

AD-A032 862

FLORIDA UNIV GAINESVILLE DEPT OF PLANT PATHOLOGY  
BIOLOGICAL CONTROL OF AQUATIC WEEDS WITH PLANT PATHOGENS. (U)  
OCT 76 T F FREEMAN, K E CONWAY, R CHARUDATTAN DACW73-73-C-0049


F/G 6/6

UNCLASSIFIED

WES-CR-A-76-2

NL

| OF |  
AD  
A032862



END

DATE  
FILMED  
1 - 77

ADA 032862

B009804 Suply  
(submitted by  
contractor)

AD-A632862

MC

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Contract Report A-76-2	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>BIOLOGICAL CONTROL OF AQUATIC WEEDS WITH PLANT PATHOGENS,</b>		5. TYPE OF REPORT & PERIOD COVERED Final report,
7. AUTHOR(s) T. E. Freeman, K. E. Conway R. Charudattan, F. W. Zettler		8. CONTRACT OR GRANT NUMBER(s) DACW73-73-C-0049
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Plant Pathology University of Florida Gainesville, Florida 32601		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office, Chief of Engineers, U. S. Army Washington, D. C. 20314		12. REPORT DATE October 1976
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) U. S. Army Engineer Waterways Experiment Station Mobility and Environmental Systems Laboratory Vicksburg, Mississippi 39180		13. NUMBER OF PAGES 25
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) Unclassified
18. <b>WEST</b>		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <b>CR-A-76-2</b>		
19. SUPPLEMENTARY NOTES Research was done in cooperation with and in support of the Florida Department of Natural Resources; U. S. Department of Interior; the Offices of Water Resources Research and Technology; and the Center for Environmental Program, University of Florida's Institute of Food and Agricultural Sciences. Dr. E. O. Gangstad, OCE, was the Contracting Officer's representative.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In the early 1970's, an effort was initiated to locate and isolate pathogenic organisms for use in the biological control of aquatic plants with special reference to waterhyacinth. This report describes the exhaustive search which has been conducted both in the United States and in several foreign countries. Information on laboratory and field research studies is presented as well as the current state of the art in this area of aquatic plant management research.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Aquatic plant control Aquatic weeds Biological control Pathogens		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

406436

VB

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

THE CONTENTS OF THIS REPORT ARE NOT TO BE USED FOR ADVERTISING, PUBLICATION, OR PROMOTIONAL PURPOSES. CITATION OF TRADE NAMES DOES NOT CONSTITUTE AN OFFICIAL ENDORSEMENT OR APPROVAL OF THE USE OF SUCH COMMERCIAL PRODUCTS.

ACCEPTED BY	
NTIS	White Star <input checked="" type="checkbox"/>
D-5	East Street <input type="checkbox"/>
UNIVERSITY	<input type="checkbox"/>
JUSTIFICATION	
BY	
LIC. NUMBER/AVAILABILITY CODES	
DATE RECEIVED BY STAL	
A	

## PREFACE

The information presented herein was performed in part under Contract No. DACW 73-73-C-0049 with the Department of Plant Pathology of the University of Florida, Gainesville, Florida, for the Office, Chief of Engineers (OCE). The research was done in cooperation with and in support of the Florida Department of Natural Resources; the U. S. Department of the Interior; the Office of Water Resources Research and Technology, as authorized under the Water Resources Research Act as amended; and the Center for Environmental Program of the University of Florida's Institute of Food and Agricultural Sciences. This study was conducted and the report prepared by Drs. T. E. Freeman, R. Charudattan, K. E. Conway, and F. W. Zettler. Dr. E. O. Gangstad, OCE, was the Contracting Officer's representative for the contract; his assistance and constructive criticism is hereby acknowledged. The authors are also grateful to the many students and assistants who have contributed to the conduct of the work and preparation of the report. The Mobility and Environmental Systems Laboratory of the U. S. Army Engineer Waterways Experiment Station (WES), Vicksburg, Mississippi, monitored the report.

This report is considered to be a summary statement of the current state of the art of biological control of aquatic plants with pathogenic organisms. It describes the baseline or point of departure for future work in this vital research area of aquatic plant management.

Director of WES during the preparation and publication of this report was COL J. L. Cannon, CE. Technical Director was Mr. F. R. Brown.

## CONTENTS

	<u>Page</u>
PREFACE .....	2
CONVERSION FACTORS, METRIC (SI) TO U. S. CUSTOMARY AND U. S. CUSTOMARY TO METRIC (SI) UNITS OF MEASUREMENT .....	4
INTRODUCTION .....	5
RELEVANCE OF RESEARCH .....	5
RESEARCH APPROACH .....	6
ENDEMIC PATHOGENS .....	7
<i>Rhizoctonia solani</i> .....	7
<i>Acremonium zonatum</i> .....	8
<i>Cercospora</i> spp .....	8
<i>Cercospora rodmanii</i> .....	9
Fungi Associated with Waterhyacinth .....	13
Other Studies .....	14
EXOTIC PATHOGENS .....	14
INTEGRATED CONTROL .....	17
CONCLUSION .....	17
RECOMMENDATIONS .....	18
REFERENCES .....	19
BIBLIOGRAPHY .....	20
TABLES 1-4	



**CONVERSION FACTORS, METRIC (SI) TO U. S. CUSTOMARY  
AND U. S. CUSTOMARY TO METRIC (SI) UNITS OF MEASUREMENT**

Units of measurement used in this report can be converted as follows:

<u>Multiply</u>	<u>By</u>	<u>To Obtain</u>
	<b><u>Metric (SI) to U. S. Customary</u></b>	
millimicrons	$0.03937 \times 10^{-6}$	inches
microns	0.00003937	inches
metres	3.28084	feet
hectares	2.47108	acres
litres	0.26417	gallons
cubic metres	35.31466	cubic feet
grams	0.002204622	pounds (mass)
Celsius degrees or Kelvins	1.8	Fahrenheit degrees*
	<b><u>U. S. Customary to Metric (SI)</u></b>	
inches	0.0254	metres
feet	0.3048	metres
acres	4046.856	square metres
gallons (U. S. liquid)	0.00378541	cubic metres
pounds (force) per square inch	6.894757	kilopascals
Fahrenheit degrees	0.555	Celsius degrees or Kelvins**

- 
- \* To obtain Fahrenheit (F) temperature readings from Celsius (C) readings, use the following formula:  $F = (1.8)(C) + 32$ . To obtain Fahrenheit readings from Kelvins (K), use:  $F = (1.8)(K - 273.15) + 32$ .
  - \*\* To obtain Celsius (C) temperature readings from Fahrenheit (F) readings, use the following formula:  $C = (0.555)(F - 32)$ . To obtain Kelvin (K) readings, use:  $K = (0.555)(F + 459.67)$ .

# BIOLOGICAL CONTROL OF AQUATIC WEEDS WITH PLANT PATHOGENS

## INTRODUCTION

1. Plant pathogens have many characteristics that make them ideal candidates as biocontrols for aquatic weeds.<sup>1,2</sup> They are numerous and diverse, frequently host specific, and easily disseminated and self-perpetuating. In addition, they will not completely eliminate a host species, and do not normally affect man or other animals. With these points in mind, a modest program was begun at the University of Florida at Gainesville in 1970, with the object of the evaluation and subsequent use of plant pathogens as biocontrol agents for aquatic weeds. The program was expanded with the aid of a matching grant from the U. S. Department of Interior's Office of Water Resources Research and subsequent support from the Florida Department of Natural Resources and the U. S. Army Corps of Engineers (Contracts DACW 73-71-C0002 and DACW 73-73-C-0049). The research described herein was conducted partly under Contract DACW 73-73-C-0049.

2. The program has progressed rapidly considering the lack of initial background information. We have developed a considerable backlog of information about diseases affecting aquatic plants. The objective of the utilization of plant pathogens in biocontrol programs for noxious aquatic plants has not yet been completely fulfilled. However, we have reached the critical stage when field evaluation of at least two pathogens of waterhyacinth is fully warranted. An additional three or four organisms should soon also reach this point. All of these efforts are aimed at waterhyacinths. We have also attempted to find and research diseases with biocontrol potential for other aquatic weeds. To do these studies, we have requested continuing support of this critical area of research by the U. S. Army Corps of Engineers. Such support is vital in order to enable us to more completely reach our objective and to begin reaping the benefits of an operational biological control program in the shortest possible time.

## RELEVANCE OF RESEARCH

3. The aquatic weed problem is one of considerable proportion that appears to be growing in magnitude rather than diminishing or even stabilizing. This is occurring despite the expenditure of considerable sums of money and human energy in the application of conventional methods of mechanical and chemical control.

4. The Florida Department of Natural Resources estimates over \$15,000,000 are expended annually in Florida for aquatic weed control. These control efforts are concentrated primarily on the estimated 100,000 ha\* of water hyacinth (*Eichhornia crassipes*) and 40,000 ha of hydrilla (*Hydrilla verticillata*) that occur in the state. Lesser attention is given to the approximately 20,000 ha of other aquatic weeds, such as Eurasian watermilfoil (*Myriophyllum spicatum*) and alligatorweed (*Alternanthera philoxeroides*) (Burkhalter, personal communication). Despite these efforts, aquatic weed infestations have increased steadily in the years since these plants were introduced. The range of these plants has also expanded. Within the last 2-3 yr, Eurasian watermilfoil was found in the St. Johns

---

\* A table of factors for converting metric (SI) units of measurement to U. S. customary units and U. S. customary units to metric (SI) units is presented on page 4.

River watershed, and hydrilla was found to infest Rodman Reservoir on the Cross Florida Barge Canal, Okeechobee, Orange, and Lochlossa Lakes.

5. Florida is by no means unique in having a tremendous aquatic weed problem. Proliferating water weed populations are of concern in the rest of the United States, Middle Europe, Africa, Asia, and South and Central America. Indeed the problem is worldwide, but is more acute in the warmer latitudes where waterhyacinth, hydrilla, watermilfoil, alligatorweed, salvinia (*Salvinia* spp.), and waterlettuce (*Pistia stratiotes*) are the major offenders. Reasons for the increasing aquatic weed problem are complex, but are definitely related to man and his activities. With the increase in population and the accompanying environmental problems, it has become apparent that new methods of aquatic weed control must be found. Conventional methods have not been entirely satisfactory because of cost, overall ineffectiveness, or environmental pollution. The energy problem as it relates to fossil fuel supply has also served to emphasize the need for low-energy methods of control.

6. In recent years, biological control methods have received considerable attention. Various species of herbivorous insects, fish, snails, and even the manatee have been, or are being, investigated for their ability to exert some control pressure on noxious aquatic plants. Some of them, such as the alligatorweed flea beetle, have been reasonably effective, especially in an integrated control program. Surprisingly, until our program was initiated, plant pathogens had been rarely considered as biocontrol agents. They have all the prerequisites of a biocontrol agent and thus offer an untapped reservoir of potential usefulness, either alone or in an integrated control program with insects and, perhaps, chemicals. Our research efforts, the first 3 yr of which are summarized in Reference 2, are aimed at bringing to fruition this use of plant pathogens in control programs for aquatic weeds.

#### RESEARCH APPROACH

7. We are using two approaches in our efforts to use plant pathogens to control aquatic weeds. They are:

- a. The use of endemic or native plant pathogens as a type of "biological herbicide" through the artificial induction of epiphytotics. We consider this to be the most rapid approach from an operational standpoint.
- b. The search for an ultimate utilization of exotic plant pathogens. This has been the classical approach successfully used by entomologists in their biological control efforts toward imported weeds. This facet involves the search for pathogens near the center of origin of the noxious species, in an area where climatic conditions are similar to those where the pest is a problem in this country. This is the slower of the two approaches from an operational standpoint.

8. During the past 3 yr, our efforts have been directed primarily toward those pathogens with definite biocontrol potential. These are: the endemic pathogens of waterhyacinth; *Acremonium zonatum*, *Cercospora piaropi*, *C. rodmanii*, *Rhizoctonia solani*, and three exotic ones; *Bipolaris stenospila*, *Uredo eichhorniae*, and *Rhizoctonia* sp. We have carried out extensive cultural studies in the laboratory; greenhouse studies, including cross-inoculation onto other plant species; host range studies; and in the case of the endemic pathogens, small-scale field tests. These latter tests have shown *A. zonatum* and *C. rodmanii* to have considerable potential as biocontrols. We are presently testing both of these at locations in Florida and in Lake Concordia in Louisiana. In this latter test, we have the two pathogens combined with two insects (*Neochetina eichhorniae* and *Arzama densa*) in all possible combinations. This test is being conducted in cooperation with the U. S. Army Corps of Engineers, U. S.

Army Engineer Waterways Experiment Station (WES), and the U. S. Department of Agriculture (USDA) with the approval of the Louisiana Department of Agriculture and the Louisiana Fish and Game Commission. We believe *C. rodmanii* to have been the cause of a spectacular decline of waterhyacinth in Rodman Reservoir in 1971. This natural decline saved the U. S. Army Corps of Engineers approximately \$35,000 in spray cost in that body of water (Zeiger, personal communication). *A. zonatum* and *N. eichhorniae* combinations are also being tested in south Florida near Ft. Lauderdale.

9. Work with the exotic pathogens is being done in our quarantine facility, which is limited in size. Therefore, the work is progressing at a slower pace than with the endemic pathogens.

## ENDEMIC PATHOGENS

### "*Rhizoctonia solani*"

10. Fungal isolate RhEa from Panama was shown to be highly pathogenic to waterhyacinths.<sup>3</sup> It was identified as a species of *Rhizoctonia* closely related to *R. solani*. We, therefore, deemed it advisable to test endemic isolates of this same fungus. Several isolates<sup>4</sup> were found to be very nearly as pathogenic to waterhyacinth as was RhEa. An isolate from bean was selected (H287) for further study as a biocontrol agent. In the meantime, studies were continued under quarantine conditions with RhEa.

11. After completion of sclerotial survival studies with RhEa, we were convinced that the fungus had potential as a biocontrol agent for aquatic plants. Sclerotia of RhEa survived for over 26 months when submerged in lake water. Survival rate was probably longer, but the supply of sclerotia for testing purposes was exhausted after 26 months. The pathogenicity of cultures derived from submerged sclerotia was equal to that of the stock culture and cultures derived from dry sclerotia.<sup>5</sup> In addition, it was shown that certain *R. solani* isolates were capable of attacking underwater portions of waterhyacinth and other aquatics.<sup>6,7</sup> These later results were further encouragement to proceed with field tests of domestic isolates of *R. solani*.

12. On the negative side, *R. solani* possesses some distinct disadvantages. It is a common soil-inhabiting parasite that affects numerous species of plants. Under ideal conditions for disease development, it will attack a large number of commercial crop plants (Table 1). Many consider this to completely preclude the use of this organism for biological control purposes. However, its use as a biological herbicide in the aquatic environment would not necessarily increase its potential as a pathogen of terrestrial plants. Indeed, when terrestrial crop plants were sprayed with inoculum of *R. solani*, fewer of them were attacked than when these same plants were inoculated under ideal disease conditions in the greenhouse (Table 1).

13. Initial field studies to assess the potential of domestic isolates of *R. solani* were begun in the fall of 1973. An isolate of *Acremonium zonatum*, formerly *Cephalosporium zonatum*, described by Rintz<sup>8</sup> was also included in these studies. These fungi were used alone and in combination on a well-established stand of waterhyacinths in an isolated area of Lake Alice on the University of Florida campus. Both fungi were mass grown in liquid culture on a relatively simple chemically defined medium (Difco Czapeks-Dox broth amended with 0.5 percent yeast extract). Both fungi were grown in this manner, and the mycelial mats collected after 14 days. These mats were ground in a large commercial Waring blender. The resulting mycelial suspensions were sprayed on waterhyacinth plots in Lake Alice on 10 October. The two fungi were applied singly and in combination. Spraying was accomplished with a 12-gal conventional Broyhill sprayer at 100 psi. Plots were approximately 1/20 of an acre.

14. Infection by *R. solani* was apparent in less than a week and with *A. zonatum* in less than two.

By the first frost in early December, secondary spread of *A. zonatum* was apparent and resulting damage was significant. Lesions caused by *R. solani* persisted, but secondary spread was negligible and damage was less than anticipated. Plots were observed throughout the winter and into the following growing season. No evidence of *R. solani* infection the following spring was noted, whereas sporadic *A. zonatum* infections occurred. These results were disappointing in the case of *R. solani*, but not totally unexpected. The lack of a spore stage was a deterrent in secondary spread on aerial portions of the plant. On the other hand, results obtained with *A. zonatum* were more promising than laboratory and greenhouse studies had indicated. Thus, *R. solani* studies were curtailed in favor of more intense study on *A. zonatum*.

#### **"Acremonium zonatum"**

15. In addition to the field test (paragraphs 13-14) and host range studies (Table 1), we have conducted a wide variety of additional studies with the fungus *A. zonatum*. For instance, during the course of our preliminary work, it became apparent that the culture medium on which *A. zonatum* was grown affected its pathogenicity to waterhyacinth. Pathogenicity as well as total inoculum production was enhanced by enriching the culture medium. Pathogenicity increased as culturing medium was changed from cornmeal agar to Czapeks agar to Czapeks agar plus yeast extract to potato-dextrose agar to potato-dextrose agar plus yeast extract.

16. The fact that pathogenicity of *A. zonatum* changed due to culturing substrate brought forth the idea that other factors may also be used to increase pathogenicity. In at least one other plant pathogen, *Cladospodium fulvum* on tomato, virulence has been increased through irradiation of cultures with ultraviolet radiation. Short wave UV radiation in the 253-m $\mu$  range was most effective. Since this wavelength is germicidal, dosages were critical. A decision was made to embark upon a study of the effectiveness of UV radiation for inducing mutations for increased virulence to waterhyacinth in populations of *A. zonatum*. Thus far, we have isolated over 400 single spore cultures derived from spores irradiated with UV light (253 m $\mu$ ) for periods ranging from 30 to 180 sec. Many of these cultures differ morphologically from the parent cultures.

17. So far, approximately 200 cultures have been tested for changes in their pathogenicity to waterhyacinth. Where changes have been noted, in most cases, there was a reduction in pathogenicity. However, in three or four cases, there appeared to be a slight increase in pathogenicity. The fact that some changes appear to have occurred is encouraging at this stage in these studies.

18. Additional field tests of *A. zonatum* were conducted during the summer of 1974 both at Gainesville and Ft. Lauderdale. In June of 1975, it was included, along with three other biotic agents, in an integrated control test in Lake Concordia.

19. Results with *A. zonatum* have been erratic. Infection appears to be related to the growth habit of the waterhyacinth. Large robust plants seem to be more readily infected than smaller ones. Where infection does take place, subsequent spread by the pathogen is slow and does not keep pace with the prolific growth of the host. *A. zonatum* is most effective when used in an integrated system with insects and other fungi.

#### **"Cercospora" spp**

20. During the winter of 1973-74, waterhyacinth plants in the area of Gainesville were found affected by a leaf-spotting disease not previously noted. The disorder was found to be incited by a species of *Cercospora* subsequently identified as *C. piaropi*.<sup>9</sup> This was only the second report of the occurrence of this organism since it was originally reported from Texas in 1914. The other occurrence was in India.

The fungus did not appear to be causing appreciable damage to the waterhyacinth plant at the time it was first noted.

21. In December 1973, Dr. Conway isolated, along with many other fungi, a *Cercospora* species from declining waterhyacinth in Rodman Reservoir.<sup>10</sup> Preliminary tests showed the fungus to be pathogenic on water hyacinth, and a secondary test showed it to inflict considerable damage on this plant. In fact, affected plants eventually died and sank to the bottom of the test vats. Therefore, this fungus was programmed for more detailed laboratory study and eventual field testing.

22. Microscopic examination revealed the fungus to be a typical *Cercospora*. However, spore measurements showed the spores to be much longer than those recorded for *C. piaropi*<sup>9</sup> which is the only previously reported *Cercospora* on waterhyacinth. Spores of *C. piaropi* rarely exceed 150 $\mu$  in reports of its occurrence, whereas those of the Rodman *Cercospora* frequently exceed 400 $\mu$ . In addition, symptoms caused by this fungus appear to differ from those recorded for *C. piaropi*. Symptoms caused by the former is a general blasting of the foliage, whereas the latter produces more discrete spots on the leaves. However, there was the distinct possibility that we were dealing with two manifestations of the same fungus and disease. When leaves exhibiting *C. piaropi* symptoms with fruiting characterized by small spores are brought into the laboratory and placed under moist conditions, long spores frequently develop on the dead tissue. Therefore, we were not sure if we had a long spore variant of *C. piaropi* or a new waterhyacinth species of *Cercospora* (*Cercospora* spp. are distinguished by the host upon which they occur). Further study revealed that the spore not only differed in size but in basal morphology. These differences along with symptomology prompted its description as a new species of *Cercospora* designated *C. rodmanii*.<sup>11</sup>

#### **"*Cercospora rodmanii*"**

23. Optimum temperature for growth of *C. rodmanii* is near 20°C; however, excellent growth occurs over the range of 20° to 30°C. Pathogenicity appears correlated with growth, but this has not been definitely determined. The fungus grows rapidly for a *Cercospora* reaching a maximum growth on potato-dextrose agar plus yeast extract in 10-14 days at 20°C. It sporulates sparsely on waterhyacinth-extract agar, but profusely on the dead leaf tissue in the laboratory, greenhouse, and field.

24. Temperature studies tended to indicate the fungus would more likely attack the plant during mild weather. Therefore, field studies were begun in the fall on established stands of waterhyacinth in an isolated area of Lake Alice. An initial application was made 4 September 1974. Infection took place within two weeks, but subsequent spread was slow and a second application was made 3 October 1974 (Figure 1). Inoculum consisted of the mycelial growth from 60 Roux bottles cultured for two weeks (approximately 1000 g wet weight).

25. By mid-November, extensive damage had occurred in the immediate area sprayed, and spread was apparent around the entire perimeter of the lake (Figure 2). By the first frost on 2 December, considerable damage was apparent in areas of secondary spread, and spore trappings of the fungus were made at considerable distance from the original inoculation site, indicating extensive secondary spread. After this first frost and subsequent ones on 9 and 17 December, it was apparent that diseased plants were more severely damaged than more healthy ones. It should be noted that all these frosts were relatively light and healthy plants were not severely damaged. Above-average temperatures occurred in the latter part of January and February 1975, and some of the waterhyacinths began sending out offshoots. Evidently the apical meristem of the plants had not been killed, and the plants were able to resume growth. However, it was apparent that a severe stress had been placed on the waterhyacinths




Figure 1. The damage resulting from inoculation of waterhyacinth with a *Cercospora* sp. in an isolated area of Lake Alice. ① Application of fungus to healthy waterhyacinth on 3 October 1974. ② Area of immediate spray (arrow) showing initial damage on 15 October 1974. ③ Overall area viewed from inoculation site on 13 November 1974. ④ Same view as ③ on 21 November 1974. ⑤ Same area on 17 December 1974, after three light frosts. ⑥ Closeup of diseased and frosted plants on 6 December 1974




Figure 2. Spread of *Cercospora* sp. on waterhyacinth in Lake Alice after 3 October 1974 application. ① Initial spread from inoculation site (arrow) through sawgrass barrier into main body of lake on 13 November 1974. ② View from sawgrass barrier looking toward main body of lake on 18 November 1974. ③ Aerial view showing spread of disease from initial inoculation site (arrow) around entire perimeter of lake on 21 November 1974. ④ Color IR photo of lake on 21 November 1974. Diseased waterhyacinths are brownish-red whereas healthy ones are bright red



when the plants in the pool area, which were normally 2-3 ft tall, were less than 6 in. high. In comparison to these plants, waterhyacinths in the main lake where the disease was less severe were their normal 3-4 ft. Thus, *C. rodmanii* fulfills the purpose of a biological control organism for waterhyacinths by increasing the stress on the plant and not necessarily eliminating the entire population.

26. We wanted to know what effect infection would have on the plants over a period of time. Four more sprays were applied on the pool area, two in March and one each in May and July.

27. In April, we were able to determine that most of the small waterhyacinths in the pool were infected again with *C. rodmanii*. However, with the approach of summer, the waterhyacinths began their rapid growth phase, and by June, the new leaves were outgrowing the disease. The infection of *C. rodmanii* during the summer was confined to the older, lower leaves. This condition prevailed throughout the summer until September when the waterhyacinth growth was slowed by cooler night temperatures. Increased infection was now apparent in the pool area. The waterhyacinths in the pool area showed a general browning by mid-September. This browning has continued into October until definite disease symptoms could be seen on the plants throughout the pool. Damage to waterhyacinth was approximately one month ahead of last year, and we were looking for increased damage this fall and winter. We will not be able to totally evaluate the extent of damage until the end of the 1976 growing season.

28. This disease that occurred in Lake Alice as a result of our inoculation with *C. rodmanii* is strikingly similar to the Rodman disorder noted in 1971 and to a lesser extent in 1972-1973. Dave Bowman, Reservoir Manager, observed the disease in Lake Alice and believes what he saw to be the same disease he had observed in Rodman. In addition, the root-rotting phase so notable in Rodman has begun to appear in Lake Alice. We now believe *C. rodmanii* to be the causal agent of the Rodman disorder.

29. Encouraged by the success in Lake Alice last fall, test plots were set up in Rodman Reservoir. A site was chosen behind tree population No. 4 to exclude outside interference with the tests. The purpose of this experiment was to reestablish the disease in the reservoir. Although the disease was very prevalent in 1971, its severity on the waterhyacinth population has lessened each year.

30. Five sprays were to be applied from the shoreline, one every two weeks. The multiple sprays were felt necessary to begin infection and to increase the inoculum to a high enough level to create an epiphytic. Spray operations were begun on 28 February 1975. The next day, the U. S. Army Corps of Engineers began to raise the water level of the reservoir from its winter draw-down depth of 4.67 m to a level of 5.49 m. The water level had risen, and the waterhyacinths were growing and floating free of the shoreline by the second spray date. By the third spray application, the spray plots were moving with the rising water; therefore, one of our objectives of increasing the inoculum in one area could not be achieved.

31. In mid-April, damage on the plants due to *C. rodmanii* was present along a gradient from the opening onto the reservoir from tree population No. 4 to our spray site onshore. By this time, there was a definite reduction in plant growth in our plots when compared to the untreated waterhyacinths that surrounded these areas. In addition to the disease in the tree population No. 4 area, there was also a heavy natural infection of *C. rodmanii* in the main part of the reservoir in the Orange Springs and Blue Springs areas.

32. During May, the disease in the main reservoir continued to stress and brown the waterhyacinths. On May 30, another spray plot was established in the Cypress tree stand in tree population No. 4.

33. In June, the waterhyacinths in the Orange and Blue Springs areas were completely browned. The symptoms on the plants were typical for *C. rodmanii* damage.

34. By July, the waterhyacinths in the Cypress stand were beginning to brown at the tips of the leaves, and in mid-July, the waterhyacinths in our original spray plots were showing the typical *C. rodmanii* symptoms. Waterlettuce was invading soon after the severely infected waterhyacinths died and sank to the bottom. By late July, the waterhyacinths in the Cypress stand were also dying out and open water was beginning to show.

35. On 7 August, aerial inspection of the original spray site showed continued browning and drop out of the waterhyacinths. In the Cypress tree area, there was now 10-20 acres of open water. By mid-August, the estimate of open water in the Cypress tree area exceeded 20 acres. Large mats of waterhyacinths showed typical symptoms, which included many dead plants with floating, spindly petioles and leaves. It was also noted that completely dead plants continued to float until broken apart by wind and water action.

36. By mid-October, the area of open water was estimated to be 35-40 acres, with an additional 20 acres invaded by waterlettuce. The area now is aesthetically pleasing.

37. Results with *C. piaropi* have been very encouraging and it is programmed for more intense study. The host range tests (Table 2) were especially encouraging as they indicate the fungus to be highly host specific.

38. The field symptoms of *C. rodmanii* suggested that toxin may be involved. Symptoms were typical of those incited by toxic by-products in other disease syndromes. However, studies conducted by a graduate student, Mr. Ray Martyn, did not verify the presence of a toxic product in culture filtrates of the fungus.

39. We have carried out limited epidemiological studies. Extensive spore trappings were conducted in Lake Alice during the fall of 1975, when temperature variations ranged from below freezing at night to day temperatures in the eighties. Results show that sporulation by *C. rodmanii* was curtailed by temperatures below 50° F. Maximum occurred when day temperatures were in the 70- to 85-deg range. On 8 December 1975, spore trappings reached a high of 193 spores per cubic metre of air volume. Laboratory studies failed to reveal a forceable discharge mechanism for spores of *C. rodmanii*. Therefore, both spore release and subsequent dispersal is probably passive in nature.

#### **Fungi Associated with Waterhyacinth**

40. In November 1973, an active research project was initiated when Dr. Conway joined the project to catalogue the mycoflora found on waterhyacinths. His specific assignment was to study the mycoflora associated with declining waterhyacinths in Rodman Reservoir. The mycoflora can be divided into two main categories: those occurring on living tissue and those occurring on declining tissue (Table 3).

41. The fungi that occur on living tissue are studied for their possible role in the control of the prolific waterhyacinths. Fungi identified and cultured in this group include (a) Ascomycetes—*Mycosphaerella* sp., *Didymella* sp., *Leptosphaerulina* sp., and *Asteromella* sp.; and (b) Fungi Imperfecti—*Ascochyta* sp., *Cercospora* sp., *Phoma* sp., *Myrothecium striatisperum*, and *Cephalosporiopsis* sp.

42. Fungi associated with dead and declining tissue include (a) Fungi Imperfecti—*Curvularia lunata* var. *aeria*, *Alternaria* sp., *Dendryphion* sp., *Nigrospora sphaerica*, *Nigrospora oryzae*, *Thysanophora longispora*, *Periconca* sp., *Scolecobasidium constrictum*, *Penicillium* spp., *Tilletiopsis* sp., *Fusarium* sp., *Pestalotia* sp., *Botryodiploidea* sp., *Acremonium zonatum*, *Memnoniella* sp., *Epicoccum* sp.,

*Cladosporium* spp., *Pithomyces chartatum*, *Aspergillus* spp., an unknown synematal fungus, *Rhizoctonia* sp. and *Pyrenochaeta* sp.; (b) Phycomycetes—*Mucor* sp. and *Circinella* sp.; and (c) Ascomycete—*Melanospora* sp.

43. The role of saprophytic fungi is often neglected in disease research; however, several important contributions to our knowledge of waterhyacinth control and utilization can be realized through their investigation. *Periconia* is a saprophytic fungus that produces a toxin which causes disease symptoms on certain plants. When the fungus is inoculated onto the waterhyacinth, a leaf spot is produced with the fungus continuing to grow on the leaf. The presence of *Aspergilli* and *Penecillia* may indicate the production of aflatoxins on the waterhyacinth. This is obviously important to researchers attempting to use waterhyacinths as a food source. The presence of *Pithomyces chartarum* is of even greater importance to nutrition researchers. It is the only fungus known to have caused a natural widespread outbreak of poisoning. The toxin, sporidesmin, causes liver damage, loss of weight, icterus, and photosensitivity to grazing animals. An additional project has been initiated through our department in cooperation with the Department of Animal Science Nutrition Laboratory and the Department of Veterinary Science to study mycotoxins present in waterhyacinths and their effect on animals. Thus, our efforts in determining mycoflora that may have biological control potential may also provide valuable information for workers concerned with other methods of control and utilization of the waterhyacinth.

#### Other Studies

44. Less extensive studies have been conducted with several other endemic pathogens of aquatic plants. These were concerned primarily with minor pathogens of waterhyacinth and the alligatorweed stunt virus.<sup>12</sup>

45. Repeated attempts to transmit the alligatorweed stunt virus by mechanical means and insects were unsuccessful. However, it was transmitted by grafting of healthy and diseased stock. Such a method of transmission limits this agent's usefulness as an effective biocontrol. It does prove conclusively that the malady is viral induced and not a genetic abnormality. Dr. Zettler is convinced that the virus belongs in the beet yellows group and may actually be a strain of beet yellows. Being in this group could present problems in clearing this virus for use as a biocontrol, since tristeza virus of citrus also belongs to the beet yellows complex.

46. Two other minor pathogens have been reported affecting waterhyacinth in Florida. There are a *Sigmoidea* sp.<sup>13</sup> and *Mycoleptodiscus terrestris*.<sup>14</sup> Both appear to be weak pathogens of little potential in biological control. A new nonpathogenic fungus of the genus *Doratomyces* was also recovered from waterhyacinth foliage.<sup>15</sup>

#### EXOTIC PATHOGENS

47. Dr. Charudattan has traveled extensively in search of exotic pathogens on waterhyacinth and hydrilla. In 1973, he spent 90 days in India (supported by the Florida Department of Natural Resources) collecting pathogens on these two plants. The regions in India surveyed included Sringar (State of Kashmir), the surroundings of Delhi and Calcutta, Dehra Dan Utter Pradesh, Kota (Rajasthan), and several areas in the state of Orissa. These areas were not covered during his previous tour in India from November 1971 to February 1972. In addition, several previously surveyed areas around Madras were revisited. One hundred and eighty-two fungal and bacterial isolates were obtained from hydrilla, waterhyacinth, waterlettuce, and miscellaneous submersed aquatic plants. Ninety-five were from hydrilla and the remainder from other plants.

48. Twenty-four of the Indian hydrilla isolates proved to be pathogenic to hydrilla. These were primarily species of *Penicillium*, *Aspergillus*, and *Trichoderma*. These were pathogenic through the production of toxic metabolites.<sup>16</sup> These toxic agents were identified as oxalic acid and penicillic acid. They are general toxic agents not specific for hydrilla. Thus, their direct use in a biocontrol program appears limited.

49. Fourteen of the remaining isolates were found to be pathogenic to waterhyacinth. Among these were isolates of *Mycrothesium roridum*, *Alternaria eichhorniae*, and *Acremonium* sp. The others have not been identified because they are weakly pathogenic and are of academic interest only. The three pathogens have been evaluated and rejected as biocontrol agents in India. Nevertheless, we are maintaining them for a possible future role here in the United States.

50. Dr. R. Charudattan undertook two survey tours during the first quarter of 1974. The first trip was to the Dominican Republic from 11 to 16 February. On this survey tour, about 82 fungal and bacterial isolates from diseased waterhyacinths were collected. The genera of fungi collected included *Helminthosporium*, *Phoma*, *Macrophoma*, *Colletotrichum*, *Fusarium*, *Chaetomium*, *Pestalotia*, and *Alternaria*. Unfortunately, the rust and smut diseases of waterhyacinth reported to occur in this country were not found. The second trip was to Venezuela between 3 and 12 March and yielded a collection of approximately 87 cultures. The majority of these are fungi, but they are yet to be identified. Pathogenicities of about 35 isolates from the Dominican Republic have been tested on waterhyacinth. So far, a species of *Phoma* and two species of *Helminthosporium* have been found to be virulent on waterhyacinth. One of the species of *Helminthosporium* is extremely pathogenic on waterhyacinth. This appears to be a hitherto unreported pathogen of waterhyacinth. Preliminary tests indicate it to be comparable in pathogenicity to *Acremonium zonatum* and *Rhizoctonia solani*. Among the diseases found in Venezuela was a necrotic leaf spot surrounding each feeding injury caused by the insect, *Neochetina*. Diseases of alligatorweed or *Pistia* were not encountered, though these plants were found in these countries. Several new and useful contacts were established in the two countries.

51. Dr. Charudattan traveled to Argentina and Uruguay during the last half of April 1974 in search of diseases on waterhyacinth and alligatorweed. On this trip, he was successful in finding the rust disease on waterhyacinth, but was unsuccessful in locating rust of alligatorweed. The rust on waterhyacinth was present in Argentina but not in Uruguay, where it had been reported to flourish. He was able to bring back adequate material for critical study in our isolation greenhouse.

52. Morphologically the rust fits the description of *Uredo eichhorniae*. This fungus was originally recorded on waterhyacinth in the Dominican Republic in 1927. The organism has been loosely referred to as *Puccinia eichhorniae*. No basis for this name has been established, since only the uredial stage of this rust is known. The material from Argentina did not have teliospores, and they have not formed in material brought back to Gainesville. We have been able to infect plants in the isolation greenhouse with the rust. Therefore, we should be able to maintain the culture and increase it for further study.

53. Considerable work remains to be done before we can determine if this disease has biocontrol potential. It did not appear to be causing significant damage on waterhyacinth in Argentina. However, this does not rule out the possibility that it will cause significant damage on the host in Florida. A different host gene pool may exist here that has not had the selection pressure of the disease exerted upon it. This we will soon determine. The big advantage of rust diseases is their host specificity. However, some rust species can affect related plant species and many have alternate hosts in nonrelated plant groups. Presently we do not have information concerning additional related hosts or alternate hosts of *U. eichhorniae*. Dr. Charudattan did find a similar rust on *Pontederia* sp. growing in close proximity to

infected waterhyacinths. A similar rust also occurs on *Pontederia* in Florida.<sup>17</sup> Cross-inoculation studies with these two rusts are presently in progress.

54. One additional exotic pathogen was added to our collection during 1974. This was a species of *Alternaria* isolated from leaf spots on alligatorweed from Puerto Rico. Diseased material was collected and transferred to us by Mr. Chuck Zeiger of the Corps of Engineers and Dr. Dave Perkins of the USDA. We had received unofficial reports of an *Alternaria* affecting waterhyacinth in California as well as other locations. This represents our first collection of the fungus. Its pathogenic potential is still under evaluation.

55. With the completion of the preceding trips, we now have a fair idea of the disease picture as it occurs on waterhyacinth. Interestingly, the widest variety of disease have been found in the area of Puerto Rico and the Dominican Republic.

56. The two exotic pathogens that appear to have the most biocontrol potential are the waterhyacinth rust and the *Helminthosporium (Bipolaris)* on waterhyacinth. These have been studied the most extensively and are still being actively investigated.

57. Initial attempts to inoculate waterhyacinth with the rust pathogen, *U. eichhorniae*, were unsuccessful, possibly due to lack of proper environment. A variety of treatments to stimulate the germination of uredospores of waterhyacinth rust were tried. They were: hydration of spores for 24 hr, freezing (-5°C for 24 hr) or heating (for 3 min at 55°C) prior to hydration; and hydration in various concentrations of a mineral salts medium, in specialized media for growing rusts (obtained from Dr. C. A. Hollis), and in 0.1 percent solution of Tween 20. All these treatments failed to stimulate spores of *U. eichhorniae* to germinate. Hydration in dilute (0.0025 to 0.025 percent) aqueous solutions of nonanol and octanol stimulated less than 1 percent of the spores to germinate.

58. After this initial failure, Charudattan again visited Argentina, Uruguay, Paraguay, and Brazil from 12 October through 8 November 1975, to collect spore samples of waterhyacinth rust, *Uredo eichhorniae*. As part of the general information he is seeking on rusts of Pontederiaceae, he also makes collections of *Uromyces pontederiae* on *Pontederia cordata*, *P. lanceolata*, and *Eichhornia azurea*. An apparently new rust on *Reussia subovata* was also discovered in the Misiones province of Argentina. Its relation to *U. eichhorniae*, *U. pontederiae*, and *Uromyces heterantherae* is being studied. The taxonomic study of these rusts would be vital to the host-specificity tests on *U. eichhorniae*.

59. A sample of *U. eichhorniae* from Dique Lujan, Campana, Argentina, was successfully inoculated onto waterhyacinth from Florida in greenhouse tests. Several additional techniques of spore germination were tried on *U. eichhorniae*. These included varying periods of hydration of uredospores, heat-shock, various temperatures, and incubation with volatile alcohols, synthetic media, or host leaf-extracts. Freezing spores for periods up to 24 hr prior to hydration seems to trigger spore germination and subsequent infection of host. Use of very freshly collected uredospores also seemed vital to successful germination and host-infection.

60. The possible lack of prolonged contact between uredospores and the waxy leaves of waterhyacinth was considered partly responsible for failure of infections. Hence, undiluted glycerine and three different oils, namely heavy paraffin oil, citrus oil, and Visko-Rhap mineral oil, were tested as sticking agents in inoculations with *U. eichhorniae*. The oils were less desirable than glycerine. The less viscous citrus oil and Visko-Rhap proved toxic to waterhyacinth leaves, causing burn damage. While mineral oil was nontoxic, it was suspected of creating a hydrophobic phase between spores and leaves and thus preventing spore germination. Glycerine proved to be a successful medium for spore inoculation.

61. Close to 100 percent relative humidity following deposition of uredospores on leaves was necessary for infections. This was achieved by maintaining spores and plants in chambers containing saturated  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$  solution.

62. Rust infection on inoculated waterhyacinth was visible in about two to three weeks after spore deposition. Development of uredosori was complete after the third week. During a two-week period following eruption of uredosori, several crops of spores have been observed.

63. Inoculations of waterhyacinths with *U. pontederiae* from *Pontederia* spp. and *E. azurea* are being attempted to determine cross infectivity among rusts of Pontederiaceae.

64. The isolate of *Bipolaris* identified as *B. stenospila*<sup>18</sup> from the Dominican Republic was compared for its virulence with an isolate of *Helminthosporium stenospilum* Drechs. (= *B. stenospila*) from the American Type Culture Collection. The former was significantly more virulent and destructive to waterhyacinth than the latter which was moderately pathogenic. In comparison, *H. cynodontis*, *H. sacchari* and *H. sativum* were weakly pathogenic while the following were nonpathogenic to waterhyacinth: *H. maydis*, *H. victoriae*, *H. setariae* and *H. carbonum*. Both *H. sacchari* and *H. stenospila* are sugarcane pathogens found in Florida and other sugarcane regions of the southeastern United States. Therefore, it is likely that we may find an isolate of *B. stenospila* in Florida that is more virulent on waterhyacinth than the type specimen of *H. stenospila* (Table 4).

#### INTEGRATED CONTROL

65. In cooperation with Dr. Perkins at Ft. Lauderdale, limited integrated controls tests using *Acremonium zonatum* and the insect *Neochetinia eichhorniae* were conducted. These tests indicated, though inconclusively, that increased damage resulted from the use of the two biotic agents together.

66. A more extensive integrated test was set in Lake Concordia, the week of 29 June 1975. This test, which is still in progress, is a cooperative effort with the Aquatic Plant Research Branch of the WES in Vicksburg and the Biocontrol Laboratory of the USDA in Gainesville. The object is to test the effectiveness of two plant pathogens and two insects, alone and in all possible combinations on 2- by 2-m aluminum frames containing waterhyacinths. The treatments are: (a) *Arzama densa* (insect), (b) *Neochetinia eichhorniae* (insect), (c) *Cercospora rodmanii* (plant pathogen), and (d) *Acremonium zonatum* (plant pathogen). Rates of treatment were: *A. densa* = 40 per cage; *N. eichhorniae* = 50 per cage; *Cercospora* = 80-g wet weight mycelium and spores; and *A. zonatum* = 160-g wet weight mycelium and spores. Plant pathogens were put on in split applications 24 and 25 June. One half of each inoculum batch was put on in approximately 2ℓ of water per plot 24 June. The following day the other half of the inoculum was sprayed on the plots. Insects were released two weeks later (10 July). At this time, infection was noted by both pathogens (personal communication with Neal Spencer, USDA, and Sam Shirley, WES).

67. The Lake Concordia experiment has been monitored at regular intervals during the growing season and will continue at least through the 1976 growing season.

68. Preliminary weighing data show that the four agents in combination cause a significant decrease in biomass accumulation. By mid-September, this weight of plants attacked by all four biotic agents was less than 50 percent of that of the check plants. Complete results of this test will be published upon its completion in 1976. The preliminary results are very encouraging.

#### CONCLUSION

69. Significant advances have been made in realizing the goal of the utilization of plant pathogens

in biocontrol programs for aquatic weeds. It is anticipated that the results of this study will help bring this goal to fruition within the next few years. The support of the U. S. Army Corps of Engineers has been a significant contributing factor in making these efforts possible.

### RECOMMENDATIONS

70. The EPA-AIPS Workshop held at the University of Arkansas from 23 to 24 June 1975 recommended that:

- a. APHIS and the Working Group on Natural Enemies evaluate proposals and make judgment in cooperation with the State Plant Regulation Agency of whether or not to import fungi for use in the Classical Biocontrol Strategy.
- b. EPA develop guidelines for registration of fungi for use as mycoherbicides.
- c. Policy be formulated to permit commercial firms to have limited term exclusivity to public use patents.
- d. Advice and council of the American Phytopathological Society and other societies as appropriate, be sought. Identify expertise needed for the development of guidelines for registration of fungal pesticide products by EPA.
- e. AIBS continue this workshop approach for development of specific registration criteria for biological pesticides.

In addition, the following study program was outlined.

- a. Identify the fungus and determine its synonymy, genus, species, strain, and race.
- b. Identify the target plant or organism and its synonymy and biology, genus, species, strain, and biotypes.
- c. Determine the life cycle of the pathogen or that of the nearest representative perfect genus of the taxon. Investigate genetic stability.
- d. Determine the disease cycle or etiology of the disease (seed borne, soil borne, wind borne, water borne, overwintering mechanism, motility, plant parts affected, chlorosis, necrosis, wilt, sterility).
- e. Determine the epidemiology of the disease (cardinal temperatures for growth sporulation and durability in storage or the environment).
- f. Determine geographic area over which the target species occurs and the distribution of the disease.
- g. Make an intensive host range study beginning with related plant specified for all stages.
- h. Characterize the components of the fungus as it will be employed in field tests (conidia, sclerotia, mycelium, sexual spores, sporangia, chlamydo spores).
- i. Characterize the material other than fungal material included in product.
- j. Examine the toxicological properties of the product:
  - (1) Acute oral (2 animal species).
  - (2) Dermal (guinea pig).
  - (3) Respiratory (2 animal).
  - (4) Potential for mycotoxin production.
- k. Determine effects on fish and wildlife.
- l. Determine efficacy in field.

## REFERENCES

1. Zettler, F. W. and Freeman, T. E., "Plant Pathogens as Bio-Controls of Aquatic Weeds," *Annual Review, Phytopathology*, Vol 10, 1972, pp 455-470.
2. Freeman, T. E., Charudattan, R., and Zettler, F. W., "Biological Control of Water Weeds with Plant Pathogens," Publication No. 23, 1973, Water Resources Research Center, University of Florida, Gainesville, Fla.
3. Freeman, T. E. and Zettler, F. W., "Rhizoctonia Blight of Waterhyacinth," *Phytopathology*, Vol 61, 1971, p 892.
4. Joyner, G. G. and Freeman, T. E., "Pathogenicity of *Rhizoctonia solani* to Aquatic Plants," *Phytopathology*, Vol 63, 1973, pp 681-685.
5. Freeman, T. E., "Survival of Sclerotia of *Rhizoctonia solani* in Lake Water," *Plant Disease Repr.*, Vol 57, 1973, pp 601-602.
6. Joyner, B. G., *Characterization of a Rhizoctonia sp. Pathogenic to Aquatic Plants*, M.S. Thesis, University of Florida, Gainesville, Fla., 1972.
7. Freeman, T. E. "Rhizoctoniosis of Aquatic Plants," *Encyclopedia of Science and Technology Yearbook*, McGraw-Hill, New York, 1975, pp 327-328.
8. Rintz, R. E., "Zonal Leafspot of Waterhyacinths," *Hyacinth Control Journal*, Vol 11, 1973, pp 41-44.
9. Freeman, T. E. and Charudattan, R., "Occurrence of *Cercospora piaropi* on Waterhyacinth in Florida," *Plant Disease Rptr.*, Vol 58, 1974, pp 277-278.
10. Conway, K. E., Freeman, T. E., and Charudattan, R., "The Fungal Flora of Waterhyacinths in Florida, Part I," Publication No. 30, 1974, Water Resources Research Center, University of Florida, Gainesville, Fla.
11. Conway, K. E., "Cercospera Rodmanii, A New Pathogen of Waterhyacinth with Biological Control Potential" (in preparation), *Canadian Journal Bot.*
12. Hill, H. R. and Zettler, F. W., "A Virus-Like Stunting Disease of Alligatorweed from Florida," *Phytopathology*, Vol 63, p 443.
13. Lin, C. Y. and Charudattan, R., "*Sigmoidea* sp., A New Pathogenic Aquatic Hyphomycete of Waterhyacinth," *Proceedings, American Phytopathological Society*, Vol 2, 1975, p 137.
14. Charudattan, R. and Conway, K. E., "*Mycocleptodiscus terrestris* Leaf-Spot on Waterhyacinth," *Plant Dis. Repr.*, Vol 66, 1975, pp 77-80.
15. Conway, K. E. and Kimbrough, J. W., "A New *Doratomyces* from Waterhyacinth," *Mycotaxon*, Vol 2, 1975, pp 127-131.
16. Charudattan, R., "Pathogenicity of Fungi and Bacteria from India to Hydrilla and Waterhyacinth," *Hyacinth Contr. Journal*, Vol 11, 1973, pp 44-48.
17. Charudattan, R. and Conway, K. E., "Comparison of *Uredo eichhorniae*, the Waterhyacinth Rust with *Uromyces pontederiae*," *Mycologia*, Vol 67, 1975, pp 653-657.
18. Charudattan, R., Conway, K. E., and Freeman, T. E., "A Blight of Waterhyacinth, *Eichhornia crassipes* Caused by *Bipolaris stenospila* (*Helminthosporium stenospilum*)," *Proceedings, Phytopathology Society*, Vol 2, 1975, p 65.



## BIBLIOGRAPHY

Charudattan, R. and Lin, C. Y., "Isolates of *Penicillium*, *Aspergillus*, and *Trichoderma* Toxic to Aquatic Plants," *Proceedings, EWRC Fourth International Symposium of Aquatic Weeds*, Vienna, 1974, pp 142-143.

\_\_\_\_\_, "Penicillium, Aspergillus, and Trichoderma Isolates Toxic to Hydrilla and Other Aquatic Plants," *Hyacinth Control Journal*, Vol 12, 1973, pp 70-73.

Charudattan, R. et al., "Studies on the Use of Plant Pathogens in Biological Control of Aquatic Weeds in Florida," *Proceedings, EWRC Fourth International Symposium on Aquatic Weeds*, Vienna, 1974, pp 144-151.

Charudattan, R., "Evaluation of Foreign Pathogens as Biocontrols of Hydrilla and Waterhyacinth in the U. S. A.," *Proceedings, Second International Congress of Plant Pathology*, 1973, Abstract 0390; also Reprint No. 3, *WSSA Newsletter*, Vol 2, 1974, p 11.

\_\_\_\_\_, "Use of Plant Pathogens to Control Aquatic Weeds," *Impact of the Use of Microorganisms on the Aquatic Environment*, Ecological Research Series, U. S. Environmental Protection Agency, Corvallis, Oreg., 1975.

\_\_\_\_\_, "Weed Control with Plant Pathogens," *Agrichemical Age*, Jan-Feb 1975.

Conway, K. E., "Procedures Used to Test Endemic Plant Pathogens for Biological Control of Waterhyacinth," *Proceedings, Phytopathology Society*, Vol 2, 1974, p 31.

Freeman, T. E. and Zettler, F. W., "A Disease of Waterhyacinth with Biological Control Potential," *Abstracts of 1972 Meeting of Weed Science Society of America*, p 61.

Freeman, T. E., Charudattan, R., and Conway, K. E., "Use of Plant Pathogens for Bioregulation of Aquatic Macrophytes," *Proceedings, EPA Conference, Biological Control for Water Quality Enhancement*, 1975.

Freeman, T. E., Zettler, F. W., and Charudattan, R., "Phytopathogens as Biocontrols for Aquatic Weeds," *PANS.*, Vol 20, 1974, pp 181-185.

\_\_\_\_\_, "Utilization of Phytopathogens as Biocontrols for Aquatic Weeds," *Third International Symposium on Biological Control of Aquatic Weeds*, Montpellier, France, 1973; also published in *Proceedings, Conference on Integrated Systems of Aquatic Plant Control*, 1974, pp 97-102.

Gangstad, E. O. et al., "Aquatic Plant Control Program; Aquatic Weed Control with Plant Pathogens," Report 8, Dec 1974, U. S. Army Engineer Waterways Experiment Station, CE, Vicksburg, Miss.

Goeden, R. E. et al., "Present Status of Projects on the Biocontrol of Weeds with Insects and Plant Pathogens in the United States and Canada," *Weed Science*, Vol 22, 1974, pp 490-495.

Hayslip, H. F. and Zettler, F. W., "Past and Current Research on Diseases of Eurasian Watermilfoil (*Myriophyllum spicatum* L.)," *Hyacinth Control Journal*, Vol 11, 1973, pp 38-40.

Hayslip, H. F., *Evaluation of Plant Pathogens as Biocontrols of Eurasian Watermilfoil (Myriophyllum spicatum L.)*, M.S. Thesis, University of Florida, Gainesville, Fla., 1972.

Hill, H. R. and Rintz, R. E., "Observations of Declining Water Lettuce Populations in Lake Izabel, Guatemala," *Proceedings, Southern Weed Science Society*, Vol 25, 1972, pp 374-380.

Hill, H. R., *Survey and Evaluation of Plant Pathogens of Alligatorweed (Alternanthera philoxeroides (Mart.) Griseb.)*, M.S. Thesis, University of Florida, Gainesville, Fla., 1972.

Hill, H. R., Zettler, F. W., and Freeman, T. E., "Plant Pathogens with Potential for Biological Control of Aquatic Weeds," *Proceedings, Southern Weed Science Society*, Vol 25, 1972, p 388.

Ridings, W. H. and Zettler, F. W., "Aphanomyces Blight of Amazon Sword Plant," *Phytopathology*, Vol 62, 1972, Abstract, p 806; also Vol 63, 1973, pp 289-295.

Rintz, R. E. and Freeman, T. E., "*Fusarium roseum* Pathogenic to Waterhyacinth in Florida," *Phytopathology*, Vol 62, 1972, p 806.

Rintz, R. E., *Location, Identification and Characterization of Pathogens of the Waterhyacinth*, - Ph. D. Dissertation, University of Florida, Gainesville, Fla., 1973.

Zettler, F. W. and Freeman, T. E., "Potential for the Use of Plant Pathogens as Biocontrol Agents of Weeds," *Proceedings, Second International Congress of Plant Pathology*, St. Paul, Minn., 1973.

**Table 1**  
**Infection of Crop Plants by "Acremonium zonatum" and "Rhizoctonia solani" After**  
**Spray Inoculation with a Mycelial and Spore Suspensions**

Crop	Variety	Infection	
		"R. solani"	"A. zonatum"
Cabbage	Charleston Wakefield	--*	--
Cantaloupe	Hales Best Jumbo	x**	--
Carrot	Imperator	--	--
Celery	Giant Pascal	--	--
Collard	Georgia	--	--
Cucumber	Poinsett	xx†	--
Eggplant	Florida Market	--	--
Endive	Green curled	--	--
Escarole	Batavian	--	--
Field corn	PAB 751	--	--
Grapefruit	Duncan	--	--
Irish potato	Sebago	--	--
Lettuce	Great Lake	--	--
Lima bean	Henderson	--	--
Mustard	Florida Broadleaf	--	--
Oats	Fulghum	--	--
Okra	Clemson Spineless	--	--
Onion	White Globe	--	--
Orange	Temple	--	--
Peanut	Florunner	--	--
Pole bean	Kentucky Wonder	xxx††	--
Radish	Scarlet Globe	--	--
Rye	Weser	--	--
Slash pine	--	--	--
Snap beans	Harvester	--	--
Soybeans	Bragg	--	--
Southern peas	Cream 40	--	--
Squash	Early Summer Crookneck	--	--
Strawberry	Florida 90	--	--
Sugarcane	CL 41-223	--	--
Sweet corn	Silver Queen	--	--
Sweet pepper	Yolo L	--	--
Sweet potato	?	--	--
Tangerine	Dancy	--	--
Tobacco	Turkish NN	--	--
Tomato	Homestead	--	--
	Manalucia	--	--
	Walter	--	--
	MH 1	--	--
Watermelon	Congo	xxx	--
	Charleston Grey	xxx	--
Wheat	Holden	--	--

\* No infection.  
 \*\* Moderate infection.  
 † Severe infection.  
 †† Mild infection.

Table 2  
Plants Included in Greenhouse and Field Host Range Test for  
Susceptibility to "Cercospera Rodmanii"

Plant	Variety or Description	Resistance
Avocado		..*
Beans	Harvester	..
Beans	Lima Fordhook bush	..
Beans	Runner Kentucky Wonder (pole type)	..
Beans	Speckled butter Jackson Wonder	..
Beans	White baby lima Henderson bush	..
Beets	Detroit Dark Red	..
Cabbage	Charleston Wakefield	..
Cantaloupe	Hale's Best	..
Carrots	Imperator	..
Celery	Giant Pascal	..
Collard	Georgia	..
<i>Colocasia</i> sp.		..
Corn	Field corn PAG 751	..
Corn	Funks 5945**	..
Corn	Funks 4762**	..
Corn	Florida Sweet	..
Corn	Golden Hybrid NK 199†	..
Corn	Silver Queen	..
Cotton	Sea Island	..
Cotton	DPL 16**	..
Cotton	Stoneville 213**	..
Cowpeas	Zipper Cream	..
Cucumber	Pickling F1 hybrid†	..
Eggplant	Florida Market	..
<i>Hydrocotyle umbellata</i> **		..
Grapefruit		..
Lemon		..
Lettuce	Great Lakes bibb (sporulated on senescent leaves)	..
Red mangrove		..
Mustard	Florida Broadleaf	..
Oats	Fulghum	..
Okra	Clemson Spineless	..
Onion	Purple	..
Onion	Red	..
Onion	White	..
Orange	Sour	..
Orange	Sweet	..
Peanuts	Florunner	..
Peas	Cream Acre	..
Peppers	Banana	..
Peppers	Hot Hungarian	..
Pine tree	Slash	..
<i>Pontederia cordata</i>		..
Potatoes	Irish Red La Soda	..
Pyracantha		..
Radishes	Scarlet Red Globe	..
Rice	Saturn**	..
Rye	Weser	..
Squash	Early Summer Crookneck	..
Strawberry	Florida 90	..
Soybeans	Bragg**	..
Soybean	Davis	..
Sugarcane		..
Sweet potatoes	Georgia Red	..
Tangerine		..
Tobacco	Turkish	..
Tomatoes	Homestead	..
Tomatoes	MH-1	..
Tomatoes	Manalucia**	..
Tomatoes	Walter	..
Waterhyacinth	Inoculum check	..
Watermelon	Charleston Grey	..
Watermelon	Crimson Sweet†	..
Wheat	Holden	..

\* Not susceptible.  
\*\* Greenhouse only.  
† Field only.

**Table 3**  
**Fungi Found Associated with Waterhyacinth**  
**During Winter and Spring Months**

Fungi	Pathogenicity
<b>Subdivision</b>	
<b>Ascomycotina</b>	
Class—Pyrenomycetes	
1. <i>Melanospora</i> sp.	None
Class—Loculoascomycetes	
2. <i>Leptosphaerulina</i> sp.	None
3a. <i>Didymella exigua</i>	None
b. <i>Ascochyta</i> (imperfect stage)	None
4. <i>Mycosphaerella</i> sp.	Slight
<b>Deuteuromycotina</b>	
Class—Coelomycetes	
5. <i>Phoma</i> spp.	Varies—moderate to none
6. <i>Botryodiploidea</i> sp.	Slight
Class—Hyphomycetes	
7. Unknown synnematal fungus	None
8. <i>Mycoleptodiscus terrestris</i>	Slight to none
9. <i>Myrothecium cinctum</i>	None
10. <i>Epicoccum purpurascens</i>	None
11. <i>Alternaria</i> spp.	None
12a. <i>Aspergillus flavus</i>	None
b. <i>Aspergillus niger</i>	None
13. <i>Acremonium zonatum</i>	Good-excellent
14. <i>Bipolaris</i> spp.	Good—under investigation
15. <i>Cercospora</i> sp.	Good-excellent
16. <i>Cladosporium</i> spp.	None
17a. <i>Curvularia brachyspora</i>	None
b. <i>Curvularia penniseti</i>	Slight
18. <i>Dendryphiella infuscans</i>	Not tested
19. <i>Exserohilum prolatum</i>	Under investigation
20. <i>Memnoniella subsimplex</i>	Not tested
21. <i>Perionia echinoclaoe</i>	Slight
22. <i>Pithomyces chartarum</i>	None
23a. <i>Nigrospora oryzae</i>	None
b. <i>Nigrospora sphaerica</i>	None
24. <i>Thysanophora longispora</i>	None
25. <i>Scolecobasidium humicola</i>	None
26. <i>Stemphylium vericarium</i>	Under investigation
27. <i>Ustilaginoidea</i> sp. (?)	Not tested

**Table 4**  
**Virulence of "Bipolaris stenospila" from the Dominican Republic (353),  
 American Type Culture Collection (13447) and Florida (14 b) to  
 Waterhyacinth and Four Gramineaceous Plants**

<u>Isolate</u>	<u>Waterhyacinth</u>	<u>Bermudagrass</u>	<u>Sugarcane</u>	<u>Corn*</u>	<u>Oat*</u>
353	4+**	3+	2+	0	0
13447	2+	2+	2+	Not tested	Not tested
14 b	2+	3+	2+	Not tested	Not tested

- 
- \* Varieties resistant and susceptible to Helminthosporium were tested.
  - \*\* Rating of pathogenicity was based on the degree of damage. Replicated three times (plant). Experiment repeated twice with grasses and four times with waterhyacinth.

In accordance with ER 70-2-3, paragraph 6c(1)(b), dated 15 February 1973, a facsimile catalog card in Library of Congress format is reproduced below.

Freeman, T            E

Biological control of aquatic weeds with plant pathogens, by T. E. Freeman, R. Charudattan, K. E. Conway, [and] F. W. Zettler, Department of Plant Pathology, University of Florida, Gainesville, Florida. Vicksburg, U. S. Army Engineer Waterways Experiment Station, 1976.

25 p. illus. 27 cm. (U. S. Waterways Experiment Station. Contract report A-76-2)

Prepared for Office, Chief of Engineers, U. S. Army, Washington, D. C., under Contract No. DACW73-73C-0049.

References: p. 19-21.

1. Aquatic plant control. 2. Aquatic weeds. 3. Biological control. 4. Pathogens. I. Charudattan, R., joint author. II. Conway, K. E., joint author. III. Zettler, F. W., joint author. IV. Florida. University, Gainesville. Dept. of Plant Pathology. V. U. S. Army. Corps of Engineers. (Series: U. S. Waterways Experiment Station, Vicksburg, Miss. Contract report A-76-2)  
TA7.W34c no.A-76-2