

AD-A052 875

ARMY INST OF DENTAL RESEARCH WASHINGTON D C

F/G 6/1

FERMENTATION CHARACTERISTICS OF STRAINS OF 'STREPTOCOCCUS MUTAN--ETC(U)

MAR 78 T C LYON

UNCLASSIFIED

NL


| OF |
AD
A052875



END
DATE
FILMED

5 -78

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER	2. GOVT ACCESSION NO.	4. RECIPIENT'S CATALOG NUMBER
6	9	11
3. TITLE (and Subtitle) Fermentation Characteristics of Strains of <i>Streptococcus mutans</i>		5. REPORT NUMBER COVERED Dec 69 - Mar 72
7. AUTHOR T. C. Lyon, Jr.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Institute of Dental Research Walter Reed Army Medical Center Washington, DC 20012		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command ATTN: (SGRD-RP) Washington, DC 20012		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS DA OD 6034
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 29 Mar 78
		13. NUMBER OF PAGES 5
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <i>Streptococcus mutans</i> , sucrose, xylitol, fermentation, dental caries		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Realization of the important role of <i>Streptococcus mutans</i> in the initiation of dental caries has resulted in extensive studies of all its biologic characteristics. The ultimate goal is to control or eliminate it from the oral cavity. One means of accomplishing this is to reduce or eliminate its carbohydrate source. This study reports those carbohydrates which <i>Streptococcus mutans</i> cannot metabolize.		

DDC
RECEIVED
APR 19 1978
F

038 670 -

alt

UDC FILE COPY AD A052875

FERMENTATION CHARACTERISTICS OF STRAINS OF *Streptococcus mutans*

T. C. Lyon, Jr.

Chief, Divisions of Preventive Dentistry
and Clinical Specialties
U. S. Army Institute of Dental Research
Walter Reed Army Medical Center
Washington, D. C. 20012

Running head: Fermentation Patterns of *S. mutans*

Synopsis

Realization of the important role of *Streptococcus mutans* in the initiation of dental caries has resulted in extensive studies of all its biologic characteristics. The ultimate goal is to control or eliminate it from the oral cavity. One means of accomplishing this is to reduce or eliminate its carbohydrate source. This study reports those carbohydrates which *Streptococcus mutans* cannot metabolize.

ACCESSION FOR	
W. H. Section	☒
Re-File Section	☐
EX	
DISTRIBUTION/AVAILABILITY CODES	
DISC.	MAIL, SIG/ or SPECIAL
A	

FERMENTATION CHARACTERISTICS OF STRAINS OF *Streptococcus mutans*

Although the role of sucrose in a number of systemic diseases is not clearly defined, there is little doubt as to its significance in the initiation and progression of dental caries (Newbrun, J. Dent. Child. 36:13-22, 1969). Among the most significant factors in this relationship is the formation of long-chain polymers of glucose (mutans) when sucrose is utilized as a carbohydrate source. The ability of *Streptococcus mutans* to produce such polymers has led to its consideration as a major etiologic agent in the initiation of a carious lesion (Loesche, et al., Inf. and Immun. 11:1252-1260, 1975). Knowledge of this particular aspect of sucrose metabolism has resulted in the search for a substitute sweetening agent. Xylitol shows great promise of being such an agent. Not only is it an effective sweetener, but recent studies suggest that it prevents initiation of the carious process (Scheimer, et al., Acta Odontol. Scand. 32:383-412, 1974). However, as the result of a report which suggested that it may have carcinogenic potential, studies within the United States have been stopped. Thus, until further evidence is available concerning this aspect of the metabolism of xylitol, it seems advisable to consider the overall carbohydrate fermentation characteristics of *S. mutans*. Such information may broaden the range of sweeteners which could be used as sucrose substitutes.

Included in this study were the following strains of *S. mutans*: HS6, AHT, and E49 (serotype a); FA-1 (serotype b); NCTC 10449 (serotype c); and S1-1 and 6715-10 (serotype d). Each liter of fermentation medium contained 20g trypticase, 2.0g K_2HPO_4 , 0.01g $MgSO_4 \cdot 7H_2O$, and 20g of sugar,

with pH adjusted to 6.8. Sugars which could not be autoclaved were filter sterilized (see Table). Control tubes contained no sugar. All tubes were incubated at 37°C for 14 days. After reaching room temperature, tubes were shaken well and pH was measured using a Beckman pH meter. Unless otherwise indicated, readings represent an average of two tubes.

The findings (Table) indicate that all strains failed to utilize arabinose, glycerol, inositol, and melezitose as carbohydrate sources. Using only strains 10449, HS-6, and FA-1, the study was expanded to include cellobiose, galactose, maltose, mannose, melibiose, rhamnose, trehalose, and xylose. These strains failed to ferment rhamnose and xylose.

Of the carbohydrates which *S. mutans* failed to ferment, only glycerol is currently used as a commercial sweetener. Consideration should be given to the use of one, or a combination of the other sugars, i.e., arbinose, inositol, melezitose, rhamnose, or xylose, as sucrose substitutes.

FERMENTATION CHARACTERISTICS OF STRAINS OF *S. mutans*

Strains of <i>S. mutans</i>	Arabinose**	Glucose**	Glycerol	Isitol	Inulin	Lactose	Mannitol	Melezitose	Raffinose	Salicin	d Sorbitol	Starch
HS 6**	6.4	4.6	7.2	7.3	5.1	5.0	4.8	6.7	5.0	5.1	5.3	5.6
AHT	6.6	4.5	6.9	6.8	4.4	4.5	4.6	6.9	4.5	4.6	4.6	5.5
E 49	6.4	4.5	7.0	6.8	4.7	4.7	4.8	7.0	4.8	4.6	4.8	6.0
FA 1	6.7	4.5	7.0	7.3	4.7	4.5	4.8	7.2	4.7	4.7	4.9	5.3
10449 NCTC	6.4	4.4	6.9	6.9	4.3	4.5	4.5	6.6	4.5	4.6	4.8	5.0
SL-1	6.5	4.3	7.0	7.0	4.5	4.5	4.8	6.9	7.0	7.1	4.7	5.2
6715-10	6.5	4.4	6.8	6.8	4.6	4.6	4.9	6.9	6.8	6.9	4.9	5.2
Control**	6.4	6.4	7.4	7.4	7.3	7.0	7.1	6.9	7.2	7.5	7.4	7.3

*Filter sterilized - cellobiose, galactose, maltose, mannose, and trehalose were also filter sterilized

**pH recorded from a single tube