AD-	A056 89	6 UNI CAL MAT	TED TE CULATI 78 H	CHNOLOG ON OF T H MICH	HE ENE	SEARCH RGETICS H HOBB	CENTER AND CO	EAST H	ARTFORD N DYNAM F 78-1187	CONN ICS OF 49620-	F/ ELECTR 77-C-00 N	G 20/5 0ETC 64 L	(U)
	OF   AD56 896				genner.	astan.						The second secon	
And the second		Concernante Concernante		ALL DATE OF THE OWNER	-						B.Ť.D		1 Pilling
				and a second sec	- - - - - - - - - - - - - - - - - - -			angeogra		END DATE FILMED 9 78			
			34										
1												3.	

UTRC-R78-922853 UNCLASSIFIED SECURITY O ASSINCATION OF THIS PAGE (When Date En EPORT DOCUMENTATION PAGE BEFORE COMPLETING FORM 2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER AFUSR 8 TYPE OF REPORT & PERIOD COVERED CALCULATION OF THE ENERGETICS AND COLLISION DYNAMICS OF ELECTRONIC TRANSITION LASERS . Interim re 50 H. Harvey Michels F49620-77-C-0064 R. H./Hobbs 9. PERFORMING ORGANIZATION NAME AND ADDRESS PROGRAM ELEMENT, PROJECT, TASK United Technologies Research CenterJ Silver Lane 61102F East Hartford, Conn 06108 23Ø3/B1 1. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NC May 178 Bolling AFB DC 20332 B ER OF PAGES 35 14. MONITORING AGENCY NAME & ADDRESS(II dillerent from 15. SECURITY CLASS. (of this report) UNCLASSIFIED DECLASSIFICATION/DOWNGRADING 154. 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, it different from Report) DDC RORUNR 18. SUPPLEMENTARY NOTES AUG 1 1978 5 SIT 9. KEY WORDS (Continue on reverse side if necessary and identify by block number) ELECTRONIC STRUCTURE POTENTIAL ENERGY CURVES Q (+) 1 sub 1 8 2 Sigma(+) subju C 2 Signatty subj & 20 ABSTRACT (Continue on reverse side Hingcessery and identify by block number) The research performed under this contract was directed toward the calculation of the electronic structure of selected chemical systems in order to determine their potential as laser systems. The systems studied included the  $0^{2}$  symmetry of Hg<sub>2</sub>, the A( $\Sigma_{U}$ ) and B( $\Sigma_{T}$ ) states for Hg<sub>2</sub>U, all low-lying  $2\Sigma$  and  $2\pi$  states of HgTl. A list is given of le the publication and talks resulting from this program. DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLET UNCLASSIFIED

# UNITED TECHNOLOGIES RESEARCH CENTER



East Hartford, Connecticut 06108

SOCOA DA

R78-922853 Approved for public release; distribution unlimited.

Calculation of the Energetics and Collision Dynamics of Electronic Transition Lasers

Interim Scientific Report

Contract No. F49620-77-C-0064 Project-Task 2303/B1

ACCESSION fo	r
NTIS DDC WANNOUNCED JUSTIFICATION	White Soction 🖄 Buff Soction 🗆
DISTRIBUTIO	N/AVAILABILITY CODES
Dist.	AVAIL and/or SPECIAL
A	

REPORTED BY	,	H.	X.	hichele	
	Н.	н.	Mic	hels	

R. H. Hobbs

APPROVED BY Brild A. Peterson

DATE May 1978

NO. OF PAGES \_\_\_\_\_\_\_

COPY NO.

78 07 26 097

[]

transma a

-

Π

1

0

0

## FOREWORD

This report was prepared for the Air Force Office of Scientific Research, United States Air Force by the United Technologies Corporation, Research Center, East Hartford, Connecticut, under Contract F49620-77-C-0064, Project-Task No. 2303/B1. The performance period for the technical program was from 1 April 1977 through 31 March 1978. The project monitor was Major Robert A. Osteryoung, USAF.

#### ACKNOWLEDGMENTS

- International Property in the second secon

1

It is a pleasure to acknowledge the advise and assistance rendered by the author's associates. We are indebted to Judith B. Addison for her help both with computer programming and with the analysis of the results of this research. Her efforts have been of real value during the course of this work. We also wish to acknowledge many fruitful discussions with research associates and the beneficial collaboration with Dr. Gerald A. Peterson on several research problems. Finally, we acknowledge the continued support and encouragement given by Lt.Col.John T. Viola of AFOSR and Major David S. Olson of AFWL whose confidence in the value of this research has been of real help.

# Calculation of the Energetics and Collision Dynamics of Electronic Transition Lasers

0

0

0

0

[]

Π

Π

I

I

I

0

0

0

0

## TABLE OF CONTENTS

Section	Page
ABSTRACT	. 1
STATEMENT OF WORK	• 2
REVIEW OF TECHNICAL EFFORT	. 3
RESEARCH PROGRESS DURING THE CONTRACT YEAR	. 5
PUBLICATIONS AND PRESENTATIONS	. 7
A. Technical Reports in Journals	. 7
<ul> <li>B. Technical Reports in Books</li> <li>C. Technical Paper and Lecture Presentations</li> </ul>	. 9
ILLUSTRATIONS	. 9
TABLES	. 15
APPENDICES	. 19

## ILLUSTRATIONS

I

]

]

0

0

1

0

0

]]

1

]

1

-

-

-

1

]]

[]

Figure		Page
1	Low-Lying Potential Energy Curves for SiF	9
2	Correlation Diagram for Intermediate Coupling of the Hg <sub>2</sub> Molecule	10
3	Ground State Potential Energy Curve for Hg <sub>2</sub>	11
4	Low-Lying Potential Energy Curves for Hg2+	12
5	Correlation Diagram for Intermediate Coupling of the HgTl Molecule	13
6	Low-Lying Potential Energy Curves for MgNa	14

Q

U

U

.

-

[

-

-

-

[]

0

[]

0

0

# TABLES

Table		Page
1	SiF/SiF <sub>2</sub> Correlations	15
II	Low-Lying Molecular States of SiF and Their Dissociation Limits	16
111	Low-Lying Molecular States of $\text{Hg}_2^+$ and Their Dissociation Limits	17
IV	Low-Lying Molecular States of MgNa and Their Dissociation Limits	18

# APPENDICES

1

1

I

I

I

Ţ

i i i

[]

1

0

Π.

IJ

]

]

]

]

Appendix		Page
A	Electronic Structure of the Noble Gas Dimer Ions	19
В	Configuration Interaction Studies of the HeH <sup>+</sup> Molecular Ion. IV. The Triplet Sigma, Pi and Delta States	20
с	Electronic Structure of the Noble Gas Dimer Ions. I. Potential Energy Curves and Spectroscopic Constants	21
D	Electronic Structure of the Noble Gas Dimer Ions. II. Absorption Spectrum for the A ${}^{2}\Sigma_{u}^{+} \rightarrow D  {}^{2}\Sigma_{g}^{+}$ System	22
E	Electronic Structure of the Noble Gas Dimer Ions. III. Absorption Specthum for the A ${}^{2}\Sigma^{+}_{u} \rightarrow B {}^{2}\Pi$ System	23
F	Electronic Structure of Excited States of Selected Atmospheric Systems	24

#### ABSTRACT

Knowledge of the partition of energy between the products of a chemical reaction is fundamental to the understanding of the dynamics of collisional reactions. Atom-atom and atom-molecule reactions that yield vibrationally excited diatomic molecules are of particular scientific and military interest. An understanding of the dynamics of the simplest of these reactions is a prerequisite for understanding more complicated rearrangement gas phase reactions. A knowledge of at least limited regions of potential energy surfaces is of importance in analyzing the vibrational and rotational relaxation rates and can be utilized in a predictive fashion. In the low energy regime there are little data available on collision processes and experimental measurements are difficult. Thus, a program to develop theoretical techniques and construct computer programs for calculating potential energy surfaces and kinetic cross sections for simple collision processes is appropriate. The potential energy surfaces will include both atom-atom and atom-diatom systems with special emphasis on molecular reactions yielding electronically excited products which have potential as visible or UV chemical lasers. These systems shall include metal oxides and halides, metal excimers and the noble gashalides. This research will develop techniques and computational schemes for a priori prediction of the kinetics and the product distribution of atomdiatom reactions. This will involve (1) improved calculation techniques for potential energy surfaces for the reactants and products, (2) quantum and classical mechanical calculations of the dynamics of the reaction and (3) detailed calculations for systems with high potential as electronic transition lasers.

## Calculation of the Energetics and Collision Dynamics of Electronic Transition Lasers

STATEMENT OF WORK

The contractor shall:

a. Perform a theoretical research investigation of the electronic structure and energetics of excimer systems and on chemical reactions in such systems which would yield a sufficient density of excited electronic species capable of lasing in the visible or UV spectrum regions. Calculate electronic wave-functions and energies as a function of internuclear separation followed by calculations of electronic transition moments for systems selected as typical of this class, and which appear to offer a high potential as candidate excimer lasers. In addition, research shall be carried out related to the development of practical computational techniques for prediction of the kinetics and the product distribution of chemical reactions involving electronically excited reactants or products. Experimental data will be iteratively examined and integrated with the results generated from this theoretical investigation. The computations shall be carried out using density functional and ab initio methods, using computer codes developed or modified at this Center. Specifically, this investigation shall comprise the following:

(1) Conduct research related to the development of practical computational techniques for defining potential energy surfaces for reactants and products with possible electronic excitation and illustrate the utility of such techniques by a study of the long-range interactions in the Si +  $F_2$ system. This portion of the research has the highest priority.

(2) Survey existing experimental and theoretical data available to the Contractor pertaining to excimer systems.

(3) Perform quantum mechanical calculations of the electronic wavefunctions, electronic transition moments, and radiative lifetimes for three (3) diatomic metal excimer systems. These systems shall be chosen from the following:  $Hg_2$ ,  $Cd_2$ , CdHg, TlHg, TlCd. If other chemical systems are subsequently identified as potential candidates these can be substituted.

1

Π

#### REVIEW OF TECHNICAL EFFORT

Calculations of the dynamics of molecular collisions are ordinarily carried out with the aid of the Born-Oppenheimer separation of the electronic and nuclear motion. One proceeds by calculating the electronic energy as a function of the positions of the nuclei, which are assumed to be stationary. This electronic energy, plus the electrostatic repulsion between the nuclei, defines a potential energy hypersurface on which the nuclei may be regarded as moving. A potential energy hypersurface defined in this way is referred to as adiabatic, and is appropriate for describing the nuclear motion in the limit of low velocity. There are many collisions for which an adiabatic potential energy hypersurface provides an adequate description. However, most reactive collisions and many collisions which do not lead to reaction are inadequately described by an adiabatic potential energy hypersurface. These collisions are characterized by velocities of nuclear motion sufficient to affect adversely the Born-Oppenheimer separation, with the result that the overall wavefunction must be described as a superposition of terms involving different electronic energy states. Under these conditions, it will be useful to consider adiabatic potential energy hypersurfaces corresponding to all electronic states relevant to the overall wavefunction.

When the different potential energy hypersurfaces are well separated in energy, the nuclear motion can ordinarily be described in terms of motion on a single hypersurface. However, when two or more hypersurfaces are close in energy, they can be expected to mix appreciably in the overall wavefunction, and it will then be necessary not only to calculate the hypersurfaces but also to calculate the quantities needed to discuss their mixing in the overall wavefunction.

The calculation of a point on a potential energy hypersurface is equivalent to calculating the energy of a diatomic or polyatomic system for a specified nuclear configuration, and therefore will present considerable practical computational difficulty. For certain problems or nuclear configurations, the maximum possible accuracy will be needed and under these conditions relatively elaborate <u>ab initio</u> methods are indicated. Under other conditions, it may be possible to use less elaborate and more rapid computational methods, and density functional or other approaches may then prove useful.

A knowledge of at least limited regions of potential energy surfaces is of importance in analyzing the electronic, vibrational and rotational relaxation rates in chemical laser systems. These relative relaxation rates govern both the feasibility of lasing and the power that is potentially available. Such relaxation rates are governed by long-range forces between atomic and

3

molecular fragments and thus knowledge of the dissociation behavior represented by the potential energy surfaces for the laser system can be used in calculations of the kinetic behavior.

Our reserach effort to date has been devoted to the theoretical development of computational techniques for the prediction of potential energy surfaces, and to studies of the energetics of prototype electronic transition laser systems, the prediction of the radiative lifetimes of electronically excited species, and studies of photon energy loss paths such as photoabsorption or ionization of the upper excited electronic states. The goal of this research program is to develop a technical information base in support of the development of a practical chemical laser system operating on an electronic transition.

## RESEARCH PROGRESS DURING THE CONTRACT YEAR

Electronic structure calculations were performed for the Si+F<sub>2</sub> system at various geometries. Studies were carried out for the ground X  ${}^{1}A_{1}$  state, the first excited A  ${}^{1}B_{1}$  state and for the separated atom-molecule states of Si ( ${}^{3}P$ ) + F<sub>2</sub>( ${}^{1}\Sigma_{g}^{+}$ ). The SiF/SiF<sub>2</sub> correlations are shown in Table I. Our calculated excitation energy is in good agreement with experiment. These calculations also indicate a maximum in the lowest  ${}^{1}A_{1}$  potential energy surface relative to Si+F<sub>2</sub> similar to that found in the diatomic molecule SiF. These results are shown in Fig. 1. The corresponding atomic correlations are given in Table II. The origin of the maximum in the A  ${}^{2}\Sigma^{+}$  state of SiF is an avoided crossing with the  ${}^{2}\Sigma^{+}$  ionic state connecting to Si<sup>+</sup> ( ${}^{2}P$ ) + F<sup>-</sup> ( ${}^{1}S$ ). The origin of the maximum in SiF<sub>2</sub> is an avoided crossing with the  ${}^{1}A_{1}$  surface arising from SiF<sup>+</sup>(X ${}^{1}\Sigma^{+}$ ) + F<sup>-</sup>( ${}^{1}S$ ). This maximum suggests that favorable branching should occur to the higher metastable a  ${}^{4}\Sigma^{-}$  state of SiF in the exothermic reaction of Si+F<sub>2</sub>. Similar chemistry should be observed in the GeF and SnF systems. Further studies are in progress.

Calculations have also been initiated on the Hg<sub>2</sub>, Hg<sub>2</sub><sup>+</sup>, MgNa and TlHg molecules. These studies include density functional calculations with explicit treatment of all core electrons. The symmetries  $0_g^+$  for Hg<sub>2</sub>, A  $2 \varepsilon_u^+$  and B  $2 \varepsilon_g^+$  for Hg<sub>2</sub><sup>+</sup>, all low-lying  $2 \varepsilon$  and  $2 \pi$  states of MgNa and the  $2 \varepsilon$  and  $2 \pi$  states of HgTl have been examined. The results obtained to date are summarized for each system below.

## Hg2

Annual S

A correlation diagram for intermediate coupling of the Hg<sub>2</sub> molecule is shown in Fig. 2. Density functional calculations have been carried out on the ground X  ${}^{1}\Sigma_{g}^{+}$  state of the system. The potential energy curve for X  ${}^{1}\Sigma_{g}^{+}$  of Hg<sub>2</sub> is presented in Fig. 3. We find a calculated dissociation energy of 0.2 eV, in good agreement with experiment.

# Hg2+

A correlation diagram for the low-lying molecular states of  $Hg_2^+$  and their dissociation limits is shown in Table III. Density functional calculations were carried out for the ground A  ${}^{2}\Sigma_{u}^{+}$  state and the excited B  ${}^{2}\Sigma_{g}^{+}$  state dissociating to the Hg  ${}^{1}S_{g}$  (6s<sup>2</sup>) + Hg  ${}^{+}S_{g}$ (6s) asymptotic limit. These potential energy curves are presented in Fig. 4. Calculations of the radiative lifetimes for the A  ${}^{2}\Sigma_{u}^{+}$  + B  ${}^{2}\Sigma_{g}^{+}$  system is in progress. Our calculated dissociation energy for the ground A  ${}^{2}\Sigma_{u}^{+}$  state of Hg ${}^{+}_{2}$  is 0.64 eV. There are no experimental data for comparison.

## HgT1

Detailed density functional calculations have been carried out on the low-lying doublet states of HgTl. A correlation diagram for the intermediate coupling of the HgTl molecule is shown in Fig. 5. Further studies on this system are in progress. 1

-----

1

-----

1

1

-

## MgNa

The low-lying doublet sigma and pi states of MgNa have been studied in detail. A correlation diagram for the low-lying molecular states of MgNa and their dissociation limits is shown in Table IV. Ab initio calculations were carried out on the  $^{2}\Sigma$  and  $^{2}\Pi$  symmetries. These states are presented in Fig. 6. Further studies, including the ionic state, are in progress.

#### PUBLICATIONS AND PRESENTATIONS

The significant research results obtained under this Contract have been prepared for publication in technical journals and books or presented at technical meetings. These papers and meetings are listed below. Abstracts of the published papers are included in the Appendices to this report.

#### A. Technical Reports in Journals

- "Electronic Structure of the Noble Gas Dimer Ions". H. H. Michels, R. H. Hobbs and L. A. Wright, to be published in the International Journal of Quantum Chemistry, 1978 Sanibel Symposium Issue, 1978.
- "Configuration Interaction Studies of the HeH<sup>+</sup> Molecular Ion. IV. The Triplet Sigma, Pi and Delta States." T. A. Green, H. H. Michels and J. C. Browne, to be published in the Journal of Chemical Physics, July 1, 1978.
- "Electronic Structure of the Noble Gas Dimer Ions. I. Potential Energy Curves and Spectroscopic Constants." H. H. Michels and R. H. Hobbs, to be published in the Journal of Chemical Physics, 1978.
- 4. "Electronic Structure of the Noble Gas Dimer Ions. II. Absorption Spectrum for the A  ${}^{2}\Sigma_{u}^{+} \rightarrow D {}^{2}\Sigma_{g}^{+}$  System." H. H. Michels and R. H. Hobbs, to be published in the Journal of Chemical Physics, 1978.
- 5. "Electronic Structure of the Noble Gas Dimer Ions. III. Absorption Spectrum for the A  ${}^{2}\Sigma_{u}^{+} \rightarrow B {}^{2}\Pi_{g}$  System." H. H. Michels and R. H. Hobbs, to be published in the Journal of Chemical Physics, 1978.

## B. Technical Articles in Books

 "Electronic Structure of Excited States of Selected Atmospheric Systems." H. H. Michels, to be published as a chapter in Volume II of the "Excited State in Chemical Physics". Edited by J. Wm. McGowan, John Wiley & Sons, New York, 1978.

## C. Technical Paper and Lecture Presentations

 "Calculations of the Energetics and Radiative Properties of Visible/ UV Chemical Laser Systems." H. H. Michels. Presented at the AFOSR/AWFL Contractors' Meeting, Kirtland Air Force Base, Albuquerque, New Mexico, May 16-20, 1977.

7

- "Photodissociation of Noble Gas Dimer Ions." H. H. Michels. Presented at the 30th Annual Gaseous Electronics Conference, Palo Alto, California, October 18-21, 1977.
- "Theoretical Studies of the Energetics of Electronic Transition Lasers." H. H. Michels. Presented at the AFOSR/AFGL Chemical Dynamics Contractors' Meeting, Air Force Geophysics Lab, Hanscom Air Force Base, Bedford, Massachusetts, October 26-27, 1977.
- "Electronic Structure of Noble Gas Dimer Ions." Presented at the 1978 Sanibel Symposium on Atomic, Molecular and Solid State Theory, Sheraton Palm Coast Inn, Daytona, Florida, March 10-16, 1978.

1

- ----

----

----

-----

1

and a

# LOW-LYING POTENTIAL ENERGY CURVES FOR SIF

.



FIG. 1

76-12-41-1



# CORRELATION DIAGRAM FOR INTERMEDIATE COUPLING OF THE Hg2 MOLECULE



UNIQUE ASSIGNMENTS FROM (J1, J2) TO (A, S) COUPLING ARE INDICATED BY SOLID LINES

10

Trent .

1

0

1

1

-

-----

1

1

1

-

T

Conception of the local division of the loca

Constanting of the



## (DENSITY-FUNCTIONAL METHOD)



78-06-50-2

FIG. 3



R (a.u.)

78-06-50-1

1

]]

-

and a

L

-

the state

[]

]

1

[]

FIG. 4

aptendent.

Francisco Constants

and a second second

I

E

# CORRELATION DIAGRAM FOR INTERMEDIATE COUPLING OF THE TIH MOLECULE



UNIQUE ASSIGNMENTS FROM (J1. J2) TO (A,S) COUPLING ARE INDICATED BY SOLID LINES



# LOW-LYING POTENTIAL ENERGY CURVES FOR MgNa

FIG. 6

0

- series

( international

-

]

-----

]

]

]

14

and the second se

-----

COMPACT.

- Charles

Π

and a second

# TABLE I

SIF/SIF<sub>2</sub> CORRELATIONS

Dissociation Limit	<u>eV</u>	Molecular States
SIF $(X^2 \pi) + F (^2 P)$	0.0	$SiF_2 (1, 3_{A_1}, 1, 3_{B_1})$
SIF (A $2z^+$ ) + F ( $2P$ )	2.83	SiF <sub>2</sub> ( <sup>1,3</sup> A <sub>1</sub> , <sup>1,3</sup> B <sub>1</sub> )
SiF (a ${}^{4}\Sigma^{-}$ ) + F ( ${}^{2}P$ )	3.70	$SiF_2 ({}^{3,5}A_1, {}^{3,5}B_1)$
$SiF^{+}(X^{1}\Sigma^{+}) + F^{-}(^{1}S)$	3.81	$siF_2$ ( <sup>1</sup> A <sub>1</sub> )

## TABLE II

0

0

The second second second second

1

the second

[]

[]

LOW-LYING MOLECULAR STATES OF SIF AND THEIR DISSOCIATION LIMITS

Dissociation Limit	eV	Molecular States
Si + F		2 - 4 - 2 +
${}^{3}P_{g} (3s^{2}3p^{2}) + {}^{2}P_{u} (2p^{5})$	0.0	$\begin{array}{c} 2\Sigma^{-}(2), & 4\Sigma^{-}(2), & 4\Sigma^{-}(2), \\ 4\Sigma^{+}(1), & 2\Pi(2), & 4\Pi(2), & 2\Delta(1), \\ 4\Delta(1) \end{array}$
${}^{1}D_{g} (3s^{2}3p^{2}) + {}^{2}P_{u} (2p^{5})$	0.781	$2 \Sigma^{-}(1), 2 \Sigma^{+}(2), 2 \Pi(2), 2 \Delta(2), 2 \Phi(1)$
${}^{1}S_{p}(3s^{2}3p^{2}) + {}^{2}P_{u}(2p^{5})$	1.908	$2 \Sigma^{+}(1), 2 \Pi(1)$
${}^{5}S_{u} (2s3p^{3}) + {}^{2}P_{u} (2p^{5})$	4.131	$4_{5}(1), 6_{5}(1), 4_{\pi}(1), 6_{\pi}(1), 6_{\pi$
Si <sup>+</sup> + F <sup>-</sup>		
${}^{2}P_{\mu} (3s^{2}3p) + {}^{1}S_{g} (2p^{6})$	4.701	$2^{2} \pm (1), 2^{\pi}(1)$

# TABLE III

# LOW-LYING MOLECULAR STATES OF Hg + AND THEIR DISSOCIATION LIMITS

$ {}^{1}s_{g}(5d^{10}6s^{2}) + {}^{2}s_{g}(5d^{10}6s) \qquad 0.0 \qquad {}^{2}z_{g}^{+}(1), {}^{2}z_{u}^{+}(1) \\ {}^{3}P_{u}(5d^{10}6s6p) + {}^{2}s_{g}(5d^{10}6s) \qquad 5.004 \qquad {}^{2}z_{g}^{+}(1), {}^{2}z_{u}^{+}(1), {}^{4}z_{g}^{+}(1), \\ {}^{4}z_{u}^{+}(1), {}^{2}\pi_{g}(1), {}^{2}\pi_{u}(1), \\ {}^{4}\pi_{g}(1), {}^{4}\pi_{u}(1) \\ \\ {}^{1}s_{g}(5d^{10}6s^{2}) + {}^{2}D_{g}(5d^{9}6s^{2}) \qquad 5.334 \qquad {}^{2}z_{g}^{+}(1), {}^{2}z_{u}^{+}(1), \\ {}^{2}\pi_{g}(1), {}^{2}\pi_{u}(1), \\ {}^{2}\Delta_{g}(1), {}^{2}\Delta_{u}(1) \\ \\ {}^{1}P_{u}(5d^{10}6s6p) + {}^{2}s_{g}(5d^{10}6s) \qquad 6.702 \qquad {}^{2}z_{g}^{+}(1), {}^{2}z_{u}^{+}(1), {}^{2}\pi_{g}(1), \\ {}^{2}\pi_{u}(1) \\ \\ {}^{1}s_{g}(5d^{10}6s^{2}) + {}^{2}P_{u}(5d^{10}6p) \qquad 6.948 \qquad {}^{2}z_{g}^{+}(1), {}^{2}z_{u}^{+}(1), \\ {}^{2}\pi_{g}(1), {}^{2}\pi_{u}(1) \\ \end{array} $	Dissociation Limits Hg + Hg <sup>+</sup>	<u>eV</u>	Molecular States
${}^{3}P_{u}(5d^{10}6s6p) + {}^{2}S_{g}(5d^{10}6s) \qquad 5.004 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{4}\Sigma_{g}^{+}(1), {}^{4}\Sigma_{g}^{+}(1), {}^{4}\Sigma_{g}^{+}(1), {}^{4}\Sigma_{g}^{+}(1), {}^{4}\Sigma_{u}^{+}(1), {}^{2}\Lambda_{u}(1), {}^{4}\Lambda_{u}(1)$ ${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}D_{g}(5d^{9}6s^{2}) \qquad 5.334 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{2}\Lambda_{u}(1), {}^{2}\Lambda_{u$	${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}S_{g}(5d^{10}6s)$	0.0	$2z_{g}^{+}(1), 2z_{u}^{+}(1)$
${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}D_{g}(5d^{9}6s^{2}) \qquad 5.334 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), \\ {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1), \\ {}^{2}\Delta_{g}(1), {}^{2}\Delta_{u}(1) \\ {}^{1}P_{u}(5d^{10}6s6p) + {}^{2}S_{g}(5d^{10}6s) \qquad 6.702 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{2}\Pi_{g}(1), \\ {}^{2}\Pi_{u}(1) \\ {}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}P_{u}(5d^{10}6p) \qquad 6.948 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), \\ {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1) \\ {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1) \end{cases}$	${}^{3}P_{u}(5d^{10}6s6p) + {}^{2}S_{g}(5d^{10}6s)$	5.004	${}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{4}\Sigma_{g}^{+}(1), \\ {}^{4}\Sigma_{u}^{+}(1), {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1), \\ {}^{4}\Pi_{g}(1), {}^{4}\Pi_{u}(1)$
${}^{1}P_{u}(5d^{10}6s6p) + {}^{2}S_{g}(5d^{10}6s) \qquad 6.702 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1)$ ${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}P_{u}(5d^{10}6p) \qquad 6.948 \qquad {}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1), {}^{2}\Pi_{u}(1), {}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1), {$	${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}D_{g}(5d^{9}6s^{2})$	5.334	${}^{2}\Sigma_{g}^{+}(1), {}^{2}\Sigma_{u}^{+}(1),$ ${}^{2}\Pi_{g}(1), {}^{2}\Pi_{u}(1),$ ${}^{2}\Delta_{g}(1), {}^{2}\Delta_{u}(1)$
${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}P_{u}(5d^{10}6p) \qquad 6.948 \qquad {}^{2}\Sigma_{g}^{+}(1), \; {}^{2}\Sigma_{u}^{+}(1), \\ {}^{2}\Pi_{g}(1), \; {}^{2}\Pi_{u}(1)$	$P_{u}(5d^{10}6s6p) + {}^{2}S_{g}(5d^{10}6s)$	6.702	$2 \Sigma_{g}^{+}(1), 2 \Sigma_{u}^{+}(1), 2 \Pi_{g}(1),$ $2 \Pi_{u}(1)$
	${}^{1}S_{g}(5d^{10}6s^{2}) + {}^{2}P_{u}(5d^{10}6p)$	6.948	$2z_{g}^{+}(1), 2z_{u}^{+}(1),$ $2\pi_{g}(1), 2\pi_{u}(1)$

# TABLE IV

0

1

1

[]

0

1

# LOW-LYING MOLECULAR STATES OF MgNa AND THEIR DISSOCIATION LIMITS

Dissociation Limits	eV	Molecular States
Mg + Na		
${}^{1}S_{g}(3s^{2}) + {}^{2}S_{g}(3s)$	0.0	$2\Sigma^{+}(1)$
${}^{1}S_{g}(3s^{2}) + {}^{2}P_{u}(3p)$	2.103	$2\Sigma^{+}(1), 2\pi(1)$
${}^{3}P_{u}(3s3p) + {}^{2}S_{g}(3s)$	2.712	$2_{\Sigma^{+}(1)}, 4_{\Sigma^{+}(1)}, 2_{\Pi}(1), 4_{\Pi}(1)$
${}^{1}S_{g}(3s^{2}) + {}^{2}S_{g}(4s)$	3.191	<sup>2</sup> <sup>2</sup> <sup>+</sup> (1)

 $Mg^{+} + Na^{-}$  ${}^{2}S_{g}(3s) + {}^{1}S_{g}(3s^{2})$ 

6.975  $2\Sigma^+(1)$ 

## APPENDIX A

## ELECTRONIC STRUCTURE OF THE NOBLE GAS DIMER IONS\*

H. H. Michels and R. H. Hobbs United Technologies Research Center East Hartford, Connecticut 06108

> L. A. Wright Kirtland Air Force Base New Mexico 87117

## ABSTRACT

A systematic study of the electronic structure and chemical binding in the dimer ion sequence, Ne<sub>2</sub><sup>+</sup>, Ar<sub>2</sub><sup>+</sup>, Kr<sub>2</sub><sup>+</sup> and Xe<sub>2</sub><sup>+</sup>, has been carried out using both density functional and <u>ab initio</u> configuration interaction computational approaches. This study includes detailed calculations of the pertinent potential energy curves and an analysis of the calculated spectroscopic properties of the bound A  ${}^{2}\Sigma_{u}^{+}$  (1/2<sub>u</sub>I) state of these ions.

Supported in part by the Air Force Office of Scientific Research under Contract F49620-77-C-0064

## APPENDIX B

CONFIGURATION INTERACTION STUDIES OF THE HeH<sup>+</sup> MOLECULAR ION. IV. THE TRIPLET SIGMA, PI AND DELTA STATES.\*

> T. A. Green Sandia Laboratories Albuquerque, New Mexico 87115

H. H. Michels United Technologies Research Center East Hartford, Connecticut 06108

J. C. Browne Department of Physics and Computer Sciences University of Texas Austin, Texas 78712

#### ABSTRACT

The method of superposition of configurations was applied to the triplet sigma, pi, and delta states of HeH<sup>+</sup> which correlate to the separated atom states of principle quantum number less than or equal to 3. The calculations were done for internuclear separations,  $0 \le R \le 65.5$  a.u., on a mesh adequate for interpolation. Similar calculations on the singlet states have already been reported. The present calculations complete the accurate evaluation of the potential energy curves for this system which are required for low- and intermediate-energy collision studies. In addition to the energy eigenvalues and eigenfunctions, dipole, gradient and radial coupling matrix elements were calculated for the sigma and pi states. Primarily, this paper presents information on the eigenvalues. The accuracy of the tripletstate calculations is comparable to that obtained for the singlet states. The similarities and differences in the pattern of avoided crossings for the triplet and singlet states are exhibited. These are mainly determined by the differences in the triplet and singlet energy-level schemes of the united and separated atoms.

I

<sup>\*</sup> Supported in part by the Air Force Office of Scientific Research under Contract F49620-77-C-0064.

#### APPENDIX C

## ELECTRONIC STRUCTURE OF THE NOBLE GAS DIMER IONS. 1. POTENTIAL ENERGY CURVES AND SPECTROSCOPIC CONSTANTS\*

H. H. Michels and R. H. Hobbs United Technologies Research Center East Hartford, Connecticut 06108

#### ABSTRACT

A systematic study of the electronic structure and chemical binding in the dimer ion sequence, Ne<sub>2</sub><sup>+</sup>, Ar<sub>2</sub><sup>+</sup>, Kr<sub>2</sub><sup>+</sup> and Xe<sub>2</sub><sup>+</sup>, has been carried out using both density functional and <u>ab initio</u> configuration-interaction computational techniques. This study includes detailed calculations of the pertinent potential energy curves and an analysis of the calculated spectroscopic properties of the bound states of these ions. A regular progression is found in the spectroscopic properties for the ground A  ${}^{2}\Sigma^{+}_{1/2u}$  state which leads to some remarkably simple conclusions concerning the nature of the binding and the size of these dimer ions. For the heavier systems, Kr<sub>2</sub><sup>+</sup> and Xe<sub>2</sub><sup>+</sup>, spinorbit coupling becomes important, resulting in a strong mixture of the  $\Lambda$ -s coupled  $\Sigma$  and  $\Pi$  states. This mixing affects the strength of the binding in the ground state. A comparison with other <u>ab initio</u> studies and an analysis of the asymptotic behavior at large internuclear separations is given.

These dimer ion species illustrate the classic Hartree-Fock symmetry dilemma arising from improper dissociation character. The nature of this problem for ionized homopolar species is discussed.

 Supported in part by the Air Force Office of Scientific Research under Contract F49620-77-C-0064.

## APPENDIX D

ELECTRONIC STRUCTURE OF THE NOBLE GAS DIMER IONS. II. ABSORPTION SPECTRUM FOR THE A  ${}^{2}\Sigma_{g}^{+} \rightarrow D {}^{2}\Sigma_{g}^{+}$  SYSTEM\*

> H. H. Michels and R. H. Hobbs United Technologies Research Center East Hartford, Connecticut 06108

## ABSTRACT

A systematic study of the electronic structure and chemical binding in the dimer ion sequence, Ne<sub>2</sub><sup>+</sup>, Ar<sub>2</sub><sup>+</sup>, Kr<sub>2</sub><sup>+</sup> and Xe<sub>2</sub><sup>+</sup>, has been carried out using both density function and <u>ab initio</u> configuration interaction computational approaches. From the results of this study, the absorption spectrum for the UV transitions of the A  ${}^{2}\Sigma_{u}^{+} \rightarrow D {}^{2}\Sigma_{g}^{+}$  system has been calculated.

\* Supported in part by the Air Force Office of Scientific Research under Contract F49620-77-C-0064. I

## APPENDIX E

ELECTRONIC STRUCTURE OF THE NOBLE GAS DIMER IONS. III. ABSORPTION SPECTRUM FOR THE A  ${}^{2}\Sigma_{u}^{+}$  + B  ${}^{2}\Pi_{g}$  SYSTEM

> H. H. Michels and R. H. Hobbs United Technologies Research Center East Hartford, Connecticut 06108

## ABSTRACT

A systematic study of the electronic structure and chemical binding of the dimer ion sequence, Ne<sub>2</sub><sup>+</sup>, Ar<sub>2</sub><sup>+</sup>, Kr<sub>2</sub><sup>+</sup> and Xe<sub>2</sub><sup>+</sup>, has been carried out using both density functional and <u>ab initio</u> configuration interaction computational approaches. From the results of this study, the absorption spectrum for the visible/IR transition of the A  ${}^{2}\Sigma_{u}^{+}$  + B  ${}^{2}\Pi_{g}$  system has been calculated.

 Supported in part by the Air Force Office of Scientific Research under Contract F49620-77-C-0064.

#### APPENDIX F

## ELECTRONIC STRUCTURE OF EXCITED STATES OF SELECTED ATMOSPHERIC SYSTEMS\*

## H. H. Michels United Technologies Research Center East Hartford, Connecticut 06108

#### I. INTRODUCTION

In order to evaluate the thermodynamic and radiation properties of a natural or perturbed state of the upper atmosphere or ionosphere, the thermal and transport properties of heated air are required. Such properties are also of particular interest in plasma physics, in gas laser systems, and in basic studies of airglow and the aurora. In the latter area, the release of certain chemical species into the upper atmosphere results in luminous clouds that display the resonance electronic-vibrational-rotation spectrum of the released species. Such spectra are seen in rocket releases of chemicals for upper atmospheric studies and upon reentry into the atmosphere of artificial satellites. Of particular interest in this connection is the observed spectra of certain metallic oxides and air diatomic species. From band intensity distribution of the spectra, and knowledge of the f-values for electronic and vibrational transitions, the local conditions of the atmosphere can be determined (Ref. 1).

Present theoretical efforts which are directed toward a more complete and realistic analysis of the transport equations governing atmospheric relaxation and the propagation of artifical disturbances require detailed information of thermal opacities and LWIR absorption in regions of temperature and pressure where molecular effects are important (Refs. 2, 3). Although various experimental techniques have been employed for both atomic and molecular systems, theoretical studies have been largely confined to an analysis of the properties (bound-bound, bound-free and free-free) of atomic systems (Refs. 4, 5). This has been due in large part to the unavailability of reliable wavefunctions for diatomic molecular systems, and particularly for excited states

and a

and a second

\* Supported in part by AFOSR under Contract No. F44620-73-C-0077.

or states of open-shell structures. More recently (Refs. 6-9), reliable theoretical procedures have been prescribed for such systems which have resulted in the development of practical computational programs.

The theoretical analysis of atmospheric reactions requires the knowledge of the electronic structure of the ground and low-lying excited states of atoms, ions and small molecular clusters of nitrogen and oxygen and, in certain regions, the interaction of water or other small molecules with these clusters. Because of the computational complexity for systems with large numbers of electrons, traditional <u>ab initio</u> theoretical methods are difficult and expensive to apply. However, no clear alternative choice is currently on the horizon, although progress continues in semiempirical and perturbative approaches to the calculation of electronic structure. Such approaches have met with limited success in applications to electronically excited states of molecules. <u>Ab initio</u> computational programs based on the variational theorem, incorporating analytic basis functions and coupled with a configuration interaction analysis, remain the backbone of our techniques, for studies to chemical accuracy, which are applicable to systems such as the atmospheric diatomic molecules and ions.

The potential curves derived from such calculations can often be empirically improved by comparison with so-called experimental curves derived from observes spectroscopic data, using RKR or other inversion procedures. It is often found, particularly for the atmospheric systems, that the remaining correlation errors in a configuration interaction (CI) calculation are similar for many excited electronic states of the same symmetry or principle molecular orbital description. Thus, it is often possible to calibrate an entire family of calculated excited state potential curves to near-spectroscopic accuracy. Such a procedure has been applied to the systems described here.

The particular choice of the atmospheric systems to be presented here proved to be somewhat of a problem. Calculations for the N<sub>2</sub> molecule, including all low-lying valence and Rydberg states, had been performed several years ago (Ref. 10) as had a fairly extensive set of calculations for  $0_2^-$ (Ref. 11). Gilmore has described the known spectroscopic states for N<sub>2</sub>,  $0_2$ , NO and several states of their corresponding positive and negative ions (Ref. 12). A complete analysis of all of the valence and low-lying Rydberg states for N<sub>2</sub>,  $0_2$  and NO including ions, seemed too formidable (and expensive) a task, so the systems N<sub>2</sub>,  $0_2$ , NO,  $0_2^-$  and NO<sup>+</sup> have been selected for inclusion in this review. This choice includes the most important atmospheric positive ion, NO<sup>+</sup>, and the chemically most interesting negative ion,  $0_2^-$ .

The general composition of this article is as follows. We first present a critical review of the current status of electronic structure calculations for molecular systems. This is followed by a compilation of the potential energy curves, derived spectroscopic analysis and pertinent discussion of the atmospheric molecules selected above.

annual a

1

-----

L

I

I

#### REFERENCES

- Harang, O.: AlO Resonant Spectrum for Upper Atmosphere Temperature Determination. AFCRL-66-314, Environmental Research Paper No. 192, 1966.
- Churchill, D. R. and R. E. Meyerott: J. Quant. Spectr. Radiative Transfer, 5, 69, 1965.
- The Airglow and the Aurorae, edited by E. B. Armstrong and A. Dalgarno, Pergamon Press, New York, 1955.
- Armstrong, B. H., R. R. Johnston and P. S. Kelly, Jr.: J. Quant. Spectr. Radiative Transfer, 5, 55, 1965.
- Johnston, R. R., B. H. Armstrong and O. R. Platas: J. Quant. Spectr. Radiative Transfer, 5, 49, 1965.
- 6. Harris, F. E. and H. H. Michels: Int. J. Quantum Chem., IS, 329, 1967.
- Krauss, M.: Compendium of <u>ab initio</u> Calculations of Molecular Energies and Properties. NBS Technical Note 438, 1967.
- Wahl, A. C., P. J. Bertoncini, G. Das and T. L. Gilbert: Int. J. Quantum Chem., <u>IS</u>, 123, 1967.
- Schaefer, H. F.: Electronic Structure of Atoms and Molecules. Addison-Wesley Publishing Co., 1972.
- 10. Michels, H. H.: J. Chem. Phys., 53, 841, 1970.
- Michels, H. H.: Electronic and Atomic Collisions, Volume 2, edited by L. M. Branscomb, North Holland Publishing Co., Amsterdam, 1170, 1971.
- 12. Gilmore, F. R.: J. Quant. Spectr. Radiative Transfer, 5, 369, 1965.