

# @LEVEL #

# DDC FILE COPY AD AO 62207

# NPS55-78-019 NAVAL POSTGRADUATE SCHOOL Monterey, California





SIMULATION METHODS FOR POISSON PROCESSES IN NONSTATIONARY SYSTEMS

by

Peter A. W. Lewis and Gerald S. Shedler

AUGUST 1978

Approved for public release; distribution unlimited.

78 12 11 173

NAVAL POSTGRADUATE SCHOOL MONTEREY, CALIFORNIA

Rear Admiral Tyler F. Dedman Superintendent Jack R. Borsting Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

Ken U.W

Peter A. W. Lewis, Professor Department of Operations Research

edler (by P.A.w.L)

Gerald S. Shedler IBM Research Laboratory

Reviewed by:

Michael G. Sovereign, Chairman // Department of Operations Research Released by:

1 olles

William M. Tolles Dean of Research

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS
REPORT NUMBER 2. GOVT A	CCESSION NO. 3. RECIPIENT'S CATALOG NUMBER
HNPS55-78-019	
TITLE (and Subilitie)	5. TYPE OF REPORT & PERIOD COVERE
Simulation Methods for Poisson Processe	s in (9) Technical pept.
Nonstationary Systems	
AUTHOR(.)	8. CONTRACT OR GRANT NUMBER(*)
Peter A. W. Lewis and G. S. Shedler	
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TAS
Naval Postgraduate School	AREA & WORK UNIT NUMBERS
Monterey, Ca. 93940	N0001478WR80035
Office of Naval Research	August 1978
Arlington, VA 22217	13. NUMBER OF PAGES
4. MONITORING AGENCY NAME & ADDRESS(It different from Contr	rolling Office) 15. SECURITY CLASS. (of this report)
APLACE AN UNIT	Unclassified
10/NSF-AF-476	15. DECLASSIFICATION/DOWNGRADING SCHEDULE
6. DISTRIBUTION STATEMENT (of this Report)	
Approved for public release; distributi	on unlimited.
13,390	
E a. L	
7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20,	, if different from Report)
16)RRAL405 / MIRRA	448804
A grunp.	TIOSOT
8. SUPPLEMENTARY NOTES	
9. KEY WORDS (Continue on reverse aide if necessary and identify h	y block number)
Nonhomogeneous Poisson processes	
Simulation; intensity function	
Thinning algorithm; gap statistics	
	block number)
0. ABSTRACT (Continue on reverse side if necessary and identify by	is a widely used model for a series
The nonhomogeneous Poisson process	
The nonhomogeneous Poisson process of events (stochastic point process) in	which the "rate" or "intensity" of
The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of po
The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables. and that t
The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde number of points in any of these interv	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables, and that als has a Poisson distribution. In
The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde number of points in any of these interv this paper we first discusseseveral gen	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables, and that als has a Poisson distribution. In eral methods for simulation of the -
The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde number of points in any of these interv this paper we first discusseseveral gen	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables, and that t als has a Poisson distribution. In eral methods for simulation of the
<ul> <li>ABSTRACT (Continue on reverse side if necessary and identity by The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde number of points in any of these interv this paper we first discusses everal gen D 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE S/N 0102-014-6601  </li> </ul>	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables, and that t als has a Poisson distribution. In eral methods for simulation of the
<ul> <li>ABSTRACT (Continue on reverse side if necessary and identity by The nonhomogeneous Poisson process of events (stochastic point process) in occurrence of points varies, usually wi characteristic properties that the numb overlapping intervals are mutually inde number of points in any of these interv this paper we first discusses everal gen D FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE S/N 0102-014-6601   5</li> </ul>	which the "rate" or "intensity" of th time. The process has the er of points in any finite set of no pendent random variables, and that t als has a Poisson distribution. In eral methods for simulation of the ECURITY CLASSIFICATION OF THIS PAGE (When Date En

# UNCLASSIFIED

# SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

# 20. ABSTRACT Continued.

one-dimensional nonhomogeneous Poisson process; these include time-scale transformation of a homogeneous (rate one) Poisson process via the inverse of the integrated rate function, generation of the individual intervals between points, and generation of a Poisson number of order statistics from a fixed density function.

We then state a particular and very efficient method for simulation of nonhomogeneous Poisson processes with log-linear rate function. The method is based on an identity relating the nonhomogeneous Poisson process to the gap statistics from a random number of exponential random variables with suitably chosen parameters. This method can also be used, at the cost of programming complexity and some memory, as the basis for a very efficient technique for simulation of nonhomogeneous Poisson processes with more complicated rate functions such as a log-quadratic rate function.

> Finally, we describe a simple and relatively efficient new method for simulation of one-dimensional and two-dimensional nonhomogeneous Poisson processes. The method is applicable for any given rate function and is based on controlled deletion of points in a Poisson process with a rate function that dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates. The thinning method is also applicable to the generation of individual intervals between points, as is required in many programs for discrete-event simulations.

MTIS	Section D
000	Contion T
L'ANNART (*	0
JUSTITICALC	1
EISTRIBUTION,	AVAN ARI ITY CODEC
BISTRIBUTION,	AVAILABILITY CODES
EISTRIBUTION Dist. AVAIL	AVAILABILITY CODES
DISTRIBUTION, Dist. AVAIL	AVAILABILITY CODES
DIST. AVAIL	AVAILABILITY CODES

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

# SIMULATION METHODS FOR POISSON PROCESSES IN NONSTATIONARY SYSTEMS

Peter A. W. Lewis Naval Postgraduate School Monterey, California

Gerald S. Shedler IBM Research Laboratory San Jose, Californía

# ABSTRACT

The nonhomogeneous Poisson process is a widely used model for a series of events (stochastic point process) in which the "rate" or "intensity" of occurrence of points varies, usually with time. The process has the characteristic properties that the number of points in any finite set of nonoverlapping intervals are mutually independent random variables, and that the number of points in any of these intervals has a Poisson distribution. In this paper we first discuss several general methods for simulation of the one-dimensional nonhomogeneous Poisson process; these include time-scale transformation of a homogeneous (rate one) Poisson process via the inverse of the integrated rate function, generation of the individual intervals between points, and generation of a Poisson number of order statistics from a fixed density function.

We then state a particular and very efficient method for simulation of nonhomogeneous Poisson processes with log-linear rate function. The method is based on an identity relating the nonhomogeneous Poisson process to the gap statistics from a random number of exponential random variables with suitably chosen parameters. This method can also be used, at the cost of programming complexity and some memory, as the basis for a very efficient technique for simulation of nonhomogeneous Poisson processes with more complicated rate functions such as a log-quadratic rate function.

Finally, we describe a simple and relatively efficient new method for simulation of one-dimensional and two-dimensional noahomogeneous Poisson processes. The method is applicable for any given rate function and is based on controlled deletion of points in a Poisson process with a rate function that dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates. The thinning method is also applicable to the generation of individual intervals between points, as is required in many programs for discrete-event simulations.

# 1. INTRODUCTION

The one-dimensional nonhomogeneous (nonstationary) Poisson process (see e.g., [5, pp. 28-29; 3, pp. 94-101]) has the characteristic properties that the numbers of points in any finite set of nonoverlapping intervals are mutually independent random variables, and that the number of points in any interval has a Poisson distribution. The most general nonhomogeneous Poisson process can be defined in terms of a monotone, nondecreasing, right-continuous function  $\Lambda(x)$  which is bounded in any finite interval. Then the number of points in any finite interval, for example  $(0, x_0)$ , has a Poisson distribution with parameter  $\mu_0 = \Lambda(\mathbf{x}_0) - \Lambda(0)$ . In this paper we assume that  $\Lambda(x)$  is continuous, but not necessarily

# POISSON PROCESSES IN NONSTATIONARY SYSTEMS...Continued

absolutely continuous. The right derivative  $\lambda(\mathbf{x})$  of  $\Lambda(\mathbf{x})$  is the rate function for the process;  $\Lambda(\mathbf{x})$  is called the integrated rate function and has the interpretation that for  $\mathbf{x} \geq 0$ ,  $\Lambda(\mathbf{x}) - \Lambda(0) =$  $E[N(\mathbf{x})]$ , where N( $\mathbf{x}$ ) is the total number of points in (0, $\mathbf{x}$ ]. Note that  $\lambda(\mathbf{x})$  may jump at points at which  $\Lambda(\mathbf{x})$  is not absolutely continuous. In contrast to the homogeneous Poisson process, i.e.,  $\lambda(\mathbf{x})$  a constant (usually denoted by  $\lambda$ ), the intervals between the points in a one-dimensional nonhomogeneous Poisson process are neither independent nor identically distributed.

Applications of the one-dimensional nonhomogeneous Poisson process include modelling of the incidence of coal-mining disasters [5], the arrivals at an intensive care unit [12], transaction processing in a data base management system [15], occurrences of major freezes in Lake Constance [23], and geomagnetic reversal data [22]. The statistical analysis of trends in a one-dimensional nonhomogeneous Poisson process, based on the assumption of an exponeatial polynomial rate function, is discussed by [4, 5, 12 and 15].

One-dimensional nonhomogeneous Poisson processes are often used as models for event streams when there is gross inhomogeneity in a system, e.g., time of day effect or long-term growth in use of a facility. It is important to be able to simulate these processes since analytic results are difficult to obtain. This is particularly true in the context of queueing systems; see e.g., [19]. The methods given here for simulation of the one-dimensional nonhomogeneous Poisson process have application, for example, to study of the length of a queue at a toll booth at a time corresponding to the peak traffic time, or to study of the arrivals at an intensive care unit where the probability of a

bed being free at a time corresponding to the peak of arrivals from afternoon operations is of interest. Note that in simulations of nonhomogeneous systems of this kind, estimates of measures of system behavior will be based on multiple replications.

The two-dimensional homogeneous Poisson process (of rate  $\lambda > 0$ ) is defined by the properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any region of area A has a Poisson distribution with mean  $\lambda A$ ; see e.g., [10, pp. 31-32]. Note that the number of points in a region R depends on its area, but not on its shape. The homogeneous Poisson process arises as a limiting twodimensional point process with respect to a number of limiting operations; cf., [7,8]. Properties of the process are given in [18]. Applications of the two-dimensional homogeneous Poisson process to problems in ecology and forestry have been discussed in [24] and [9]. The model also arises in connection with naval search and detection problems.

The two-dimensional nonhomogeneous Poisson process is characterized by a (continuous) positive rate function  $\lambda(x,y)$ . Applications of the two-dimensional nonhomogeneous Poisson process include problems in forestry as well as naval search and detection. The detection and statistical analysis of trends in the two-dimensional nonhomogeneous Poisson process is discussed in [21].

# 2. <u>SIMULATION OF THE ONE-DIMENSIONAL</u> NONHOMOGENEOUS POISSON PROCESS

There are a number of methods for simulating one-dimensional nonhomogeneous Poisson process which we review briefly. Time-scale transformation of a homogeneous (rate one) Poisson process via the inverse of the integrated rate function  $\Lambda(x)$  constitutes a first general method; cf. [3, pp. 96-97]. This method is based on the result that X1, X2, ... are the points in a nonhomogeneous Poisson process with continuous integrated rate functions  $\Lambda(x)$ if and only if  $X'_1 = \Lambda(X_1)$ ,  $X'_2 = \Lambda(X_2)$ ,..., are the points in a homogeneous Poisson process of rate one. The time-scale transformation method is a direct analogue of the inverse probability integral transformation method for generating (continuous) nonuniform random numbers. For many rate functions, inversion of  $\Lambda(x)$  is not simple and must be done numerically; cf., [6] and [20]. The resulting algorithm for simulation of the nonhomogeneous Poisson process may be far less efficient than simulation based on other methods.

A second general method for simulating a one-dimensional nonhomogeneous Poisson process with integrated rate function  $\Lambda(\mathbf{x})$ is to generate the intervals between points individually, an approach which may seem more natural in the event scheduling approach to simulation. Thus, given the points  $X_1 = x_1, X_2 = x_2, ..., X_i = x_i, with$  $X_1 < X_2 < \cdots < X_i$ , the interval to the next point, X<sub>i+1</sub> - X<sub>i</sub>, is independent of x1,..., xi-1 and has distribution function  $F(x) = 1 - exp[-\{\Lambda(x_i + x) - \Lambda(x_i)\}].$  It is possible to find the inverse distribution function  $F^{-1}(\cdot)$ , usually numerically, and generate  $X_{i+1} - X_i$  according to  $X_{i+1} - X_i = F^{-1}(U_i)$ , where  $U_i$  is a uniform random number on the interval (0,1). Note, however that this not only involves computing the inverse distribution function for each interval  $X_{i+1} - X_i$ , but that each distribution has different parameters and possibly a different form. An additional complication is that  $X_{i+1} - X_i$  is not necessarily a proper random variable, i.e. there may be positive probability that X<sub>i+1</sub> - X<sub>i</sub> is infinite. It is necessary to take this into account for each interval X<sub>i+1</sub> - X<sub>i</sub> before the inverse probability

integral transformation is applied. The method is therefore very inefficient with respect to speed, more so than the time-scale transformation method.

In a third method, simulation of a non-homogeneous Poisson process in a fixed interval  $(0, x_0]$  can be reduced to the generation of a Poisson number of order statistics from a fixed density function by the following result; cf., [5, p. 45]. If  $X_1, X_2, \ldots, X_n$  are the points of the nonhomogeneous Poisson process in  $(0, x_0]$ , and if  $N(x_0) = n$ , then conditional on having observed n ( > 0) points in  $(0, x_0]$ , the X, are distributed as the order statistics from a sample of size n from the distribution function  $\{\Lambda(\mathbf{x}) - \Lambda(\mathbf{0})\}/\{\Lambda(\mathbf{x}_0) - \Lambda(\mathbf{0})\},\$ defined for  $0 < x \leq x_0$ . Simulation of the nonhomogeneous Poisson process based on order statistics is in general more efficient (with respect to speed) than either of the previous two methods. Of course, a price is paid for this greater efficiency. First, it is necessary to be able to generate Poisson variates, and second, more memory is needed than in the interval-byinterval method to store the sequence of points. Enough memory must be provided so that with very high probability the random number of points generated in the interval can be stored. Recall that the number of points in the interval (0,x0] has a Poisson distribution with mean  $\mu_0 = \Lambda(x_0) - \Lambda(0)$ . Memory of size, e.g.,  $\mu_0 + 4\mu_0^{1/2}$  will ensure that overflow will occur on the average in only 1 out of approximately 40,000 realizations. This probability is small enough so that in case of overflow, the realization of the process generally can be discarded.

We now summarize several recently developed methods for simulating one- and two-dimensional nonhomogeneous Poisson processes. These methods are discussed in greater detail in [14, 16, 17].

# POISSON PROCESSES IN NONSTATIONARY SYSTEMS...Continued

# 3. SIMULATION USING GAP STATISTICS

In a previous paper [14], we have considered the simulation of nonhomgeneous Poisson processes with degree-one exponential polynomial rate function, i.e.,  $\lambda(\mathbf{x})$  of the form

 $\lambda(\mathbf{x}) = \exp\{\gamma_0 + \gamma_1 \mathbf{x}\} = \lambda \exp\{\gamma_1 \mathbf{x}\}, \gamma_1 \neq 0.$  (1)

The rate function (1) is the simplest of a general family of log-linear rate functions, i.e., rate functions whose logarithms are linear in the coefficients [4; 12] which are useful in analyzing nonhomogenerous Poisson processes. The rate function (1) represents a situation in which the rate is monotonically increasing or decreasing depending on whether  $\gamma_1$  is greater than or less than zero, with  $\gamma_1$ equal to zero giving a homogeneous Poisson process. The case where  $\gamma_1$  is less than zero and the case where  $\gamma_1$  is greater than zero are quite distinct; in the first situation  $\lambda(x) \neq 0$  as  $x \neq \infty$ , and in the second,  $\lambda(x) + \infty$  as  $x + \infty$ . Moreover, when  $\gamma_{1}$  is less than 0, the intervals between events are not proper random variables since there is a nonzero probability that there is no event after any fixed point x.

In [14] a method for simulating the nonhomogeneous Poisson process is given based on an identity relating the nonhomogeneous Poisson process with rate function (1) to the gap statistics from a random number of exponential random variables with suitably chosen parameters. This method avoids costly ordering and taking of logarithms required by direct simulation methods and is more efficient than time-scale transformation of a homogeneous Poisson process via the inverse of the integrated rate function  $\Lambda(x)$ .

Simulation of the one-dimensional nonhomogeneous Poisson process in a fixed interval  $(0, x_0]$  is more natural than simulation for a fixed

number of events since time, not serial number, is the basic parameter of the inhomogeneity discussed here. Thus the gap statistics algorithm for simulation of the nonhomogeneous Poisson process generates the sequence of times-to-events in a fixed interval. Although such a method requires more memory than successive generation of individual times until the next event, it is far more efficient.

We now state the method of Lewis and Shedler [14] for simulation via gap statistics of the one-dimensional nonhomogeneous Poisson process with rate function (1). This scheme, which is particular to the degree-one exponential polynomial rate function, can use standard packages for exponential random numbers (e.g., [13]) and obviates the need for ordering of the random numbers. It is based on the result (see [25]) that the gap process associated with a Poisson distributed (parameter  $-\lambda/\gamma_1 > 0$ ) number of exponential (parameter  $\beta = -\gamma_1 > 0$ ) gap statistics is a nonhomogeneous Poisson process with rate function  $\lambda(\mathbf{x}) = \lambda \exp(\gamma, \mathbf{x})$ on (0, ~). Efficient methods for generation of Poisson random numbers for which the generation time does not increase proportionally with the mean are given by [1,2] and [11].

Assuming the availability of a source of unit exponential random numbers  $E_1, E_2$ , ..., obtained by logarithms or by other methods, the resulting algorithm for generating the events in the nonhomogeneous Poisson process is as follows.

# <u>Algorithm 1</u>. Gap Statistics Technique $(\gamma_1 < 0)$ .

- 1. Generate m as a Poisson random number with parameter  $-\lambda/\gamma_1$ . If m = 0, exit; there are no events in  $(0,x_0]$ .
- 2. For m > 0, if  $E_1/(\beta m)$  is greater than  $x_0$ , exit; there are no events in  $(0,x_0]$ . Otherwise, set  $E_1/(\beta m)$  equal to  $X_1$ .

- 3. If  $E_2/\{\beta(m-1)\} + X_1 > x_0$ , then return  $X_1$  and exit. Otherwise, set it equal to  $X_2$ .
- 4. Continue, possibly for m times. If  $E_m/\beta + X_{m-1} > x_0$ , return  $X_1, X_2, \dots, X_{m-1}$ and exit. Otherwise, set this equal to  $X_m$ , return  $X_1, X_2, \dots, X_m$  and exit.

The case  $\gamma_1 > 0$  is handled in the same way as  $\gamma_1 < 0$  by using a time-reversal technique, as follows. Simulate according to Algorithm 1 with  $\lambda(x) = \lambda^{\#}$  $\times \exp{\{\gamma_1^{\#}x\}}$ , where  $\lambda^{\#} = \exp{\{\gamma_0 + \gamma_1 x_0\}}$ and  $\gamma_1^{\#} = -\gamma_1$ . The output of Algorithm 1 is a sequence  $x_1^{\#}, x_2^{\#}, \dots, x_n^{\#}$ . Then set  $x_1 = x_0 - x_n^{\#}, x_2 = x_0 - x_{n-1}^{\#}, \dots, x_n = x_0 - x_1^{\#}$ . These  $X_i$  are the required events in the nonhomogeneous Poisson process for  $\gamma_1 > 0$ .

Lewis and Shedler [16] consider the simulation in a fixed interval of the nonhomogeneous Poisson process with degreetwo exponential polynomial rate function

 $\lambda(\mathbf{x}) = \exp\{\alpha_0 + \alpha_1 \mathbf{x} + \alpha_2 \mathbf{x}^2\}, \quad \alpha_2 \neq 0$  (2)

the case  $\alpha_2 = 0$  giving the degree-one exponential polynomial rate function. Again, the case where  $\alpha_2$  is less than zero and the case where  $\alpha_2$  is greater than zero are distinct. The simulation method give: is based on representation of the process as a superposition of two independent nonhomogeneous Poisson processes, one of which has a fitted rate function of the form (1); simulation of the latter process is accomplished via the gap statistics algorithm [14]. A rejection-acceptance technique is used to generate the other, more complex, nonhomogeneous Poisson process. The resulting algorithm is more efficient than time-scale transformation of a homogeneous Poisson process; see [20]. This method can be improved by using the thinning algorithm given in the next section to simulate the second nonhomogeneous Poisson process. The method can also be

extended to more complex rate functions than the degree-two exponential polynomial.

# 4. <u>SIMULATION OF NONHOMOGENEOUS</u> POISSON PROCESSES BY THINNING

In this section we describe a new method [17] for simulating a nonhomogeneous Poisson process. The method is not only conceptually simple, but is also computationally simple and relatively efficient. In fact, at the cost of some efficiency, the method can be applied to simulate the given nonhomogeneous Poisson process without the need for numerical integration or routines for generating Poisson variates. Used in conjunction with the special methods given by Lewis and Shedler [14,16], the method can be used to simulate quite efficiently nonhomogeneous Poisson processes with rather complex rate function, in particular, combinations of long-term trends and fixed-cycle effects. The method is also easily extended to the problem of simulating the two-dimensional nonhomogeneous Poisson process (see Section 5), and of simulating conditional and doubly stochastic Poisson processes.

Simulation of a nonhomogeneous Poisson process with general rate function  $\lambda(x)$  in a fixed interval  $(0, x_0]$  can be based on thinning of a nonhomogeneous Poisson process with rate function  $\lambda^*(x) \ge \lambda(x)$ . The main result [17, Theorem 1] is that if  $x_1^*, x_2^*, \ldots, x_{N^*}(x_0)$  are the points of the process with rate function  $\lambda^*(x)$  in the interval  $(0, x_0]$  and if the point  $x_1^*$  is deleted with (independent) probability  $1 - \lambda(x_1^*)/\lambda(x_1^*)$ , then the remaining points form a nonhomogeneous Poisson process with rate function  $\lambda(x)$  in the interval  $(0, x_0]$ .

This result is the basis for the method of simulating one-dimensional nonhomogeneous Poisson processes on an interval  $(0, x_0)$ given by Algorithm 2.

# POISSON PROCESSES IN NONSTATIONARY SYSTEMS...Continued

# Algorithm 2. One-Dimensional Nonhomogeneous Poisson Process (Thinning)

- 1. Generate points in the nonhomogeneous Poisson process  $\{N^{*}(x)\}$  with rate function  $\lambda^{*}(x)$  in the fixed interval  $(0,x_{0}]$ . If the number of points generated, n<sup>\*</sup>, is such that n<sup>\*</sup> = 0, exit; there are no points in the process  $\{N(x)\}$ .
- 2. Denote the (ordered) points by  $X_1^*, X_2^*$ , ...,  $X_{n*}^*$ . Set i = 1 and k = 0.
- 3. Generate  $U_i$ , uniformly distributed between 0 and 1. If  $U_i \leq \lambda(X_i^*) / \lambda^*(X_i^*)$ , set k equal to k+1 and  $X_k = X_i^*$ .
- 4. Set i equal to i+1. If  $i \leq n^*$ , go to 3.
- 5. Return  $X_1, X_2, \dots, X_n$ , where n = k, and also n.

The method of thinning of Algorithm 2 is essentially the obverse of the conditional method of Section 2 using conditioning and acceptance-rejection techniques to generate the random variables with density function  $\lambda(\mathbf{x})/\{\Lambda(\mathbf{x}) - \Lambda(0)\}$  [16, Algorithm 3]. The differences are subtle, but computationally important. In the acceptancerejection method, it is first necessary to generate a Poisson variate with mean  $\mu_0 = \Lambda(\mathbf{x}_0) - \Lambda(0)$ , and this involves an integration of the rate function  $\lambda(\mathbf{x})$ . Then the Poisson( $\mu_0$ ) number, n, of variates generated by acceptance-rejection must be ordered to give  $X_1, X_2, \dots, X_n$ .

In the simplest form of the method of thinning,  $\lambda^*(\mathbf{x})$  is taken equal to  $\lambda^* = \max_0 \leq \mathbf{x} \leq \mathbf{x}_0^{\lambda}(\mathbf{x})$ , so that, for instance the points  $\mathbf{x}_1^*, \mathbf{x}_2^*, \dots, \mathbf{x}_{n^*}^*$  can be generated by cumulating exponential ( $\lambda^*$ ) variates until the sum is greater than  $\mathbf{x}_0$  (cf. [14, Algorithm 1]. The generated points are then thinned. No ordering, no integration of  $\lambda(\mathbf{x})$  and no generator of Poisson variates is required. Of course for both algorithms to be efficient, computation of  $\lambda(\mathbf{x})$  and  $\lambda^{*}(\mathbf{x})$  must be easy relative to computation of the inverse of  $\Lambda(\mathbf{x})$ .

For the thinning algorithm (as well as the algorithm based on conditioning and acceptance-rejection) efficiency as measured by the number of points deleted is proportional to

$$\mu_0/\mu_0^* = \{\Lambda(\mathbf{x}_0) - \Lambda(0)\}/\{\Lambda^*(\mathbf{x}_0) - \Lambda^*(0)\};$$

this is the ratio of the areas between 0 and  $\mathbf{x}_0$  under  $\lambda(\mathbf{x})$  and  $\lambda^*(\mathbf{x})$ . Thus,  $\lambda^*(\mathbf{x})$  should be as close as possible to  $\lambda(\mathbf{x})$  consistent with ease of simulating the nonhomogeneous Poisson process  $\{\mathbf{N}^*(\mathbf{x}):\mathbf{x} \geq 0\}$ .

It is important to note that the method of thinning can be used to generate individual intervals between events occurring in  $(0, x_0]$  if  $\lambda(x)$  is bounded on  $(0, x_0]$ . The resulting algorithm is not only useful in the event scheduling approach to simulation, but also in the generation of conditional Poisson processes. Informally, the "one at a time" thinning algorithm is as follows. If  $\lambda^*(x) = \lambda^* \ge \max_0 \le x \le x_0^{-\lambda}(x)$ , then the ith interval  $x_i - x_{i-1}$  is obtained by generating and cumulating exponential( $\lambda^*$ ) random numbers  $F_{i,1}^*, F_{i,2}^*, \ldots$ until, for the first time,

$$U_{i,j} \leq \lambda (X_{i-1} + E_{i,1}^{*} + \dots + E_{i,j}^{*}) / \lambda^{*} ,$$
  
i = 1,2,...; j = 1,2,...,

where the  $U_{i,j}$  are independent, uniform (0,1) random numbers.

# 5. <u>SIMULATION OF TWO-DIMENSIONAL</u> HOMOGENEOUS POISSON PROCESSES

Recall that the two-dimensional homogeneous Poisson process (of rate  $\lambda > 0$ ) has the characteristic properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any region of area A has a Poisson distribution with mean  $\lambda A$ .

In considering the two-dimensional homogeneous Poisson process, projection properties of the process depend quite critically on the geometry of the regions considered. These projection properties are simple for rectangular and circular regions, and make simulation of the homogenecus process quite easy. We consider here the case of a rectangular region. The following result forms the basis for simulation of the two-dimensional homogeneous **Poisson** process of rate  $\lambda$  in a fixed rectangle  $R = \{(x,y): 0 \le x \le x_0, 0 \le y \le y_0\}.$ If  $(X_1, Y_1)$ ,  $(X_2, Y_2)$ ,...,  $(X_N, Y_N)$  denote the positions of the points of the process in R, labelled so that  $X_1 < X_2 < \cdots$ , then  $x_1, x_2, \ldots, x_N$  form a one-dimensional homogeneous Poisson process on  $0 \le x \le x_0$  of rate  $\lambda y_0$ ; if the points are relabelled  $(X'_1, Y'_1), (X'_2, Y'_2), \dots, (X'_N, Y'_N)$  so that  $Y'_1 < Y'_2 < \cdots < Y'_N$ , then  $Y'_1, Y'_2, \dots, Y'_N$  form a one-dimensional homogeneous Poisson process on  $0 < y \leq y_0$  of rate  $\lambda x_0$ .

We state next conditional properties of the Poisson process in a rectangle. The important thing to note is that although the processes obtained by projection of the points onto the x and y axes are not independent, there is a type of conditional independence which makes it easy to simulate the two-dimensional process. Thus, conditional on having observed n > 0 points  $(X_1, Y_1)$ ,  $(X_2, Y_2)$ ,..., $(X_n, Y_n)$  in R, labelled so that  $X_1 < X_2 < \cdots < X_n$ , the  $X_1, X_2, \ldots, X_n$ are uniform order statistics on  $0 \le x \le x_0$ , and the  $Y_1, Y_2, \ldots, Y_n$  are independent and uniformly distributed on  $0 < y \le y_0$ , independent of the  $X_i$ .

From these two results, the following simulation procedure is obtained.

# Algorithm 3. Two-Dimensional Homogeneous Poisson Process in a Rectangle

- 1. Generate points in the one-dimensional homogeneous Poisson process of rate  $\lambda y_0$  on  $(0, x_0]$ . If the number of points generated, n, is such that n = 0, exit; there are no points in the rectangle.
- 2. Denote the points generated by  $x_1 < x_2 < \cdots < x_n$ .
- 3. Generate  $Y_1, Y_2, \ldots, Y_n$  independent, uniformly distributed random numbers on  $(0, y_0)$ .
- 4. Return  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$  as the coordinates of the two-dimensional homogeneous Poisson process in the rectangle, and n.

Note that generation of the points  $x_1, x_2$ , ..., $x_n$  in steps 1 and 2 can be accomplished by cumulating exponential  $(\lambda y_0)$  random numbers. Alternatively, after generating a Poisson random number N = n (with parameter  $\lambda x_0 y_0$ ), n independent, uniformly distributed random numbers on  $(0, x_0]$  can be ordered; see [14, p. 502].

The basis for another algorithm for simulation of the two-dimensional homogeneous Poisson process in a rectangle is the following corollary. Denote the Poisson points by  $(X_1, Y_1), (X_2, Y_2), \ldots$ , where the index does not necessarily denote an ordering on either axis. Conditionally, the pairs  $(X_1, Y_1),$  $(X_2, Y_2), \ldots, (X_N, Y_N)$  are independent random variables, and furthermore, for each pair  $(X_i, Y_i), X_i$  is distributed uniformly between 0 and  $x_0$ , independently of  $Y_i$ , which is uniformly distributed between 0 and  $y_0$ .

Direct generation of homogeneous Poisson points in noncircular or nonrectangular regions is difficult. The processes obtained by projection of the points on the two axes are nonhomogeneous Poisson processes with complex rate functions determined by the geometry of the region. However, the conditional independence which is found in

### POISSON PROCESSES IN NONSTATIONARY SYSTEMS...Continued

circular and rectangular regions for the processes on the two axes is not present. In particular, given that there are n points  $(X_1, Y_1), \ldots, (X_n, Y_n)$  in a nonrectangular region, the pairs  $(X_i, Y_i)$  are mutually independent, but  $X_i$  is in general not independent of  $Y_i$ ,  $i = 1, 2, \ldots, n$ . Therefore it is simpler to enclose the region in either a circle or a rectangle, generate a homogeneous Poisson process in the enlarged area, and subsequently exclude points outside of the given region.

# 6. <u>SIMULATION OF TWO-DIMENSIONAL</u> NONHOMOGENEOUS POISSON PROCESSES

The two-dimensional nonhomogeneous Poisson process  $\{N(x,y):x \ge 0, y \ge 0\}$  is specified by a positive rate function  $\lambda(x,y)$  which for simplicity is assumed here to be continuous. Then the process  $\{N(x,y)\}$  has the characteristic properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any such region R has a Poisson distribution with mean  $\Lambda(R)$ ; here  $\Lambda(R)$  denotes the integral of  $\lambda(x,y)$  over R, i.e., over the entire area of R.

The basic result of Section 4 on thinning of one-dimensional nonhomogeneous Poisson processes generalizes to two-dimensional nonhomogeneous Poisson processes. Thus, suppose that  $\lambda(\mathbf{x},\mathbf{y}) \leq \lambda^*(\mathbf{x},\mathbf{y})$  in a fixed rectangular region of the plane. If a nonhomogeneous Poisson process with rate function  $\lambda^*(\mathbf{x},\mathbf{y})$  is thinned according to  $\lambda(\mathbf{x},\mathbf{y})/\lambda^*(\mathbf{x},\mathbf{y})$  (i.e., each point  $(\mathbf{X}_i,\mathbf{Y}_i)$ ) is deleted independently if a uniform (0,1) random number  $U_i$  is greater than  $\lambda(\mathbf{X}_i,\mathbf{Y}_i)/\lambda^*(\mathbf{X}_i,\mathbf{Y}_i)$ , the result is nonhomogeneous Poisson process with rate function  $\lambda(\mathbf{x},\mathbf{y})$ .

The nonhomogeneous Poisson process with rate function  $\lambda(\mathbf{x},\mathbf{y})$  in an arbitrary but

fixed region R can be generated by enclosing the region R either in a circle [17], or a rectangle, and applying Algorithm 3. The following procedure assumes that the region R has been enclosed in a rectangle R<sup>\*</sup>, and that  $\lambda^* = \max\{\lambda(x,y):x, y \in R\}$  has been determined; here the bounding process is homogeneous with rate  $\lambda^*$  in the rectangle R<sup>\*</sup>.

Algorithm 4. Two-Dimensional Nonhomogeneous Poisson Process (Thinning)

- 1. Using Algorithm 2, generate points in the homogeneous Poisson process of rate  $\lambda^*$  in the rectangle R<sup>\*</sup>. If the number of points, n<sup>\*</sup>, is such that n<sup>\*</sup> = 0, exit; there are no points in the nonhomogeneous Poisson process.
- 2. From the n<sup>\*</sup> points generated in 1, delete the points that are not in R, and denote the remaining points by  $(x_1^*, y_1^*)$ ,  $(x_2^*, y_2^*)$ ,...,  $(x_m^*, y_m^*)$  with  $x_1^* < x_2^* < \cdots < x_m^*$ . Set i = 1 and k = 0.
- 3. Generate  $U_i$  uniformly distributed between 0 and 1. If  $U_i \leq \lambda(X_i^*, Y_i^*) / \lambda^*$ , set k = k+1,  $X_k = X_i^*$  and  $Y_k = Y_i^*$ .
- Set i equal to i+1. If i ≤ m<sup>\*</sup>, go to 3.
- 5. Return  $(X_1, Y_1)$ ,  $(X_2, Y_2)$ ,...,  $(X_n, Y_n)$ , where n = k and n.

# 7. CONCLUSION

We have summarized previously known general methods for simulating nonhomogeneous Poisson processes in one dimension. In addition, we have described the simple and efficient new methods of Lewis and Shedler for simulating nonhomogeneous Poisson processes in one and two dimensions. Extensions of the thinning algorithm to the simulation of homogeneous or nonhomogeneous conditional or doubly stochastic Poisson processes will be described elsewhere.

# REFERENCES

1. Ahrens, J. and Dieter, U. (1974a). Computer methods for sampling from gamma, beta, Poisson and binomial distributions. Computing, Vol. 12, pp. 223-46.

2. Ahrens, J. and Dieter, U. (1974b). Non Uniform Random Numbers, Graz, Austria: Technische Hoschule in Graz.

3. Çinlar, E. (1975). Introduction to Stochastic Processes. Englewood Cliffs, New Jersey: Prentice-Hall.

4. Cox, D. R. (1972). The statistical analysis of dependencies in point processes. In Stochastic Point Processes, ed.
P. A. W. Lewis, pp. 55-56, New York: Wiley.

5. Cox, D. R. and Lewis, P. A. W. (1966). The Statistical Analysis of Series of Events. London: Methuen.

6. Gilchrist, R. (1977). Some problems encountered in generating realisations from a useful nonhomogeneous Poisson process. In <u>Proceedings of the European</u> <u>Congress of Statisticians</u> 1976, Ed. J. R. Barra et al., Amsterdam: North Holland.

7. Goldman, J. R. (1967a). Stochastic point processes: limit theorems. <u>Ann.</u> <u>Math. Stat.</u>, Vol. 38, pp. 771-79.

 Goldman, J. R. (1967b). Infinitely divisible point processes in R<sup>n</sup>. J. Math. Anal. Appl. Vol. 17, pp. 133-46.

9, Holgate, F. (1972). The use of distance methods for the analysis of spatial distributions of points. In <u>Stochastic</u> Point Processes, Ed. P. A. W. Lewis, pp. 122-35. New York: Wiley.

10. Karlin, S. and Taylor, H.M. (1975). A First Course in Stochastic Processes. New York: Academic Press.

11. Knuth, D. E. (1969). The Art of Computer Programming, Vol. 2, Semi-Numerical Algorithms, Reading, Massachusetts: Addison-Wesley.

12. Lewis, P. A. W. (1972). Recent advances in the statistical analysis of univariate point processes. In <u>Stochastic</u> <u>Point Processes</u>, Ed. P. A. W. Lewis, pp. <u>1-54.</u> New York: Wiley.

13. Lewis, P. A. W. and Learmonth, G. (1973). Naval Postgraduate School random number generator package LLRANDOM. Naval Postgraduate School Report NPS55Lw73061A. Monterey, California.

14. Lewis, P. A. W. and Shedler, G. S. (1976a). Simulation of nonhomogeneous processes with log-linear rate function. Biometrika, Vol. 63, pp. 501-05.

15. Lewis, P. A. W. and Shedler, G. S. (1976b). Statistical analysis of a nonstationary series of events in a data base system. IBM J. Res. Devel., Vol. 20, pp. 465-82. 16. Lewis, P. A. W. and Shedler, G. S. (1977). Simulation of nonhomogeneous Poisson processes with degree-two exponential polynomial rate function. IBM Research Report RJ 1953. San Jose, California.

 Lewis, P. A. W. and Shedler, G. S. (1978). Simulation of nonhomogeneous Poisson processes by thinning. IBM Research Report RJ 2286. San Jose, California.

18. Miles, R. E. (1970). On the homogeneous planar Poisson point process. Mathematical Biosciences, Vol. 6, pp, 85-127.

19. Newell, G. F. (1968). Queues with time-dependent arrival rates (I, II, III). J. Appl. Prob., Vol. 5, pp. 436-451, 479-490, 591-606.

20. Patrow, M. L. (1977). A comparison of two algorithms for the simulation of nonhomogeneous Poisson processes with degree-two exponential polynomial intensity function. M.S. Thesis, Department of Operations Research. Naval Postgraduate School. Monterey, California (National Technical Information Service, Springfield, Virginia; AD-A047164.)

21. Rantschler, R. D. (1973). The detection and estimation of trends in two-dimensional Poisson processes. M.S. Thesis. Department of Operations Research. Naval Postgraduate School. Monterey, California. (National Technical Information Service, Springfield, Virginia; AD-A76136.)

22. Reyment, R. A. (1976). Geomagnetic reversal data statistically appraised. Tectonophysics, Vol. 31, pp. 73-91.

23. Steinijans, V. (1972). A stochastic point process model of the occurrence of major freezes in Lake Constance. <u>Appl.</u> <u>Statistics</u>, Vol. 25, pp. 58-61.

24. Thompson, H. R. (1955). Spatial point processes with applications to ecology. Biometrika, Vol. 42, pp. 102-15.

25. Vere-Jones, D. (1970). Stochastic models for earthquake occurrence. J. Roy. Statist. Soc. B, Vol. 32, pp. 1-62.

Acknowledgments. Subbort under Office of Naval Research Grant NRG42-343 and National Science Foundation Grant AF476 is gratefully acknowledged by P.A.W. Lewis

# DISTRIBUTION LIST

		NO.	OF	COPIES
STATISTICS AND FRCBABILITY CFFICE OF NAVAL RESEARCH CODE 436 ARLINGION	PROGRAM			
VA	22217		1	
CFFICE OF NAVAL RESEARCH NEW YERK AFEA CFFICE 715 BREACWAY - 5TH FLOOR ATTN: DR. RJEER GRAFTON NEW YORK, NY	10003		1	
DEFICE OF NAVAL RESEARCH B	RANCH OFF			
536 SCUTH CLARK STREET ATTN: DEPUTY AND CHIEF SC CHICAGO, IL	IENTIST 60605		1	
LIBRARY NAVAL OCEAN SYSTEMS CENTER				
CA	92152		1	
NAVY LIBPARY NATIONAL SPACE TECHNOLOGY	LAB			
ATTN: NAVY LIBRARIAN CAY ST. LCUIS MS	39522		1	
NAVAL ELECTRONIC SYSTEMS C	GM MA NO			
NATIENAL CENTER NO. 1 Arlington VA	20360		1	
CIPECTOR NAVAL REASARCH LA	EURATORY			
CODE 2029 WASHINGTON, C.C.	20375		1	
DEFENSE COCUMENTATION CENT CAPEFON STATION	ER		ź	
ALEXANDRIA VIFGINIA 22314				
TECHNICAL INFORMATION DIVI NAVAL RESEARCH LABORATORY	SION		1	
WASHINGTON, C. C.	20375			

CFFICE OF NAVAL RESEARCH SAN FRANCISCE AREA OFFICE	NO.	OF	COPIES
760 MARKET STREET SAN FRANCISCE CALIFORNIA 94102		1	
TECHNICAL LIBRARY NAVAL ORENANCE STATICN		1	
TRUTAR HEFE HEREEARD - 20040			
NAVAL SHIP ENGINEERING CENTER PHILADELFHIA DIVISION TECHNICAL LIBRARY FHILADELFHIA PENNSYIVANIA 19112		1	
		-	
BUREAU CF NAVAL PRESONNEL DEPARTMENT OF THE NAVY TECHNICAL LIERARY WASHINGTON C. C. 20370		1	
LIBRARY CODE 3212 NAVAL POSTGRACLATE SCHOOL		1	
		-	
DEFARTMENT OF MATHEMATICS LNIVERSITY CF NORTH CARCLINA CHARLOTTE			
NC 28223		1	
PROF. T. W. ANCERSON DEFARTMENT OF STATISTICS STANFORC UNIVERSITY		,	
STANFORD , CALIFCRNIA 94305		-	
FROF. F. J. AN SCOMBE CEFARTMENT OF STATISTICS YALE UNIVERSITY NEW FAVEN			
CCNNECTICUT 06520		1	
PROF. L. A. AFCIAN INSITILTE CF INCUSTRIAL ADMINISTRATION			
SCHENECTADY NEW YORK 12308		1	
PROF. C. R. BAKER DEFARTMENT OF STATISTICS UNIVERSITY (F. NCTRH CAROLINA CHAPEL FILL.			
NORTH CARCLINA 27514 11		1	

PROF. R. E. BECHOFER DEFARTMENT OF OPERATIONS RESEARCH CCRNELL UNIVERSITY ITHACA NEW YORK 1485C	1
FRCF. N. J. BERSHAD SCHOOL CF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE , CALIFORNIA S2664	1
P. J. BICKEL CEPARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA BEFKELEY, CALIFORNIA 94720	1
PROF. F. W. BLCCK DEFARTMENT OF MATHEMATICS UNIVERSITY CF FITTSBURGH PITTSBURGH PA 15260	1
FREF. JOSEPH BLUM DEFT. OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE THE AMERICAN UNIVERSITY WASHINGTON DC 20016	1
PFOF. R. A. BRADLEY DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE , FLORIDA 32306	1
PROF. R. E. BARLOW CFERATICNS RESEARCH CENTER CCLLEGE OF ENGINEERING LNIVERSITY OF CALIFORNIA BEFKLEY CALIFORNIA 54720	1
MR. C. N BENNETT NAVAL COASTAL SYSTEMS LABEFATORY CODE P761 PANAMA CITY, FLERIDA 32401	1
PROF. U. N. BHAT CCMPUTER SCIENCE / OPERATIONS RESEARCH CENTER SOUTHERN METHODIST UNIVERSITY CALLAS TEXAS 75275	1
PRCF. W. R. BLISCHKE DEPT. OF QUANTITATIVE BUSINESS ANALYSIS UNIVERSITY OF SOUTHERN CALIFOFNIA LOS ANGELES, CALIFORNIA S0007	1

DR. DERRILL J. BORDELCN NAVAL UNCERWATER SYSTEMS CENTER COCE 21 NEWPORT	1
C2 84 0	
J. E. BOYER JR DEPT. OF STATISTICS SCUTHERN METHODIST UNIVERSITY DALLAS TX 75275	1
DR. J. CHANERA U. S. ARMY RESEARCH P. G. BEX 12211 RESEARCH TRIANGLE PARK, NEFTH CAROLINA 27706	1
PROF. F. CHERNOFF DEPT. OF MATHEMATICS MASS INSTITUTE OF TECHNOLOGY CAMBRILGE, MASSACHUSETTS 02139	1
PROF. C. DERMAN DEFARTMENT OF CIVIL ENGINEERING AND ENGINEERING MECHANICS COLUMBIA UNIVERSITY NEW YORK NEW YORK 10027	1
FRCF. R. L. EISNEY VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEPT. CF INDUSTRIAL ENGINEERING AND OPERATIONS RESEARCH BLACKSBUFG, V/ 24061	1
WR. J. DOWLING DEFENSE LOGISTICS STUDIES INFORMATION EXCHANGE ARMY LOGISTICS MANAGEMENT CENTER FORT LEE VIRGINIA 20390	1
PRCF J. D. ESARY CEPT. CF OPERATIONS RESEARCH AND ADMINISTRATIVE SCIENCES NAVAL POSTGRADUATE SCHOOL MONTEREY. CALIFORNIA 53540	1
DR. M. J. FISCHER DEFENSE COMMUNICATIONS AGENCY 1860 WIEHLE AVENUE RESTON VIRGINIA 22070	1

PROF. C. P. GAVER DEFT. OF OPERATIONS RESEARCH NAVAL FESTEFADUATE SCHEEL MONTEREY CA	1
MR. GENE F. GLEISSNER APPLIED MATHEMATICS LABORATORY CAVID TAYLOF NAVAL SHIP RESEARCH AND DEVELOPMENT CENTER EETHESCA MD 20084	1
PROF. S. S. GUPTA DEPARTMENT OF STATISTICS FURDUE UNIVERSITY LAFAYETTE INDIANA 47907	1
PROF. D. L. HANSON DEFT OF MATH. SCIENCES STATE UNIVERSITY OF NEW YORK, EINGHAMTCN BINGHAMTCN NY 13901	1
PROF. F. J. HARRIS DEFT. OF ELECTRICAL ENGINEERING SAN CIEGC STATE UNIVERSITY SAN DIEGC CA 52182	1
FRCF. L. H. HERBACH DEFT. CF CFERATICNS RESEARCH AND SYSTEMS ANALYSIS POLYTECHNIC INSTITUTE CF NEW YORK BRCOKLYN NY 11201	1
FRCF. N. J. HINICH DEPARTMENT OF ECONOMICS VIRGINIA POLYTECHNIC INSTITUTE ANC STATE UNIVERSITY BLACKSEURG VIRGINIA 24C61	1
PROF. W. M. HIRSCH INSTITUTE OF MATHEMATICAL SCIENCES New York University New York 10453 New York 10453	1
PROF. C. L. IGLEHART DEFARTMENT OF OPERATIONS RESEARCH STANFERD UNIVERSITY STANFERD, CALIFORNIA 94350	1

PRCF. J. B. KACANE DEPARTMENT CF STATISTICS CARNEGIE-MELLCN FITTSBURGH FENNSYLVANIA 15213	1
DR. RICHARE LAU DIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET FASADENA CA 91101	1
DR . A. R. LAUFER DIRECTOR OFFICE OF NAVAL RESEARCH ERANCH OFF 1030 EAST GREEN STREET PASADENA CA S1101	1
FROF. M. LEACRETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NORTH CAROLINA 27514	1
CR. J. S. LEE J. S. LEE ASSCCIATES, INC. 2001 JEFFERSON DAVIS HIGHWAY SUITE EC2 ARLINGTON VA 22202	1
PROF. L. C. LEE DEPARTMENT OF STATISTICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSEURG VA 24061	1
PROF. R. S. LEVENWERTH DEPT. OF INCUSTRIAL AND SYSTEMS ENGINEERING UNIVERSITY OF FLORIDA GAINSVILLE . FLORIDA 32611	1
PRCF G. LIEPERMAN STANFORD INIVERSITY DEPARTMENT OF CPERATIONS RESEARCH STANFORD CALIFORNIA 94305	1

DR. JAMES F. MAAR NATIONAL SECURITY AGENCY FORT MEACE, MARYLAND 20755 1 PROF. R. W. MADSEN DEFARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA 1 NC 65201 DR. N. R. MANN SCIENCE CENTER RCCKWELL INTERNATIONAL CORPORATION F.C. BOX 1085 HCUSAND CAKS, CALIFORNIA 91360 1 DR. W. H. MARLOW PREGRAM IN LOGISTICS THE GECRGE WASHINGTON UNIVERSITY 707 22ND STREET, N. W. WASHINGTON, E. C. 20037 1 PROF. E. MASRY DEFT. APPLIEC PHYSICS AND INFCRMATICN SERVICE UNIVERSITY OF CALIFERNIA LA JOLLA CALIFORNIA 1 \$2093 DF. BRUCE J. MCCDNALD SCIENTIFIC CIRECTOR SCIENTIFIC LIAISCN GROUF CFFICE OF NAVAL RESEARCH AMERICAN EMBASSY - TOKYO AFC SAN FRANCISCO 1 96503 FRCF. J. A. MUCKSTADT CEPT. CF CFERATIONS RESEARCH CCRNELL UNIVERSITY ITHACA, NEW YORK 19850 1 DF. JANET M. MYFRE THE INSTITUTE OF DECISION SCIENCE FOR BLEINESS AND PUBLIC FOLICY CLAREMONT MEN'S COLLEGE CLAREMONT CA 91711 1 WR. H. NISSELSCN BLREAU OF THE CENSUS PCCM 2C25 FRCERAL FUILCING 3 WASHINGTUN . C. C. 2C33 1

1

1

1

1

1

1

1

1

MISS B. S. ORLEANS NAVAL SEA SYSTEMS COMMAND (SEA 03F) RM 10508 ARLINGTON VIRGINIA 20360 FRCF. C. B GWEN CEFARTMENT OF STATISTICS SCUTHERN METHCEIST UNIVERSITY DALLAS TEXAS 75222 DR. A. PETRASOVITS ROOM 2078 , FCCD AND DRUG ELDG. TUNNSY'S PASTURE OTTOWA , CNTARIC KIA-GL2 , CANADA PROF. S. L. PHCENIX SIELEY SCHOOL OF MECHANICAL AND AEROSPACE ENGINEERING CCFNELL UNIVERSITY ITHACA NY 14850 14850 DR. A. L. POWELL CIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 495 SUMMER STREET BOSTON 00010 C2210 MA MR. F. R. PRIORI CCCE 224 CPERATIONSL TEST AND ONRS EVALLATION FORCE (OPTEVFOR) NORFOLK , VIEGINIA 20360 PROF. M. L. FUFI DEFT. CF MATHEMATICS F.C. BOX F INCIANA UNIVERSITY FOUNCATION BLOOMING TON IN 47401 PROF. H RC6BINS DEFARTMENT OF MATHEMATICS COLUMBIA UNIVERSITY NEW YORK, NEW YORK, 10027

FROF. M ROSENBLATT DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA \$2093
FROF. S. M. ROSS COLLEGE CF ENGINEERING UNIVERSITY CF CALIFORNIA BEFKELEY CA 94720
PRCF. I RUBIN SCHOOL OF ENGINEERING AND APPLIEC SCIENCE UNIVERSITY OF CALIFCRNIA LCS ANGELES CALIFCRNIA 9C024
PROF. 1. R. SAVAGE DEFARTMENT OF STATISTICS YALE UNIVERSITY NEL HAVEN. CONNECTICUT (6520)
PROF. L. L., JR. SCHARF DEFARTMENT OF ELECRICAL ENGINEERING COLORACO STATE UNIVERSITY FT. COLLINS, COLORADO 80521
FRCF. R. SERFLING DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE FLORIDA 32306
PROF. W. R. SCHUCANY DEPARTMENT OF STATISTICS SCUTHERN METHODIST UNIVERSITY DALLAS, TEXAS 75222
FRCF. D. D. SIEGMUND DEFT. OF STATISTICS STANFORD UNIVERSITY STANFORD CA 94305
FRCF. M. L. SHOOMAN DEFT. CF FLECTRICAL ENGINEERING FOLYTECHNIC INSTITUTE OF NEW YORK BROOKLYN, NEW YORK 11201
PRCF. N. SINGPURWALLA DEPT. CF CFERATIONS RESEARCH THE GEORGE WASHINGTON UNIVERSITY 707 22ND ST. N. W. WASHINGTON, C. C. 18

DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANCANT OF THE FARINE CORPS WASHINGTON , C. C. 20380 1 MR. CHARLES S. SMITH CASD (IGL), FENT AGON MASHINGTON DC 1 20301 DR. D. E. SMITH DESMATICS INC. P.C. BCX 618 STATE COLLEGE PENNSYLVANIA J6801 1 FRCF. W. L. SMITH CEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL NOFTH CAROLINA 27514 1 FRCF. H SOLOMON DEFARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORC, CALIFERNIA 94305 1 MR. GLENN F. STAHLY NATIONAL SECURITY AGENCY FORT MEADE MARYLANC 20755 1 MR. EAVIE A. SWICK AEVANCED FREJECTS GROUP CODE 8103 MAVAL FESEARCH LAB. WASHINGTON DC 1 20375 WR. WENDELL G. SYKES ARTHUR D. LITTLE, INC. ACCRN PARK CAMBRIEGE MA 1 C2140 FROF. J. F. THOMPSON DEPARTMENT OF MATHEMATICAL SCIENCE RICE UNIVERSITY HCUSTON, TEXAS 77C01 1

NO. OF COPIES FRCF. W. A. THOMPSON DEPARTMENT CF STATISTICS UNIVERSITY OF MISSCURI COLUMBIA , MISSOLRI 65201 1 FREF. F. A. TILLMAN DEPT. OF INCUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN KS 1 66506 1 FREF J. W. TUKEY DEFAFTMENT OF STATISTICS PRINCETON UNIVERSITY PRINCETON , N. J. C854C FROF. A . F . VEINOTT CEFARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSTITY STANFORD CALIFORNIA \$4305 1 CANIEL H. WAGNER Station Square CNE Faqli, Pennsylvan IA 19301 1 PROF. GRACE WALBA DEPT. OF STATISTICS UNIVERSITY OF WISCONSIN MADISON 1 53706 . FRCF. K. T. WA DEPARTMENT OF MATHEMATICAL SCIENCES CLEMSON UNIVERSITY CLEMSON, SOLTH CAROLINA 29631 1 PROF. G. S. LATSON CEPARTMENT OF STATISTICS FRINCETON , N. J. 08540 1 FRCF. BERNARE WIEROW STANFERD ELECTRONICS LAB STANFORD UNIVERSITY STANFORD CA 1 \$43 05 FRCF. G. E. WHITEHOUSE DEPT. OF INDUSTRIAL ENGINEERING LEFIGH UNIVERSITY BETHLEHEM FA 1 18015 20

PROF. S. ZACKS CEFT. OF MATHEMATICS AND STATISTICS CASE WESTERN FESERVE UNIVERSITY CLEVELAND., CHIC 44106	1
PRCF. M. ZIA-HASSAN DEFARTMENT OF INDUSTRIAL AND SYSTEMS ENGINEERING ILLINCIS INSTITUTE OF TECHNOLOGY CHICAGO 60616	1
CR. DAVID BRILLINGER STATISTICS CEPT. UNIVERSITY OF CALIFORNIA BEFKELEY, CALIFORNIA 54720	1
CR. CHIN CHEN CCMMONWEAL TH ASSOCIATES, INC. 205 E. WASHINGTON JACKSON, MI 49201	1
FRCFESSOR PETER CHEN ALFRED P. SLCAN SHCOOL CF MGNT MASS INSTITUTE CF TCHNCLOGY 50 MEMOFIAL DRIVE CAMBRIEGE MASSACHUSETTS 02139	1
DR. E. CINLAR O.R. DEPT. MCRTHWESTERN UNIVERSITY EVANSTON, ILLINDIS 60201	1
DR. GUY C CORYNEN L. 156 UNIV OF CALIFORNIA LAWRENCE LIVERMORE LABORATORY F. D. EOX EOE LIVEFMCRE, CALIFORNIA 94550	1
DR. MICHAEL A. CRANE CONTROL AN/LYSIS CORPORATION EOC WELCH PEAC FALD ALTE , CALIFORNIA 94304	1
CR. D. F. CALEY STATISTICS DEPT. (I A S) AUSTRALIAN NATIONAL UNIVERSITY CANBERRA, A. C. T. 2606	1

HSU DER-ANN SCH OF ENGF./APPLIED SCI. CEFT. CF CIVIL & GEOLOGICAL ENGR. THE ENGINEERING QUACRANGLE FFINCEION NEL JERSEY C8540	1
ASSISTANT CEAN CAV ELDREDGE SCHOOL OF EUSINESS ADMINISTRATION UNIVERSITY OF EVANSVILLE EVANSVILLE INDIANA 47702	1
CR. G. W. FISHMAN CIFRICULUM IN O. R. UNIVERSITY OF NORTH CAFCLINA PHILLIPS ANNEX , CHAPPEL HILL NORTH CAROLINA 27514	1
CR. A. V. FOLCEN DEPARTMENT CF PHYSIOLOGY THE UNIVERSITY LEEDS LS2 SJT ENGLANC	1
DR. EUGENE C. FOMER MANAGEMENT SCI. & ENGINEERING DEPT. C.W. POST COLLEGE CF LONG ISLAND UNIVERSITY GREENVALE, NEW YORK 11548	1
DR. D. VERE JONES DEPT. OF MATHEMATICS VICTORIA UNIVERSITY OF WELLINGTON , P. D. BCX 196 WELLINGTON , NEW ZEALAND	1
J. M. LIITTSCHWAGER SYSTEMS CIVISICN CCLLEGE OF ENGINEERING THE UNIVERSITY OF IOWA 10WA CITY , ICWA 52242	1
Y. W. CE KHAADSTENIET DEFATRMENT OF EIULOGY FREE UNIVERSITY CE BOELELAAN 1087 AMSTERCAM-EUITENVELDER THE NETHERLANES	1
JACK P. LANFOLT DEFENCE & CIVIL INST. OF ENVIR. MED 1133 SFEPART AVE. WEST PO BUX 2000 DOWNSVIEW ONTARIO, CANADA	1

MF. FRANK MCNOLTY 1142 PENE AVE. SUNNYVALE, CA 94087	1
CR. P. A. F. CRAN STATISTICS CEPT. (I A S) AUSTRALIAN NATIONAL UNIVERSITY CANBERRF, A. C. T 2606 AUSTRALIA	1
G. W. MCRRISON DAK RIDGE NATIONAL LABORATORY CAK RICGE, TENNESSE 27830	1
MR. BRADLEY NICKEY LSCA - FCREST SERVICE P. O. BOX 5007 FIVERSICE CALIFCRNIA 92507	1
CR. DAVID CAKES DEFT. OF STATISTICS HARVARE UNIVERSITY SCIENCE CENTER, 1 OXFORD ST. COMBRIDGE, MA. O2132	1
Professor Emanuel Parzen, Director Institute of Statistics Teaxs A&M University College Station, TX 77843	1
MR ART PETERSON DEFT. OF EIGSTATISTICS SC-32 UNDVERSITY OF WASHINGTON SEATTLE, WASHINGTON 58195	1
CAVID * RCBINSON CONTROL ANALYSIS CCRPORATION 800 WELCH REAC PALO ALTO CALIFORNIA 94304	1
C. H. SAUER(21-129) THOMAS J. WATSCN RESEARCH CENTER F.C. BCX 218 YCRKTOWN HEIGHTS, NEW YCRK 10598	1

23

MR. C. A. STEELE JR F.C. BCX 45 MAGNGLIA, MA C1930 1 FRCF P. SLLLO SCHOOL OF MANAGEMENT FENSSELAER POLYTECHNIC INSTITUTE TROY, N.Y. 12181 1 DR. FOBEFT TRAPPL INSITIUT FUR ALLGEMEINE UND VERGLEICHENDE PHYSIOLOGIE DER UNIVERSITAT WIEN SCHWARZSFANIERSTRASSE 17 A-1090 WIEN OSTERREICH (ALSTRIA) 1 PALL A. HILLIS PCLY TECHNICA 2824 W. GECRGE MASON FALLS CHURF , VA. 22042 1 TUNCEL M. YEGULALP KRUMB SCHCCL CF MINES 918 S.H. MLDD CCLUMBIA UNIVERSITY 520 H. 12CTH ST. NEW YROK, N.Y. 10027 1 SHIEGKI YOKOYAMA RESEARCH CENTER OF APPLIED INFORMATION SCIENCES TOHOKU UNIVERSITY KATAHIRA-CFO 2-1, SENDAI 980 JAPAN 1 Dean of Research 1 Code 012 Naval Postgraduate School Monterey, Ca. 93940 Library (Code 55) 1 Naval Postgraduate School Monterey, CA 93940

50

Professor Peter Lewis Code 55Lw Naval Postgraduate School Monterey, Ca. 93949

Mr. G. Shedler K53-282 IBM Research Lab Cottle Road San Jose, Ca. 95193

R. J. Stampfel Code 55 Naval Postgraduate School Monterey, Ca. 93940 25