

AD-A068 903

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
STRONGLY STABLE STATIONARY SOLUTIONS IN NONLINEAR PROGRAMS. (U)
FEB 79 M KOJIMA

F/G 12/1

DAAG29-75-C-0024


UNCLASSIFIED

MRC-TSR-1920

NL

| OF |

AD
A068903



END
DATE
FILMED
7-79
DDC

LEVEL

2

MRC Technical Summary Report #1920

**STRONGLY STABLE STATIONARY SOLUTIONS
IN NONLINEAR PROGRAMS**

Masakazu Kojima

*See back page
for 1473*

AD A068903

**Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706**

February 1979

(Received December 19, 1978)

DDC
RECEIVED
MAY 23 1979
C

**Approved for public release
Distribution unlimited**

Sponsored by

U.S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D.C. 20550

DDC FILE COPY

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

STRONGLY STABLE STATIONARY SOLUTIONS IN NONLINEAR PROGRAMS

Masakazu Kojima

Technical Summary Report #1920
February 1979

ABSTRACT

For each continuously twice differentiable map $f = (f_0, f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^{1+m}$, we define a nonlinear program:

$$\begin{aligned} P1(f) \quad & \text{minimize } f_0(x) \\ & \text{subject to } f_i(x) = 0 \quad (1 \leq i \leq l), \\ & \quad \quad \quad f_j(x) \leq 0 \quad (l+1 \leq j \leq m). \end{aligned}$$

A (Kuhn-Tucker's) stationary solution x to $P1(f)$ is said to be strongly stable if there exists an open neighborhood U of x such that each open neighborhood $V \subset U$ of x contains a stationary solution to a perturbed problem $P1(f+g)$ which is unique in U whenever $g_i(x)$, $\partial g_i(x)/\partial x_j$ and $\partial^2 g_i(x)/\partial x_j \partial x_k$ ($0 \leq i \leq m$, $1 \leq j \leq n$, $1 \leq k \leq n$) are sufficiently small for all x in U . We will give conditions on the gradient vectors and the Hessian matrices of f_i ($0 \leq i \leq m$) which characterize the strong stability. These conditions are then applied to a parametric nonlinear program:

$$\begin{aligned} P2(t) \quad & \text{minimize } h_0(x,t) \\ & \text{subject to } h_i(x,t) = 0 \quad (1 \leq i \leq l), \\ & \quad \quad \quad h_j(x,t) \leq 0 \quad (l+1 \leq j \leq m), \end{aligned}$$

where t is a parameter vector varying in a closed subset T of \mathbb{R}^q . Let Σ^S be the set of points (x,t) in $\mathbb{R}^n \times T$ such that x is a strongly stable stationary solution to $P2(t)$. Under a certain constraint qualification and the continuity and the differentiability of the map $h : \mathbb{R}^n \times T \rightarrow \mathbb{R}^{1+m}$, we will establish that if S is a connected subset of Σ^S and if x^* is a local minimum solution to $P2(t^*)$ for some $(x^*, t^*) \in S$ then x is a local minimum solution to $P2(t)$ for all $(x,t) \in S$. Finally this result is applied to showing some interesting properties of a class of methods developed in the fixed point and complementarity theory.

AMS(MOS) Subject Classification - 90C30, 49B50

Key Words - Nonlinear Program, Stability, Parametric Program, Fixed Point and Complementarity Theory

Work Unit Number 5 - Mathematical Programming and Operations Research

Sponsored by the United States Army under Contract No. DLAG29-75-C-0024. This material is based upon work supported by the National Science Foundation under Grant No. MCS78-09525.

SIGNIFICANCE AND EXPLANATION

When we construct mathematical models from practical problems in the field of operations research, economics, engineering, etc., the data which we can utilize usually have uncertainty. We may not get exact data or the data may be changing as time goes. In such a model it is important to take account of the stability of the solution. Here we say that a solution to a model is stable if any slight perturbation to the data yields a small change of the solution.

In this paper we study stability of a mathematical programming model, which involves an objective function to be minimized (or maximized) under certain constraints. We give conditions on the data of the model which characterize the stability. Applications to a mathematical programming model having parameters and a class of computational methods are also discussed.

ACCESSION for

NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		<input type="checkbox"/>

BY

DISTRIBUTION/AVAILABILITY CODES

Dis. or SPECIAL

M

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

Masakazu Kojima

1. INTRODUCTION.

Let R^n be the n -dimensional Euclidean space. Throughout the paper we use the l_∞ norm $\|x\| = \max_i |x_i|$ for each $x = (x_1, \dots, x_n) \in R^n$. Let F be the class of maps $f = (f_0, f_1, \dots, f_m)$ from R^n into R^{1+m} such that each f_i is continuously twice differentiable, where m is a nonnegative integer. For each $f \in F$ we define the nonlinear program:

$$P(f) \quad \begin{aligned} & \text{minimize } f_0(x) \\ & \text{subject to } x \in X(f), \end{aligned}$$

where

$$X(f) = \{x \in R^n : f_i(x) = 0 \quad (1 \leq i \leq k), \\ f_j(x) \leq 0 \quad (k+1 \leq j \leq m)\}.$$

We call $X(f)$ the constraint set to the nonlinear program $P(f)$. For each positive number δ and $x \in R^n$, let

$$B_\delta(x) = \{x' \in R^n : \|x' - x\| \leq \delta\}.$$

An $x \in X(f)$ is said to be a local minimum solution to $P(f)$ if there is a positive number δ such that

$$f_0(x) \leq f_0(x') \quad \text{for all } x' \in X(f) \cap B_\delta(x).$$

It is well-known that under an assumption which is called a constraint qualification every local minimum solution to $P(f)$ satisfies the Kuhn-Tucker stationary condition (Kuhn and Tucker [20], see also Fiacco and McCormick [15], Mangasarian [23]). We will write the Kuhn-Tucker stationary condition in the form of a system of equations.

For each $\alpha \in R$, let

$$\alpha^+ = \max(0, \alpha) \quad \text{and} \quad \alpha^- = \min(0, \alpha).$$

For each continuously twice differentiable map $\psi : R^n \rightarrow R$ we use the notations $\psi(x)$

and $\nabla^2 \psi(x)$ for the gradient vector and the $n \times n$ Hessian matrix of ψ at $x \in R^n$. Then the Kuhn-Tucker stationary condition to $P(f)$ can be written as the system of equations

$$(1-1) \quad F(x, y) = 0,$$

where

$$(1-2) \quad F(x, y) = \begin{pmatrix} \nabla f_0(x) + \sum_{i=1}^k y_i \nabla f_i(x) + \sum_{j=k+1}^m y_j \nabla f_j(x) \\ -f_1(x) \\ \vdots \\ -f_k(x) \\ y_{k+1} - f_{k+1}(x) \\ \vdots \\ y_m - f_m(x) \end{pmatrix}$$

for all $(x, y) \in R^{n+m}$. If $x \in R^n$ satisfies (1-1) for some $y \in R^m$ then we call x a stationary solution to $P(f)$, y a Lagrange multiplier vector associated with x and (x, y) a stationary point to $P(f)$. The constraint qualification which we will impose on a stationary (or local minimum) solution x^* to $P(f)$ is:

Condition 1.1. The set $\{\nabla f_i(x^*) : f_i(x^*) = 0, 1 \leq i \leq k\}$ is linearly independent.

If a local minimum solution x^* to $P(f)$ satisfies Condition 1.1 then it is a stationary solution, i.e., there exist a Lagrange multiplier vector $y^* \in R^m$ such that (x^*, y^*) satisfies (1.1). See Fiacco and McCormick [15]. A stationary (or local minimum) solution x^* to $P(f)$ is said to be isolated if there is a positive number such that $B_\delta(x^*)$ contains no stationary (or local minimum) solution to $P(f)$ except x^* .

Let x^* be an isolated stationary solution to $P(f)$. We will study the existence and uniqueness of a stationary solution to a perturbed problem $P(f+g)$ in a small open neighborhood of x^* , where $g \in F$. For each $g \in F$ and a subset U of R^n , let

Sponsored by the United States Army under Contract No. DAMG29-75-C-0024. This material is based upon work supported by the National Science Foundation under Grant No. MCS78-09525.

(1-3)
$$F^* = (g \in F : g_0(x) = \frac{1}{2} x^T D x + c^T x, g_1(x) = d_1 (1 \leq i \leq l))$$
 for some $n \times n$ symmetric matrix D ,
 $c \in R^n$ and $d = (d_1, \dots, d_m) \in R^m$.

In Section 5, we will show that a s -stable (w.r.t. F) stationary solution of $P1(f)$ is a local minimum if and only if it satisfies the strong second-order sufficient condition (Robinson [27]) for an isolated local minimum solution.

In Section 6, we will be concerned with the parametric nonlinear program:

$$P2(t) (t \in T) \quad \text{minimize } h_0(x, t)$$
 subject to $x \in Y(t)$,

where

$$Y(t) = \{x \in R^n : h_1(x, t) = 0 \quad (1 \leq i \leq k)$$

$$h_j(x, t) \leq 0 \quad (k+1 \leq j \leq m)\},$$

t is a parameter vector varying in a closed subset T of R^q and
 $h = (h_0, h_1, \dots, h_m) : R^n \times T \rightarrow R^{1+m}$. We assume that $h(\cdot, t) \in F$ for each $t \in T$ and that $h_1(x, t), \partial h_1(x, t)/\partial x_j$ and $\partial^2 h_1(x, t)/\partial x_j \partial x_k$ ($0 \leq i \leq m, 1 \leq j \leq n, 1 \leq k \leq n$) are continuous with respect to $(x, t) \in R^n \times T$. Let I be the set $\{(x, t) \in R^n \times T : x$ is a stationary solution of $P2(t)\}$, and I^s the set $\{(x, t) \in I : x$ is a s -stable (w.r.t. F) stationary solution of $P2(t)\}$. Under the constraint qualification that the set $\{\nabla h_1(x, t) : h_1(x, t) = 0, 1 \leq i \leq m\}$ is linearly independent for all $(x, t) \in I^s$, where $\nabla h_1(x, t) = (\partial h_1(x, t)/\partial x_1, \dots, \partial h_1(x, t)/\partial x_n)^T$, we will prove that if S is a connected set in I^s and if x^* is a local minimum solution to $P2(t^*)$ for some $(x^*, t^*) \in S$ then x is a local minimum solution to $P2(t)$ for every $(x, t) \in S$.

In Section 7, we will focus our attention to the case where $T = [0, 1] \subset R$. In a class of computational methods (Eaves [9, 11], Kojima [17, 19], Lemke [21], Saigal [30], Todd [33]) developed in the fixed point and complementarity theory (see, for example, Allgower and Geory [6], Cottle and Dantzig [7], Eaves [10], Lemke [22], Saigal [30], Todd [32]), the structure of the one dimensional parameter family $P2(t)$ ($t \in T$) of nonlinear programs is utilized. To solve $P1(f)$, each method of the class artificially

norm (g, U)

$$= \sup_{0 \leq i \leq m} \sup_{x \in U} \max \{ |g_i(x)|, \| \nabla g_1(x) \|, \| \nabla^2 g_1(x) \| \},$$
 where for each $n \times n$ matrix

$$\|A\| = \max \{ \|Ax\| : x \in R^n, \|x\| = 1 \}.$$

Definition. Let x^* be a stationary solution to $P1(f)$. x^* is strongly stable with respect to a subclass F' of F (abbreviated by s -stable (w.r.t. F')) if for some $\delta^* > 0$ and each $\delta \in (0, \delta^*)$ there exists an $\alpha > 0$ such that whenever $g \in F'$ and $\text{norm}(g, B_{\delta^*}(x^*)) \leq \alpha$, $B_\delta(x^*)$ contains a stationary solution to $P1(f+g)$ which is unique in $B_{\delta^*}(x^*)$.

Obviously, if $F_2 \subset F_1 \subset F$ and if a stationary solution x^* to $P1(f)$ is s -stable (w.r.t. F_1) then it is s -stable (w.r.t. F_2). By the definition, we also see that if a stationary solution x^* to $P1(f)$ is not s -stable (w.r.t. a subclass F' of F) then for any positive numbers δ^*, δ, α with $\delta < \delta^*$ there exists a map $g \in F'$ such that $\text{norm}(g, B_{\delta^*}(x^*)) \leq \alpha$ and that $B_\delta(x^*)$ contains either no stationary solution to $P1(f+g)$ or more than one stationary solutions to $P1(f+g)$. In this paper, we derive necessary and sufficient conditions for the s -stability (w.r.t. F) and then apply them to showing some topological properties of the stationary solution set to a parametric nonlinear program.

In our discussions, the map $F : R^{n+m} \rightarrow R^{n+m}$ defined by (1-2) plays an essential role. Generally, the map F is not continuously differentiable on R^{n+m} but is piecewise continuously differentiable (abbreviated by PC^1) on R^{n+m} . Some definitions and properties on PC^1 maps will be given in Section 2.

In Sections 3 and 4, we will give conditions on the gradients vectors and the Hessian matrices of f_i ($0 \leq i \leq m$) which characterize the s -stability (w.r.t. F). We will also show that the s -stability w.r.t. the class F is equivalent to the s -stability with respect to its proper subclass

construct a map $h : \mathbb{R}^n \times T \rightarrow \mathbb{R}^{1+m}$ such that $h(x,1) = f(x)$ for all $x \in \mathbb{R}^n$, i.e., $P2(t) = P1(f)$ and that the program $P2(0)$ has a trivial stationary solution x^0 . Then starting from $(x^0, 0) \in I$, the method traces the connected component $S^0 \subset I$ containing $(x^0, 0)$. Under certain conditions, the method attains an approximation $(\hat{x}, 1)$ of $(x^*, 1) \in S^0$. By the definition, x^* is a stationary solution to $P1(f)$. However, it is not always true that x^* is a local minimum solution to $P1(f)$. Furthermore, it can happen that the value of the objective function f_0 at x^* is greater than the one at the initial solution x^0 even if $x \in X(f)$ for all $(x,t) \in S^0$. As a corollary to the results given in Section 6, we will see that if x^0 is a local minimum solution of $P2(0)$ and if x is a s -stable (w.r.t. F) stationary solution of $P2(t)$ for every $(x,t) \in S^0$ then S^0 is a one dimensional curve with the form $\{(x(t), t) : t \in [0,1]\}$ and $x(t)$ is a local minimum solution to $P2(t)$ for every $t \in [0,1]$. It should be noted that if x was not s -stable (w.r.t. F) for some $(x,t) \in S^0$, any open neighborhood of x might contain no stationary solution or more than one stationary solution to $P2(t + \epsilon)$ for every sufficiently small $\epsilon > 0$. If, in addition, we employ a map $h : \mathbb{R}^n \times T \rightarrow \mathbb{R}^{1+m}$ such that

$$h_0(x,t) = (1-t)f_0(x) + tf_0(x)$$

$$h_1(x,t) = f_1(x) \quad (1 \leq i \leq m)$$

for all $(x,t) \in \mathbb{R}^n \times T$, then we can prove that $f_0(x(t))$ is monotone nonincreasing with respect to $t \in [0,1]$.

Many studies have been made on the stability or the sensitivity of (local) minimum solutions to parametric programs (Berge [5], Dantzig, Folkman and Shapiro [8], Evans and Gould [13], Fiacco [14], Fiacco and McCormick [15], Robinson [25, 26, 27], etc.). They mainly discussed the continuity of the minimum value of the objective function, the continuity of the set of minimum solutions and/or the continuity of an isolated local minimum (or stationary) solution with respect to a small change of the parameter vector. In Robinson [25], a quantitative bound on a variation of an isolated local minimum solution caused by a small change of the parameter vector was given. Under an

additional assumption on the differentiability with respect to the parameter vector, Fiacco [14] provided a method for estimating the variation.

Let $(x,y) \in \mathbb{R}^{n+m}$ be a stationary point to $P2(t)$. If $(i : y_i > 0, 1 \leq i \leq m) = (i : h_i(x,t) = 0, 1 \leq i \leq m)$, we say that (x,y) satisfies strict complementarity. In the papers [14,25] above, the strict complementarity was assumed at a stationary point whose variation would be investigated so that they could apply the standard implicit function theorem to a system of equations associated with the Kuhn-Tucker stationary condition. If we replace $\nabla f_1(x)$ by $\nabla h_1(x,t) = (\partial h_1(x,t)/\partial x_1, \dots, \partial h_1(x,t)/\partial x_n)^T$ ($0 \leq i \leq m$) in (1-2), we will have the map $H(x,y,t) : \mathbb{R}^{n+m+1} \rightarrow \mathbb{R}^{n+m}$ and the system of equations $H(x,y,t) = 0$ which is equivalent to the Kuhn-Tucker stationary condition to $P2(t)$ (see (6-2)). Note that if the strict complementarity holds at a solution (x,y) to $H(x,y,t) = 0$ then the map $H(\cdot, \cdot, t) : \mathbb{R}^{n+m} \rightarrow \mathbb{R}^{n+m}$ is continuously differentiable in some open neighborhood of (x,y) . In the case where the strict complementarity does not hold, however, the approach based on the standard implicit function theorem for C^1 maps can not be used. Throughout this paper we do not assume the strict complementarity. Without the strict complementarity assumption, Robinson [27] recently gave a sufficient condition for the continuity of the variation of an isolated stationary solution to $P2(t)$ with respect to the parameter vector t . His approach is based on the study of generalised equations which include the Kuhn-Tucker stationary condition as a special case (Robinson 26)). On the other hand, our main tools are the degree theory of the continuous maps (see, for example, Ortega and Rheinboldt [24]) and some fundamental results on PC^1 maps (Sotomayor [16], Whitehead [34]).

2. PRELIMINARIES.

Let P be a subset of \mathbb{R}^n and K a collection of a finite number of closed convex polyhedral sets with nonempty interior. K is said to be a subdivision of \mathbb{R}^n if

(2-1) the union of all σ in K is P

(2-2) $\sigma_1 \cap \sigma_2$ is a common face of σ_1 and σ_2 for every pair σ_1 and σ_2

σ_2 in K with $\sigma_1 \cap \sigma_2 \neq \emptyset$.

We call each $\sigma \in K$ a piece of K . When K is a subdivision of P , we will write $P = |K|$. A PL (piecewise linear) map $\psi : |K| \rightarrow \mathbb{R}^m$ is a continuous map such that

the restriction $\psi|_\sigma$ of the map ψ to each piece $\sigma \in K$ is affine, i.e., there exist a $P \times P$ matrix $A(\sigma)$ and an $a(\sigma) \in \mathbb{R}^m$ such that

$$\psi(z) = A(\sigma)z + a(\sigma) \text{ for every } z \in \sigma.$$

A PC^1 (piecewise continuously differentiable) map $\psi : |K| \rightarrow \mathbb{R}^m$ is a continuous map such that for each piece σ of K there exist an open set $U \supset \sigma$ and a continuously differentiable map $\psi' : U \rightarrow \mathbb{R}^m$ such that $\psi(z) = \psi'(z)$ for all $z \in \sigma$. Obviously, a PL map $\psi : |K| \rightarrow \mathbb{R}^m$ is PC^1 . We will use the symbol $D\psi(z; \sigma)$ for the Jacobian matrix of the restriction $\psi|_\sigma$ of a PC^1 map $\psi : |K| \rightarrow \mathbb{R}^m$ to a piece σ of K at $z \in \sigma$.

Now we shall show that the map $F : \mathbb{R}^{n+m} \rightarrow \mathbb{R}^{n+m}$ defined by (1-2) is PC^1 . Let J^0 denote the index set $\{i+1, \dots, m\}$. For each $J \subset J^0$, define

$$(2-3) \quad \tau(J) = \{(x, y) \in \mathbb{R}^{n+m} : y_j \geq 0 (j \in J), y_j \leq 0 (j \in J^0 \setminus J)\}$$

where $J^0 \setminus J = \{j \in J^0 : j \notin J\}$. We include the two cases $J = \emptyset$ and $J = J^0$.

$$\tau(\emptyset) = \{(x, y) \in \mathbb{R}^{n+m} : y_j \leq 0 (j \in J^0)\}$$

$$\tau(J^0) = \{(x, y) \in \mathbb{R}^{n+m} : y_j \geq 0 (j \in J^0)\}.$$

Let

$$(2-4) \quad F^0 = \{\tau(J) : J \subset J^0\}.$$

Then F^0 is a subdivision of \mathbb{R}^{n+m} , i.e., $\mathbb{R}^{n+m} = |F^0|$. It is easily verified that F is PC^1 on the subdivision F^0 .

Let $\psi^k : |K| \rightarrow \mathbb{R}^p$ be PC^1 ($k=1,2$), where $P = |K| \subset \mathbb{R}^p$, and ϵ, ρ be positive numbers. If

$$\|\psi^2(z) - \psi^1(z)\| \leq \epsilon \text{ for all } z \in |K|$$

and

$$\|(D^2(z; \sigma) - D\psi^1(z; \sigma))u\| \leq \rho \|D\psi^1(z; \sigma)u\|$$

for all $z \in \sigma$, $\sigma \in K$ and $u \in \mathbb{R}^p$ with $\|u\| = 1$,

we say that ψ^2 is an (ϵ, ρ) -approximation of ψ^1 . See Kojima [18] or Whitehead [34] for more general definition.

Let $\psi : |K| \rightarrow \mathbb{R}^p$ be PC^1 , where $P = |K| \subset \mathbb{R}^p$. For each $z^0 \in |K|$, let

$$(2-5) \quad M(z^0, K) = \{\sigma \in K : z^0 \in \sigma\}.$$

It is easily verified that

$$D\psi(z^0; \tau)(z - z^0) = D\psi(z^0; \sigma)(z - z^0)$$

if $\tau, \sigma \in M(z^0, K)$ and $z \in \tau \cap \sigma$.

Hence we can consistently define a PL map $\psi^{z^0} : |M(z^0, K)| \rightarrow \mathbb{R}^p$ by

$$\psi^{z^0}(z) = \psi(z^0) + D\psi(z^0; \sigma)(z - z^0) \text{ for every } z \in \sigma \text{ and } \sigma \in M(z^0, K).$$

The map $\psi^{z^0} : |M(z^0, K)| \rightarrow \mathbb{R}^p$ can be regarded as a PL approximation of $\psi : |K| \rightarrow \mathbb{R}^p$ at the point $z^0 \in |K|$ with the use of the Jacobian matrix. A PC^1 map $\psi : |K| \rightarrow \mathbb{R}^p$ is said to be locally nonsingular at $z^0 \in |K|$ if the PL map $\psi^{z^0} : |M(z^0, K)| \rightarrow \mathbb{R}^p$ is one-to-one. A nonsingular PC^1 map $\psi : |K| \rightarrow \mathbb{R}^p$ is a PC^1 map which is locally nonsingular at every $z^0 \in |K|$.

Suppose (x^0, y^0) is a stationary point to $F(f)$ (i.e., a solution to (1-1)) and that Condition 1.1 holds. In Section 3, we will show that $F : |F^0| \rightarrow \mathbb{R}^{n+m}$ is locally nonsingular at $z^0 = (x^0, y^0)$ if and only if

$$\det DF(z^0; \sigma) > 0 \text{ for all } \sigma \in M(z^0, F^0)$$

or

$$\det DF(z^0; \sigma) < 0 \text{ for all } \sigma \in M(z^0, F^0).$$

In Section 4, we will establish that x^0 is σ -stable (w.r.t. F) if and only if $z^0 = (x^0, y^0)$ is a locally nonsingular point of the map $F : |F^0| \rightarrow \mathbb{R}^{n+m}$.

We now state some fundamental lemmas on PC^1 maps which we will use later.

Throughout the lemmas below, K denotes a subdivision of a subset P of \mathbb{R}^p .

Lemma 2.1. Assume that $P = |K|$ is compact and that $\psi : |K| \rightarrow \mathbb{R}^p$ is a one-to-one and nonsingular PC^1 map. Then there exist positive numbers ϵ, ρ such that every

(ϵ, ρ) -approximation to $\psi : |K| \rightarrow \mathbb{R}^p$ is one-to-one and nonsingular.

Proof. See Theorem 3 of Whitehead [34].

Lemma 2.2. Let $\psi : |K| \rightarrow \mathbb{R}^p$ be PC^1 . Assume that

(2-6) a point z^* lies in the interior of P ,

(2-7) $\psi : |K| \rightarrow \mathbb{R}^p$ is locally nonsingular at z^* .

Then there is a positive number δ such that

(2-8) $\psi : |K| \rightarrow \mathbb{R}^p$ is locally nonsingular at every $z \in B_\delta(z^*) \cap P$,

(2-9) ψ is one-to-one in $B_\delta(z^*) \cap P$.

Proof. See Lemmas 2-10 and 2-11 of Kojima [18].

For each subset U of \mathbb{R}^p , we use the notations $\text{int } U$, $\text{cl } U$ and $\text{bd } U$ for the interior of U , the closure of U and the boundary of U , respectively. Let U be an open subset of \mathbb{R}^p and $\psi : \text{cl } U \rightarrow \mathbb{R}^p$ be continuous. Suppose that $\psi(z) \neq c$ for all $z \in \text{bd } U$. We define $\text{deg}(\psi, U, c)$ to be the degree of the map ψ at c with respect to U (see, for example, Ortega and Rheinboldt [24]).

Lemma 2.3. Let $\psi : |K| \rightarrow \mathbb{R}^p$ be PC^1 and z^* an interior point of $P = |K|$. Assume that $B_\delta(z^*) \subset |K|$ for some positive δ^* and that ψ is one-to-one in $B_{\delta^*}(z^*)$.

Then either

(2-10) $\text{deg}(\psi, \text{int } B_\delta(z^*), \psi(z^*)) = +1$ for every $\delta \in (0, \delta^*)$

or

(2-11) $\text{deg}(\psi, \text{int } B_\delta(z^*), \psi(z^*)) = -1$ for every $\delta \in (0, \delta^*)$.

If (2-10) (or (2-11)) holds then

$$\det D\psi(z) \geq 0 \quad (\text{or } \leq 0)$$

for all $z \in \sigma \cap \text{int } B_\delta(z^*)$ and $\sigma \in K$.

Proof. Since $B_{\delta^*}(z^*)$ is a compact subset of \mathbb{R}^p , ψ maps $B_{\delta^*}(z^*)$ onto $\psi(B_{\delta^*}(z^*))$ homeomorphically. Hence either (2-10) or (2-11) holds (see, for example, Chapter XVI,

14 of Alexandroff [3]). On the other hand, by the invariance theorem of domain (Lemma 3.9 of Eilenberg and Steenrod [1]), we see that $\psi(\text{int } B_\delta(z^*))$ is an open subset of \mathbb{R}^p for each $\delta \in (0, \delta^*)$. We assume that (2-10) holds, and show that

$$\det D\psi(z) \geq 0 \quad \text{for all } z \in \sigma \cap \text{int } B_\delta(z^*) \text{ and } \sigma \in K.$$

Let $z \in \sigma \cap \text{int } B_\delta(z^*)$ and $\det D\psi(z) < 0$. Then there is a $z' \in \text{int } \sigma \cap \text{int } B_\delta(z^*)$ such that $\text{sign } \det D\psi(z; \sigma) = \text{sign } \det D\psi(z'; \sigma) \neq 0$. Thus it suffices to show that

$\det D\psi(z'; \sigma) > 0$. Note that ψ is continuously differentiable in some open neighborhood

of z' . Since $\text{int } B_\delta(z^*)$ is a convex subset of \mathbb{R}^p and ψ is one-to-one in $B_\delta(z^*)$,

the image of the line segment $\{(1-t)z^* + tz' : 0 \leq t \leq 1\} \subset \text{int } B_\delta(z^*)$ under the map

ψ does not intersect with $\psi(\text{bd } B_\delta(z^*))$. By the homotopy invariance theorem (6.2.2

of Ortega and Rheinboldt [24]), we obtain

$$\begin{aligned} 1 &= \text{deg}(\psi, \text{int } B_\delta(z^*), \psi(z^*)) \\ &= \text{deg}(\psi, \text{int } B_\delta(z^*), \psi(z')) \\ &= \text{sign } \det D\psi(z'; \sigma). \end{aligned}$$

Q.E.D.

3. LOCAL NONSINGULARITY OF THE MAP $F : \mathbb{R}^k \rightarrow \mathbb{R}^{n+m}$.

In this section we give a necessary and sufficient condition for the \mathbb{R}^k map $F : \mathbb{R}^k \rightarrow \mathbb{R}^{n+m}$ defined by (1-2) to be locally nonsingular at a $z^0 \in |\mathbb{R}^k|$.

If w^1, \dots, w^k are p -dimensional vectors, we denote by (w^1, \dots, w^k) the closed convex polyhedral cone spanned by them, i.e.,

$$(w^1, \dots, w^k) = \left\{ \sum_{i=1}^k \lambda_i w^i : \lambda_i \geq 0 \ (1 \leq i \leq k) \right\}.$$

Lemma 3.1 (Samelson, Thrall and Waslar [3]).

Let $v^1, \dots, v^p, w^1, \dots, w^p$ be p -dimensional vectors. Define

$$\Pi = \{(w^1, \dots, w^p) : w^i = v^i \text{ or } v^i \ (i = 1, \dots, p)\}.$$

Suppose that $\det(v^1, \dots, v^p) > 0$. Then Π is a subdivision of \mathbb{R}^p if and only if for every $(w^1, \dots, w^p) \in \Pi$

$$\text{sign } \det(w^1, \dots, w^p) = (-1)^s,$$

where s is the number of v^i 's among w^1, \dots, w^p .

Corollary 3.2. Let Π be the collection of all the orthants of \mathbb{R}^p , i.e.,

$$\Pi = \{(w^1, \dots, w^p) : w^i = e^i \text{ or } -e^i \ (1 \leq i \leq p)\},$$

where e^i denotes the i -th unit vector in \mathbb{R}^p . Let $\psi : |\Pi| \rightarrow \mathbb{R}^p$ be PL. Then ψ is one-to-one if and only if either

$$\det D\psi(0; \sigma) > 0 \text{ for all } \sigma \in \Pi$$

or

$$\det D\psi(0; \sigma) < 0 \text{ for all } \sigma \in \Pi.$$

Proof. We first show the "only if" part. Suppose that ψ is one-to-one on $|\Pi| = \mathbb{R}^p$.

Since $\psi : |\Pi| \rightarrow \mathbb{R}^p$ is piecewise linear, we have

$$\psi(x) = D\psi(0; \sigma)x + \psi(0) \text{ for every } x \in \sigma \in \Pi.$$

Hence

$$\det D\psi(0; \sigma) \neq 0 \text{ for all } \sigma \in \Pi,$$

and the desired result follows from Lemma 2.3.

We now show the "if" part. Performing an appropriate linear transformation to the map ψ if necessary, we may assume that $\psi(0) = 0$,

$$\det D\psi(0; \sigma) > 0 \text{ for all } \sigma \in \Pi$$

and that

$$D\psi(0; \sigma) = [e^1, \dots, e^p],$$

where $\sigma = (-e^1, \dots, -e^p)$. Then we have

$$\psi(z) = D\psi(0; \sigma)z \text{ for every } z \in \sigma \in \Pi.$$

Let $\sigma^+ = (e^1, \dots, e^p)$ and $D\psi(0; \sigma^+) = (u^1, \dots, u^p)$. Then, for every $\sigma \in \Pi$, we have

$$\psi(e^1) = D\psi(0; \sigma)e^1 = D\psi(0; \sigma^+)e^1 = u^1 \text{ if } e^1 \in \sigma$$

and

$$\psi(-e^1) = D\psi(0; \sigma)(-e^1) = D\psi(0; \sigma^+)(-e^1) = -e^1 \text{ if } -e^1 \in \sigma.$$

Hence, for every $\sigma \in \Pi$,

$$D\psi(0; \sigma)e^1 = \begin{cases} u^1 & \text{if } e^1 \in \sigma \\ -e^1 & \text{if } -e^1 \in \sigma. \end{cases}$$

Recall that $\det D\psi(0; \sigma) > 0$ for all $\sigma \in \Pi$. By Lemma 3.1, the collection

$$\{(w^1, \dots, w^p) : w^i = u^1 \text{ or } -e^1 \ (1 \leq i \leq p)\}$$

subdivides \mathbb{R}^p . This is equivalent to that $\psi : |\Pi| \rightarrow \mathbb{R}^p$ is one-to-one. Q.E.D.

Now we are ready to derive a necessary and sufficient condition for the local

nonsingularity of the map $F : \mathbb{R}^k \rightarrow \mathbb{R}^{n+m}$ at a $z^0 \in |\mathbb{R}^k|$. Let

$$z^0 = (x^0, y^0) \in |\mathbb{R}^k| = \mathbb{R}^{n+m}.$$

Let

$$J_0 = \{j : y_j^0 > 0, 1 + 1 \leq j \leq m\},$$

and

$$J_1 = \{j : y_j^0 \geq 0, 1 + 1 \leq j \leq m\}.$$

Obviously, we see $J_0 \subset J_1$ and

$$\Pi(z^0, \mathbb{R}^0) = \tau(J) : J_0 \subset J \subset J_1,$$

where $\Pi(z^0, \mathbb{R}^0)$ denotes the collection of pieces $\tau(J) \in \mathbb{R}^0$ which contain z^0 and $\tau(J)$ is defined by (2-3). By the definition of the local nonsingularity,

Obviously Q^* is symmetric. By a simple calculation we see that Q^* satisfies (3-4) for every $\gamma > 0$. The existence of an $n \times n$ symmetric matrix Q^* satisfying (3-4) can be shown similarly if we replace (3-5) by

$$y_1 \dots y_k y_{k+1} \dots y_p < 0 \text{ and } y_j = 0 \text{ (} k+1 \leq j \leq p \text{)} .$$

Q.E.D.

We introduce some notations. For each $x = (x, y) \in R^n$, let

$$(3-6) \quad L(x) = \sum_{i=1}^k y_i^2 f_i(x) + \sum_{i=k+1}^p y_i^2 f_i(x) .$$

We can express the Jacobian matrix $DF(x; \tau(J))$ for each $\tau(J) \in N(x, R^p)$ in terms of $L(x)$ and $\nabla f_i(x)$ ($1 \leq i \leq m$). For simplicity, we assume $J = (k+1, \dots, k)$ for some $k \leq m$. Then

$$(3-7) \quad DF(x; \tau(J)) = \begin{array}{c|ccc|ccc} L(x) & \nabla f_1(x) & \dots & \nabla f_k(x) & 0 & \dots & 0 \\ \hline -\nabla f_1(x)^T & \dots & \dots & \dots & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ -\nabla f_k(x)^T & \dots & \dots & \dots & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ -\nabla f_m(x)^T & \dots & \dots & \dots & 0 & \dots & 0 \end{array} \begin{array}{c} n \\ k \\ m-k \\ m-k \\ m-k \\ m-k \\ m-k \end{array}$$

where E denotes the $(m-k) \times (m-k)$ identity matrix. Hence

$$(3-8) \quad \det DF(x; \tau(J)) = \det \begin{bmatrix} L(x) & A \\ -A^T & 0 \end{bmatrix} ,$$

where

$$(3-9) \quad A = [\nabla f_1(x), \dots, \nabla f_k(x)] .$$

Let D be an $n \times n$ symmetric matrix and W a subspace of R^n with $\dim W = p \geq 1$. We say that the matrix D has a positive (negative or zero) determinant if $\det B^T D B > 0$ (< 0 or $= 0$), where B is an $n \times p$ matrix whose columns form a basis of W . We also say that the matrix D is positive definite (or positive semi-definite) on the subspace W if the $p \times p$ matrix $B^T D B$ is positive definite (or positive semi-definite). For convenience, we assume that each $n \times n$ matrix has a positive determinant and is positive definite on the zero dimensional subspace $W = \{0\}$.

Theorem 3.5. Let $z^0 = (z^0, y^0) \in \tau(J)$ for some $J \subset \{1, \dots, m\}$. If the set (3-10)

$$(3-10) \quad \{\nabla f_i(z^0) : i \in \{1, \dots, k\} \cup J\}$$

is linearly dependent then $\det DF(z^0; \tau(J)) = 0$. In the case where the set (3-10) is linearly independent,

$$\det DF(z^0; \tau(J)) > 0 \text{ (} = 0 \text{ or } < 0 \text{)}$$

if and only if the $n \times n$ matrix $L(z^0)$ has a positive (zero or negative) determinant on the space $W = \{w \in R^n : \nabla f_i(z^0)^T w = 0 \text{ (} i \in \{1, 2, \dots, k\} \cup J \text{)}\}$.

Proof. For simplicity, we assume that $J = \{k+1, \dots, m\}$. Then (3-8) holds where A is the $n \times k$ matrix defined by (3-9). The first assertion of the theorem follows immediately. Suppose that A has rank k . If $k = n$ then the dimension of the subspace W is zero and $\det DF(z^0; \tau(J)) = (\det A)^2 > 0$; hence we obtain the desired result. Assume that $k < n$. Choose an $n \times (n-k)$ matrix B such that $B^T A$ is the $(n-k) \times k$ zero matrix and that $B^T B$ is the $k \times k$ identity matrix. Then the set of columns of B forms a basis of the subspace W . Let $D = L(z^0)$ and

$$C = [A \quad B] .$$

We first deal with the case $\det D \neq 0$. It follows from the identity

$$\begin{bmatrix} D^{-1} & 0 \\ A^T D^{-1} & B^T \end{bmatrix} \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} = \begin{bmatrix} E & D^{-1} A \\ 0 & A^T D^{-1} A \end{bmatrix}$$

that

$$(3-11) \quad \det D^{-1} \det \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} = \det A^T D^{-1} A ,$$

where E denotes the identity matrix with an appropriate size. We also see

$$C^T D^{-1} C = \begin{bmatrix} A^T D^{-1} A & 0 \\ 0 & B^T D B \end{bmatrix} .$$

Hence

$$(\det C)^2 (\det D^{-1}) = (\det A^T D^{-1} A) (\det B^T D B) .$$

Substituting (3-11) into the above equality, we have

$$(3-12) \quad (\det C)^2 = \det \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} (\det B^T D B) .$$

when $\det D = 0$ we have

$$(3-12)' \quad (\det C(\epsilon))^2 = \det \begin{bmatrix} D(\epsilon) & A \\ -A^T & 0 \end{bmatrix} (\det B^T D(\epsilon) B)$$

for every sufficiently small $\epsilon > 0$, where

$$D(\epsilon) = D + \epsilon Z, \quad C(\epsilon) = (A D(\epsilon) B)$$

Note that $\det D(\epsilon) \neq 0$ for every sufficiently small $\epsilon > 0$. Taking the limit as $\epsilon \rightarrow 0$ in (3-12)', we obtain (3-12). Therefore (3-12) holds even if $\det D = 0$.

On the other hand, we see

$$\begin{bmatrix} A^T \\ B^T \end{bmatrix} [A \quad DB] = \begin{bmatrix} A^T A & A^T DB \\ 0 & B^T DB \end{bmatrix},$$

which implies

$$(\det [A \quad B]) (\det C) = (\det A^T A) (\det B^T DB).$$

Since $\det [A \quad B] \neq 0$ and $(\det A^T A) \neq 0$, we have that $\det C = 0$ if and only if $\det B^T DB = 0$. Therefore, by (3-12), we obtain

$$(3-13) \quad \det \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} > 0 \text{ (or } < 0) \text{ if } \det B^T DB > 0 \text{ (or } < 0).$$

What we have left to show is

$$\det \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} = 0 \text{ if } \det B^T DB = 0.$$

Assume $\det B^T DB = 0$. By Lemma 3.4, there exist $n \times n$ symmetric matrices Q^+ and Q^- satisfying (3-4) and (3-5) for every $\gamma > 0$. Replacing D by $D + \gamma Q^+$ (or $D + \gamma Q^-$) in (3-13) and taking the limit as $\gamma \rightarrow 0$, we obtain

$$\det \begin{bmatrix} D & A \\ -A^T & 0 \end{bmatrix} \geq 0 \text{ (or } \leq 0). \quad \text{Q.E.D.}$$

As a direct consequence of Theorems 3.3 and 3.5, we obtain the following result:

Corollary 3.6. Let $z^* = (x^*, y^*)$, J_0 and J_1 be as in Theorem 3.3. Then

$F : |K^*| \rightarrow R^{n+m}$ is locally nonsingular at z^* if and only if

(3-14) the set $\{\nabla_j^T(x^*) : j \in \{1, \dots, l\} \cup J_1\}$ is linearly independent

and

(3-15) for all J such that $J_0 \subset J \subset J_1$ the $n \times n$ matrix $L(x^*)$ has a positive (or negative) determinant on the subspace $\{w \in R^n : \nabla_j^T(x^*)^T w = 0 \text{ (} j \in \{1, 2, \dots, l\} \cup J)\}$.

4. MAIN RESULTS.

For each perturbation $g \in F$ to the problem $PI(f)$, we define the PC^1 map

$G : |K^*| \rightarrow R^{n+m}$ as follows:

$$(4-1) \quad G(x, y) = \begin{pmatrix} v_{g_0}(x) + \sum_{i=0}^k y_i v_{g_1}(x) + \sum_{j=1}^m y_j^* v_{g_2}(x) \\ -g_1(x) \\ \vdots \\ -g_k(x) \\ -g_{k+1}(x) \\ \vdots \\ -g_m(x) \end{pmatrix}$$

for every $(x, y) \in R^{n+m}$. Then the Kuhn-Tucker stationary condition to the perturbed

problem $PI(f+g)$ can be written as

$$(4-2) \quad F(x, y) + G(x, y) = 0$$

Theorem 4.1. Let $z^* = (x^*, y^*)$ be a stationary point to $PI(f)$. Assume that F is locally nonsingular at z^* . Then x^* is a s -stable (w.r.t. F) stationary solution to $PI(f)$.

Proof. Let $I^* = \{i : f_i(x^*) = 0, 1 \leq i \leq m\}$. By Corollary 3.6, the set $\{f_i(x^*) : i \in I^*\}$ is linearly independent i.e., Condition 1.1 holds. For simplicity, we assume that $I^* = \{1, 2, \dots, k\}$. By the continuity of f_i ($1 \leq i \leq m$) at x^* , we can take positive numbers δ_0 and α_0 such that if $x \in B_{\delta_0}(x^*)$,

$$|g_1(x)| \leq \alpha_0, \quad \|v_{g_1}(x)\| \leq \alpha_0 \quad (1 \leq i \leq m)$$

then

$$(4-3) \quad \{i : f_i(x) + g_1(x) = 0, 1 \leq i \leq m\} \subset \{1, \dots, k\}.$$

On the other hand, by Lemma 2.2, there is a positive number $\delta_1 \leq \delta_0$ such that F is locally nonsingular at every $z \in B_{\delta_1}(z^*)$ and that F is one-to-one in $B_{\delta_1}(z^*)$. If $x \in B_{\delta_1}(x^*)$, $|g_1(x)| \leq \alpha_0, \|v_{g_1}(x)\| \leq \alpha_0$ ($1 \leq i \leq m$) and (x, y) satisfies (4-2) then (4-3) and

$$-(v_{f_0}(x) + v_{g_0}(x)) = \sum_{i=1}^k y_i (v_{f_1}(x) + v_{g_1}(x)) + \sum_{i=k+1}^m y_i^* (v_{f_2}(x) + v_{g_2}(x)).$$

$$y_1^* = (v_{f_1}(x) + v_{g_1}(x)) < 0 \quad (k+1 \leq i \leq m)$$

hold. Since the set of vectors $v_{f_1}(x^*)$ ($1 \leq i \leq k$) is linearly independent, there is a positive numbers $\delta_2 \leq \delta_1$ and $\alpha_1 \leq \alpha_0$ such that if $x \in B_{\delta_2}(x^*)$,

$|g_1(x)| \leq \alpha_1, \|v_{g_1}(x)\| \leq \alpha_1$ ($0 \leq i \leq m$) and (x, y) satisfies (4-2) then $y \in B_{\delta_1}(y^*)$. Therefore we have shown the existence of positive numbers δ_1, δ_2 and α_1 such that

$$(4-4) \quad F \text{ is one-to-one on } B_{\delta_2}(x^*) \times B_{\delta_1}(y^*),$$

$$(4-5) \quad F \text{ is locally nonsingular at every } z \in B_{\delta_2}(x^*) \times B_{\delta_1}(y^*),$$

$$(4-6) \quad \text{if } \text{norm}(g, B_{\delta_2}(x^*)) \leq \alpha_1, x \in B_{\delta_2}(x^*) \text{ and } (x, y) \text{ is a solution to (4-2) then } y \in B_{\delta_1}(y^*).$$

What we have left to show is that for every $\delta \in (0, \delta_2]$ there is an $\alpha \in (0, \alpha_1]$ such that whenever $\text{norm}(g, B_{\delta}(x^*)) \leq \alpha$, $B_{\delta}(x^*) \times B_{\delta}(y^*)$ contains a solution to (4-2)

which is unique in $B_{\delta}(x^*) \times B_{\delta}(y^*)$. Let $P = B_{\delta}(x^*) \times B_{\delta}(y^*)$. Note that the restriction $F|_P$ of the map F to the polyhedral set P is PC^1 . By (4-4) and (4-5), and by Lemma 2.1, there are positive numbers ϵ and ρ such that if the map G given by (4-1) satisfies

$$(4-7) \quad \|G(z)\| \leq \epsilon \text{ for every } z = (x, y) \in P$$

and

$$(4-8) \quad \|DG(z; \sigma)\| \leq \rho \|DG(z; \sigma)\|$$

for every $z \in \sigma \cap P$, $\sigma \in K^*$ and $w \in R^{n+m}$ with $\|w\| = 1$.

then $F + G$ is one-to-one in P . Define

$$\gamma = \min\{\|DG(z; \sigma)\| : z \in \sigma \cap P, \sigma \in K, \|w\| = 1\}.$$

It follows from (4-5) that $\gamma > 0$. Choose a positive number $\alpha_2 \leq \alpha_1$ such that if $\text{norm}(g, B_{\delta}(x^*)) \leq \alpha_2$ then the corresponding $G : |K^*| \rightarrow R^{n+m}$ defined by (4-1) satisfies (4-7) and

$$\|DG(z; \sigma)\| \leq \rho \gamma \text{ for all } z \in \sigma \cap P, \sigma \in K \text{ and } w \in R^{n+m} \text{ with } \|w\| = 1.$$

hence G satisfies (4-8) and $F + G$ is one-to-one in P . Thus $\text{norm}(g, B_{\delta_2}(x^*)) \leq \alpha$ implies that P contains at most one solution to (4-2).

Finally we prove that for each positive number $\delta \leq \delta_2$ there is a positive number $\alpha \leq \alpha_2$ such that $B_{\delta}(z^*) = B_{\delta}(x^*) \times B_{\delta}(y^*)$ contains a solution to (4-2) whenever $\rho(g, B_{\delta_2}(x^*)) \leq \alpha$. By Lemma 2.3 and the homotopy invariance theorem (6.2.2) of Ortega and Rheinboldt [24]), we can find a positive number $\alpha < \alpha_2$ such if $\text{norm}(g, B_{\delta_2}(x^*)) \leq \alpha$ then

$$\deg(F + G, B_{\delta}(z^*), 0) = \deg(F, B_{\delta}(z^*), 0) = +1 \text{ or } -1,$$

which implies that $B_{\delta}(z^*)$ contains a solution to (4-2) (see Kronecker Theorem, 6.3.1 of Ortega and Rheinboldt [24]).

Q.E.D.

Theorem 4.2. Let $z^* = (x^*, y^*)$ be a stationary point to $P(f)$. Assume that the Condition 1.1 holds and that x^* is s -stable (w.r.t. F^*) stationary solution to $P(f)$, where F^* is given by (1-3). Then F is locally nonsingular at $z^* = (x^*, y^*)$.

Proof. For simplicity of notations, we assume that $\{i : f_i(x^*) = 0, 1 \leq i \leq m\} = \{1, 2, \dots, k_1\}$ for some k_1 . It follows from the first assumption that the set of vectors $\nabla f_i(x^*)$ ($1 \leq i \leq k_1$) is linearly independent. By the continuity of f_i and ∇f_i at x^* ($1 \leq i \leq m$), we can take positive numbers δ_0 and α_0 such that if $x \in B_{\delta_0}(x^*)$, $\|g_1(x)\| \leq \alpha_0$ and $\|\nabla g_1(x)\| \leq \alpha_0$ ($1 \leq i \leq m$) then

$$\{i : f_i(x) + g_1(x) = 0, 1 \leq i \leq m\} \subset \{1, \dots, k_1\}$$

and the set of the vectors $\nabla f_i(x) + \nabla g_1(x)$ ($1 \leq i \leq k_1$) is linearly independent. By the second assumption, for some $\delta^* > 0$ and each positive $\delta \leq \min\{\delta_0, \delta^*\}$ there is a positive number $\alpha \leq \alpha_0$ such that if $\text{norm}(g, B_{\delta_0}(x^*)) \leq \alpha$ then $B_{\delta}(x^*) \times \mathbb{R}^m$ contains a solution to (4-2) which is unique in $B_{\delta_0}(x^*) \times \mathbb{R}^m$. Taking

$$g_0(x) = -c^T x, \\ g_1(x) = d_1 \quad (1 \leq i \leq m)$$

where $c \in \mathbb{R}^m$ and $d = (d_1, \dots, d_m) \in \mathbb{R}^m$, we see that for every sufficiently small $(c, d) \in \mathbb{R}^{2m}$ the system of equations

$$F(x, y) = (c, d), \quad (x, y) \in B_{\delta_0}(x^*) \times \mathbb{R}^m$$

has a unique solution. This implies that for some positive $\delta_1 \leq \min\{\delta_0, \delta^*\}$ F is one-to-one in $B_{\delta_1}(z^*)$. Hence, by Lemma 2.3, for some $s \in (-1, +1)$ and any $\delta \in (0, \delta_1)$, we have

$$\deg(F, B_{\delta}(z^*), 0) = s$$

and

$$\text{sign det } DF(z^*) = s \text{ or } 0 \text{ for every } z \in \sigma \cap B_{\delta_1}(z^*) \text{ and } \sigma \in K^*.$$

If $\text{sign det } DF(z^*) = s$ for every $\sigma \ni z^*$ then, by Theorem 3.3, F is locally non-singular at z^* .

We assume on the contrary that $\text{det } DF(z^*, \sigma^*) = 0$ for some $\sigma^* \in K^*$ containing z^* , and show that for any positive numbers $\delta \leq \min\{\delta_0, \delta^*\}$ and $\alpha \leq \alpha_0$ there exists $g \in P^*$ with $\text{norm}(g, B_{\delta_0}(x^*)) \leq \alpha$ such that $F + G$ is not one-to-one in $B_{\delta}(z^*)$.

Then we can see that x^* is not s -stable (w.r.t. F^*) because if two distinct $(x^j, y^j) \in B_{\delta}(z^*)$ ($j = 1, 2$) satisfy

$$F(x^j, y^j) + G(x^j, y^j) = (c, d)$$

for some sufficiently small $(c, d) \in \mathbb{R}^{2m}$ and if we define

$$g_0^j(x) = g_0(x) - c^T x, \\ g_1^j(x) = g_1(x) + d_1 \quad (1 \leq i \leq m)$$

for all $x \in \mathbb{R}^m$ then x^1 and x^2 are two distinct stationary solutions to $P(f + g^j)$.

Let $0 < \delta \leq \min\{\delta_0, \delta^*\}$ and $\alpha > 0$. Recall that $z^* \in \sigma^* \in K^*$ and

$\text{det } DF(z^*, \sigma^*) = 0$. Let $J \subset \{1, \dots, k_1\}$ be such that $\sigma^* = \tau(J)$, and k_0 be the dimension of the space $W = \{v \in \mathbb{R}^m : \nabla f_i(x^*)^T v = 0, i \in \{1, \dots, k\} \cup J\}$. If $k_0 = 0$ then, by Theorem 3.5, we have $\text{det } DF(z^*, \tau(J)) > 0$, a contradiction. Hence $k_0 \geq 1$.

Let B be an $n \times k_0$ matrix whose columns form a basis of the space W . By Theorem 3.5, we see

$$\text{det } B^T L(z^*) B = 0,$$

where $L(z^*)$ is defined by (3-6). Let $\bar{z} = (-1, 1) \setminus \{z^*\}$. By Lemma 3.4, there exists an $n \times n$ symmetric matrix Q such that

$$\text{sign det } B^T (L(z^*) + \gamma Q) B = \bar{z} \text{ for all } \gamma > 0.$$

Define, for all $x \in \mathbb{R}^n$,

$$g_0(x) = \frac{1}{2} (x - x^*)^T \nabla^2 \varphi(x - x^*)$$

and

$$g_1(x) = 0 \quad (1 \leq i \leq m).$$

Then, by Theorem 3.5, we see

$$\begin{aligned} (4-9) \quad & \text{sign det}[\nabla^2 L(x^*, 0^*) + DG(x^*, 0^*)] \\ & = \text{sign det } B^T(L(x^*) + \nabla \Phi)B \\ & = \bar{\alpha} \quad \text{for every } \gamma > 0. \end{aligned}$$

Since F is one-to-one in $B_\delta(x^*)$ and $\deg(F, \text{int } B_\delta(x^*), 0) = \alpha$, using the homotopy invariance theorem (6.2.2 of Ortega and Rheinbold [24]), we have

$$\deg(F + G, \text{int } B_\delta(x^*), 0) = \alpha \quad \text{for every sufficiently small } \gamma,$$

which together with (4-9) implies that $F + G$ is not one-to-one in $B_\delta(x^*)$ for every sufficiently small $\gamma > 0$ (see Lemma 2.3). Q.E.D.

Corollary 4.3. Let $x^* = (x^*, y^*)$ be a stationary point to $\Pi(f)$. Assume that

Condition 1.1 holds. Then the following three are equivalent:

(4-10) F is locally nonsingular at x^* .

(4-11) x^* is a s -stable (w.r.t. F) stationary solution to $\Pi(f)$.

(4-12) x^* is a s -stable (w.r.t. F^0) stationary solution to $\Pi(f)$.

Proof. We have shown that (4-10) implies (4-11) (Theorem 4.1) and that (4-12) implies (4-10) (Theorem 4.2). By the definition of the strong stability, (4-11) obviously

implies (4-12). Q.E.D.

5. S-STABLE (w.r.t. F) LOCAL MINIMUM SOLUTIONS.

In this section, we focus our attention to the set of local minimum solutions to $\Pi(f)$ which are s -stable (w.r.t. F). We will be concerned with the following two conditions:

Condition 5.1. $L(x^*)$ is positive semi-definite on the space $\{v \in \mathbb{R}^n : \nabla f_i(x^*)^T v = 0 \text{ for all } i \in \{1, \dots, m\}\}$ such that $f_1(x^*) = 0$.

Condition 5.2. $L(x^*)$ is positive definite on the space $\{v \in \mathbb{R}^n : \nabla f_i(x^*)^T v = 0 \text{ for all } i \in \{1, 2, \dots, l\} \cup \{j : y_j^* > 0, 1 \leq j \leq m\}\}$.

Lemma 5.3. Suppose that $x^* = (x^*, y^*) \in \mathbb{R}^{n+m}$ is a stationary point to $\Pi(f)$. If x^* is a local minimum solution to $\Pi(f)$ satisfying Condition 1.1 then $x^* = (x^*, y^*)$ satisfies Condition 5.1. If $x^* = (x^*, y^*)$ satisfies Condition 5.2 then x^* is an isolated local minimum solution to $\Pi(f)$.

Proof. See Sections 2.2 and 2.3 of Piacco and McCormick [15].

The following two theorems characterize s -stable (w.r.t. F) local minimum solutions to $\Pi(f)$.

Theorem 5.4. Let $x^* = (x^*, y^*)$ be a stationary point to $\Pi(f)$. Assume that Conditions 1.1 and 5.2 hold. Then x^* is s -stable (w.r.t. F).

Proof. In view of Theorem 3.3 and Corollary 4.3, it suffices to show

$$\det \nabla^2 L(x^*; J) > 0 \quad \text{for all } J \text{ such that } J_0 \subset J \subset J_1,$$

where J_0 and J_1 are given by (3-1). Let $J_0 \subset J \subset J_1$. If the index set

$\{1, \dots, l\} \cup J$ has n elements, Theorem 3.5 ensures that $\det \nabla^2 L(x^*; J) > 0$. Suppose that the set $\{1, \dots, l\} \cup J$ has $k < n$ elements. Then the subspace W of \mathbb{R}^n defined by

$$W = \{v \in \mathbb{R}^n : \nabla f_i(x^*)^T v = 0 \text{ for all } i \in \{1, \dots, l\} \cup J\}$$

has the dimension $n - k \geq 1$. Let B be an $n \times (n - k)$ matrix whose columns form a basis of the space W . Since W is a subspace of

$$\{v \in \mathbb{R}^n : \nabla f_i(x^*)^T v = 0 \text{ for all } i \in \{1, \dots, l\} \cup J_0\},$$

it follows from Condition 5.2 that the $(n - k) \times (n - k)$ matrix $B^T L(x^*) B$ is positive

definite, which implies $\det B^T L(z^*) B > 0$. Therefore, by Theorem 3.5, we obtain $\det DF(z^*; r(J)) > 0$. Q.E.D.

Theorem 5.5. Let $z^* = (x^*, y^*)$ be a stationary point to $P(f)$. Assume that z^* is s-stable (w.r.t. F) and that Conditions 1.1. and 5.1 hold. Then $z^* = (x^*, y^*)$ satisfies Condition 5.2.

Proof. By Corollary 4.3, $F : |K^e| \rightarrow \mathbb{R}^{n+m}$ is locally nonsingular at z^* . Let J_0 and J_1 be the index set defined by (3-1). For simplicity of notations, we assume that $J_0 = \{1, \dots, k_0\}$ and $J_1 = \{k_0+1, \dots, k_1\}$ for some k_0 and k_1 such that $z \leq k_0 \leq k_1 \leq n$. If $k_0 = k_1 = n$ then the dimension of the space

$$\{w \in \mathbb{R}^n : \forall f_i (z^*)^T w = 0 \text{ for all } i \in \{1, \dots, t\} \cup J_0\}$$

is zero and Condition 5.2 holds. Suppose $k_0 \leq n-1$. For each $k \in \{k_0, \dots, k_1\}$, define

$$W_k = \{w \in \mathbb{R}^n : \forall f_i (z^*)^T w = 0 \text{ (} 1 \leq i \leq k \text{)}\}.$$

Then each space W_k has dimension $n-k$ and $W_{k+1} \subset W_k$ ($k_0 \leq k \leq k_1 - 1$). Choose an $n \times (n-k_0)$ matrix C such that the first $(n-k)$ columns of C form a basis of W_k ($k_0 \leq k \leq \min\{k_1, n-1\}$). We shall show that the $(n-k_0) \times (n-k_0)$ symmetric matrix $C^T L(z^*) C$ is positive definite. Let B^k be the $n \times (n-k)$ matrix consisting of the first k columns of C ($k_0 \leq k \leq \min\{k_1, n-1\}$). Assume that $k_1 \leq n-1$. It follows from Condition 5.1 that the $(n-k_1) \times (n-k_1)$ symmetric matrix

$$(5-1) \quad \begin{matrix} k_1 \\ (B^k)^T L(z^*) B^k \end{matrix}$$

is positive semi-definite. On the other hand, by the local nonsingularity, the matrix (5-1) is nonsingular (see Corollary 3.6). Hence the matrix (5-1) is positive definite.

By Corollary 3.6, for every $k \in \{k_0, \dots, k_1\}$

$$(5-2) \quad \det(B^k)^T L(z^*) B^k > 0$$

Thus we have shown that all the leading principal minors of the symmetric matrix $C^T L(z^*) C$ are positive. Hence $C^T L(z^*) C$ is positive definite.

Finally we deal with the case where $k_1 = n$, i.e., $J_1 = \{1, \dots, n\}$. In this case we see $\dim W_{k_1} = 0$. By Theorem 3.5, $\det DF(z^*; r(J_1)) > 0$, and by Theorem 3.3, $\det DF(z^*; r(\{1, \dots, k\})) > 0$ for all $k \in \{k_0, \dots, n\}$. Thus (5-2) holds for every $k \in \{k_0, \dots, n-1\}$ (Theorem 3.5). Therefore all the leading principal minors of $C^T L(z^*) C$ are positive. Q.E.D.

As a direct consequence of Lemma 5.3 and Theorem 5.5, we have:

Corollary 5.6. Let $z^* = (x^*, y^*)$ be a stationary point to $P(f)$. Suppose that z^* is s-stable (w.r.t. F) and that Condition 1.1 holds. Then the following three are equivalent:

(5-3) z^* is an isolated local minimum solution to $P(f)$.

(5-4) $z^* = (x^*, y^*)$ satisfies Condition 5.1.

(5-5) $z^* = (x^*, y^*)$ satisfies Condition 5.2.

6. AN APPLICATION TO A PARAMETRIC PROGRAM.

Let T be a closed subset of R^n , h_i a map from $R^n \times T$ into R ($0 \leq i \leq n$) and $h = (h_0, h_1, \dots, h_m)$. Throughout this and next sections, we assume that $h(\cdot, t) \in F$ for each $t \in T$ and that $\partial h_i(x, t)/\partial x_j, \partial^2 h_i(x, t)/\partial x_j \partial x_k$ ($0 \leq i \leq m, 1 \leq j \leq k \leq n$) are continuous with respect to $(x, t) \in R^n \times T$. We will deal with the following parametric nonlinear program:

$$P2(t) \quad \begin{aligned} & \text{minimize} && h_0(x, t) \\ & \text{subject to} && x \in Y(t), \end{aligned}$$

where

$$Y(t) = \{x \in R^n : h_1(x, t) = 0 \quad (1 \leq i \leq l), \\ h_j(x, t) \leq 0 \quad (l+1 \leq j \leq m)\}.$$

and $t \in T$ is a parameter vector. The Kuhn-Tucker stationary condition to $P2(t)$ can be written as

$$(6-1) \quad \begin{aligned} & H(x, y, t) = 0, \\ & \text{where} \\ & H(x, y, t) = \begin{pmatrix} \nabla h_0(x, t) + \sum_{i=1}^l y_i \nabla h_i(x, t) + \sum_{j=l+1}^m y_j \nabla h_j(x, t) \\ -h_1(x, t) \\ \vdots \\ -h_l(x, t) \\ y_{l+1} - h_{l+1}(x, t) \\ \vdots \\ y_m - h_m(x, t) \end{pmatrix} \end{aligned}$$

for every $(x, y, t) \in R^{n+m} \times T$, where $\nabla h_i(x, t) = (\partial h_i(x, t)/\partial x_1, \dots, \partial h_i(x, t)/\partial x_n)^T$. Note that for each $t \in T$ the map $H(\cdot, \cdot, t) : R^{n+m} \times R^{n+m}$ is PC^1 on the subdivision R^n defined by (2-3) and (2-4). Let

$$(6-3) \quad \begin{aligned} I &= \{(x, t) \in R^n \times T : x \text{ is a stationary solution to } P2(t)\}, \\ I^s &= \{(x, t) \in R^n \times T : x \text{ is a s-stable (w.r.t. } F) \text{ stationary} \\ & \quad \text{solution to } P2(t)\}, \\ I^* &= \{(x, t) \in R^n \times T : x \text{ is a local minimum solution to } P2(t)\}. \end{aligned}$$

obviously $I^* \subset I$. We will assume the following constraint qualification which ensures that $I^* \subset I$.

Condition 6.1. If $(x, t) \in I \cup I^*$ then the set of vectors $\nabla h_i(x, t)$ such that $h_i(x, t) = 0$ ($1 \leq i \leq m$) is linearly independent.

Theorem 6.2. Let $z^* = (x^*, y^*)$ be a stationary point to $P2(t^*)$. Assume that the map $H(\cdot, \cdot, t^*) : R^{n+m} \rightarrow R^{n+m}$ is locally nonsingular at $z^* = (x^*, y^*)$. Then there exist open neighborhoods U of z^* , V of t^* and a continuous map $z : V \cap T \rightarrow U$ such that $z(t)$ is a stationary point to $P2(t)$ for all $t \in V \cap T$ and that if $z^* \in U$ is a stationary point to $P2(t)$ for some $t \in V \cap T$ then $z^* = z(t)$.

Proof. By Theorem 4.1., x^* is a s-stable (w.r.t. F) stationary solution to $P2(t^*)$. Hence there exist positive numbers δ and β such that if $t \in B_\beta(t^*)$ then $B_\delta(x^*)$ contains a unique stationary solution to $P2(t)$, which we will denote by $x(t)$. Let $y(t) \in R^m$ be the Lagrange multiplier vector associated with the stationary solution $x(t)$ to $P2(t)$; $(x(t), y(t), t)$ satisfies (6-1). Let $z(t) = (x(t), y(t))$. The continuity of the map z for every t sufficiently close to t^* follows from the fact $z(t)$ is a locally nonsingular point of $H(\cdot, \cdot, t) : R^{n+m} \rightarrow R^{n+m}$, which implies that $x(t)$ is a s-stable (w.r.t. F) stationary solution to $P2(t)$, for every t sufficiently close to t^* . Q.E.D.

Let $V^1 \subset V^2 \subset R^p$. We say that V^1 is open (or closed) relative to V^2 if $V^1 = V^2 \cap U$ for some open (or closed) subset U of R^p .

Theorem 6.3. Assume that Condition 6.1 holds. Then $I^* \cap I^*$ is open and closed relative to I^* .

Proof. First we show $I^* \cap I^*$ is open relative to I^* . Let $(x^*, t^*) \in I^* \cap I^*$. By Condition 6.1, there is a unique $y^* \in R^m$ such that (x^*, y^*, t^*) satisfies (6-1) and that the map $H(\cdot, \cdot, t^*) : R^{n+m} \rightarrow R^{n+m}$ is locally nonsingular at $z^* = (x^*, y^*)$. It suffices to show that if a solution (x, y, t) of (6-1) is sufficiently close to (x^*, y^*, t^*) then $(x, t) \in I^*$. Let $I_0 = \{(1, \dots, 1) \cup \{j : y_j^* > 0, 1 \leq j \leq m\}$. By Corollary 5.6, the $n \times n$ symmetric matrix $H(z^*, t^*)$ is positive definite on the space W , where

$$w = (w \in \mathbb{R}^n : \nabla h_1(x^0, t^0)^T w = 0, i \in I_0).$$

$$M(x, t) = \nabla^2 h_0(x, t) + \sum_{i=1}^k \lambda_i \nabla^2 h_i(x, t) + \sum_{j=1}^m \mu_j \nabla^2 h_j(x, t)$$

for all $(x, t) = (x, y, t) \in \mathbb{R}^{n+m} \times \mathbb{T}$ and $\nabla^2 h_1(x, t)$ denote the Hessian matrix of the map $h_1(\cdot, \cdot, t) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ at x . Hence if a solution $(s, t) = (x, y, t)$ to (6-1) is sufficiently close to (x^0, y^0, t^0) then $I_0 \subset I$ and the matrix $M(s, t)$ is positive definite on the space $\{w \in \mathbb{R}^n : \nabla h_1(x, t)^T w = 0 (i \in I)\}$, where

$I = \{1, \dots, k\} \cup \{j : y_j > 0, 1 + 1 \leq j \leq m\}$; hence $(x, t) \in I^0$ by Lemma 5.3. Thus we have shown that $I^0 \cap I^*$ is open relative to I^0 .

To prove that $I^0 \cap I^*$ is closed relative to I^0 , we consider a sequence $\{(x^p, t^p) \in I^0 \cap I^*\}$ which converges to some $(x^0, t^0) \in I^0$, and we shall show $(x^0, t^0) \in I^*$. Corresponding to each $(x^p, t^p) \in I^0 \cap I^*$, there is a unique $y^p \in \mathbb{R}^m$ such that (x^p, y^p, t^p) satisfies (6-1). Also for some unique $y^0 \in \mathbb{R}^m$, (x^0, y^0, t^0) satisfies (6-1). Let $J_1 = \{i : h_1(x^0, t^0) = 0, 1 + 1 \leq i \leq m\}$. Taking an appropriate subsequence if necessary, we may assume that

$$J = \{j : h_j(x^p, t^p) = 0, 1 + 1 \leq j \leq m\}$$

for some $J \subset \{1, \dots, m\}$ and all p . By the continuity of the maps $h_1 (1 \leq i \leq m)$, we see $J_1 \supset J$. Hence Condition 6.1 ensures that $\{y^p\}$ converges to y^0 .

For every p , define

$$w_p = (w \in \mathbb{R}^n : \nabla h_1(x^p, t^p)^T w = 0, i \in \{1, \dots, k\} \cup J)$$

and

$$w_0 = (w \in \mathbb{R}^n : \nabla h_1(x^0, t^0)^T w = 0, i \in \{1, \dots, k\} \cup J)$$

By Lemma 5.3, the $n \times n$ symmetric matrix $M(x^p, t^p)$, where $x^p = (x^p, y^p)$, is positive semi-definite on the space w_p . Taking the limit as $p \rightarrow \infty$, we obtain that $M(x^0, t^0)$ is positive semi-definite on w_0 , which together with $J \subset J_1$ implies that $M(x^0, t^0)$ is positive semi-definite on the space

$$\{w \in \mathbb{R}^n : \nabla h_1(x^0, t^0)^T w = 0, i \in \{1, \dots, k\} \cup J_1\}.$$

The desired result follows from Corollary 5.6.

Q.E.D.

It is easily verified that if Condition 6.1 holds then I^0 is open relative to I . Hence $I^0 \cap I^*$ is open relative to I under Condition 6.1.

Corollary 6.4. Suppose that Condition 6.1 holds. Let S be a connected subset of I^0 which contains (x^0, t^0) such that x^0 is a local minimum solution to $P2(t^0)$.

Then x is a local minimum solution to $P2(t)$ for all $(s, t) \in S$.

Proof. By Theorem 6.3, there exist an open subset U of \mathbb{R}^{n+k} and a closed subset V of \mathbb{R}^{n+k} such that $I^0 \cap I^* = I^0 \cap U = I^0 \cap V$. Hence

$$(x^0, t^0) \in S \cap I^* = S \cap I^0 \cap I^* \cap U = S \cap U = S \cap V,$$

which implies that the nonempty subset $S \cap I^0$ of S is open and closed relative to S .

Since S is connected, we obtain $S = S \cap I^0$.

Q.E.D.

7. AN APPLICATION TO A CLASS OF CONTINUOUS DEFORMATION METHODS.

To solve $P1(f)$, we artificially construct another nonlinear program $P1(g)$ which has a trivial stationary solution x^0 and a parametric nonlinear program $P2(t)$ ($t \in [0,1]$) which continuously deforms $P1(g)$ to $P1(f)$, i.e., $h(\cdot, 0) = g$ and $h(\cdot, 1) = f$. Then we start from $(x^0, 0)$ and trace the connected component $S^0 \subset I$ which contains $(x^0, 0)$ to attain a stationary solution x^1 to $P2(1)$. Some methods (Laves [9, 11], Kojima [17, 19], Lemke [21], Saigal [30], Todd [33]) which were developed in the fixed point and complementarity theory are based on the above ideas.

We can prove:

Theorem 7.1. Let $T = [0,1]$. Suppose Condition 6.1 holds. Let x^0 be a σ -stable (w.r.t. F) local minimum solution to $P2(0)$ and $S^0 \subset I$ be a compact connected set which contains $(x^0, 0)$. Assume that $S^0 \subset I^s$. Then S^0 can be written as

$$S^0 = \{(x(t), t) : 0 \leq t \leq t^*\}$$

for some $t^* \in [0,1]$ and some continuous map $x : [0, t^*] \rightarrow R^n$. Furthermore, each $x(t)$ is an isolated local minimum solution to $P2(t)$.

Proof. The first part directly follows from Theorem 6.2 and the second from Corollary 6.4.

Remark. By using Sard's theorem (5.2.5 of Ortega and Rheinboldt [24]), we can easily verify that for almost all $(c,d) \in R^{n+m}$ the set

$$\{(x,y,t) \in R^{n+m} \times [0,1] : H(x,y,t) = (c,d)\}$$

consists of disjoint piecewise smooth one manifolds. See also Alexander [1], Chow, Mallet-Paret and York [6], and Garcia and Gould [16].

Corollary 7.2. Let T, x^0, s^0, t^* and x be as in Theorem 7.1. If $h : R^{n+1} \rightarrow R^{1+m}$ has the form

$$\begin{aligned} h_0(x,t) &= (1-t)g_0(x) + tf_0(x) \\ h_1(x,t) &= f_1(x) \quad (1 \leq i \leq m) \end{aligned}$$

for every $(x,t) \in R^n \times T$, then $x(t) \in X(f)$ for all $t \in [0, t^*]$ and $f_0(x(t))$ is monotone nonincreasing with respect to $t \in [0, t^*]$.

Proof. It suffices to show that if $\epsilon > 0$ is sufficiently small and $0 \leq t < t + \epsilon \leq t^*$ then $f_0(x(t + \epsilon)) \leq f_0(x(t))$. Let $t \in [0, t^*]$. Since $x(t)$ is an isolated local minimum solution to $P2(t)$, there exists a positive number δ such that

$$h_0(x(t), t) < h_0(x', t) \text{ for all } x' \in U,$$

where $U = \{x' \in X(f) : x' \in B_\delta(x(t)), x' \neq x(t)\}$. Specifically, we have

$$h_0(x(t), t) < h_0(x', t) \text{ for all } x' \in V,$$

where $V = \{x' \in U : \|x' - x(t)\| = \delta\}$. By the continuity of h_0 , if $\epsilon > 0$ is sufficiently small then

$$h_0(x(t), t + \epsilon) < h_0(x', t + \epsilon) \text{ for all } x' \in V$$

Hence for every sufficiently small $\epsilon > 0$, the program

$$\begin{aligned} &\text{minimize } h_0(x', t + \epsilon) \\ &\text{subject to } x' \in X(f) \cap B_\delta(x(t)) \end{aligned}$$

has a minimum solution in the interior of $B_\delta(x(t))$, which must coincide with $x(t + \epsilon)$. Therefore, we obtain that, for every sufficiently small $\epsilon > 0$,

$$\begin{aligned} h_0(x(t), t) &\leq h_0(x(t + \epsilon), t) \\ h_0(x(t + \epsilon), t + \epsilon) &\leq h_0(x(t), t + \epsilon) \end{aligned}$$

from which the desired result follows.

Q.E.D.

8. CONCLUDING REMARKS.

As stated in the Introduction, the Kuhn-Tucker stationary condition can be also formulated as a system of generalized equations. The local nonsingularity of the map $F : |x| + g^{n+m}$ defined by (1-2) is equivalent to the strong regularity (see Robinson [26, 27]) for the system of generalized equations associated with the Kuhn-Tucker stationary condition. Theorem 6.2 follows from Theorem 2.1 of Robinson [27]. Also Theorems 3.3 and 5.4 have close relations with Theorems 4.1 and 3.1 of [27], respectively.

The result in Theorem 7.1 was recently shown by Saigal [30] for a special case where $P_2(t)$ has no constraints (i.e., $m = 0$). See also Saari and Saigal [28].

REFERENCES

- [1] J. C. Alexander, "The topological theory of an embedding method", Symposium on Embedding Methods for Nonlinear Problems, Linz, Austria, Oct. 1977.
- [2] J. C. Alexander and J. A. York, "The homotopy continuation method: Numerically implementable topological procedures", Transactions of the American Mathematical Society 242 (1978) 271-284.
- [3] P. S. Alexandroff, Combinatorial Topology, Graylock Press, New York, 1960.
- [4] E. Allgower and E. Georg, "Simplicial and continuation methods for approximating fixed points", Tech. Report, Colorado State University, Fort Collins, Colorado, August, 1978.
- [5] C. Berge, Topological Spaces, Macmillan, New York, 1963.
- [6] S. W. Chow, J. Mallet-Paret and J. A. York, "Finding zeros of maps: Homotopy methods that are constructive with probability one", to appear in Journal of Non-linear Analysis - Theory, Methods and Applications.
- [7] R. W. Cottle and G. B. Dantzig, "Complementary pivot theory of mathematical programming", Linear Algebra and Its Application 1 (1968) 103-125.
- [8] G. B. Dantzig, J. Folkman and M. Shapiro, "On the continuity of the minimum set of a continuous function", Journal of Mathematical Analysis and Applications 17 (1967) 519-548.
- [9] B. C. Eaves, "On quadratic programming", Management Science 17 (1971) 698-711.
- [10] B. C. Eaves, "A short course in solving equations with PL homotopies", SIAM-NMS Proceedings 9 (1976) 73-143.
- [11] B. C. Eaves, "Computing stationary points", Mathematical Programming Study 7 (1976) 1-14.
- [12] S. Eilenberg and N. Steenrod, Foundations of Algebraic Topology, Princeton University Press, New Jersey, 1952.
- [13] J. P. Evans and P. J. Gould, "Stability in nonlinear programming", Operations Research 18 (1970) 107-118.
- [14] A. V. Fiacco, "Sensitivity analysis for nonlinear programming using penalty methods", Mathematical Programming 10 (1976) 287-311.

- [15] A. V. Fiacco and G. P. McCormick, Nonlinear Programming: Sequential Unconstrained Minimization Technique, Wiley, New York, 1968.
- [16] C. B. Garcia and F. J. Gould, "A theorem on homotopy paths", Mathematics of Operations Research 3 (1978) 282-289.
- [17] M. Kojima, "Computational methods for solving the nonlinear complementarity problem", Kyoto Engineering Reports 27 (1974) 1-41.
- [18] M. Kojima, "Studies of PL approximations of piecewise- C^1 mappings in fixed point and complementarity theory", Mathematics of Operations Research 3 (1978) 17-36.
- [19] M. Kojima, "A complementary pivoting approach to parametric nonlinear programming", to appear in Mathematics of Operations Research.
- [20] H. W. Kuhn and A. W. Tucker, "Nonlinear programming", in J. Neyman ed., Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, University of California Press, California, 1951.
- [21] C. E. Lemke, "Bimatrix equilibrium points and mathematical programming", Management Science 11 (1965) 681-689.
- [22] C. E. Lemke, "Recent results on complementarity problems", in J. B. Rosen, O. L. Mangasarian and K. Ritter ed., Nonlinear Programming, Academic Press, New York, 1970.
- [23] O. L. Mangasarian, "Nonlinear Programming", McGraw-Hill, New York, 1969.
- [24] J. M. Ortega and W. Rheinboldt, Iterative Solutions of Nonlinear Equations in Several Variables, Academic Press, New York, 1970.
- [25] S. M. Robinson, "Perturbed Kuhn-Tucker points and rates of convergence for a class of nonlinear programming algorithms", Mathematical Programming 7 (1974) 1-16.
- [26] S. M. Robinson, "Generalized equations and their solutions, Part I: Basic theory", to appear in Mathematical Programming Study 10.
- [27] S. M. Robinson, "Strongly regular generalized equations", Technical Summary Report #1877, Mathematics Research Center, University of Wisconsin, Sept. 1978.
- [28] G. Saari and R. Saigal, "Generic properties of paths generated by fixed point algorithms", in preparation.
- [29] R. Saigal, "Fixed point computing methods", in Encyclopedia of Computer Science and Technology, Vol. 6, Marcel Dekker Inc., 1977.

- [30] R. Saigal, "The fixed point approach to nonlinear programming", ORSA/TIMS Meeting in Atlanta, November, 1977; also to appear in Mathematical Programming Study.
- [31] M. Samelson, R. M. Thrall and O. Weslar, "A partition theorem for Euclidean n -space", Proceedings of American Mathematical Society 9 (1958) 805-807.
- [32] M. J. Todd, The Computation of Fixed Points and Applications, Springer, New York, 1976.
- [33] M. J. Todd, "New Fixed-point algorithms for economic equilibria and constrained optimization", Tech. Report 362, School of Operations Research and Industrial Engineering, Cornell University, Ithaca, New York, 1977.
- [34] J. H. C. Whitehead, "On C^1 -complexes", Annals of Mathematics 41 (1940) 809-824.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1920	2. GOVT ACCESSION NO. 14 / MRC-75R-1920	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) STRONGLY STABLE STATIONARY SOLUTIONS IN NONLINEAR PROGRAMS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) Masakazu Kojima		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) 15 / DAAG29-75-C-0924, ✓ NSF-MCS78-09525
11. CONTROLLING OFFICE NAME AND ADDRESS see item 18 below		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 5 - Mathematical Programming and Operations Research
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 / 23p.		17. REPORT DATE February 1979
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 9 / Technical summary report		18. NUMBER OF PAGES 36
18. SUPPLEMENTARY NOTES U.S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709 National Science Foundation Washington, D.C. 20550		19. SECURITY CLASS. (of this report) UNCLASSIFIED
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear Program, Stability, Parametric Program, Fixed Point and Complementarity Theory		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) For each continuously twice differentiable map $f = (f_0, f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^{1+m}$, we define a nonlinear program: $P1(f) \quad \text{minimize } f_0(x)$ $\text{subject to } f_1(x) = 0 \quad (1 \leq i \leq l),$ $f_j(x) \leq 0 \quad (l+1 \leq j \leq m). \quad (\text{continued})$		

Yuu

A (Kuhn-Tucker's) stationary solution x to $P1(f)$ is said to be strongly stable if there exists an open neighborhood U of x such that each open neighborhood $V \subset U$ of x contains a stationary solution to a perturbed problem $P1(f + g)$ which is unique in U whenever $g_i(x)$, $\partial g_i(x)/\partial x_j$ and $\partial^2 g_i(x)/\partial x_j \partial x_k$ ($0 \leq i \leq m$, $1 \leq j \leq n$, $1 \leq k \leq n$) are sufficiently small for all x in U . We will give conditions on the gradient vectors and the Hessian matrices of f_i ($0 \leq i \leq m$) which characterize the strong stability. These conditions are then applied to a parametric nonlinear program:

$$\begin{aligned}
 P2(t) \quad & \text{minimize } h_0(x, t) \\
 & \text{subject to } h_i(x, t) = 0 \quad (1 \leq i \leq l) , \\
 & \quad \quad \quad h_j(x, t) \leq 0 \quad (l + 1 \leq j \leq m) .
 \end{aligned}$$

where t is a parameter vector varying in a closed subset T of R^q . Let Σ^S be the set of points (x, t) in $R^n \times T$ such that x is a strongly stable stationary solution to $P2(t)$. Under a certain constraint qualification and the continuity and the differentiability of the map $h : R^n \times T \rightarrow R^{l+m}$, we will establish that if S is a connected subset of Σ^S and if x^* is a local minimum solution to $P2(t^*)$ for some $(x^*, t^*) \in S$ then x is a local minimum solution to $P2(t)$ for all $(x, t) \in S$. Finally this result is applied to showing some interesting properties of a class of methods developed in the fixed point and complementarity theory.