

AD-A071 946


COLORADO STATE UNIV FORT COLLINS DEPT OF MATHEMATICS F/G 12/1
CONTINUITY OF BEST RECIPROCAL POLYNOMIAL APPROXIMATION ON < 0 , --ETC(U)
APR 79 C B DUNHAM, G D TAYLOR AFOSR-76-2878

UNCLASSIFIED


AFOSR-TR-79-0856

NL

| OF |
AD
A071946




END
DATE
FILMED
8 -79
DDC



CONTINUITY OF BEST RECIPROCAL POLYNOMIAL APPROXIMATION ON $[0, \infty)$

by

Charles B. Dunham and G. D. Taylor¹**LEVEL**Introduction

Over the past ten years considerable progress has been made in studying various questions concerning rational approximation on unbounded sets. To a large extent the starting point of this effort was the paper of Cody, Meinardus and Varga [6] and this has led to investigations of best approximation properties in various settings [1-4, 8-9] and studies of the error of best approximation [10-12].

In this paper we wish to study the best approximation properties of strong uniqueness and continuity of the best approximation operator for reciprocal polynomial approximation on $[0, \infty)$ of continuous positive functions tending to 0 as $x \rightarrow \infty$. Thus, we define

- (1) $C_0^+[0, \infty) = \{f \in C[0, \infty): f(x) > 0, x \in [0, \infty) \text{ and } \lim_{x \rightarrow \infty} f(x) = 0\}$,
and
(2) $R_n = \{\frac{1}{p}: p \in \Pi_n, p(x) > 0, x \in [0, \infty)\}$, $n \geq 1$,

where Π_n denotes the class of all algebraic real polynomials of degree $\leq n$. Furthermore, define $\|f\| = \sup\{|f(x)|: x \in [0, \infty)\}$ in what follows. In this setting, it is known that best approximations exist and are unique [3, 4] and that the following characterization theorem holds:

¹Research sponsored by the Air Force Office of Scientific Research, Air Force Systems Command, USAF, under Grant AFOSR-76-2878C and by National Science Foundation under Grant MCS-78-05847.

Approved for public release;
distribution unlimited.

79 07 27 038

A071946

DDC FILE COPY

THEOREM 1 [4]. Let $f \in C_0^+[0, \infty) \sim R_n$ with $n \geq 1$. Then $\frac{1}{p^*}$ is the best approximation to f from R_n on $[0, \infty)$ iff

(i) (standard alternation) there exist $\{x_i\}_{i=0}^{n+1}$, $0 \leq x_0 < x_1 < \dots < x_{n+1}$, such that $|f(x_i) - \frac{1}{p^*(x_i)}| = \|f - \frac{1}{p^*}\|$, $i = 0, \dots, n+1$ and

$$f(x_i) - \frac{1}{p^*(x_i)} = -(f(x_{i+1}) - \frac{1}{p^*(x_{i+1})}), \quad i = 0, \dots, n;$$

or

(ii) (nonstandard alternation) $\partial p^* \leq n-1$ and there exist $\{x_i\}_{i=0}^n$,

$$0 \leq x_0 < x_1 < \dots < x_n \text{ such that } f(x_i) - \frac{1}{p^*(x_i)} = (-1)^{n-i} \|f - \frac{1}{p^*}\|.$$

In both cases the points $\{x_i\}$ are called extreme points. Also, we wish to note that for $n \geq 1$, p^* cannot be a constant. Indeed, since $f(x) > 0$ for all $x \in [0, \infty)$ and $\lim_{x \rightarrow \infty} f(x) = 0$, then in order for the reciprocal of a constant, $\frac{1}{c^*}$, to be a best approximation to f , we must have that $c^* = \frac{2}{M}$ where $M = \max_{x \geq 0} f(x)$. Since $f(x) \rightarrow 0$ as $x \rightarrow \infty$ we can find $x_0 > 0$ such that $f(x) < \frac{1}{M}$ for $x \geq x_0$. It is then easily seen that for $p^*(x) = \epsilon(x - x_0) + c^*$ with $\epsilon > 0$ and sufficiently small that

$$\|f - \frac{1}{p^*}\| < \|f - \frac{1}{c^*}\| \text{ by a straightforward continuity-compactness}$$

argument.

In addition, it has been shown in [3] that if $\frac{1}{p^*} \in R_n$ is the best approximation to $f \in C_0^+[0, \infty)$ from R_n with $\partial p^* = n$ then both strong uniqueness (i.e., $\|f - \frac{1}{p}\| - \|f - \frac{1}{p^*}\| \geq \gamma \|\frac{1}{p} - \frac{1}{p^*}\|$, $\gamma = \gamma(f) > 0$ for all $\frac{1}{p} \in R_n$) and Lipschitz continuity of the best approximation operator at f (i.e., $\|\frac{1}{p^*} - \frac{1}{p_g}\| \leq \beta \|f - g\|$, $\beta = \beta(f) > 0$, $g \in C_0^+[0, \infty)$ and $\frac{1}{p_g}$ the best approximation to g from R_n) hold. Furthermore, it was shown in [3]

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)

NOTICE OF TRANSMITTAL TO DDC

This technical report has been reviewed and is approved for public release IAW AFR 190-12 (7b).

Distribution is unlimited.

A. D. BLOSE

Technical Information Officer

that for each f whose corresponding best approximation from R_n , $\frac{1}{p^*}$, satisfies $ap^* \leq n - 2$ the strong uniqueness theorem cannot hold. In this present paper we shall prove that if $f \in C_0^+[0, \infty)$ has $\frac{1}{p^*} \in R_n$ as its best approximation then (i) if $ap^* \leq n - 2$ (i.e. $\frac{1}{p^*}$ is deficient of order two or more) then the best approximation operator is discontinuous at f and (ii) if $ap^* = n - 1$ then the best approximation operator is continuous at f . It remains open as to whether or not a strong uniqueness theorem holds in the case that $ap^* = n - 1$.

Main Results

In this section we state and prove our main results. The first result establishing the discontinuity of the best approximation operator is given in two parts. The first theorem will treat this problem for the case that either $\frac{1}{p^*}$ is deficient of order three or more, or when $\frac{1}{p^*}$ is deficient of order two and $f - \frac{1}{p^*}$ possesses a standard alternating sequence. In this case we can prove even stronger results concerning the discontinuous behavior of the best approximation operator. The second theorem will treat the discontinuity of the best approximation operator when $\frac{1}{p^*}$ is deficient of order two with only nonstandard alternation holding for $f - \frac{1}{p^*}$. Our final result will be to prove that the best approximation operator is continuous whenever $\frac{1}{p^*}$ is deficient of order one.

THEOREM 2. Let $f \in C_0^+[0, \infty) \sim R_n$ and $\frac{1}{p^*} \in R_n$ be the best approximation to f from R_n . Further, assume that $ap^* \leq n - 2$ and that if $ap^* = n - 2$

then $f - \frac{1}{p^*}$ possesses a standard alternating set. Then, given $\epsilon > 0$ there exists $\delta > 0$, $\left\{\frac{1}{p_k}\right\}_{k=1}^{\infty} \subset R_n$ and $\{g_k\}_{k=1}^{\infty} \subset C_0^+[0, \infty)$ such that each g_k has $\frac{1}{p_k}$ as its best approximation from R_n , g_k converges uniformly to f on $[0, \infty)$ and $\delta \leq \left\|\frac{1}{p^*} - \frac{1}{p_k}\right\| \leq \epsilon$ for all k .

REMARK. This theorem establishes that not only is the best approximation operator discontinuous at f but, in fact, that it is also not possible for a local (relative to $\frac{1}{p^*}$) strong uniqueness result to hold.

Proof. Set $E = \|f - \frac{1}{p^*}\| > 0$ and assume without loss of generality that $\epsilon \leq \frac{E}{4}$. Set $\delta = \frac{\epsilon}{8}$. Since we are assuming throughout this paper that $n \geq 1$, we have that $p^*(x)$ is not identically equal to a constant which implies that $\lim_{x \rightarrow \infty} p^*(x) = \infty$. Select $\beta > 0$ such that $f(x) \leq \epsilon$ and $p^*(x) > \frac{4}{\epsilon}$ for all $x \geq \beta$. Set $e_k = (\frac{\epsilon}{4} + \frac{2}{p^*(k)})^{-1}$ and note that for $k \geq \beta$, $p^*(k) > e_k$. Define $p_k \in \Pi_n$ by

$$p_k(x) = e_k + (p^*(x) - e_k)\left[\left(\frac{x}{k} - 1\right)^2 + \frac{e_k}{p^*(k) - e_k}\right], \quad k \geq \beta.$$

Since for all $x \geq \beta$, $p^*(x) > e_k$, we have that $(p^*(x) - e_k)\left[\left(\frac{x}{k} - 1\right)^2 + \frac{e_k}{p^*(k) - e_k}\right] > 0$ implying that $p_k(x) > e_k > \frac{4}{3\epsilon}$ for $x \geq \beta$.

Next, observe that $e_k \rightarrow \frac{1}{\epsilon}$ as $k \rightarrow \infty$ and $\left(\frac{x}{k} - 1\right)^2$ converges uniformly to 1 on $[0, \beta]$ as $k \rightarrow \infty$. Thus, p_k converges uniformly to p^* on $[0, \beta]$ as $k \rightarrow \infty$. Now, let

$$\eta = \min\left(\min_{x \in [0, \beta]} f(x), \min_{x \in [0, \beta]} \frac{1}{p^*(x)}, \epsilon\right)$$

Accession For	
ADIS Gen&I	☒
DDC TAB	☐
Unannounced	☐
Justification	☐
By _____	
Distribution/ _____	
Availability Codes	
Dist	Availand/or special
PA	

and select $\mu \geq \beta$ such that for $k \geq \mu$, $\max_{x \in [0, \beta]} \left| \frac{1}{p_k(x)} - \frac{1}{p^*(x)} \right| \leq \frac{\eta}{2}$. This

implies that $\frac{1}{p_k(x)} \geq \frac{1}{p^*(x)} - \frac{\eta}{2} \geq \frac{\eta}{2} > 0$ for all $x \in [0, \beta]$ and $k \geq \mu$.

Hence $\frac{1}{p_k} \in R_n$ for $k \geq \mu$ as $p_k(x) > 0$ for all $x \in [0, \infty)$.

Now, since $p_k(k) = 2e_k$ we have that $\frac{1}{p_k(k)} - \frac{1}{p^*(k)} = \frac{\varepsilon}{8} = \delta$. Thus,

$\left\| \frac{1}{p_k} - \frac{1}{p^*} \right\| \geq \delta$ for all $k \geq \mu$. Also, for $k \geq \mu$ we have that

$\left| \frac{1}{p_k(x)} - \frac{1}{p^*(x)} \right| \leq \frac{\eta}{2} < \varepsilon$ for $x \in [0, \beta]$. In addition, for $x \geq \beta$ we have

that $\frac{1}{p^*(x)} < \frac{\varepsilon}{4}$ and $\frac{1}{p_k(x)} \leq \frac{3\varepsilon}{4}$ for $k \geq \mu$. Hence $\left\| \frac{1}{p_k} - \frac{1}{p^*} \right\| \leq \varepsilon$ as claimed.

Finally, define g_k , for $k \geq \mu$, by

$$g_k(x) = \begin{cases} f(x) + \frac{1}{p_k(x)} - \frac{1}{p^*(x)}, & x \in [0, \beta] \\ f(x), & x \geq \beta + \frac{1}{k} \\ \text{linear with endpoint values } f(\beta) + \frac{1}{p_k(\beta)} - \frac{1}{p^*(\beta)} \text{ and } f(\beta + \frac{1}{k}), & x \in [\beta, \beta + \frac{1}{k}]. \end{cases}$$

Clearly, $g_k \in C[0, \infty)$, $k \geq \mu$ and since $f(x) + \frac{1}{p_k(x)} - \frac{1}{p^*(x)} \geq f(x) - \frac{\eta}{2} > 0$

for $x \in [0, \beta]$ we have that $g_k \in C_0^+[0, \infty)$ for $k \geq \mu$. Since $f(\beta) + \frac{1}{p_k(\beta)} - \frac{1}{p^*(\beta)}$

$< f(\beta) + \frac{\eta}{2} \leq f(\beta) + \frac{\varepsilon}{2} \leq f(\beta) + \frac{\varepsilon}{8} \leq \frac{3\varepsilon}{8}$ and $f(\beta + \frac{1}{k}) \leq \frac{\varepsilon}{4}$ we have that

$g_k(x) \leq \frac{3\varepsilon}{8}$ for $x \in [\beta, \beta + \frac{1}{k}]$ implying $g_k(x) \leq \frac{3\varepsilon}{8}$ for $x \geq \beta$. Also,

$\left\| \frac{1}{p_k} - \frac{1}{p^*} \right\| < \varepsilon \leq \frac{\varepsilon}{4}$ implies that $\frac{1}{p_k(x)} < \frac{1}{p^*(x)} + \frac{\varepsilon}{4} \leq \frac{\varepsilon}{2}$ for $x \geq \beta$. From

this it follows that $\left| g_k(x) - \frac{1}{p_k(x)} \right| \leq \frac{\varepsilon}{8}$ for $x \geq \beta$. In addition, for

for $x \in [0, \beta]$ we have that $g_k(x) - \frac{1}{p_k(x)} = f(x) - \frac{1}{p^*(x)}$ and this implies

that $g - \frac{1}{p_k}$ exhibits the same alternating behavior as $f - \frac{1}{p^*}$ on $[0, \infty)$.

Thus, if $f - \frac{1}{p^*}$ has a standard alternating set so does $g_k - \frac{1}{p_k}$ implying

that $\frac{1}{p_k}$ is the best approximation to g_k from R_n on $[0, \infty)$. If $f - \frac{1}{p^*}$

possesses only a nonstandard alternating set then so does $g_k - \frac{1}{p_k}$. Since

in this case we must have that $\alpha p_k \leq n - 3$, we must have that $\alpha p_k \leq n - 1$

implying once again that $\frac{1}{p_k}$ is the best approximation to g_k from R_n on

$[0, \infty)$. Since it is clear that g_k converges uniformly to f on $[0, \infty)$,

the proof is completed by relabeling the sequences $\left\{\frac{1}{p_k}\right\}_{k=\mu}^{\infty}$ and $\{g_k\}_{k=\mu}^{\infty}$

as $\left\{\frac{1}{p_k}\right\}_{k=1}^{\infty}$ and $\{g_k\}_{k=1}^{\infty}$, respectively.

For the case that $\alpha p^* = n - 2$ and $f - \frac{1}{p^*}$ has only a nonstandard alternating sequence we have the slightly weaker theorem:

THEOREM 3. Let $f \in C_0^+[0, \infty) \sim R_n$ and $\frac{1}{p^*} \in R_n$ be its best approximation

from R_n . Further, assume that $\alpha p^* = n - 2$ and $f - \frac{1}{p^*}$ possesses only a

nonstandard alternating set. Then there exists $\left\{\frac{1}{p_k}\right\}_{k=1}^{\infty} \subset R_n$ and

$\{g_k\}_{k=1}^{\infty} \subset C_0^+[0, \infty)$ such that for each k , $\frac{1}{p_k}$ is the best approximation to

g_k from R_n on $[0, \infty)$, g_k converges uniformly to f on $[0, \infty)$ and

$$\left\| \frac{1}{p_k} - \frac{1}{p^*} \right\| \geq \frac{7}{8}E \text{ where } E = \left\| f - \frac{1}{p^*} \right\| > 0.$$

Proof. Select $\beta > 0$ such that $p^*(x) \geq \frac{8}{E}$, $f(x) \leq \frac{E}{8}$ and $p^*(x)$ is monotone increasing for $x \geq \beta$. For $k \geq \beta$, define

$$p_k(x, t) = t + (p^*(x) - t) \left[\left(\frac{x}{k} - 1 \right)^2 + \frac{t}{p^*(k) - t} \right], \quad 0 \leq t \leq \frac{1}{E}, \quad x \geq 0.$$

Note that $p_k(x, t)$ is continuous on $[0, \infty) \times [0, \frac{1}{E}]$. Define

$h(t) = \min\{p_k(x, t) : x \in [\beta, 2k]\}$ and observe that h is a continuous

function of t , $0 \leq t \leq \frac{1}{E}$. In addition, $h(0) = \min\{p^*(x) \left(\frac{x}{k} - 1 \right)^2 : x \in [\beta, 2k]\} = 0$

as $k \geq \beta$ and that $h(\frac{1}{E}) = \min\{\frac{1}{E} + (p^*(x) - \frac{1}{E}) \left[\left(\frac{x}{k} - 1 \right)^2 + \frac{1}{E p^*(k) - 1} \right] : x \in [\beta, 2k]\} > \frac{1}{E}$

as $p^*(x) > \frac{1}{E}$ on $[\beta, 2k]$. Select $e_k \in (0, \frac{1}{E})$ so that $h(e_k) = \frac{1}{E}$. Thus,

$p_k(x, e_k) \geq \frac{1}{E} > 0$ for $x \in [\beta, 2k]$. Observe that $p_k(x, e_k)$ converges

uniformly to $p^*(x)$ on $[0, \beta]$ as $k \rightarrow \infty$ since $0 < e_k \leq \frac{1}{E}$, $\frac{e_k}{p^*(k) - e_k} \rightarrow 0$ and

$\left(\frac{x}{k} - 1 \right)^2$ converges uniformly to 1 on $[0, \beta]$ as $k \rightarrow \infty$.

Next, let

$$\eta = \min \left(\min_{x \in [0, \beta]} f(x), \min_{x \in [0, \beta]} \frac{1}{p^*(x)}, \frac{E}{4} \right) > 0.$$

Select $\mu \geq \beta$ such that $k \geq \mu$ implies that $\sqrt{k} \geq \beta$, $k > 1$,

$\max \left\{ \left| \frac{1}{p_k(x, e_k)} - \frac{1}{p^*(x)} \right| : x \in [0, \beta] \right\} \leq \frac{\eta}{2}$. Thus, for $k \geq \mu$, $\frac{1}{p_k(x, e_k)} \geq \frac{\eta}{2} > 0$,

for all $x \in [0, \beta]$. This implies that for $k \geq \mu$, $\frac{1}{p_k(x, e_k)}$ is positive

and converges uniformly to $\frac{1}{p^*(x)}$ on $[0, \beta]$. In addition, for $k \geq \mu$ and

$x \in [\beta, \sqrt{k}]$ we have $p_k(x, e_k) \geq e_k + (p^*(x) - e_k) \left[\left(\frac{1}{\sqrt{k}} - 1 \right)^2 \right]$

$\geq e_k + \frac{1}{2}(p^*(x) - e_k) \geq \frac{1}{2}p^*(x) \geq \frac{4}{E}$ as $p^*(x) \geq \frac{8}{E}$ for $x \geq \beta$. Since

$\max\{\frac{1}{p_k(x, e_k)} : x \in [\beta, 2k]\} = E$ we have that if $t_k \in [\beta, 2k]$ is such that $\frac{1}{p_k(t_k, e_k)} = E$ then $t_k > \sqrt{k}$ for $k \geq \mu$.

Next, note that for $x \geq k \geq \mu$, $p_k(x, e_k)$ is a monotone increasing function of x and that $p_k(2k, e_k) = e_k + (p^*(2k) - e_k)(1 + \frac{e_k}{p^*(2k) - e_k}) \geq p^*(2k) \geq \frac{8}{E}$. Thus, $\frac{1}{p_k(x, e_k)} \leq \frac{E}{8}$ for $x \geq 2k$. Summarizing, we have shown that $\frac{1}{p_k(x, e_k)} \leq \frac{E}{4}$ for $x \in [0, \sqrt{k}]$, $\frac{1}{p_k(x, e_k)} \leq \frac{E}{8}$ for $x \geq 2k$ and $\frac{1}{p_k(x, e_k)} \leq E$ for $x \in [\sqrt{k}, 2k]$ with $t_k \in [\sqrt{k}, 2k]$ a point at which the value E is attained.

Next, define α_k by $E - \alpha_k = \max\{(\frac{1}{p_k(x, e_k)} - f(x)) : x \in [\beta, 2k]\}$.

Since $f(x) \leq \frac{E}{8}$ for $x \geq \beta$ and $\frac{1}{p_k(t_k, e_k)} = E$ we have that $E - \alpha_k \geq E - f(t_k) \geq \frac{7E}{8}$ implying that $\frac{E}{8} \geq f(t_k) \geq \alpha_k$. Let $y_k \in [\beta, 2k]$ be such that

$\frac{1}{p_k(y_k, e_k)} - f(y_k) = E - \alpha_k$ for each $k \geq \mu$. Since $\frac{1}{p_k(x, e_k)} \leq \frac{E}{8}$ for

$x \in [\beta, \sqrt{k}]$ we have that $y_k \in [\sqrt{k}, 2k]$. Also, since $f(t_k) \rightarrow 0$ as $k \rightarrow \infty$

(as $t_k \rightarrow \infty$) it follows that $\alpha_k \rightarrow 0$ as $k \rightarrow \infty$. Noting that $f(x) \leq \frac{E}{8}$ for

$x \in [\beta, \infty)$ and that $\frac{1}{p_k(x, e_k)} \leq \frac{E}{8}$ for $x \geq 2k$ we have that $|f(x) - \frac{1}{p_k(x, e_k)}|$

$\leq E - \alpha_k$ for $x \in [\beta, \infty)$ and $k \geq \mu$. Also, since $\frac{1}{p_k(t_k, e_k)} = E$ and $\frac{1}{p^*(t_k)} \leq \frac{E}{8}$

we have that $\|\frac{1}{p_k} - \frac{1}{p^*}\| \geq |\frac{1}{p_k(t_k, e_k)} - \frac{1}{p^*(t_k)}| \geq \frac{7E}{8}$.

Now define g_k by (for $k \geq u$)

$$g_k(x) = \begin{cases} f(x) + \frac{1}{p_k(x, e_k)} - \frac{1}{p^*(x)}, & x \in [0, \beta], \left| f(x) - \frac{1}{p^*(x)} \right| \leq E - \alpha_k \\ E - \alpha_k + \frac{1}{p_k(x, e_k)}, & x \in [0, \beta], f(x) - \frac{1}{p^*(x)} > E - \alpha_k \\ -E + \alpha_k + \frac{1}{p_k(x, e_k)}, & x \in [0, \beta], f(x) - \frac{1}{p^*(x)} < -E + \alpha_k \\ f(x), & x \geq \beta + \frac{1}{k} \\ \text{linear on } [\beta, \beta + \frac{1}{k}] \text{ with endpoint values } f(\beta) + \frac{1}{p_k(\beta, e_k)} - \frac{1}{p^*(\beta)} \\ \text{and } f(\beta + \frac{1}{k}). \end{cases}$$

Observe that $g_k(x) > 0$ for all $x \geq 0$. Indeed, for $x \in [0, \beta]$ with

$$\left| f(x) - \frac{1}{p^*(x)} \right| \leq E - \alpha_k \text{ we have that } g_k = f(x) + \frac{1}{p_k(x, e_k)} - \frac{1}{p^*(x)} \geq f(x) - \frac{n}{2} \geq \frac{n}{2} > 0. \text{ For } x \in [0, \beta] \text{ with } f(x) - \frac{1}{p^*(x)} > E - \alpha_k, g_k(x) = E - \alpha_k + \frac{1}{p_k(x, e_k)} \geq \frac{7E}{8} + \frac{1}{p_k(x, e_k)} > 0 \text{ and for } x \in [0, \beta] \text{ with } f(x) - \frac{1}{p^*(x)} < -E + \alpha_k,$$

$$g_k(x) = -E + \alpha_k + \frac{1}{p_k(x, e_k)} = -E + \alpha_k + \frac{1}{p^*(x)} + \frac{1}{p_k(x, e_k)} - \frac{1}{p^*(x)} \geq f(x) - \frac{n}{2} \geq \frac{n}{2} > 0. \text{ Since } f(\beta) + \frac{1}{p_k(\beta, e_k)} - \frac{1}{p^*(\beta)} \geq f(\beta) - \frac{n}{2} \geq \frac{n}{2} > 0 \text{ and } f(\beta + \frac{1}{k}) > 0$$

we have that $g_k(x) > 0$ on $[\beta, \beta + \frac{1}{k}]$ and finally g_k is positive on $[\beta + \frac{1}{k}, \infty)$

as f is. To see that $g_k(x)$ is continuous on $[0, \infty)$ one must only check on $[0, \beta]$ as for $x > \beta$ it is clearly continuous. However, on $[0, \beta]$, $g_k(x)$ is simply the truncation of $f(x) - \frac{1}{p^*(x)}$ to the range $[-E + \alpha_k, E - \alpha_k]$ plus the continuous function $\frac{1}{p_k(x, e_k)}$ showing that $g_k \in C_0^+[0, \infty)$.

Next, let us consider $\left|g_k(x) - \frac{1}{p_k(x, e_k)}\right|$. Note that by construction $\left|g_k(x) - \frac{1}{p_k(x, e_k)}\right| \leq E - \alpha_k$ for $x \in [0, \beta]$ and that, if $\{x_i\}_{i=0}^n$ with $x_0 < x_1 < \dots < x_n$ is a nonstandard alternating set for $f - \frac{1}{p^*}$ then we must have that $x_n < \beta$ and

$$g_k(x_i) - \frac{1}{p_k(x_i, e_k)} = \operatorname{sgn}(f(x_i) - \frac{1}{p^*(x_i)})(E - \alpha_k) = (-1)^{n-i}(E - \alpha_k).$$

Next, on $[\beta, \beta + \frac{1}{k}]$ we have that $f(\beta) \leq \frac{E}{8}$, $f(\beta + \frac{1}{k}) \leq \frac{E}{8}$ and $\left|\frac{1}{p_k(\beta, e_k)} - \frac{1}{p^*(\beta)}\right| \leq \frac{E}{8}$ so that $g_k(\beta) \leq \frac{E}{4}$ and $g_k(\beta + \frac{1}{k}) \leq \frac{E}{8}$. Thus, $g_k(x) \leq \frac{E}{4}$ on $[\beta, \beta + \frac{1}{k}]$. Also, recall that $\frac{1}{p_k(x, e_k)} \leq \frac{E}{4}$ on $[0, \sqrt{k}]$ so that $\left|g_k(x) - \frac{1}{p_k(x, e_k)}\right| \leq \frac{E}{4}$ on $[\beta, \beta + \frac{1}{k}]$. Finally, we noted earlier that $\left|f(x) - \frac{1}{p_k(x, e_k)}\right| \leq E - \alpha_k$ on $[\beta, \infty)$ so that $\left|g_k(x) - \frac{1}{p_k(x, e_k)}\right| \leq E - \alpha_k$ on $[0, \infty)$. Since there exists $y_k \in [\sqrt{k}, 2k]$ at which $f(y_k) - \frac{1}{p_k(y_k, e_k)} = -(E - \alpha_k)$ we have that $g_k - \frac{1}{p_k}$ possesses a standard alternating set at the points $x_0 < x_1 < \dots < x_n < y_k$ and thus $\frac{1}{p_k}$ is the best approximation to g_k from R_n on $[0, \infty)$. Finally, it is a straightforward argument to prove that g_k converges uniformly to f . Thus, once again reindexing the sequence $\{\frac{1}{p_k}\}_{k=\mu}^\infty$ gives the desired result. I

Next, we wish to show that if $f \in C_0^+[0, \infty)$ has $\frac{1}{p^*}$ as its best approximation from R_n with $\partial p^* = n - 1$ then the best approximation operator is continuous. This we do in the following theorem.

THEOREM 4. Let $f \in C_0^+[0, \infty) \sim R_n$ and let $\frac{1}{p^*}$ be its best approximation from R_n on $[0, \infty)$ with $\alpha p^* = n - 1$. Then, the best approximation operator is continuous at f .

Proof. Let $\{g_k\}_{k=1}^\infty \subset C_0^+[0, \infty)$ with $g_k \rightarrow f$ uniformly on $[0, \infty)$. Further,

let $\frac{1}{p_k} \in R_n$ be the best approximation to g_k on $[0, \infty)$ for each k . Then,

we must prove that $\|\frac{1}{p_k} - \frac{1}{p^*}\| \rightarrow 0$ as $k \rightarrow \infty$. Let us first note that

$$\|g_k - \frac{1}{p_k}\| \leq \|g_k - \frac{1}{p^*}\| \text{ implying that } \limsup_{k \rightarrow \infty} \|g_k - \frac{1}{p_k}\| \leq \limsup_{k \rightarrow \infty} \|g_k - \frac{1}{p^*}\|$$

$$= \|f - \frac{1}{p^*}\| = E. \text{ Also, } E = \|f - \frac{1}{p^*}\| \leq \|f - \frac{1}{p_k}\| \leq \|f - g_k\| + \|g_k - \frac{1}{p_k}\|$$

implying that $E = \liminf_{k \rightarrow \infty} (E - \|f - g_k\|) \leq \liminf_{k \rightarrow \infty} \|g_k - \frac{1}{p_k}\|$. Combining

these results gives that $\lim_{k \rightarrow \infty} \|g_k - \frac{1}{p_k}\| = E$. In addition, since

$$E \leq \|f - \frac{1}{p_k}\| \leq \|f - g_k\| + \|g_k - \frac{1}{p_k}\| \text{ we also have that } \lim_{k \rightarrow \infty} \|f - \frac{1}{p_k}\| = E.$$

Next, fix $y \in [0, \infty)$ such that $f(y) = \max\{f(x) : x \in [0, \infty)\}$. Then

since a constant cannot be a best approximation to f from R_n on $[0, \infty)$ we must have that $2E < f(y)$. Select $\delta > 0$ such that for $x \in I = [y - \delta, y + \delta] \cap [0, \infty)$ we must have $f(x) \geq \frac{1}{2}(2E + f(y)) > 2E$. Choose β such that $k \geq \beta$ implies that $\|f - \frac{1}{p_k}\| \leq \frac{3}{2}E$. Then for $k \geq \beta$ and $x \in I$, we have that

$$0 < m = 2E - \frac{3}{2}E \leq f(x) - \frac{3}{2}E \leq \frac{1}{p_k(x)} \leq f(x) + \frac{3}{2}E \leq \|f\| + \frac{3}{2}E = M.$$

In addition, observe that the inequality $\frac{1}{p_k(x)} \leq M$ holds for all $x \in [0, \infty)$

and $k \geq \beta$. Let $\{p_v\}$ be a subsequence of $\{p_k\}$. Then, since $\frac{1}{M} \leq p_v(x) \leq \frac{1}{m}$

for all $x \in I$, there exists a subsequence $\{p_\mu\}$ of $\{p_\nu\}$ such that p_μ converges uniformly to some $\bar{p} \in R_n$ on I . This implies that the coefficients of p_μ converge to the coefficients of $\bar{p}$ which in turn implies that for each $x \in [0, \infty)$, $p_\mu(x) \rightarrow \bar{p}(x)$. Thus, we must have $\frac{1}{M} \leq \bar{p}(x) \leq \frac{1}{m}$ on I and $\frac{1}{M} \leq \bar{p}(x)$ on $[0, \infty)$. This last inequality shows that $\bar{p} \in R_n$. Furthermore, for $x \in [0, \infty)$ fixed, $\left| f(x) - \frac{1}{\bar{p}(x)} \right|$

$$= \lim_{n \rightarrow \infty} \left| f(x) - \frac{1}{p_\mu(x)} \right| \leq \lim_{n \rightarrow \infty} \left\| f - \frac{1}{p_\mu} \right\| = E. \text{ Thus, } \left\| f - \frac{1}{\bar{p}} \right\| \leq E \text{ implying}$$

that $\bar{p} \equiv p^*$ by the uniqueness of best approximations from R_n . Since this is true for any subsequence $\{p_\nu\}$ of $\{p_k\}$ we must have that this is also true for the full sequence $\{p_k\}$. That is, that p_k converges uniformly to p^* on I and pointwise on $[0, \infty)$. To complete this argument we must prove that $\frac{1}{p_k}$ converges uniformly to $\frac{1}{p^*}$ on $[0, \infty)$.

From the above discussion we have that $\frac{1}{p_k}$ converges pointwise to $\frac{1}{p^*}$ on $[0, \infty)$ and, in fact, on any fixed closed interval $[0, \alpha]$, $\alpha > 0$, $\frac{1}{p_k}$ converges uniformly to p^* (due to the coefficient convergence).

In order to establish this final fact, we must examine the coefficient convergence in more detail. Thus, let $p^*(x) = a_{n-1}^* x^{n-1} + \dots + a_0^*$ with $a_{n-1}^* > 0$ (here we are using our hypothesis that $\deg p^* = n - 1$ and $p^* \in R_n$) and let $p_k(x) = a_n^k x^n + \dots + a_0^k$ where we know that the leading nonzero coefficient of p_k must be positive. In addition, we have that $a_j^k \rightarrow a_j^*$ as $k \rightarrow \infty$ for $j = 0, 1, \dots, n$ where $a_n^* = 0$. Thus, there exists $\gamma \geq \beta$ such that $k \geq \gamma$ implies that $a_{n-1}^k \geq \frac{a_{n-1}^*}{2} > 0$ and $|a_j^k - a_j^*| \leq 1$ for $j = 0, \dots, n - 2$. Thus, given $\epsilon > 0$ there exists $\delta > 0$ such that

$$\hat{p}(x) = \frac{a_{n-1}^*}{2}x^{n-1} + (a_{n-2}^* - 1)x^{n-2} + \dots + (a_0^* - 1) \geq \frac{2}{\epsilon}. \text{ Since } p_k(x) \geq \hat{p}(x)$$

for $k \geq \gamma$ and $p^*(x) > \hat{p}(x)$ for all $x \geq \delta$ we have that

$$\left| \frac{1}{p_k(x)} - \frac{1}{p^*(x)} \right| \leq \left| \frac{1}{p_k(x)} \right| + \left| \frac{1}{p^*(x)} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

for $k \geq \gamma$ and $x \geq \delta$. On $[0, \delta]$ we have that $\frac{1}{p_k}$ converges uniformly to $\frac{1}{p^*}$.

Thus, we may select $K \geq \gamma$ such that $k \geq K$ implies $\left| \frac{1}{p_k(x)} - \frac{1}{p^*(x)} \right| < \epsilon$ for

all $x \in [0, \delta]$. Hence, for $k \geq K$ we have that $\left\| \frac{1}{p_k} - \frac{1}{p^*} \right\| < \epsilon$ implying

the desired result.

Concluding Remarks

Observe that the question of whether or not a strong uniqueness result holds for the case that $f \in C_0^+[0, \infty)$ with its best approximation $\frac{1}{p^*}$ from R_n satisfying $ap^* = n-1$ remains open. Likewise, the question of Lipschitz continuity of the best approximation operator remains open in this case.

A second item of interest is that in ordinary rational approximation on a finite interval, nonstandard (i.e., fewer) alternation due to degeneracy of the best approximation may be unimportant as the set of f with degenerate best approximations is nowhere dense [5, 7]. If the corresponding result that $\{f: \text{the best approximation } \frac{1}{p^*} \in R_n \text{ has } ap^* < n\}$ was nowhere dense then we could expect to be able to usually employ the simpler theory of [8] for this problem. However, the continuity result for degree $n-1$ implies that every f with nonstandard alternation and best approximation $\frac{1}{p^*} \in R_n$ with $ap^* = n-1$ has all g sufficiently close with nonstandard alternation and best approximations of degree $n-1$. In

this regard, an interesting question is to characterize those f for which nonstandard alternation will occur. Some initial results in this direction have been obtained by the second author and D. Leeming.

REFERENCES

1. H.-P. Blatt, Rationale Approximation auf $[0, \infty)$, ZAMM, 53 (1973), 182-183.
2. H.-P. Blatt, Rationale Tschebyscheff-Approximation über unbeschränkten Intervallen, Habilitationsschrift der Universität, Erlangen-Nürnberg, 1974.
3. D. Brink, Tchebycheff approximation by reciprocals of polynomials on $[0, \infty)$, Ph.D. Thesis, Michigan State University, 1972.
4. D. Brink and G. D. Taylor, Chebyshev approximation by reciprocals of polynomials on $[0, \infty)$, J. Approximation Theory, 16 (1976), 142-148.
5. E. W. Cheney and H. L. Loeb, Generalized rational approximation, SIAM J. Numer. Anal., Ser. B1 (1964), 11-25.
6. W. J. Cody, G. Meinardus and R. S. Varga, Chebyshev rational approximation to e^{-x} in $[0, +\infty)$ and applications to heat-conduction problems, J. Approx. Theory, 2 (1969), 50-65.
7. C. B. Dunham, Alternating Chebyshev approximation, TAMS, 178 (1973), 95-109.
8. C. B. Dunham, Chebyshev approximation by interpolating rationals on $[\alpha, \infty)$, Math. Comput., 29 (1975), 549-551.
9. E. H. Kaufman, Jr. and G. D. Taylor, Uniform approximation with rational functions having negative poles, J. Approximation Theory, to appear.
10. A. R. Reddy and Oved Shisha, Rational approximation on the positive real axis - a survey, preprint.
11. E. B. Saff, R. S. Varga and W.-C. Ni, Geometric convergence of rational approximations to e^{-z} in infinite sectors, preprint.
12. E. B. Saff, A. Schönhage and R. S. Varga, Geometric convergence to e^{-z} by rational functions with real poles, Numer. Math., 25 (1976), 307-322.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR-79-0856	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CONTINUITY OF BEST RECIPROCAL POLYNOMIAL APPROXIMATION ON $[0, \infty)$	5. TYPE OF REPORT & PERIOD COVERED Interim $\neq$ Rept.	
6. AUTHOR(s) Charles B. Dunham and G. D. Taylor	7. CONTRACT OR GRANT NUMBER(s) AFOSR-76-2878 NSF-MC578-05847	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Colorado State University Department of Mathematics Fort Collins, CO 80523	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A3	
10. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, D. C. 20332	11. REPORT DATE Apr 1979	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) $[12] 15p.$	13. NUMBER OF PAGES 14	
	14. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Reciprocal polynomial approximation on $[0, \infty)$, constrained rational approximation.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Continuity of the best approximation operator is studied for approximating continuous positive functions tending to zero as $x \rightarrow \infty$ by reciprocals of polynomials on $[0, \infty)$ is studied. Both negative and positive results are obtained. $\nearrow a$ infinity. approaches infinity		