

AD-A075 834

OKLAHOMA UNIV NORMAN DEPT OF CHEMISTRY

F/G 7/3

THE X-RAY CRYSTAL STRUCTURE OF TRIMETHYLTIN (IV) AZIDE. A UNIQUE--ETC(U)

SEP 79 D CUNNINGHAM , M B HOSSAIN

N00014-77-C-0432

UNCLASSIFIED TR-12

NL

| OF |

AD  
A075834



END  
DATE  
FILMED  
11-79  
DDC

AD A 075834

DDC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

LEVEL II

REPORT DOCUMENTATION PAGE		REVISIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 12	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The X-Ray Crystal Structure of Trimethyltin(IV) Azide. A Unique One-Dimensional $[(CH_3)_3Sn]^{δ+}[(CH_3)_3Sn(N_3)_2]^{δ-}$ Polymer. <i>Delta</i>		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Desmond/Cunningham, M.B./Hossain, K.C./Molloy, D. van der/Helm and J. J./Zuckerman		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Oklahoma Department of Chemistry Norman, Oklahoma 73019		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0432
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of Navy Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-636
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <i>12 12</i>		12. REPORT DATE 24 September 1979
		13. NUMBER OF PAGES 8
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited <i>9 Technical rept's</i>		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for Publication in J. Chem. Soc., Chem. Commun.		DDC RECEIVED OCT 30 1979 B
18. SUPPLEMENTARY NOTES <i>14 TR-12</i>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Trimethyltin Azide, Organotins, Azides, X-ray, Crystal and Molecular Structure, Low Temperature X-ray Study, Polymer, One-Dimensional Polymer, Linear Polymer, Associated Structure, Bridging Azide Groups, $\alpha$ -Nitrogen Azide Bridging.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Our 1978 review of the tin structural literature (ref. 11) reveals many examples of axially-most electronegative, trigonal bipyramidal triorganotin moieties in associated polymer chains, but the structure of trimethyltin azide is unique in consisting of $SnC_3$ moieties in alternating $[(CH_3)_3Sn]^{δ+}$ cationic and $[N_3(CH_3)_3Sn]^{δ-}$ anionic environments in a one-dimensional array with the tin atoms bridged at unequal distances by the $\alpha$ -nitrogen atoms of the azide group at which site the chains are bent. <i>alpha</i> <i>(delta+)</i>		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)*(delta-) 403 980*

OFFICE OF NAVAL RESEARCH

Contract N00014-77-C-0432

Task No. NR 053-636

TECHNICAL REPORT NO. 12

The X-Ray Crystal Structure of Trimethyltin(IV) Azide. A Unique  
One-Dimensional  $\{[(\text{CH}_3)_3\text{Sn}]^{\delta+}[(\text{CH}_3)_3\text{Sn}(\text{N}_3)_2]^{\delta-}\}_n$  Polymer

by

Desmond Cunningham, M. B. Hossain, K. C. Molloy,  
D. van der Helm and J. J. Zuckerman

Prepared for Publication

in

J. Chem. Soc., Chem. Commun.

University of Oklahoma  
Department of Chemistry  
Norman, Oklahoma 73019

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

\*This document has been approved for public release  
and sale; its distribution is unlimited

79 10 29 139

# STATEMENT

Our 1978 review of the tin structural literature (ref. 11) reveals many examples of axially-most-electronegative, trigonal bipyramidal triorganotin moieties in associated polymer chains, but the structure of trimethyltin azide is unique in consisting of  $\text{SnC}_3$  moieties in alternating  $[(\text{CH}_3)_3\text{Sn}]^{\delta+}$  cationic and  $[\text{N}_3(\text{CH}_3)_3\text{SnN}_3]^{\delta-}$  anionic environments in a one-dimensional array with the tin atoms bridged at unequal distances by the  $\alpha$ -nitrogen atoms of the azide group at which site the chains are bent.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist	AIL and/or SPECIAL
A	

The X-Ray Crystal Structure of Trimethyltin(IV) Azide at 138K. A Unique,  
One-Dimensional  $\{[(\text{CH}_3)_3\text{Sn}]^{\delta+}[(\text{CH}_3)_3\text{Sn}(\text{N}_3)_2]^{\delta-}\}_n$  Polymer

By Desmond Cunningham<sup>+</sup>, M. B. Hossain, K. C. Molloy, D. van der Helm\* and J. J. Zuckerman\*

(Department of Chemistry, University of Oklahoma, Norman, OK  
73019 USA)

Summary

The structure of trimethyltin(IV) azide, as determined at 138K by X-ray diffraction, consists of one-dimensional polymeric chains in which planar  $\text{SnC}_3$  moieties in alternating  $[(\text{CH}_3)_3\text{Sn}]^{\delta+}$  and  $[\text{N}_3(\text{CH}_3)_3\text{SnN}_3]^{\delta-}$  environments are bridged through the  $\alpha$ -nitrogen atoms of the azide group.

Organic and inorganic derivatives of the azide group exhibit a variety of properties which have made them the subject of continuing interest. While organic azides are of importance in the drug and dye industry, it is among the metal azides that the most subtle variations of bonding and stability occur, thus placing this class of compounds both at the forefront of research in the explosives industry and as model systems for investigation by the structural chemist<sup>1</sup>.

Triorganometal(IV) azides are unique insofar as they have relatively high thermal stability and appear to be shock insensitive<sup>2</sup>. Trimethylsilyl- and germyl azides are liquids at room

---

<sup>+</sup>On leave from the Department of Chemistry, University College, Galway, Ireland.

\*Author(s) to whom correspondence should be addressed.

temperature and apparently monomeric, while the corresponding azide derivatives of tin and lead are relatively high melting solids ( $121^\circ$  and  $>200^\circ\text{C}$ , decomp., respectively).<sup>2</sup> Although infrared and Mössbauer studies strongly suggest a polymeric structure for trimethyltin azide,<sup>3-8</sup> the intriguing question of how the azide bridging actually occurs, i.e., through the  $\alpha$ -nitrogen only or, as has been suggested,<sup>3,8,9</sup> through both the  $\alpha$ - and  $\gamma$ -nitrogens, remains uncertain. The situation has prompted us to make a preliminary report of the crystal structure of this complex.

Suitable crystals were obtained by slow vacuum sublimation at ca.  $70^\circ\text{C}$ . Trimethyltin(IV) azide crystallizes in the orthorhombic space group  $Pnc2$ ,  $a = 8.684(3)$ ,  $b = 11.727(4)$ , and  $c = 6.612(1)\text{\AA}$ ,  $Z = 4$ ,  $V = 673.35\text{\AA}^3$ ,  $D_{\text{calc}} = 2.03\text{ g cc}^{-3}$ . Diffraction data were collected on an Enraf-Nonius CAD-4 automatic diffractometer at  $138\pm 2\text{K}$  and were corrected for Lorentz, polarization and absorption effects.<sup>10</sup> The structure was solved by Patterson and difference Fourier methods. The tin and non-hydrogen light atoms were refined anisotropically and isotropically, respectively, to an R-value of 5.26% for 922 unique reflections or 6.77% for all data (1249 reflections) using a block-diagonal least squares program.

The structure is shown in Figure 1 along with pertinent

bond lengths and angles. The chain polymeric nature of the complex is confirmed, but there are two other noteworthy features associated with the structure. Firstly, bridging is through the  $\alpha$ -nitrogen of the azide group at which the chains are bent with an Sn(1)-N(1)-Sn(2) angle of  $133.2(4)^\circ$  in an analogous manner to that found in similar species,<sup>11</sup> especially trimethyltin chloride<sup>12</sup> and methoxide.<sup>13</sup> The  $\text{SnC}_3$  moieties are eclipsed in the former, but staggered as in our azide in the latter. In trimethyl-<sup>14</sup> and triphenyltin<sup>15</sup> isothiocyanate, on the other hand, the bridging occurs through both nitrogen and sulfur atoms.

Secondly, as a direct consequence of the space group symmetry there are two non-equivalent tin sites, alternate tin atoms making two short [2.290(6)Å] or two long [2.441(6)Å] bonds to nitrogen (see Figure 1). The geometry at each tin is trigonal bipyramidal with the  $\text{SnC}_3$  moieties planar, and the more electronegative nitrogen atoms occupying apical positions with an N(1)-Sn-N(1') angle of  $175.8(3)^\circ$  associated with the short Sn-N bonds, and an angle of  $178.4(3)^\circ$  associated with the long Sn-N bonds. This the only example of an associated organotin polymer in which the chains consist of alternating tin environments.<sup>11</sup> This arrangement is probably best described as a combination of trimethyltin(IV) cations and trimethyltin(IV) diazide anions,  $\{[(\text{CH}_3)_3\text{Sn}]^{\delta+} [\text{N}_3(\text{CH}_3)_3\text{SnN}_3]^{\delta-}\}_n$ , in a one-dimensional array. The tin atoms are not sufficiently different to be distinguished by <sup>119m</sup>Sn Mössbauer

spectroscopy.<sup>16</sup> This and other organotin(IV) azides are currently being subjected to solid state tin-119 nmr study.<sup>17</sup>

Acknowledgement. Our work is supported by the U.S. Office of Naval Research (J.J.Z.) and the National Cancer Institute, U.S. D.H.E.W.  
\* through Grant No. CA-17562 (D.v.d.H.).


# References and Notes

- (1) See, for example, Energetic Materials, Vol. 1, ed. by D. H. Fair and R. F. Walker, Plenum Press, New York (1977).
- (2) See, for example: J. S. Thayer and R. West, Advan. Organometal. Chem., 1967, 5, 149; J. S. Thayer, Organometal. Chem. Revs., 1966, 1, 157; M. F. Lappert and H. Pyszora, Advan. Inorg. Chem. Radiochem., 1966, 9, 133.
- (3) J. G. A. Luijten, M. J. Janssen and G. J. M. van der Kerk, Rec. Trav. Chim., 1962, 81, 202.
- (4) G. J. M. van der Kerk, J. G. A. Luijten and M. J. Janssen, Chimia, 1962, 14, 10.
- (5) J. S. Thayer and R. West, Inorg. Chem., 1964, 3, 889.
- (6) J. S. Thayer and D. P. Strommen, J. Organometal. Chem., 1966, 5, 383.
- (7) H.-S. Cheng and R. H. Herber, Inorg. Chem., 1970, 9, 1686.
- (8) J. Müller, Z. Naturforsch., 1979, 34b, 536.
- (9) Comprehensive Inorganic Chemistry, ed. by J. C. Bailar, H. J. Emeléus, R. Nyholm and A. F. Trotman-Dickenson. Pergamon Press, London (1975), Vol. 2, 93.
- (10) Room temperature Weissenberg photographs show that reflections of the type  $hkl$ ,  $h = 2n+1$  are very weak and diffuse, indicative of some disorder in the crystal at ambient temperature which is apparently absent at  $138 \pm 2$  K.
- (11) J. A. Zubieta and J. J. Zuckerman, Prog. Inorg. Chem., 1978, 24, 251.
- (12) M. B. Hossain, J. L. Lefferts, K. C. Molloy, D. van der Helm and J. J. Zuckerman, Inorg. Chim. Acta, 1979, 36, L409.

- (13) A. M. Domingos and G. M. Sheldrick, Acta Crystallogr., 1974, B30, 519.
- (14) R. A. Forder and G. M. Sheldrick, J. Chem. Soc., Chem. Commun., 1969, 1125, J. Organomet. Chem., 1970, 21, 115.
- (15) A. M. Domingos and G. M. Sheldrick, J. Organometal. Chem., 1974, 67, 257.
- (16) Even the tin atoms in the five- and six-coordinated dimethyltin(IV) double salt  $[(CH_3)_2SnCl \cdot terpy]^+$   $[(CH_3)_2SnCl_3]^-$  are barely distinguishable by Mössbauer spectroscopy [N. W. G. Debye, E. Rosenberg and J. J. Zuckerman, J. Am. Chem. Soc., 1968, 90, 3234].
- (17) G. E. Maciel and J. J. Zuckerman, unpublished results.

Figure Caption

Figure 1. A segment of the polymeric structure of trimethyltin azide, showing atomic labelling. Atoms related by a two-fold symmetry along  $\underline{c}$  are primed. Atoms in special positions are: Sn(1), C(1)[0,0,z; Wykoff notation:c]; Sn(2), C(4)[1/2,0,z;b]. The remaining atoms are in general positions.



#### STATEMENT

Our 1978 review of the tin structural literature (ref. 11) reveals many examples of axially-most-electronegative, trigonal bipyramidal triorganotin moieties in associated polymer chains, but the structure of trimethyltin azide is unique in consisting of  $\text{SnC}_3$  moieties in alternating  $[(\text{CH}_3)_3\text{Sn}]^{\delta+}$  cationic and  $[\text{N}_3(\text{CH}_3)_3\text{SnN}_3]^{\delta-}$  anionic environments in a one-dimensional array with the tin atoms bridged at unequal distances by the  $\alpha$ -nitrogen atoms of the azide group at which site the chains are bent.

