

OFFICE OF NAVAL RESEARCH Contract N00014-76-C-0826 Task No. NR 056-625 1.125 TECHNICAL REPORT NO. 81-1 AD A 09542 ί, REACTIONS OF METAL-METAL MULTIPLE BONDS. 8. EORMING MO-MO QUADRUPLE BONDS BY REDUCTIVE ELIMINATION (ALKYL GROUP DISPROPORTIONATION) tin leid IN THE REACTIONS OF 1,2-Mo2R2 (NMe2) (COMPOUNDS (MAM) WITH CARBON DIOXIDE AND 1,3-DIARYLTRIAZENES . bν M.H. Chisholm, K. Folting, D.A. Haitko and J.C. Huffman Prepared for Publication in the Journal of the American Chemical Society FEB 2 3 1981 Department of Chemistry Indiana University Bloomington, IN 47405 January 30,7 1981 FILE COP Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited. 1-16 (9: 81 2 23 04 8

<b>REPORT DOCUMENTATION PAGE</b>	READ INSTRUCTIONS BEFORE COMPLETING FORM
REPORT NUMBER 2. GOVT ACCESSION N	0. 3. RECIPIENT'S CATALOG NUMBER
AD - AC 954:	· 1
TITLE (and Subjitie) Reactions of Metal-Metal Multiple	5. TYPE OF REPORT & PERIOD COVERED
onds. 8. Forming Mo-Mo Quadruple Bonds by Reduc-	
ive Elimination (Alkyl Group Disproportionation)	Technical Report 1981
n the Reactions of 1, $R_2(NMe_2)$ , Compounds (M	ENd. PERFORMING ORG. REPORT NUMBER
vith Carbon Dioxide and 1, -Diaryltriazenes.	TR-81-1
AUTHOR()	8. CONTRACT OR GRANT NUMBER(.)
I.H. Chisholm, K. Folting, D.A. Haitko and	
.C. Huffman	N00014-76-C-0826
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Department of Gnemistry Indiana University	
Riognation IN 47405	
CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research	I IADUARY 30 1001
Department of the Navy	January JU, 1701
	34
4. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	15. SECURITY CLASS. (of this report)
	154. DECLASSIFICATION/DOWNGRADING
This document has been approved for public rele is unlimited.	ase and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, 11 different i	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the ebstrect entered in Block 20, 11 different in	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract ontered in Block 20, 11 different i	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, 11 different i	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the ebetrect entered in Block 20, 11 different i 8. SUPPLEMENTARY NOTES	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the ebstrect entered in Block 20, 11 different 8. SUPPLEMENTARY NOTES	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the ebetrect entered in Block 20, 11 different in 8. SUPPLEMENTARY NOTES	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different i 8. SUPPLEMENTARY NOTES	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different i 8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse side if necessary and identify by block number molybdenum metal metal bond quadruple bond re	rom Report)
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different in 8. SUPPLEMENTARY NOTES 1. KEY WORDS (Continue on reverse side if necessary and identify by block number molybdenum, metal-metal bond, quadruple bond, re carbon dioxide, dimethylamide	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different in 8. SUPPLEMENTARY NOTES 1. KEY WORDS (Continue on reverse elde 11 necessary and identify by block number molybdenum, metal-metal bond, quadruple bond, re carbon dioxide, dimethylamide	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different in 8. SUPPLEMENTARY NOTES 1. KEY WORDS (Continue on reverse elde 11 necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re carbon dioxide, dimethylamide	ease and sale; its distributi
This document has been approved for public rele is unlimited. 7. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different in 8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse olds 11 necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re carbon dioxide, dimethylamide	ease and sale; its distributi
This document has been approved for public rele is unlimited. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different in SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse oldo if necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re carbon dioxide, dimethylamide	ease and sale; its distributi
This document has been approved for public rele is unlimited. T. DISTRIBUTION STATEMENT (of the obstract ontered in Block 20, 11 different is S. SUPPLEMENTARY NOTES S. SUPPLEMENTARY NOTES Note: Supplementary note: Second of the second second dentify by block number nolybdenum, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Addition of CO, and 1,3-diaryltriazenes	ease and sale; its distributi
This document has been approved for public release is unlimited. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different is SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES Supplement, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Addition of CO <sub>2</sub> and 1,3-diaryltriazenes (M=M), where R = C <sub>0</sub> H <sub>c</sub> , CH <sub>0</sub> CD <sub>0</sub> , CD <sub>0</sub> CH <sub>0</sub> , CH(CH <sub>0</sub> )	ease and sale; its distributi from Report) eductive elimination, insertio -0 1,2-Mo <sub>2</sub> R <sub>2</sub> (NMe <sub>2</sub> ) <sub>4</sub> compound - and n-, sec- and tert-C, H <sub>o</sub>
This document has been approved for public release is unlimited. The DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different is Supplementary notes KEY WORDS (Continue on reverse olds if necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide ABSTRACT (Continue on reverse olds II necessary and identify by block number Addition of CO <sub>2</sub> and 1,3-diaryltriazenes (M=M), where $R = C_2H_5$ , $CH_2CD_3$ , $CD_2CH_3$ , $CH(CH_3)$ yield Mo-Mo quadruply bonded compounds Mo <sub>2</sub> (0,CN)	Pase and sale; its distributi from Report) rom Report) eductive elimination, insertic $r_{2}$ and n-, sec- and tert-C4Hg (e_1), and Mo <sub>2</sub> (ArN <sub>2</sub> Ar), respectively
This document has been approved for public release is unlimited. 7. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different is 8. SUPPLEMENTARY NOTES 1. KEY WORDS (Continue on reverse elde 11 necessary and identify by block number molybdenum, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide 2. ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Addition of CO <sub>2</sub> and 1,3-diaryltriazenes (M=M), where $R = C_2H_5$ , $CH_2CD_3$ , $CD_2CH_3$ , $CH(CH_3)$ yield Mo-Mo quadruply bonded compounds $Mo_2(O_2CN)$ tively, by the reductive elimination (alkyl	Prom Report) (rom Report) eductive elimination, insertion $_{2}$ on $1,2-Mo_{2}R_{2}(NMe_{2})_{4}$ compound $_{2}$ and $n-$ , sec- and tert-C <sub>4</sub> Hg $_{3}(e_{2})_{4}$ and $Mo_{2}(ArN_{3}Ar)_{4}$ , respectively group disproportionation) of
This document has been approved for public release is unlimited. DISTRIBUTION STATEMENT (of the obstract entered in Black 20, If different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide ABSTRACT (Continue on reverse side If necessary and identify by block number Addition of $CO_2$ and 1,3-diaryltriazenes (MEM), where $R = C_2H_5$ , $CH_2CD_3$ , $CD_2CH_3$ , $CH(CH_3$ yield Mo-Mo quadruply bonded compounds $Mo_2(O_2CN)$ tively, by the reductive elimination (alkyl alkene and alkane. The reactions of the label	Prove and sale; its distributi from Report) $(-0 \ 1, 2 - Mo_2R_2(NMe_2)_4 \ compound_2 \ and \ n-, \ sec- \ and \ tert-C_4Hg$ $(e_2)_4 \ and \ Mo_2(ArN_3Ar)_4, \ respective group \ disproportionation) \ old \ ethyl \ compounds \ show \ that the section \ the sect$
This document has been approved for public release This unlimited. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different is SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde 11 necessary and identify by block number nolybdenum, metal-metal bond, quadruple bond, re- carbon dioxide, dimethylamide ABSTRACT (Continue on reverse elde 11 necessary and identify by block number Addition of CO <sub>2</sub> and 1,3-diaryltriazenes (M=M), where R = C <sub>2</sub> H <sub>5</sub> , CH <sub>2</sub> CD <sub>3</sub> , CD <sub>2</sub> CH <sub>3</sub> , CH(CH <sub>3</sub> yield Mo-Mo quadruply bonded compounds Mo <sub>2</sub> (O <sub>2</sub> CN tively, by the reductive elimination (alkyl alkene and alkane. The reactions of the label	Prom Report) Trom Report) eductive elimination, insertion $_{2}$ and n-, sec- and tert-C <sub>4</sub> Hg $_{2}$ $_{3}$ Ar $_{4}$ , respectively $_{4}$ $_{1}$ $_{2}$ $_{2}$ $_{3}$ $_{4}$ $_{1}$ $_{2}$ $_{4}$ $_{1}$ $_{2}$ $_{2}$ $_{3}$ Ar $_{1}$ $_{4}$ , respectively $_{3}$ $_{4}$ $_{1}$ $_{1}$ $_{1}$ $_{2}$ $_{2}$ $_{3}$ $_{1}$ $_{2}$ $_{2}$ $_{3}$ $_{1}$ $_{2}$ $_{3}$ $_{1}$ $_{2}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{4}$ $_{2}$ $_{2}$ $_{3}$ $_{1}$ $_{2}$ $_{2}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{1}$ $_{3}$ $_{2}$ $_{3}$ $_{3}$ $_{1}$ $_{3}$ $_$

#### LUIRITY CLASSIFICATION OF THIS PAGE (Whon Date Entered)

the elimination of the hydrocarbons is intramolecular and that the  $\beta$ -hydroger atom of one ethyl group is transferred to the  $\alpha$ -carbon atom of the other. These reactions proceed through intermediates of the form  $Mo_2R_2(NMe_2)_2L_2$ , where  $L = O_2CNMe_2$  or  $ArN_3Ar$ . The compound  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ , where  $C_7H_8 =$ p-tolyl, has been isolated and structurally characterized: space group P21/a,  $\alpha = 16.503(4)$  Å, b = 23.116(7) Å, c = 11.318(3) Å,  $\beta = 88.15(1)^\circ$  and Z = 4. The molecule has virtual, but not crystallographically imposed  $C_2$  symmetry. The  $Mo \equiv Mo$  bond, 2.171(1) Å, is bridged by a cis-pair of triazenido ligands which afford sufficient flexibility to allow a non-eclipsed geometry. Each molybdenun atom is coordinated to three nitrogen atoms and a carbon atom which lie roughly in a plane. Pertinent averaged bond distances are Mo-C = 2.21(1) Å, Mo-N(dimethylamide) = 1.95(1) Å and Mo-N (triazenido) = 2.17(1) and 2.25(1) Å with the longer bond being trans to the  $Mo-NMe_2$  group. Plausible mechanisms for the reductive elimination sequence are presented and discussed in the light of related reactions.

Accession Tor MTIS GRA&I DIEC INB. Uppmionicand Justific time. By\_\_\_\_\_ Distribution/ Aveils Clivy Codes (1, 2) = 1

SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

### Introduction

The preparation<sup>2</sup> of an extensive series of  $1,2-M_2R_2(NMe_2)_4$  compounds (M=M), where M = Mo and W, and R = Me, Et, i-Pr, n-Bu, sec-Bu, tert-Bu, CH<sub>2</sub>CMe<sub>3</sub> and CH<sub>2</sub>SiMe<sub>3</sub>, provides the opportunity of studying the reactivity patterns of  $\sigma$ -alkyl groups<sup>3</sup> bonded to dimetal centers. Of particular note in the above series are the thermally stable compounds which contain  $\sigma$ -alkyl groups that have  $\beta$ -hydrogen atoms. A common thermal decomposition pathway in mononuclear transition metal complexes containing such ligands involves the so-called  $\beta$ -hydrogen elimination reaction: M-alkyl + M-H + alkene. For dial-kylmetal complexes, this is often the step which precedes reductive elimination by carbon-hydrogen bond formation, leading to the overall reaction shown in equation 1 below. This may be called reductive alkyl group disproportionation of di-n-butyl compounds of platinum(II),  $^4$  cobalt(III)<sup>5</sup> and titanium(IV).<sup>6</sup> It seemed reasonable to suppose that reductive alkyl group disproportionation might also occur for 1,2-dialkyldimetal compounds.

$$L_n M^{x+} R_2 \rightarrow L_n M^{(x-2)+} + alkane + 1-alkene$$

Previously, we suggested that the thermal stability of compounds of the type  $1,2-Mo_2(i-Pr)_2(NMe_2)_4$  arises from two factors: (i) B-hydrogen elimination is not favored because metal atomic orbitals are not available (or suited) to form C-H-Mo 3 center bonds<sup>7</sup> and (ii) reductive elimination would generate a coordinatively unsaturated and presumably highly energetic metallic species<sup>8</sup> (Me\_2N)\_2Mo = Mo(NMe\_2)\_2. Both of these factors are controllable, in principle, by suitable reactions involving the dimethylamido ligands and previous work had shown that  $W_2Me_2(NEt_2)_4$  and  $W_2(NMe_2)_6$  react with carbon

dioxide to give  $W_2Me_2(O_2CNEt_2)_4$  and  $W_2(O_2CNMe_2)_6$ , respectively.<sup>9</sup> The structures of these two molecules, which are shown in I and II below, led us to believe that the ethyl compound  $Mo_2Et_2(O_2CNMe_2)_4$  would be labile to reductive elimination. (1) The carbamato compound  $Mo_2(O_2CNMe_2)_4$ , formed upon reductive elimination, would be analogous to scores of well characterized  $Mo_2(O_2CR)_4$  (MEM) compounds.<sup>10</sup> (2) The  $\beta$ -hydrogen elimination step would satisfy the 16-18 electron rule<sup>11</sup> and formation of the Mo-H bond along the Mo-Mo axis would be stereochemically analogous to formation of the axial W-O bonds found in  $W_2(O_2CNMe_2)_6$ . We report here the details of our studies which were prompted by these lines of reasoning. A preliminary report of some aspect of this work has appeared.<sup>12</sup>



Central skeleton of the  $W_2Me_2(O_2CNEt_2)_4$  molecule.

I



Central skeleton of the  $W_2(O_2CNMe_2)_6$  molecule. Dotted lines indicate the long, quasi-axial W-O bonds.

Π

### Results and Discussion

<u>Reactions with Carbon Dioxide</u>.  $Mo_2Me_2(NMe_2)_4$  reacts with  $CO_2$  in hydrocarbon solvents to give a finely divided yellow crystalline solid which is only sparingly soluble in hydrocarbon solvents. The compound shows a molecular ion,  $Mo_2Me_2(O_2CNMe_2)_4^+$ , in the mass spectrometer which, together with other spectroscopic and analytical data, lead us to believe it is analogous to the tungsten compound  $W_2Me_2(O_2CNEt_2)_4$ , which was structurally characterized and shown to have a metal-to-metal triple bond.<sup>9</sup>

In contrast to the above, the l,2-dialkyl dimolybdenum compounds  $Mo_2R_2^-$  (NMe<sub>2</sub>)<sub>4</sub>, where the alkyl group contains  $\beta$ -hydrogen atoms react according to equation 2 below.

# $2 Mo_2R_2(NMe_2)_4 + CO_2(excess) + Mo_2(O_2CNMe_2)_4 + Alkane + Alkene$

The compound  $Mo_2(O_2CNMe_2)_4$  is formed as a finely divided, yellow, microcrystalline solid which is virtually insoluble in hydrocarbon solvents and only very sparingly soluble in polar solvents, such as pyridine. The <sup>1</sup>H nmr spectrum shows only one resonance and the infrared spectrum shows bands associated with the  $O_2CNMe_2$  ligands.<sup>13</sup> In the mass spectrometer, the most intense ion was the ion of highest mass,  $Mo_2(O_2CNMe_2)_4^+$ ; there was also an intense doubly charged ion corresponding to  $Mo_2(O_2CNMe_2)_4^{2+}$ . There is therefore little doubt that  $Mo_2(O_2CNMe_2)_4$  is closely related to scores of other well characterized compounds of general formula  $Mo_2(O_2CR)_4$ , which contain metal-to-metal quadruple bonds.<sup>10</sup>

Studies of reaction 2 were carried out in sealed nmr tubes and the reaction course was followed by nmr spectroscopy. Within the limits of detection by the nmr method, reaction 2 is stoichiometric: equal molar quantities of alkane and alkene are liberated. At room temperatures, the reaction is

rapid and essentially complete within 30 min. Reactions involving R = n-Bugave 1-butene and n-butane (no 2-butenes were detected); for R = sec-Bu, n-butane, 1-butene and a trace of 2-butenes were evolved, while for R =tert-Bu, the eliminated hydrocarbons were iso-butylene and iso-butane. These observations are consistent with an irreversible  $\beta$ -hydrogen atom to metal transfer reaction. This was further supported by the studies of the reactions between  $CO_2$  and each of the labelled ethyl compounds  $MO_2(CH_2CD_3)_2(NMe_2)_4$  and  $MO_2(CD_2CH_3)_2(NMe_2)_4$ . In both of these reactions, the liberated ethylene was  $CH_2=CD_2$ .

The labelling pattern in the three  $C_2H_2D_2$  isomers can be deduced by <sup>2</sup>H nmr spectroscopy. See Figure 1. Each of the isomers of dideuteroethylene show a central three-line pattern with the only apparent distinguishing characteristic being the separation between the outer lines, which is 4.8 Hz for  $CH_2=CD_2$ , 3.3 Hz for <u>cis</u>-CHD=CHD and 2.2 Hz for <u>trans</u>-CHD=CHD.

The ethane liberated in these reactions is either  $CH_2DCD_3$ , when R =  $CH_2CD_3$  or  $CHD_2CH_3$ , when R =  $CD_2CH_3$ . The <sup>2</sup>H nmr spectra of the evolved ethanes are compared with the spectra obtained for commercially available samples of the specifically labelled ethanes in Figure 2. In a separate experiment, an equimolar mixture of  $Mo_2(CH_2CH_3)_2(NMe_2)_4$  and  $Mo_2(CH_2CD_3)_2(NMe_2)_4$  was allowed to react with  $CO_2$  in an nmr tube. The <sup>2</sup>H nmr spectrum of the liberated ethane was indistinguishable from that shown in Figure 2 for  $CH_2DCD_3$ . We conclude that, within the limits of detection by nmr spectroscopy, the elimination of ethane occurs by an intramolecular process: an intermolecular process would generate  $CH_3CH_2D$  and  $CH_3CD_3$  along with  $CH_2DCD_3$  and  $C_2H_6$ .

The reaction between  $Mo_2Et_2(NMe_2)_4$  and  $CO_2$  in toluene-d<sub>8</sub> was followed by <sup>1</sup>H nmr spectroscopy at -30 °C. At this temperature, the reaction is quite

slow, taking several hours to go to completion. Certain significant observations arose from these studies. (1) The ratio of ethylene and ethane remained constant throughout the course of the reaction: ethylene and ethane are formed at apparently the same rate and it is not possible to determine which is eliminated first. (2) No signal, which might have been assigned to a metal-hydride, was detected in the range -20 to +30 ppm. [None would be expected, given (1) above]. (3) The reaction rate varied slightly from sample to sample, even when concentrations of  $Mo_2Et_2(NMe_2)_4$  and  $CO_2$  were kept virtually constant. This is similar to previous observations of  $CO_2$  insertion reactions involving  $W(NMe_2)_6$  and  $W_2Me_2(NEt_2)_4$ , which were shown to require the presence of trace amounts of free amine. 14 (4) In all reactions, the build up in concentration of a reactive intermediate was seen. The <sup>1</sup>H nmr spectrum of this intermediate is shown in Figure 3 and is consistent with that expected for  $Mo_2Et_2(NMe_2)_2(O_2CNMe_2)_2$ , providing that the molecule has virtual C<sub>2</sub> symmetry. A plausible structure is shown schematically in III below. This proposed structure is similar to that found  $^{15}$  for Mo $_2(0Bu^t)_4$ - $(0_{2}COBu^{t})_{2}$  and that reported in this paper for  $Mo_{2}Et_{2}(NMe_{2})_{2}(C_{7}H_{8}N_{3}C_{7}H_{8})_{2}$ , where  $C_7H_8 = p$ -tolyl.



III

The rate determining step must involve a reaction of  $Mo_2Et_2(NMe_2)_2^{-1}(O_2CNMe_2)_2$ , but we have not been able to determine the nature of this step. It seemed reasonable to suppose that other reactions, in which dimethylamido ligands were replaced by bidentate ligands, would show similar facility in inducing reductive elimination of alkane and alkene.

<u>Reactions with 1,3-Diaryltriazenes</u>. Previously, we have shown that  $Mo_2(NMe_2)_6$  and  $Mo_2Me_2(NMe_2)_4$  react with 1,3-diaryltriazenes according to equations  $3^{16}$  and 4, <sup>17</sup> respectively.

$$3 Mo_2(NMe_2)_5 + ArNNNHAr (excess) - 3 Mo_2(NMe_2)_5 + ArNNNHAr (excess)_5 + 3 Mo_2(NMe_2)_5 + 3 Mo_2(N$$

 $Mo_2(NMe_2)_4(ArN_3Ar)_2 + 2HNMe_2$ 

4 Mo<sub>2</sub>Me<sub>2</sub>(NMe<sub>2</sub>)<sub>4</sub> + ArNNNHAr (excess) →

 $Mo_2Me_2(NMe_2)_2(ArN_3Ar)_2 + 2HNMe_2$ 

Reactions 3 and 4 appear quantitative and irreversible. Even in the presence of excess triazene, only two dimethylamido groups are substituted. Single crystal X-ray studies revealed that, in both compounds, the triazenido ligands are bidentate: in  $Mo_2Me_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ ,<sup>17</sup> they bridge the Mo=Mo bond, while in  $Mo_2(NMe_2)_4(C_6H_5N_3C_6H_5)_2$ ,<sup>16</sup> there is an unbridged Mo=Mo bond.

In contrast to the above, 1,3-diaryltriazenes react with  $Mo_2R_2(NMe_2)_4$  compounds, where R = Et and i-Pr, in hydrocarbon solvents according to equation 5.

$$5 Mo_2R_2(NMe_2)_4 + ArNNNHAr (excess)$$

 $Mo_2(ArN_3Ar)_4 + 4HNMe_2 + Alkane + Alkene$ The compound  $Mo_2(C_6H_5N_3C_6H_5)_4$  has been synthesized previously and shown by Cotton and coworkers<sup>18</sup> to have a Mo-to-Mo quadruple bond.

We were unable to obtain satisfactory elemental analyses for our compounds: indeed, analytical data varied in a seemingly erratic manner. Thus, crystals of the compound  $Mo_2(C_7H_8N_3C_7H_8)_4$ , grown from toluene/hexane solution, were studied by X-ray diffraction. This yielded a solution<sup>19</sup> having a structure essentially identical to that reported by Cotton, <u>et al.</u>,<sup>18</sup> for the  $Mo_2(PhN_3Ph)_4$ . The problem associated with obtaining good elemental analytical results probably arises from the partial solvation observed in the crystals. The blades of the aryl groups are packed in such a way as to generate channels and holes. In the crystal which we examined, toluene molecules were located and refined in the channels, but a large hole, <u>ca</u>. 12 Å in diameter, was apparently partially occupied by some molecule of inclusion. Full structural details are given in the supplementary data. The characterization of  $Mo_2(C_7H_8N_3C_7H_8)_4$  is not pertinent to the thrust of this paper beyond establishing that, in reaction 5, the metal-to-metal bond order changes from three to four.

By the careful addition of 2 equivalents of 1,3-di-p-tolyltriazene to a hydrocarbon solution of  $Mo_2Et_2(NMe_2)_4$ , we have been able to isolate  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  as an orange crystalline compound. In the solid state, it is quite stable and like  $Mo_2Me_2(NMe_2)_2(ArN_3Ar)_2$  and  $Mo_2(NMe_2)_4(ArN_3Ar)_2$  compounds may be exposed to the atmosphere for short periods of time without significant decomposition. [This contrasts with the high reactivity of  $Mo_2Et_2$ -( $NMe_2$ )<sub>4</sub>, which smolders when exposed to the atmosphere]. On the other hand,  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  slowly decomposes in solution at room temperature and eliminates ethane (no ethylene is evolved). Significantly, the addition of either  $C_7H_8NNHC_7H_8$  or  $CO_2$  causes a more rapid evolution of ethane and ethylene. Addition of  $C_7H_8NNHC_7H_8$  yields  $Mo_2(C_7H_8N_3C_7H_8)_4$ ; reaction with  $CO_2$  presumably generates  $Mo_2(C_7H_8N_3C_7H_8)_2(O_2CNMe_2)_2$  - but this remains to be established.

<u>Solid State Structure of  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ </u>. An ORTEP view of the molecule giving the atom numbering scheme is shown in Figure 4. A vie, down the Mo-Mo bond is shown in Figure 5. Final atomic coordinates and

thermal parameters are given in Table 1 and 2, respectively. Complete listings of bond distances and bond angles are given in Tables 3 and 4, respectively.

The molecule has virtual, but not crystallographically imposed, C2 symmetry and is, in this and most other respects, analogous to that observed in  $Mo_2Me_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ . The Mo-C bond distance, 2.21(1) Å (average), is longer than that in the methyl analogue, 2.193(4) Å and that in the parent compound,  $Mo_2Et_2(NMe_2)_4$ , which is 2.16(1) Å (averaged).<sup>2</sup> Also, unlike in Mo<sub>2</sub>Et<sub>2</sub>(NMe<sub>2</sub>)<sub>4</sub>, the ethyl groups are directed over the Mo≡ Mo bond. As we predicted for this situation, the Mo-BH distances across the MoEMo bond are shorter than are those to the molybdenum atoms to which the ethyl group is directly bonded. See Table 5. However, in the present case, no significant C-H---Mo intraction is seen in the ground state. The orientation of the ethyl group over the Mo=Mo moiety can readily be understood in terms of steric considerations. The tolyl groups effectively block the axial positioning of the methyl groups and direct them inwards. The obtuse Mo-C-C angles = 126.1(5) and 125.7(5)°, which may be compared with Mo-C-C =  $112.0(3)^{\circ}$  and 113.0(3)° found in  $Mo_2Et_2(NMe_2)_4$ , reflect the internal crowding of the molecule. This may also be the cause of the lengthening of the Mo-C (ethyl) bond distances.<sup>20</sup>

<u>Concluding Remarks</u>. Addition of either  $CO_2^{\prime}$  or 1,3-diaryltriazines to 1,2-Mo'<sub>2</sub>R'<sub>2</sub>(NMe<sub>2</sub>)<sub>4</sub> compounds promotes reductive elimination from the dimetal center by alkyl group disproportionation with a concomittant metal-metal bond order change from three to four when R = an alkyl group having B-hydrogen atoms. Both reactions proceed through intermediates of the form 1,2-Mo<sub>2</sub>R<sub>2</sub>-

(NMe<sub>2</sub>)<sub>2</sub>(chelate)<sub>2</sub>. The elimination of alkane and alkene is intramolecular and has a formal analogy with that found in mononuclear chemistry, namely B-hydrogen elimination followed by reductive elimination by carbon-hydrogen bond formation. The detailed pathway leading to this is, however, uncertain. The presence of the dimetal center introduces additional possibilities. For example, the H-Mo bond could be formed across the Mo=Mo bond. This would be akin to a cyclometallation reaction involving a 5-membered ring<sup>21</sup> and the structure of  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  clearly shows the geometric plausibility of this reaction pathway. The observation that excess  $C_7H_8NNNHC_7H_8$ promotes a reaction with  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ , but not with  $Mo_2R_2$ - $(NMe_2)_2$  compounds, where  $R = CH_3$  or  $CH_2CMe_3$ , could be correlated with a reversible protonation of a  $Mo-NMe_2$  nitrogen lone pair. In the  $Mo_2R_2(NMe_2)_2$ - $(C_7H_8N_3C_7H_8)_2$  compounds, the NMe<sub>2</sub> ligands are clearly strong  $\pi$ -donors to one of the five in-plane molybdenum atomic orbitals (s,  $p_x$ ,  $p_y$ ,  $d_{xy}$ ,  $d_{x^2-y^2}$ ). Upon protonation of the amide lone pair, one of these orbitals would be freed to form a CH---Mo 3-center, 2-electron bond. The structure of Mo<sub>2</sub>Et<sub>2</sub>(NMe<sub>2</sub>)<sub>2</sub>- $(C_7H_8N_3C_7H_8)_2$  suggests that the B-hydrogens of the ethyl ligands are poised for such an opportunity. Alternatively, a 1,2- to 1,1-Mo $_2$ Et $_2$  isomerization reaction, involving an alkyl group transfer across the Mo≣Mo bond, may be required, and promoted in the association reaction between 1,2-Mo<sub>2</sub>Et<sub>2</sub>(NMe<sub>2</sub>)<sub>2</sub>- $(C_7H_8N_3C_7H_8)_2$  and excess triazine, prior to B-hydrogen elimination. For a 1,1-Mo<sub>2</sub>Et<sub>2</sub>(NMe<sub>2</sub>)<sub>2</sub>( $C_7H_8N_3C_7H_8$ )<sub>2</sub> molecule, one molybdenum atom would have one in-plane atomic orbital available for B-hydrogen elimination in the more conventional sense of this terminology (i.e. mononuclear). Within this context, it should be noted that (i) there is a high kinetic barrier to 1,1- and 1,2-isomerization of  $Mo_2(NMe_2)_2(CH_2SiMe_3)_4$  compounds<sup>22</sup> and (ii) Bergman, et

<u>al</u>.,<sup>23</sup> have proposed, with good experimental evidence, that reductive elimination of acetone from 1,2-Cp<sub>2</sub>Co<sub>2</sub>Me<sub>2</sub>(CO)<sub>2</sub> proceeds via an initial methyl group transfer from one cobalt atom to the other. Finally, it should be recognized that it is not necessary to invoke the formation of an intermediate metal-hydride in these reactions. The  $\beta$ -hydrogen atom of an ethyl ligand could be transferred to the  $\alpha$ -carbon of the other ethyl ligand through the agency of one of the nitrogen atoms, which could be either a triazenido or dimethyl-amido ligand. In a formal sense this would have analogy with E<sub>2</sub> reactions of substituted alkanes. These, and other possibilities, provide us with the stimulus for further studies of reductive eliminations of hydrocarbons from dinuclear metal centers.

## Experimental Section

General procedures, including the preparations of  $1,2-Mo_2R_2(NMe_2)_4$  compounds, have been described previously.<sup>2</sup> Note the use of dry and oxygen-free solvents and atmospheres.

 $^{1}$ H and  $^{2}$ H nmr spectra were recorded on Varian HR220 MHz spectrometer equipped with a variable temperature probe. Infrared spectra were recorded on a PE-283 spectrophotometer. Mass spectra were obtained on a Varian MS-902 by the method of direct insertion, courtesy of Mr. Peter Cook, Queen Mary College, London.

<u>Preparation of  $Mo_2(0_2CNMe_2)_4$ </u>. To a 25 mL flask equipped with a stir-bar and  $N_2$  inlet, was added 0.100 g (0.23 mmoles) of  $Mo_2Et_2(NMe_2)_4$  in 15 mL of toluene. The flask was attached to a calibrated manifold, evacuated at liquid  $N_2$  temperature, and 6 equivalents of  $CO_2$  were added. As the flask gradually warmed to room temperature, the color changed from a golden yellow to orange with the precipitation of a pale yellow powder. The solution above the powder was virtually colorless at the completion of the reaction. Infrared spectra of the pale yellow powder using Nujol mull and CsI plates gave a strong absorption at 1560 cm<sup>-1</sup> ( $\vee NCO_2$  of bridging carbamate). The yellow powder was extremely insoluble in most solvents; however, a sufficient quantity was soluble in pyridine-d<sup>5</sup> to record its nmr spectrum, which showed a singlet at 2.93 ppm relative to TMS corresponding to  $O_2CNMe_2$ . Elemental analyses: Found (calculated) - C, 33.55 (33.79); H, 7.89 (8.04); N, 12.99 (13.15).

 $\frac{\text{Preparation of Mo}_2(C_7H_8N_3C_7H_8)_4(\text{toluene})_x}{\text{moles}}$ . A 25 mL round-bottomed flask was charged with 0.25 g (0.58 mmoles) of Mo $_2\text{Et}_2(\text{NMe}_2)_4$  and 0.78 g (3.5 mmoles) of di-p-tolyliriazene. Approximately 15 mL of toluene were added at

room temperature in a Vacuum Atmospheres Dri Lab. The color was intensely red-orange. Hexanes (5 mL) were added to the flask before placing it in the cool storage refrigerator of the Dri Lab at  $-35^{\circ}$ C. Dark orange crystals of  $Mo_2(C_7H_8N_3C_7H_8)_4$  were prepared in this manner. Satisfactory analysis could not be obtained for this complex as was the case for many of the p-tolyltria-zene derivatives of analogous complexes.<sup>16,17</sup> In all respects, the complex showed properties similar to the unsubstituted aromatic derivative prepared by Cotton, et al.<sup>18</sup>

Analyses of Gases Liberated. The gases liberated upon reaction of  $CO_2$  with  $MO_2(C_2H_5)_2(NMe_2)_4$ ,  $MO_2(n-C_4H_9)_2(NMe_2)_4$ ,  $MO_2(sec-C_4H_9)_2(NMe_2)_4$ ,  $MO_2(i-C_3H_7)_2(NMe_2)_4$ , and  $MO_2(t-C_4H_9)_2(NMe_2)_4$  in toluene-d<sub>8</sub> were all compared with authentic samples of the respective alkenes and alkanes by nmr spectros-copy.

The deuterated gases  $CD_3CH_2D$ , 1,1- $CD_2CH_2$ , and  $CH_3CD_2H$  were purchased from Merck, Sharp, and Dohme and compared identically with those generated in the reactions given in Figure 2 at 33.77 MHz in benzene. Calculations using  ${}^{1}H-{}^{1}H$  coupling constants given in R.M. Lynden-Bell and N. Sheppard, Proc. R. Soc. A., <u>269</u>, 1385 (1965) for ethylene and a comparison of the magnetogyric ratios  ${}_{Y}$  ( ${}^{1}H$ ) = 2.675 x 10<sup>4</sup> radians/(s·G),  ${}_{Y_{2}H}$  = 4.107 x 10<sup>3</sup> radians/(s·G) allowed the simulation of the  ${}^{2}H$  nmr at 33.77 MHz of the various dideuteroethylenes shown in Figure 1. The calculated splittings were obtained by using the program entitled NMRCAL, a part of the Nicolet 80 computer system. Having purchased the various dideuteroethylenes, the line shapes and splittings observed for the commercial samples were found to be identical to those predicted by simulation.

<u>Synthesis of  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ </u>. In a 50 mL round-bottomed flask were placed 0.50 g (1.17 mmoles) of  $Mo_2Et_2(NMe_2)_4$  and 0.53 g (2.35 mmoles) of p-tolyltriazene. To this flask 30 mL of toluene were added in the Dri Lab. The color of the solution was intensely red. After 3 days in the refrigerator of the Dri Lab at  $-35^{\circ}C$ , red crystals precipitated. Addition of 10 mL of hexane to the reaction flask induced crystallization. A crystal of appropriate dimensions was chosen for X-ray analysis.

<u>X-Ray Structural Determination of  $Mo_2(NMe_2)_2Et_2(C_7H_8N_3C_7H_8)_2$ </u>. A crystal of dimensions 0.20 x 0.20 x 0.40 mm was mounted in a nitrogen-filled glovebag and transferred to the liquid-nitrogen boil-off system of the diffractometer.

The cell dimensions obtained from 46 reflections at  $-175^{\circ}$ C with MoKa ( $\lambda = 0.71069$  Å) were a = 16.503(4) Å, b = 23.116(7) Å, c = 11.318(3) Å, B = 88.15(1)^{\circ}, V = 4315.6(5) Å<sup>3</sup>, Z = 4, d<sub>calcd</sub> = 1.353 g cm<sup>-3</sup> and space group P21/a.

A total number of 6970 reflections were collected using standard movingcrystal, moving-detector techniques with the following values: scan speed = 4.0 deg min<sup>-1</sup>, scan width = 1.8 + dispersion, single background time at extremes of scan = 5 sec, aperture size =  $3.0 \times 4.0$  mm. Of the 6970 reflections collected, the number with unique amplitudes was 5667 and the number with F > 2.33 sigma(F) was 4622. The limits of data collection were  $6^{\circ} < 2\theta < 45^{\circ}$ .

The structure was solved using a combination of direct methods and Fourier techniques. A molecule of solvent, toluene, was located. All nonhydrogen atoms were refined anisotropically. All hydrogens were located in

the difference map, but were not refined due to space limitations in the least-squares program. The final residuals were R(F) = 0.057 and Rw(F) = 0.0795. The goodness of fit for the last cycle, 1.795 and the maximum delta/-sigma, 0.01.

Acknowledgement. We thank the Office of Naval Research, the Marshall H. Wrubel Computing Center and the taxpayers of the State of Indiana for financial support of this work.

#### References

- 1. Chisholm, M.H.; Huffman, J.C.; Kirkpatrick, C.C. Inorg. Chem., submitted.
- Chisholm, M.H.; Haitko, D.A.; Folting, K.; Huffman, J.C. J. Am. Chem. Soc., submitted.
- '-Alkyl groups bonded to transition metals are important in a large number of catalytic and stoichiometric reactions. See, for example, Kochi, J.K. in "Organometallic Mechanisms and Catalysis, Academic Press, New York, San Francisco and London, 1978.
- Whitesides, G.M.; Gaasch, J.F.; Stedronsky, E.R. J. Am. Chem. Soc. 1972, 94, 5258.
- 5. Ikariya, T.; Yamamoto, A. J. Organometal. Chem. 1976, 120, 257.
- McDermott, J.X.; Wilson, M.E.; Whitesides, G.M. J. Am. Chem. Soc. 1976, 98, 6529.
- 7. Cotton, F.A.; Day, V.W. <u>J.C.S. Chem. Commun.</u> 1974, 415; Cotton, F.A.; LaCour, T.; Stanislowski, A.G. J. Am. Chem. Soc. 1974, 96, 754.
- 8. Norton, J.R. Acc. Chem. Res. 1979, 12, 139.
- 9. Chisholm, M.H.; Cotton, F.A.; Extine, M.W.; Stults, B.R. <u>Inorg. Chem.</u> 1978, 17, 603.
- 10. Cotton, F.A. Chem. Soc. Rev. 1975, 4, 27; Acc. Chem. Res. 1978, 11, 225.
- 11. Tolman, C.A. Chem. Soc. Rev. 1972, 1, 337.
- 12. Chisholm, M.H.; Haitko, D.A. J. Am. Chem. Soc. 1979, 101, 6784.
- 13. Chisholm, M.H.; Extine, M.W. J. Am. Chem. Soc. 1977, 99, 782.
- 14. Chisholm, M.H.; Extine, M.W. J. Am. Chem. Soc. 1977, 99, 802.
- Chisholm, M.H.; Cotton, F.A.; Extine, M.W.; Reichert, W.W. J. Am. Chem. Soc. 1978, 100, 1727.
- Chisholm, M.H.; Haitko, D.A.; Folting, K.; Huffman, J.C. <u>Inorg. Chem. in</u> press.

- 17. Chisholm, M.H.; Haitko, D.A.; Folting, K.; Huffman, J.C. <u>Inorg. Chem.</u>, in press.
- 18. Cotton, F.A.; Rice, G.W.; Sekutowski, J.C. Inorg. Chem. 1979, 18, 1143.
- 19. Crystal data for  $Mo_2(C_7H_8N_3C_7H_8)_4(toluene)_x$ : ref. 19. Cell dimensions at -142°C from 32 reflections: a = 10.443(7) Å, b = 17.159(8) Å, c = 19.771(13) Å,  $\beta$  = 108.08(2)°, Z = 4,  $d_{calcd}$  = 1.674 g cm<sup>-3</sup> and space group = P21/n. A total of 5168 reflections were collected, 4426 of which were unique; the number with F > 2.33 sigma(F) was 3425. The final residuals are R(F) = 0.0951 and Rw(F) = 0.0948. For the last cycle, the goodness fit was 1.450 and the maximum delta/sigma = 0.05. The Mo-Mo distance, 2.090(2) Å, the Mo-N distances, 2.14(2) Å (averaged) and the Mo-Mo-N angles 91.5(5) Å (averaged) are all essentially the same as those reported by Cotton, et al. (ref. 18) for the phenyl analogue.
- 20. For a discussion of the lengthening of the Co-C bond distances in cobaloximes as a function of steric and conformational factors, see: Marzilli, L.G.; Toscano, P.J.; Randaccio, L.; Bresciani-Pahor, N.; Calligaris, M.J. J. Am. Chem. Soc. 1979, 101, 6754; idem, ibid 1980, 102, 7372.
- 21. Parshall, G.W. Acc. Chem. Res. 1970, 3, 139.
- 22. Chisholm, M.H.; Rothwell, I.P. J.C.S. Chem. Commun. 1980, 985.
- 23. Bergman, R.G. Acc. Chem. Res. 1980, 13, 113.

Table 1. Fractional Coordinates for the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  Molecule.

<u>Note 1</u>: The isotropic thermal parameter listed for those atoms refined aniostropically are the isotropic equivalent.

Note 2: Number in parentheses in this and all following tables refer to the error in the least significant digits.

Atom	10 <sup>4</sup> x	10 <sup>4</sup> y	10 <sup>4</sup> z	10 B <sub>iso</sub>
Mo(1)	4878.8(4)	2293.1(3)	8199(1)	14
Mo(2)	5997.0(4)	2268.3(3)	9156(1)	14
N(3)	4518(4)	1480(3)	91/4(5)	15
N(4) N(5)	48/6(4)	1306(3)	10097(5)	1/
N(5)	6281(3)	1019(2)	10358(5)	14
N(0)	5909(3)	1435(3)	7267(5)	15
N(8)	5306(3)	1622(3)	7006(5)	15
N(9)	4848(3)	3014(3)	7310(5)	16
C(10)	5389(5)	3512(3)	7144(7)	23
C(11)	4063(4)	3132(3)	6769(7)	22
C(12)	3959(4)	2579(3)	9529(6)	20
C(13)	<b>399</b> 0(5)	2485(4)	10891(7)	25
N(14)	6089(3)	3001(3)	9998(5)	18
C(15)	5582(5)	3523(3)	10109(7)	23
C(16)	6894(5)	3108(4)	10505(7)	24
C(17)	6941(4)	2516(4)	/80/(6)	19
(18)	6902(5)	2406(4)	6469(7)	29
(19)	3826(4)	1141(3)	8880(0) 7012(7)	15
C(20)	2714(5)	1312(3)	7913(7)	20
(21)	2466(5)	532(3)	8205(7)	24
(22)	1741(5)	147(A)	7812(8)	29
(24)	2906(5)	340(4)	9197(7)	27
C(25)	3588(5)	653(3)	9529(7)	22
C(26)	5880(4)	1409(3)	11391(6)	18
C(27)	5872(4)	828(3)	11714(7)	22
C(28)	6243(5)	653(4)	12750(7)	27
C(29)	6649(5)	1046(4)	13468(7)	28
C(30)	7027(5)	840(4)	14583(7)	34
C(31)	6665(5)	1619(4)	13130(7)	30
C(32)	6294(5)	1811(4)	12084(7)	25
(33)	6974(4) 7456(5)	1086(3)	8417(6)	10
C (34)	7450(5) 9195(5)	1243(3)	9333(7)	20
(35)	7456(5)	1245(3)	9353(7)	20
C(35)	8185(5)	947(3)	9580(7)	20
C(36)	8417(5)	493(3)	8864(7)	22
C(37)	9208(5)	175(4)	9082(8)	28
C(38)	7911(5)	315(4)	7949(7)	25
C(39)	7193(5)	607(3)	7731(7)	21
C(40)	4911(4)	1435(3)	5961(6)	17
C(41)	4659(5)	1860(4)	5177(7)	24
C(42)	4269(5)	1701(4)	4138(7)	28
C(43)	4102(5)	1119(4)	3910(7)	26
(44)	3692(5)	944(4)	2/89(7)	31
U(45) C(46)	4350(5)	090(4)	4/14(/)	20
C(40)	-752(R)	1307(7)	3265(10)	74
C(48)	85(7)	1215(5)	3654(8)	45
C(49)	625(7)	1672(5)	3635(8)	52
C (50)	1425(8)	1588(8)	3991(9)	78
C(51)	1681(8)	1022(9)	4366(10)	83
C(52)	1108(11)	588(8)	4355(11)	98
C(53)	322(9)	683(5)	4015(9)	65

Table 2. Anisotropic Thermal Parameters for the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ Molecule. The form of the exponent is:  $exp[-2\pi^2(U_{11}h^2a^{*2} + U_{22}k^2b^{*2} + U_{33}kc^{*2} + 2U_{12}hka^{*}b^{*} + 2U_{13}h^{k}a^{*}c^{*} + 2U_{23}k^{k}b^{*}c^{*})].$ 

Atom	ווט	U22	U33	U12	U13	U23
Mo(1) Mo(2) N(3) N(4) N(5) N(5) N(6) N(7) N(8) C(10) C(12) C(12) C(12) C(12) C(13) C(12) C(13) C(13) C(14) C(15) C(22) C(23) C(22) C(23) C(22) C(22) C(23) C(22) C	15.6(4) $15.4(4)$ $20(4)$ $20(4)$ $21(4)$ $16(3)$ $21(4)$ $18(3)$ $18(4)$ $28(5)$ $24(4)$ $15(4)$ $30(5)$ $18(4)$ $31(5)$ $22(4)$ $13(4)$ $29(5)$ $19(4)$ $11(4)$ $23(5)$ $27(5)$ $20(4)$ $17(4)$ $25(5)$ $23(5)$ $43(6)$ $40(6)$ $25(5)$ $17(4)$ $30(5)$ $24(5)$ $27(5)$ $20(4)$ $17(4)$ $25(5)$ $23(5)$ $43(6)$ $40(6)$ $25(5)$ $17(4)$ $30(5)$ $24(5)$ $27(5)$ $31(5)$ $26(5)$ $11(4)$ $29(5)$ $31(5)$ $35(5)$ $25(5)$ $18(4)$ $88(10)$ $77(8)$ $87(9)$ $79(10)$	$15.7(4) \\ 16.2(4) \\ 16.2(4) \\ 18(4) \\ 22(4) \\ 15(3) \\ 19(4) \\ 19(4) \\ 19(4) \\ 26(4) \\ 14(4) \\ 30(5) \\ 37(5) \\ 39(5) \\ 22(4) \\ 20(5) \\ 37(5) \\ 39(5) \\ 22(4) \\ 20(5) \\ 37(5) \\ 39(5) \\ 22(4) \\ 20(5) \\ 27(5) \\ 26(5) \\ 41(6) \\ 30(5) \\ 24(5) \\ 28(5) \\ 33(5) \\ 49(6) \\ 54(7) \\ 63(7) \\ 51(7) \\ 41(6) \\ 30(5) \\ 24(5) \\ 28(5) \\ 33(5) \\ 49(6) \\ 54(7) \\ 63(7) \\ 51(7) \\ 41(6) \\ 30(5) \\ 28(5) $	$\begin{array}{c} 20.9(4)\\ 21.3(4)\\ 20(4)\\ 22(4)\\ 18(3)\\ 23(4)\\ 17(3)\\ 21(4)\\ 17(3)\\ 44(5)\\ 28(5)\\ 23(4)\\ 25(5)\\ 27(4)\\ 36(5)\\ 32(5)\\ 19(4)\\ 31(5)\\ 21(4)\\ 37(5)\\ 42(5)\\ 32(5)\\ 19(4)\\ 32(5)\\ 32(5)\\ 19(4)\\ 32(5)\\ 25(5)\\ 24(5)\\ 23(5)\\ 25(5)\\ 25(4)\\ 33(5)\\ 25(5)\\ 24(4)\\ 28(4)\\ 37(5)\\ 25(5)\\ 24(4)\\ 28(4)\\ 37(5)\\ 25(5)\\ 24(4)\\ 28(4)\\ 37(5)\\ 22(4)\\ 25(5)\\ 32(5)\\ 32(5)\\ 32(5)\\ 32(5)\\ 32(5)\\ 24(4)\\ 28(4)\\ 37(5)\\ 22(4)\\ 25(5)\\ 32(5)$	$\begin{array}{c} 0.8(3)\\ 0.2(3)\\ -1(3)\\ 3(3)\\ 1(3)\\ 3(3)\\ -2(3)\\ 1(3)\\ 0(3)\\ -2(4)\\ 4(4)\\ 2(4)\\ 4(4)\\ -2(3)\\ 7(4)\\ -11(4)\\ -2(3)\\ 7(4)\\ -11(4)\\ -3(4)\\ 2(4)\\ 3(3)\\ 1(3)\\ 6(4)\\ 3(4)\\ -19(4)\\ 0(4)\\ -3(4)\\ 3(4)\\ -19(4)\\ 0(4)\\ -3(4)\\ 3(4)\\ 7(4)\\ 10(4)\\ 9(4)\\ 14(5)\\ 1(5)\\ 4(4)\\ -1(3)\\ -3(4)\\ 3(4)\\ -0(4)\\ 14(5)\\ 1(5)\\ 4(4)\\ -1(3)\\ -3(4)\\ 3(4)\\ -0(4)\\ 14(4)\\ 2(4)\\ 1(4)\\ -2(3)\\ -4(4)\\ -7(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -3(4)\\ -2(7)\\ -27(7)\\ -26(11)\\ \end{array}$	$\begin{array}{c} 0.4(3) \\ -0.1(3) \\ 3(3) \\ 3(3) \\ 2(3) \\ 1(3) \\ 0(3) \\ -1(3) \\ 4(3) \\ 2(4) \\ -6(4) \\ 7(3) \\ 4(4) \\ 9(3) \\ 4(4) \\ -6(4) \\ 3(3) \\ 6(4) \\ 2(4) \\ 4(4) \\ -3(4) \\ 4(4) \\ 3(3) \\ 6(4) \\ 2(4) \\ 4(4) \\ -3(4) \\ 4(4) \\ 3(3) \\ 3(4) \\ 11(4) \\ 2(4) \\ -1(4) \\ 0(4) \\ 4(4) \\ 1(3) \\ 1(4) \\ 2(4) \\ -1(4) \\ 0(4) \\ 4(4) \\ 1(3) \\ 1(4) \\ 5(4) \\ 1(4) \\ 5(4) \\ 1(4) \\ 5(4) \\ 1(4) \\ 5(4) \\ 1(4) \\ -3(4) \\ 5(4) \\ 3(3) \\ 3(4) \\ -6(4) \\ 6(4) \\ -6(4) \\ 3(4) \\ -7(7) \\ 7(5) \\ 15(6) \\ 24(7) \end{array}$	1.1(3) -0.6(3)  1(3) -2(3)  2(3)  0(3) -0(3)  -1(3)  5(3)  0(4)  6(4)  -3(4) -0(4)  -3(3) -5(4)  -1(4)  6(4)  8(5) -1(3)  -1(4)  -5(4)  -4(4)  -4(4)  -5(4)  -4(4)  0(4)  9(4)  7(4)  3(5)  11(5)  3(4)  3(4)  12(4)  12(4)  12(4)  12(4)  12(4)  12(4)  3(4)  -5(4)  -5(4)  -7(4)  3(4)  -15(4)  -7(4)  5(5)  -15(5)  -25(9)  -25(9)  -25(9)  -1(3)  -1(3)  -1(4)  -3(3)  -5(4)  -1(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(4)  -5(5)  -25(9)  -15(5)  -25(9)  -15(5)  -25(9)  -15(5)
C (52) C (53)	143(17) 147(14)	181(19) 68(10)	45(8) 29(6)	29(12) 57(15) 13(9)	9(7) 21(11) 15(7)	-15(10) -33(10) -6(6)

All U's are x10\*\*3

# Table 3. Bond Distances, in Angstroms,

Α	В	Distance
Mo(1) Mo(1) Mo(1)	Mo(2) N(3) N(8) N(8)	2.171(1) 2.251(6) 2.160(6)
Mo(1) Mo(2) Mo(2) Mo(2)	C(12) N(5) N(6)	2.205(7) 2.171(6) 2.253(6)
Mo(2) N(3) N(3) N(4)	C(17) N(4) C(19) N(5)	2.221(7) 1.281(8) 1.431(9) 1.316(8)
N(5) N(6) N(6) N(7)	C(26) N(7) C(33) N(8)	1.412(9) 1.286(8) 1.428(9) 1.314(8)
N(8) N(9) N(9) N(14)	C(40) C(10) C(11) C(15)	1.437(9) 1.464(9) 1.476(9) 1.471(9)
N(14) C(12) C(17) C(19)	C(16) C(13) C(18) C(20)	1.484(9) 1.559(10) 1.539(11) 1.384(10)
C(19) C(20) C(21) C(22)	C(25) C(21) C(22) C(23)	1.393(10) 1.420(11) 1.384(11) 1.540(11)
C(22) C(24) C(26) C(26) C(26)	C(24) C(25) C(27) C(32)	1.411(11) 1.400(11) 1.393(11) 1.406(11)
C(27) C(28) C(29) C(29) C(29)	C(28) C(29) C(30) C(31)	1.400(11) 1.403(12) 1.502(11) 1.380(12)
C (31) C (33) C (33) C (34) C (35)	C(32) C(34) C(39) C(35) C(36)	1.421(11) 1.395(10) 1.394(10) 1.417(10)
C(36) C(36) C(38) C(40)	C(38) C(38) C(39) C(41)	1.526(10) 1.413(11) 1.391(11)
C(40) C(41)	C(46) C(42)	1.384(11) 1.407(11)

for the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  Molecule.

÷

A	В	Distance
A C (42) C (43) C (43) C (43) C (45) C (45) C (47) C (48) C (48)	B C(43) C(44) C(45) C(46) C(46) C(48) C(49) C(53)	Distance 1.400(12) 1.511(11) 1.402(11) 1.400(11) 1.479(16) 1.382(14) 1.358(16)
C (49) C (50) C (51) C (52)	C(50) C(51) C(52) C(53)	1.407(16) 1.445(21) 1.379(22) 1.383(19)

٠

Table 3. Continued.

1 400

Ì

# Table 4. Bond Angles, in Degrees,

A	B	C	Angle
Mo(2)	Mo(1)	N(3)	87.1(2)
Mo(2)	Mo(1)	N(8)	91.6(2)
Mo(2)	Mo(1)	N(9)	108.5(2)
Mo(2)	Mo(1)	C(12)	104.2(2)
N(3)	MO(1) Mo(1)	N(8)	1/.4(2)
N(3)	MO(1) Mo(1)	N(9) C(12)	75 5(2)
N(S)	Mo(1)	N(Q)	107 8(2)
N(8)	Mo(1)	$\Gamma(12)$	147.8(3)
N(9)	Mo(1)	C(12)	93,9(3)
Mo(1)	Mo(2)	N(5)	91.7(2)
Mo(1)	Mo(2)	N(6)	87.1(1)
Mo(1)	Mo(2)	N(14)	107.5(2)
Mo(1)	Mo(2)	C(17)	103.8(2)
N(5)	Mo(2)	N(6)	77.4(2)
N(5)	Mo(2)	N(14)	109.0(2)
N(5)	Mo(2)	C(17)	148.2(3)
N(6)	Mo(2)	N(14)	163.5(2)
N(D) N(JA)	MO(2) Mo(2)	( 17)	76.0(3)
$M_{0}(14)$	MU(2)	U(17)	$\frac{92.0(3)}{122.7(5)}$
Mo(1)	N(3)	C(19)	122.7(5) 123 1(5)
N(4)	N(3)	C(19)	114 0(6)
N(3)	N(4)	N(5)	114,1(6)
Mo(2)	N(5)	N(4)	120,9(4)
Mo(2)	N(5)	C(26)	126.4(5)
N(4)	N(5)	C(26)	111.8(6)
Mo(2)	N(6)	N(7)	122.6(4)
Mo(2)	N(6)	C(33)	123.6(5)
N(7)	N(6)	C(33)	113.0(6)
N(6)	N(7)	N(8)	113.4(6)
MO(1) Mo(1)	N(8)	N(7) C(40)	121.9(4)
MO(1) N(7)	N(O) N(B)	C(40)	125.7(5) 111 7(6)
Mo(1)	N(0)	C(10)	$135 \ 4(5)$
Mo(1)	N(9)	C(11)	114.2(4)
C(10)	N(9)	C(11)	110.0(6)
Mo(2)	N(14)	C(15)	134.5(5)
Mo(2)	N(14)	C(16)	114.7(5)
C(15)	N(14)	C(16)	110.1(6)
Mo(1)	C(12)	C(13)	126.1(5)
Mo(2)	C(17)	C(18)	125.7(5)
N(3)	C(19)	C(20)	117.0(6)
N(3)	C(19)	C(25)	122.6(7)

for the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_5)_2$  Molecule.

Table 4. Co	ontinued.
-------------	-----------

A	В	C	Angle
C(20) C(19)	C(19) C(20)	C(25) C(21)	120.4(7)
C(20)	C(21)	Č(22)	119.9(8)
C(21)	C(22)	C(23)	120.4(7)
C(21) C(23)	C(22)	C(24)	119.4(8)
C(22)	C(24)	C(25)	120.6(8)
C(19) N(5)	C(25)	C(24)	119.6(8)
N(5)	C(26)	C(27)	123.0(7
C(27)	C(26)	C(32)	119.4(7)
C(26)	C(27)	C(28)	119.9(8)
C(27) C(28)	C(28)	C(29)	121.9(8)
C(28)	C(29)	C(31)	117.8(8)
C(30)	C(29)	C(31)	122.1(8)
C(29)	(31)	(32)	121.7(8)
N(6)	C(33)	C(34)	117.7(7)
N(6)	C(33)	C(39)	123.1(7)
2 (34) 2 (33)	(33)	C(39)	19.2(7)
C(34)	C(35)	C(36)	119.2(7)
C(35)	C(36)	C(37)	119.8(7)
2(35) 2(37)	C(36)	C(38) C(38)	119.7(7)
C(36)	C(38)	C(39)	120.5(7)
2(33)	C(39)	C(38)	119.6(7)
1(8) 1(8)	C(40)	C(41) C(46)	117.6(7)
C(41)	C(40)	C(46)	119.9(7)
C(40)	C(41)	C(42)	120.0(7)
(4) (42)	C(42) C(43)	C(43)	120.3(8)
(42)	C(43)	C(45)	118.9(8)
(44)	C(43)	C(45)	120.5(8)
(43)	C(45) C(46)	C(46) C(45)	120.6(8)
(47)	C(48)	C(49)	119.6(11
(47)	C(48)	C(53)	120.0(12
2(49) 2(48)	U(48) C(49)	C(53) C(50)	120.4(12
(49)	C(50)	C(51)	119.7(13
(50)	C(51)	C(52)	116.8(13
()))	U(52) C(52)	L(53) C(52)	122.4(17

•

ļ

Table 5. Distances, in Angstroms,

from B-Hydrogens of the Ethyl Ligands to the Molybdenum Atoms in the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$ Molecule, as Located in the Difference Map. The numbering scheme for the hydrogen

atoms is given in Figure 5.

Atom	Atom	Distance <sup>a</sup>
 Mo(l)	H(1)	3.5(1)
Mo(1)	H(2)	3.6(1)
Mo(1)	H(3)	4.2(1)
Mo(2)	H(4)	3.5(1)
Mo(2)	H(5)	3.7(1)
Mo(2)	H(6)	4.1(1)
Mo(1)	H(4)	3.4(1)
Mo(1)	H(5)	4.2(1)
Mo(1)	H(6)	4.7(1)
Mo(2)	н(1)	3.4(1)
Mo(2)	H(2)	4.0(1)
Mo(2)	Н(3)	4.8(1)

<sup>a</sup>Errors are estimated based on the internal consistancy of the C-H distances and angles, as located in the difference synthesis.

## Captions to Figures

- Figure 1. <sup>2</sup>H nmr spectra for the isomers of C<sub>2</sub>H<sub>2</sub>D<sub>2</sub> recorded in benzene (top) are compared with the predicted spectra (bottom) based on <sup>1</sup>H-<sup>1</sup>H coupling constants for ethylene and (<sup>1</sup>H) taken from R.M. Lynden-Bell and M. Sheppard, Proc. R. Soc. A. <u>269</u>, 1385 (1965). (See Experimental Section).
- Figure 2. <sup>2</sup>H nmr spectra of the ethanes liberated in the reactions between  $Mo_2(CH_2CD_3)_2(NMe_2)_4$  and  $CO_2$  (A) and  $Mo_2(CD_2CH_3)_2(NMe_2)_4$  and  $CO_2$ (B) are compared with spectra associated with commercially supplied samples of  $CD_3CH_2D$  and  $CH_3CD_2H$ , respectively.
- Figure 3. <sup>1</sup>H nmr spectra showing the formation of  $Mo_2Et_2(NMe_2)_2(0_2CNMe_2)_2$  at 100 MHz in the reaction between  $Mo_2Et_2(NMe_2)_4$  and  $CO_2$  in toluene-d<sub>8</sub> at -30°C. The signals denoted by \* correspond to the protio impurities in toluene-d<sub>8</sub>. The signal denoted by \*\* corresponds to the methyl resonance of  $Mo_2(0_2CNMe_2)_4$  which, because of its low solubility, is precipitated as it is formed.
- Figure 4. An ORTEP view of the  $Mo_2Et_2(NMe_2)_2(C_7H_8N_3C_7H_8)_2$  molecule showing the atom numbering scheme used in the tables.
- Figure 5. An ORTEP view of the central skeleton of the  $MO_2Et_2(NMe_2)_2(C_7H_8N_3-C_7H_8)_2$  molecule viewed down the Mo=Mo bond showing the arrangement of the ethyl groups.

# ISOMERS OF DIDEUTEROETHYLENE

-











