

AD-A105 492

BROWN UNIV PROVIDENCE RI LEFSCHETZ CENTER FOR DYNAMI--ETC F/G 12/1
ON A GRADIENT-LIKE INTEGRO-DIFFERENTIAL EQUATION.(U)

JUN 81 J K HALE, K P RYBAKOWSKI

DAAG29-79-C-0161


UNCLASSIFIED

LCDS-81-9

AFOSR-TR-81-0660

NL

1 of 1
AD-A
ON-492



END
DATE
FILMED
-B-
DTIC

The logo consists of a large white circle centered on a black square background. Inside the white circle are five smaller black circles arranged in a pentagonal pattern. The text "Lefschetz Center for Dynamical Systems" is printed in a serif font across the center of the white circle.

Lefschetz Center for Dynamical Systems

OTIC
ELECT
OCT 14

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. AFOSR TR-81-0660		3. RECIPIENT'S CATALOG NUMBER	
2. GOVT ACCESSION NO. ADA105492		4. TYPE OF REPORT & PERIOD COVERED TECHNICAL rept	
5. TITLE (and Subtitle) ON A GRADIENT-LIKE INTEGRO-DIFFERENTIAL EQUATION.		6. PERFORMING ORG. REPORT NUMBER LCDS-81-9	
7. AUTHOR(s) Jack K. Hale and Krzysztof Rybakowski		8. CONTRACT OR GRANT NUMBER(s) AFOSR-76-3092	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Lefschetz Center for Dynamical Systems Division of Applied Mathematics Brown University, Providence RI 02912		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F, 2304/A4	
11. CONTROLLING OFFICE NAME AND ADDRESS Directorate of Mathematical and Information Sciences, Air Force Office of Scientific Research Bolling AFB DC 20332		12. REPORT DATE 1 JUN 1981	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 18	
15. SECURITY CLASS. (of this report) UNCLASSIFIED		15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $b: [-1,0] \rightarrow \mathbb{R}$ be a C^2 -function, $b(\theta) > 0$, $\theta \in (-1,0]$, $b(1) = 0$, $b'(\theta) \geq 0$, $b''(\theta) \geq 0$, $\theta \in [-1,0]$ and there is a θ_0 such that $b''(\theta_0) > 0$. Suppose $g: \mathbb{R} \rightarrow \mathbb{R}$ is a C^1 -function such that $\int_0^x g(s)ds \rightarrow \infty$ as $ x \rightarrow \infty$ and consider the equation (CONT.)			

ITEM #20, CONT.:

$$\dot{x}(t) = - \int_{-1}^0 b(\theta)g(x(t+\theta))d\theta$$

Every solution of this equation approaches a zero of g . If the zeros of g are bounded, there is a maximal compact invariant set $A_{b,g}$ of this equation in $C([-1,0], \mathbb{R})$ which is one dimensional and consists only of the zeros of g and the unstable manifolds of these zeros. If g has only one zero, then $A_{b,g}$ is a point. If g has no more than three simple zeros, then the set $A_{b,g}$ is simply an arc with the unstable zero connected to the stable ones. In the class of g which have five simple zeros, we show that there are five distinct ways that the zeros of g can be connected by orbits in $A_{b,g}$. Only one of these preserves the order of the zeros on the reals. This shows clearly the importance of considering the set $A_{b,g}$ and the structure of the flow on this set rather than just asserting that every solution approaches a zero of g .

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ON A GRADIENT-LIKE INTEGRO-DIFFERENTIAL
EQUATION

by

Jack K. Hale and Krzysztof P. Rybakowski

June 1, 1981

LCDS Report 81-9

DTIC
ELECTE
OCT 14 1981

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

ON A GRADIENT-LIKE INTEGRO-DIFFERENTIAL EQUATION

by

Jack K. Hale* and Krzysztof P. Rybakowski**
Division of Applied Mathematics
Lefschetz Center for Dynamical Systems
Brown University
Providence, R. I. 02912

June 1, 1981

* This research has been supported in part by the Air Force Office of Scientific Research under contract #AFOSR-76-3092D, in part by the Army Research Office under contract #ARO-DAAG-29-79-C-0161 and in part by the National Science Foundation under contract #MCS-79-05774-02.

⁺ This research was also supported by the Deutsche Forschungsgemeinschaft.

1980 Mathematics Subject Classification: 34 K 15.

[illegible]

On a gradient-like integro-differential equation

by

Jack K. Hale and Krzysztof P. Rybakowski

Abstract

Let $b: [-1,0] \rightarrow \mathbb{R}$ be a C^2 -function, $b(\theta) > 0$, $\theta \in (-1,0]$, $b(1) = 0$, $b'(\theta) \geq 0$, $b''(\theta) \geq 0$, $\theta \in [-1,0]$ and there is a θ_0 such that $b''(\theta_0) > 0$. Suppose $g: \mathbb{R} \rightarrow \mathbb{R}$ is a C^1 -function such that $\int_0^x g(s)ds \rightarrow \infty$ as $|x| \rightarrow \infty$ and consider the equation

$$\dot{x}(t) = - \int_{-1}^0 b(\theta) g(x(t+\theta)) d\theta$$

Every solution of this equation approaches a zero of g . If the zeros of g are bounded, there is a maximal compact invariant set $A_{b,g}$ of this equation in $C([-1,0], \mathbb{R})$ which is one dimensional and consists only of the zeros of g and the unstable manifolds of these zeros. If g has only one zero, then $A_{b,g}$ is a point. If g has no more than three simple zeros, then the set $A_{b,g}$ is simply an arc with the unstable zero connected to the stable ones. In the class of g which have five simple zeros, we show that there are five distinct ways that the zeros of g can be connected by orbits in $A_{b,g}$. Only one of these preserves the order of the zeros on the reals. This shows clearly the importance of considering the set $A_{b,g}$ and the structure of the flow on this set rather than just asserting that every solution approaches a zero of g .

1. Introduction

Let $b: [-1,0] \rightarrow \mathbb{R}$ be a C^2 -function such that $b(\theta) > 0$, $\theta \in (-1,0]$, $b(-1) = 0$, $b'(\theta) \geq 0$, $b''(\theta) \geq 0$ for $\theta \in [-1,0]$ and there exists a $\theta_0 \in [-1,0]$ such that $b''(\theta_0) > 0$. Suppose $g: \mathbb{R} \rightarrow \mathbb{R}$ is a C^1 -function and consider the equation

$$(1.1)(b,g) \quad \dot{x}(t) = - \int_{-1}^0 b(\theta)g(x(t+\theta))d\theta$$

For any $\phi \in C = C([-1,0], \mathbb{R})$, there is a unique solution $x(\phi)$ through ϕ at $t = 0$. If $T_f(t): C \rightarrow C$ is defined by $[T_f(t)\phi](\theta) = x(\phi)(t+\theta)$, then $T_f(t)$, $t \geq 0$, is a strongly continuous semigroup on C . We think of the solution $x(\phi)$ of (1.1)(b,g) as defining a curve $\{T_f(t)\phi, t \geq 0\}$ in C and, therefore, can consider the geometric concepts of ω -limit sets, α -limit set and invariant set (see [1]).

Suppose $\int_0^x g(s)ds \rightarrow \infty$ as $|x| \rightarrow \infty$. It is known (see [1,2]) that the ω -limit set of every solution of (1.1)(b,g) is a zero of g and, also, the α -limit set of any nonconstant bounded solution of (1.1)(b,g) is an unstable zero of g . If a is a zero of g then a is hyperbolic if and only if $g'(a) \neq 0$, uniformly asymptotically stable if $g'(a) > 0$ and unstable if $g'(a) < 0$. Furthermore, the unstable manifold $W^u(a)$ of a is one dimensional if a is unstable.

If the set of zeros of g is bounded, then there is a bounded set B such that every solution eventually enters B , that is, (1.1)(b,g) is point dissipative. It follows (see [1,2]) that there is a maximal compact invariant set $A_{b,g}$ for (1.1)(b,g) which is uniformly asymptotically stable and attracts bounded sets of C .

From the fact that the α -limit set of any nonconstant bounded solution is an unstable zero of g , it follows that $A_{b,g} = \bigcup \{W^u(a) : g(a) = 0\}$ and $A_{b,g}$ is one dimensional. The purpose of this paper is to discuss in some detail the structure of the set $A_{b,g}$ for a fixed b and a certain class of g .

Let G_k be the class of all C^1 -functions g satisfying the following conditions:

- 1) $\int_0^x g(s)ds \rightarrow \infty$ as $|x| \rightarrow \infty$.
- 2) g has exactly $2k+1$ zeros $a_1 < a_2 < \dots < a_{2k+1}$ all of which are simple.

Let the topology on G_k be that generated by the seminorms $\|g\|_M = \sup_{x \in M} (|g(x)| + |g'(x)|)$, where M is a compact set in $\mathbb{R}$. For any $g \in G_k$, all zeros of g are hyperbolic and the zeros a_{2j} , $j = 1, 2, \dots, k$, are saddle points with unstable manifolds $W^u(a_{2j})$ one dimensional. Thus, for each a_{2j} , there are exactly two distinct orbits defined for $t \in (-\infty, \infty)$ whose α -limit sets are a_{2j} . We call these orbits emanating from a_{2j} . Fix b as above. Let $g, \tilde{g} \in G_k$ have $a_1 < \dots < a_{2k+1}$ and $\tilde{a}_1 < \dots < \tilde{a}_{2k+1}$, resp., as their zeros. Call g and $\tilde{g}$ equivalent ($g \sim \tilde{g}$) if for all $i, j \in \{1, \dots, 2k+1\}$, there is an orbit $x(t)$ of (1.1)(b, g) emanating from a_i and tending to a_j as $t \rightarrow \infty$ if and only if there is an orbit $\tilde{x}(t)$ of (1.1)($b, \tilde{g}$) emanating from $\tilde{a}_i$ and tending to $\tilde{a}_j$ as $t \rightarrow \infty$. This clearly defines an equivalence relation on G_k . We say $g \in G_k$ is $\sim$ -stable if the equivalence class of g is a neighborhood of g in G_k .

It is not difficult to show that g is $\sim$ -stable if the ω -limit set of every orbit in $A_{b,g}$ which is not a point is a stable zero of g ;

that is, a point a_n , n odd, $1 \leq n \leq 2k+1$. Since $A_{b,g}$ is a global attractor and uniformly asymptotically stable, this is equivalent to saying that g is $\sim$ -stable if the ω -limit set of every orbit of (1.1)(b,g) defined and bounded on $(-\infty, \infty)$ is a stable zero of g . If it were known that the map $T_{b,g}(t)$ is one-to-one on $A_{b,g}$, this latter statement would be equivalent to the following: there is a neighborhood V of g such that, for any $\tilde{g} \in V$, there is a homeomorphism of $A_{b,g}$ onto $A_{b,\tilde{g}}$ which preserves orbits and sense of direction in time; that is, g is structurally stable when the map $T_{b,g}(t)$ is restricted to $A_{b,g}$. We have not been able to prove that $T_{b,g}(t)$ is one-to-one on $A_{b,g}$ and this is the reason for taking the weaker definition of equivalence. If g is analytic, then $T_{b,g}(t)$ is one-to-one (see [1]).

The ultimate objective would be to describe the equivalence classes in G_k . The cases $k = 0, 1$ are trivial. Suppose $k = 2$; that is, each $g \in G_2$ has five zeros $a_1 < a_2 < a_3 < a_4 < a_5$ with a_2, a_4 being saddle points, and a_1, a_3, a_5 being uniformly asymptotically stable. If a_j is an unstable equilibrium point with a_k, a_l being the corresponding ω -limit sets of the orbits emanating from a_j , we designate this by $j[k, l]$. The structure of the flow on $A_{b,g}$ and the equivalence classes in G_2 are then determined by a pair $\{2[1, j], 4[k, l]\}$ expressing the fact that the unstable manifold through a_2 has ω -limit set $\{a_1, a_j\}$ and the one through a_4 has ω -limit set $\{a_k, a_l\}$.

Our main result states there are exactly five equivalence classes in G_2 ; namely $\{2[1, 3], 4[3, 5]\}$, $\{2[1, 4], 4[3, 5]\}$, $\{2[1, 5], 4[3, 5]\}$, $\{2[1, 3], 4[2, 5]\}$, $\{2[1, 3], 4[1, 5]\}$. The only class that preserves the

natural order of the reals on $A_{b,g}$ is $\{2[1,3], 4[3,5]\}$. The first, third and fifth cases are $\sim$ -stable. The second and fourth cases have a connection between the saddle points a_2 and a_4 . It seems plausible that these cases are not $\sim$ -stable, but no proof is available.

The fact that five equivalence classes can occur indicates clearly the importance of studying the structure of the flow on $A_{b,g}$ rather than merely asserting that every solution of (1.1)(b,g) approaches a zero of g .

We have not characterized the equivalence classes in G_k for $k \geq 3$, but it should be possible to adapt the methods below to this case.

2. This section is devoted to the statement and proof of several lemmas.

Lemma 2.1. For every $\rho > 0$, $K > 0$, there is a $C_0 = C_0(K, \rho) > 0$ such that, for all C , $0 < C \leq C_0$, and all a, Δ , $|a| \leq K$, $|a| - 2|\Delta| \geq \rho$, the solution $y(n)$ of the difference equation

$$(2.1) \quad \begin{aligned} y(n) &= y(n-1) - Cy(n-2), \quad n \geq 1 \\ y(0) &= a + \Delta, \quad y(-1) = a \end{aligned}$$

satisfies $\text{sgn } y(n) = \text{sgn } a$ for $n \geq -1$.

Proof: Let $\lambda_1 = \lambda_1(C)$ be the roots of the characteristic equation $\lambda^2 - \lambda + C = 0$. If $0 \leq C < 1/4$, then λ_1, λ_2 are real, nonnegative and distinct, say $\lambda_1(C) < \lambda_2(C)$. Moreover, $\lambda_1(C)$ is continuous in C with $\lambda_1(0) = 0$, $\lambda_2(0) = 1$. For the initial data specified in (1.2) and $n \geq -1$,

$$(\lambda_2 - \lambda_1)y(n) = (a(1 - \lambda_1) + \Delta)\lambda_2^{n+1} + (a(\lambda_2 - 1) - \Delta)\lambda_1^{n+1}$$

The remainder of the argument follows easily from the fact that $\lambda_1(0) = 0$, $\lambda_2(0) = 1$ and the hypotheses on a, Δ .

Lemma 2.2. For every $\rho > 0$, $K > 0$, there is an $m_0 = m_0(\rho, K)$ such that, for all $0 < m \leq m_0$ and $f(x) = mx$, the following properties hold:

1) If $\phi(\theta)$ is continuous, positive and nonincreasing on $[-1, 0]$, $\phi(-1) \leq K$ and $\phi(-1) - 2[\phi(-1) - \phi(0)] \geq \rho$, then the solution $x(t)$ of (1.1)(b, f) through ϕ satisfies $0 < x(t) < \phi(0)$ for $t \in (0, \infty)$ and $x(t) \rightarrow 0$ as $t \rightarrow \infty$ monotonically.

2) If $\phi(\theta)$ is continuous, negative and nondecreasing on $[-1, 0]$, $-\phi(-1) \leq K$ and $-\phi(-1) - 2(\phi(0) - \phi(-1)) \geq \rho$, then the solution $x(t)$ of (1.1)(b, f) through ϕ satisfies $\phi(0) < x(t) < 0$ for $t \in (0, \infty)$ and $x(t) \rightarrow 0$ as $t \rightarrow \infty$ monotonically.

Proof: Fix $\rho > 0$, $K > 0$ arbitrarily. With $C_0 = C_0(\rho, K)$ as in Lemma 2.1, define $m_0 = C_0/M$, $M = \int_{-1}^0 b(\theta) d\theta$. Let $0 < m \leq m_0$ be arbitrary and define $C = mM$. Suppose ϕ satisfies the hypotheses in 1) and let $y(n)$ be the solution of (1.2) with $a = \phi(-1)$, $a + \Delta = \phi(0)$. From Lemma 2.1, $y(n)$ is decreasing, $0 < y(n) < \phi(0)$, $n \geq 1$, $y(n) \rightarrow 0$ as $n \rightarrow \infty$. Let $t_n \geq 0$ be the unique minimal $t_n \geq 0$ such that $x(t_n) = y(n)$ and set $t_{-1} = -1$, $t_0 = 0$. To prove the claim $0 < x(t) < \phi(0)$ in 1), it is sufficient to show that $t_{n-1} - t_{n-2} \geq 1$ for every $n \geq 1$. This inequality is true for $n = 1$. Assuming that it holds for some n , we obtain

$$\begin{aligned} -Cy(n-2) &= y(n) - y(n-1) = x(t_n) - x(t_{n-1}) \\ &= -\int_{t_{n-1}}^{t_n} \left(\int_{-1}^0 b(\theta) mx(s+\theta) d\theta \right) ds \\ &\geq -(t_n - t_{n-1}) Cx(t_{n-1} - 1) \\ &\geq -(t_n - t_{n-1}) Cx(t_{n-2}) = -(t_n - t_{n-1}) Cy(n-2) \end{aligned}$$

This implies $t_n - t_{n-1} \geq 1$ and proves $0 < x(t) < \phi(0)$, $t > 0$. Thus $x(t)$ is decreasing for $t > 0$ and approaches zero as $t \rightarrow \infty$. Case 2) is proved by replacing x by $-x$.

Lemma 2.3. Let α_1, α_2, m_2 and K be real numbers, $\alpha_1 \neq \alpha_2$, $m_2 < 0$, $K > 0$. Let $I = [\alpha_1, \alpha_2]$ if $\alpha_1 < \alpha_2$ and $I = [\alpha_2, \alpha_1]$ if $\alpha_2 < \alpha_1$. Then there is a mapping $f: I \rightarrow \mathbb{R}$, $f \in C^1(I)$, $|f| \leq K$ on I , $f'(\alpha_2) = m_2$, $f^{-1}\{0\} = \{\alpha_1, \alpha_2\}$, f is affine in a neighborhood of α_1 and α_2 and there is a function $x: \mathbb{R} \rightarrow \text{Int } I$, $x \in C^1(\mathbb{R})$, such that $\lim_{t \rightarrow \infty} x(t) = \alpha_1$, $\lim_{t \rightarrow -\infty} x(t) = \alpha_2$, and x satisfies (1.1)(b, f).

Proof: Assume that the lemma is true for $\alpha_1 < \alpha_2$ and suppose $\alpha_2 < \alpha_1$. Let $\bar{\alpha}_1 = \alpha_2, \bar{\alpha}_2 = \alpha_1$. Then $\bar{\alpha}_1 < \bar{\alpha}_2$. Choose $\bar{f} = \bar{f}(\bar{\alpha}_1, \bar{\alpha}_2, m_2)$, $\bar{x} = \bar{x}(\bar{\alpha}_1, \bar{\alpha}_2, m_2)$ as in the statement of the Lemma. Define $f: [\alpha_2, \alpha_1] \rightarrow \mathbb{R}$, $f(x) = -\bar{f}(\alpha_1 + \alpha_2 - x)$ and $x(t) = \alpha_1 + \alpha_2 - \bar{x}(t)$. Then it is obvious that f and x satisfy the Lemma. Hence it is enough to prove the Lemma for $\alpha_1 < \alpha_2$.

Let $f_2(x) = m_2(x - \alpha_2)$. Then by the assumptions on b , there is a unique $\lambda_2 > 0$ such that $x(t) = \alpha_2 - e^{-\lambda_2 t}$ is a solution of (1.1)(b, f_2) on $(-\infty, \infty)$. Let $K = 2(\alpha_2 - \alpha_1)$, $\rho = (1/3)(\alpha_2 - \alpha_1)$. Choose $m_0 = m_0(\rho, K)$ as in Lemma 2.2. For $0 < m \leq m_0$, $f_1(x) = f_1(m)(x) = m(x - \alpha_1)$ let




Figure 1

$\hat{x} = \hat{x}(m)$ be the unique coincidence point of f_1 and f_2 (cf. Fig. 1). Let $\hat{t} = \hat{t}(m)$ be the unique point such that $\alpha_2 - e^{-\lambda_2 \hat{t}} = \hat{x}$. Define $h = h_m: \mathbb{R} \rightarrow \mathbb{R}$ so that $h = f_1$ on $(-\infty, \hat{x}]$, $h = f_2$ on $[\hat{x}, \alpha_2]$ and h is

arbitrary on (α_2, ∞) but such that $h \in G_1$ and the zeros of h are: $\alpha_1 < \alpha_2 < \beta$ where β is some real number. Fix $\epsilon > 0$ and α such that $\alpha_1 < \alpha < \alpha_2 - \epsilon$, and $(\alpha_2 - \epsilon) - \alpha_1 \geq 2\rho + 2(\alpha_2 - \alpha)$.

If $y(t)$, $t \geq 0$, is the solution of (1.1)(b,h) through $\phi(\theta) = \lambda_2(\hat{t} + \theta)$, define $z(t) = \alpha_2 - e^{\lambda_2 t}$ for $t < \hat{t}$, $z(t) = y(t - \hat{t})$ for $t \geq \hat{t}$. It follows that $z(t)$ solves (1.1)(b,h) on $(-\infty, \infty)$. Also, it is clear that there is a unique minimal $t' = t'(m)$ such that $z(t') = \alpha$. For m small enough $\hat{x} > \alpha_2 - \epsilon$ and hence $t' > \hat{t}$. Moreover,

$$z(t') - \hat{x} = z(t') - z(\hat{t}) = - \int_{\hat{t}}^{t'} \left(\int_{-1}^0 b(\theta) h(z(s+\theta)) d\theta ds \right).$$

Since $\hat{x} \geq z(t) \geq \alpha$ for $t \in [\hat{t}, t']$, we have $0 \leq g(z(t)) = m(z(t) - \alpha_1)$ for such t , and this implies

$$|z(t') - \hat{x}| \leq M \cdot m(t' - \hat{t})(\hat{x} - \alpha_1),$$

where $M = \int_{-1}^0 b(\theta) d\theta$. Hence,

$$0 < \alpha_2 - \epsilon - \alpha < (\hat{x} - z(t')) \leq M \cdot m(t' - \hat{t})(\hat{x} - \alpha_1),$$

i.e., $t' - \hat{t} \rightarrow \infty$ as $m \rightarrow 0$. Hence, for all m small enough, and all $t \in [\hat{t}, \hat{t} + 1]$, $\alpha < z(t) < \alpha_2$. Moreover, taking m smaller, if necessary,

$$\begin{aligned} (2.2) \quad & z(\hat{t} - 1) - \alpha_1 - 2(z(\hat{t} - 1) - z(\hat{t})) \\ & = \alpha_2 - e^{\lambda_2(\hat{t} - 1)} - \alpha_1 - 2(e^{\lambda_2(\hat{t} - 1)} - e^{\lambda_2 \hat{t}}) > (1/2)(\alpha_2 - \alpha_1). \end{aligned}$$

Fix $m \leq K$ so that (2.2) is satisfied. For any $\delta > 0$ let $g_\delta(x)$ be a C^1 -function such that $g_\delta(x) = h(x)$ for $x \notin (\hat{x} - \delta, \hat{x} + \delta)$ and

$0 \leq h(x) - g_\delta(x) < \delta$ for $x \in \mathbb{R}$. If $s' = s'(\delta)$ is such that $\alpha_2 - e^{\lambda_2 s'} = \hat{x} + \delta$, then $s'(\delta) \rightarrow \hat{t}$ as $\delta \rightarrow 0$. If $\phi_\delta(\theta) = \alpha_2 - e^{\lambda_2(s'(\delta)+\theta)}$, then $\phi_\delta \rightarrow \phi$ as $\delta \rightarrow 0$ in $C([-1,0], \mathbb{R})$. Let $y_\delta(t)$, $t \geq 0$, be the solution of (1.1)(b, g_δ) through ϕ_δ and define $z_\delta(t) = \alpha_2 - e^{\lambda_2 t}$, $t < s'(\delta)$, $z_\delta(t) = y_\delta(t - s'(\delta))$, $t \geq s'(\delta)$. Then z_δ satisfies (1.1)(b, g_δ) on $(-\infty, \infty)$. By continuity, $z_\delta(t) \rightarrow z(t)$ as $\delta \rightarrow 0$, uniformly on compact intervals. Let $s'' = s''(\delta)$ be the smallest number such that $z_\delta(s''(\delta)) = \hat{x} - \delta$. Then $s''(\delta)$ exists and $s''(\delta) > s'(\delta)$. Moreover, since $z(\hat{t}+1) < \hat{x}$, it follows that $s''(\delta) < s'(\delta) + 1$ for $\delta > 0$ small enough. Hence, $s''(\delta) \rightarrow \hat{t}$ as $\delta \rightarrow 0$. For $\delta > 0$ small, it follows that $z_\delta(s''(\delta)+\theta) - \alpha_1$ is positive and nonincreasing. Moreover, $z_\delta(s''(\delta)-1) - \alpha_1 < \alpha - \alpha_2 < K$. Finally, by (2.2) and for δ small enough,

$$z_\delta(s''(\delta)-1) - \alpha_1 - 2(z_\delta(s''(\delta)-1) - z_\delta(s''(\delta))) \geq \rho.$$

Fixing such a small $\delta > 0$ and letting $f = g_\delta$ on $[\alpha_1, \alpha_2]$, $x(t) = z_\delta(t)$, we see from Lemma 2.2 that $\lim_{t \rightarrow -\infty} x(t) = \alpha_1$, $\lim_{t \rightarrow \infty} x(t) = \alpha_2$. This proves the lemma.

Corollary 2.4. Let $a_1 < \dots < a_{2k+1}$ be arbitrary real numbers, and, for every $i = 1, \dots, k$, let m_{2i} be any negative number. Let $M > 0$ be arbitrary. Then there exists a $g \in G_k$, such that $g(a_1) = 0$ for $i = 1, \dots, 2k+1$, $g'(a_{2i}) = m_{2i}$ for $i = 1, \dots, k$, g is affine on $(-\infty, a_1) \cup (a_{2k+1}, \infty)$ and in a neighborhood of each a_i , $|g(x)| \leq M$ for $x \in [a_1, a_{2k+1}]$. Moreover, for every $i = 1, \dots, k$ there are solutions $x_i(t)$ and $\bar{x}_i(t)$ of (1.1)(b, g) on $(-\infty, \infty)$ such that $a_{2i-1} < x_i(t) < a_{2i}$,

$$a_{2i} < \bar{x}_i(t) < a_{2i+1}, t \in \mathbb{R}, x_i(+\infty) = a_{2i-1}, x_i(-\infty) = a_{2i}, \bar{x}_i(+\infty) = a_{2i+1}, \\ \bar{x}_i(-\infty) = a_{2i}.$$

We also need some lemmas on approximation of elements in G_k by analytic functions.

Lemma 2.5. For every $g \in G_k$ and every positive, continuous function $\phi: \mathbb{R} \rightarrow \mathbb{R}$ there exists an analytic function $h \in G_k$ such that

$$|g(x) - h(x)| + |g'(x) - h'(x)| < \phi(x), \quad x \in \mathbb{R}.$$

Proof: Let $a_1 < \dots < a_{2k+1}$ be the zeros of g . There are numbers $b_1^-, b_1^+, i = 1, \dots, 2k+1, b_i^- < a_i < b_i^+$ such that $g(x) < 0$ for $x \leq b_1^-$, $g(x) > 0$ for $x \geq b_{2k+1}^+$, $g'(x) \neq 0$ for $x \in [b_i^-, b_i^+]$, and the intervals $[b_i^-, b_i^+]$ are pairwise disjoint.

Let $\psi: \mathbb{R} \rightarrow \mathbb{R}$ be a positive, continuous, function such that $\psi(x) < \phi(x)$ for $x \in \mathbb{R}$, ψ is integrable, $\psi(x) < \frac{1}{2}|g(x)|$ for $x \notin \bigcup_{i=1}^{2k+1} (b_i^-, b_i^+)$, $\psi(x) < \frac{1}{2}|g'(x)|$ for $x \in \bigcup_{i=1}^{2k+1} [b_i^-, b_i^+]$. By Whitney's Lemma ([3]), there is an analytic function $h: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$|g(x) - h(x)| + |g'(x) - h'(x)| < \psi(x) \quad \text{for } x \in \mathbb{R}.$$

We will show that $h \in G_k$. If $H(x) = \int_0^x h(s)ds$, $G(x) = \int_0^x g(s)ds$, then

$$\begin{aligned} H(x) &= \int_0^x g(s)ds + \int_0^x (h(s) - g(s))ds \\ &\geq G(x) - \int_0^x |h(s) - g(s)|ds \\ &\geq G(x) - \int_0^x \psi(s)ds \geq G(x) - \int_{\mathbb{R}} \psi(s)ds \rightarrow \infty \end{aligned}$$

as $|x| \rightarrow \infty$.

If $x \notin \bigcup_{i=1}^{2k+1} (b_i^-, b_i^+)$, then $|g(x) - h(x)| < \psi(x) < \frac{1}{2}|g(x)|$ and, hence,

$h(x) \neq 0$. It follows that all zeros of h must be contained in $\bigcup_{i=1}^{2k+1} (b_i^-, b_i^+)$. If $x \in \bigcup_{i=1}^{2k+1} [b_i^-, b_i^+]$, then $|g'(x) - h'(x)| < \psi(x) < \frac{1}{2}|g'(x)|$ and, hence, $h'(x) \neq 0$. This means that h has at most one zero in $[b_i^-, b_i^+]$. Since $\text{sign } h(b_i^-) = \text{sign } g(b_i^-) \neq \text{sign } g(b_i^+) = \text{sign } h(b_i^+)$, it follows that h has exactly one zero in $[b_i^-, b_i^+]$ and this zero is simple. The Lemma is proved.

Lemma 2.6. Let $a_1 < \dots < a_n$ and $\tilde{a}_1 < \dots < \tilde{a}_n$ be real numbers.

Then there are $q, M, 0 < q < M$ and there is an analytic mapping

$\alpha: \mathbb{R} \rightarrow \mathbb{R}$ such that α is a diffeomorphism of $\mathbb{R}$ onto $\mathbb{R}$, $q < \alpha'(x) < M$ for $x \in \mathbb{R}$, $\alpha(a_i) = \tilde{a}_i$, $i = 1, \dots, n$.

Proof: Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a C^1 -function such that $f'(x) > 0$ for $x \in \mathbb{R}$, f is affine on $(-\infty, a_1] \cup [a_n, \infty)$ and $f(a_i) = \tilde{a}_i$, $i = 1, \dots, n$. Let $\gamma = \inf_{x \in \mathbb{R}} f'(x) > 0$ and $v_k(x) = \prod_{i \neq k} (x - a_i)$, $k = 1, \dots, n$. By a simple calculation,

$$\begin{aligned} M &:= n \sup_k \{(\tanh(v_k(a_k)))^{-1}\} + 1 \\ &+ \sup_k \sup_{x \in \mathbb{R}} \left\{ (\tanh(v_k(a_k)))^{-1} \cdot \frac{d(\tanh \circ v_k)(x)}{dx} \right\} < \infty. \end{aligned}$$

By Whitney's Lemma, there is an analytic function $h: \mathbb{R} \rightarrow \mathbb{R}$ such that for $x \in \mathbb{R}$,

$$|h(x) - f(x)| + |h'(x) - f'(x)| < \gamma/3\tilde{M}.$$

If $\alpha = h + k$, where

$$k(x) = \sum_{k=1}^n \tanh(v_k(x)) \cdot \frac{f(a_k) - h(a_k)}{\tanh(v_k(a_k))}$$

then α is analytic,

$$|\alpha(x) - f(x)| \leq |h(x) - f(x)| + |k(x)| \leq \gamma/3\tilde{M} + \gamma/3.$$

Hence $\alpha(x) \rightarrow \pm\infty$ as $x \rightarrow \pm\infty$ and α is surjective. Also,

$$|\alpha'(x) - f'(x)| \leq |h'(x) - f'(x)| + |k'(x)| \leq \gamma/3\tilde{M} + \gamma/3$$

which implies $\sup_{x \in \mathbb{R}} |\alpha'(x)| < \infty$. Furthermore,

$$\frac{\gamma}{3} \leq f'(x) - \frac{\gamma}{3\tilde{M}} - \frac{\gamma}{3} \leq \alpha'(x).$$

Thus, $q < \alpha'(x) < M$, $x \in \mathbb{R}$ and α is a diffeomorphism. Finally, it is obvious that $\alpha(a_i) = \tilde{a}_i$, $i = 1, \dots, n$, and the lemma is proved.

Corollary 2.7. Let g and $\tilde{g}$ be two analytic functions belonging to G_k . Then there is a continuous mapping $H: [0,1] \rightarrow G_k$, such that $H(0) = \tilde{g}$, $H(1) = g$, and, for every $t \in [0,1]$, $H(t)$ is analytic.

Proof: Let $a_1 < \dots < a_n$ and $\tilde{a}_1 < \dots < \tilde{a}_n$ be the zeros of g and $\tilde{g}$, resp. Let α be as in Lemma 2.6. Define for $t \in [0,1]$ and $x \in \mathbb{R}$

$$(H(t))(x) = \begin{cases} \tilde{g}((1-2t)x + 2t\alpha(x)) & , \text{ if } t \leq 1/2 \\ (2-2t)\tilde{g}(\alpha(x)) + (2t-1)g(x), & \text{ if } t > 1/2 \end{cases}$$

It is easy to show that H satisfies all requirements of the Corollary.

3. The main result.

In this section, we state and prove the main result.

Lemma 3.1. Let $g \in G_k$ and a be an unstable zero of g . If $x(t)$, $t \in (-\infty, \infty)$, is a solution of (1.1)(b,g) emanating from a , then either $x(t) > a$ for all t or $x(t) < a$ for all t .

Lemma 3.1 gives a limitation on the maximal number of equivalence classes that can occur in G_k . The remainder of the discussion is concerned with $k = 2$. There are at most five equivalence classes in G_2 ; namely, those described in the introduction. The main result is

Lemma 3.2. There is an analytic function in each of the equivalence classes $\{2[1,5], 4[3,5]\}$, $\{2[1,3], 4[1,5]\}$ in G_2 .

Theorem 3.3. There are exactly five equivalence classes in G_2 . Each equivalence class contains an analytic function.

Proof of Lemma 3.1:

$$\text{Let } V(\phi) = G(\phi(0)) + \frac{1}{2} \int_{-1}^0 b'(\theta) \left[\int_{\theta}^0 g(\phi(s)) ds \right]^2 d\theta,$$

$$\text{where } G(x) = \int_0^x g(s) ds.$$

Then the derivative $\dot{V}(\phi)$ of V along solutions of (1.1) (b,g) is given by

$$(3.1) \quad \dot{V}(\phi) = -\frac{1}{2} b'(-1) \left[\int_{-1}^0 g(\phi(\theta)) d\theta \right]^2 - \frac{1}{2} \int_{-1}^0 b''(\theta) \left[\int_{\theta}^0 g(\phi(s)) ds \right]^2 d\theta,$$

(see [1]).

Since $\lim_{t \rightarrow -\infty} x(t) = a$, it follows that $\lim_{t \rightarrow -\infty} V(x_t) = G(a)$.

If the Lemma is not true, there exists a $t_0 > 0$ such that $x(t_0) = a$. By (3.1), $V(x_{t_0}) < G(a)$. Hence $\frac{1}{2} \int_{-1}^0 b'(\theta) \left[\int_{\theta}^0 g(\phi(s)) ds \right]^2 d\theta < 0$, a contradiction which proves the Lemma.

Proof of Lemma 3.2: We will show that there is a $g \in G_2$ representing class $\{2[1,5], 4[3,5]\}$. Symmetry in the proof to follow implies that there is a $g' \in G_2$ representing class $\{2[1,3], 4[1,5]\}$. Since these classes are stable classes, Lemma 2.5 implies that g and g' can be chosen analytic, which proves the lemma. Choose $a_1 < a_2$ arbitrarily. Define $f \in C^1$ such that f has exactly two zeros, a_1 and a_2 , f is affine in a neighborhood of a_1, a_2 and on $(-\infty, a_1) \cup (a_2, \infty)$, $f'(a_1) > 0$, $f'(a_2) = m_2 < 0$. There is a unique $\lambda_2 > 0$ such that $x(t) = a_2 + e^{\lambda_2 t}$, $t \in (-\infty, \infty)$ is a solution of (1.1)(b,f). Let $t_0 > 0$ be arbitrary and let $y(t)$, $t \in [0,1]$, be defined as




Figure 2

$$y(t) = x(t_0) - \int_0^t \left(\int_{-1}^{-s} b(\theta) f(x(s+\theta)) d\theta \right) ds$$

Hence, there is a $0 < t_1 < 1$ such that $\dot{y}(t) > 0$ for $t \in [0, t_1]$ and, so, $y(t_1) > x(t_0)$. Define $\tilde{f}$ such that $\tilde{f} = f$ on $(-\infty, x(t_0)]$, $\tilde{f}$ is affine on $[x(t_0), y(t_1)]$, $\tilde{f}(y(t_1)) = 0$, and $\tilde{f}$ is affine on $[y(t_1), \infty)$ with negative slope. If $\phi(\theta) = x(t_0 + \theta)$, $\theta \in [-1, 0]$, let $\tilde{y}$ be the solution of (1.1)(b, $\tilde{f}$) through ϕ , $t \geq 0$. Then obviously $\tilde{y}(t) > y(t)$ for $t \in [0, t_1]$. If we define $\tilde{x}(t) = x(t)$, $t \leq t_0$, $\tilde{x}(t) = \tilde{y}(t - t_0)$, $t > t_0$, then $\tilde{x}$ satisfies (1.1)(b, $\tilde{f}$) on $(-\infty, \infty)$. Moreover, there is an $s_1 > s_0$ such that $\tilde{x}(s_1 + \theta) > y(t_1)$ for $\theta \in [-1, 0]$, and $\tilde{x}(s_1 + \theta)$ is nondecreasing on $[-1, 0]$. Now we perturb $\tilde{f}$ a little to obtain a C^1 -map

h which coincides with $\tilde{f}$ on $[y(t_1), \infty)$, and h has four simple zeros, $a_1 < a_2 < a_3 < a_4$, where $a_3 \in (x(t_0), y(t_1))$, $a_4 = y(t_1)$ and there is a solution $z(t)$ of (1.1)(b, h) on $(-\infty, \infty)$ such that $z(t) = a_2$ as $t \rightarrow -\infty$ and, for some $s_2 \in \mathbb{R}$, $z(s_2 + \theta) > y(t_1)$, $\theta \in [-1, 0]$, $z(s_2 + \theta)$ is non-decreasing on $[-1, 0]$ (cf. Fig. 2). Now the application of Lemma 2.2 and the argument from the proof of Lemma 2.3 easily completes the proof of the theorem.

Proof of Theorem 3.3: Let g be an analytic function representing class $\{2[1,5], 4[3,5]\}$. By Corollary 2.4, there exists an $f \in G_2$ representing class $\{2[1,3], 4[3,5]\}$. Since the class $\{2[1,3], 4[3,5]\}$ is $\sim$ -stable, we infer from Lemma 2.5 that there is an analytic function $\tilde{g} \in G_2$ representing $\{2[1,3], 4[3,5]\}$. By Corollary 2.7, there is a continuous map $H: [0,1] \rightarrow G_2$ such that $H(0) = \tilde{g}$, $H(1) = g$ and $H(s)$ is an analytic function for every fixed s . Let $a_1(s) < a_2(s) < a_3(s) < a_4(s) < a_5(s)$ be the zeros of $H(t)$. It follows that $a_i(s)$ is continuous for $i = 1, \dots, 5$.

Let $s_0 = \sup\{s \in [0,1]: \text{the solution } x(s', t), t \in (-\infty, \infty) \text{ of (1.1)(b, } H(s')) \text{ emanating from } a_2(s') \text{ and staying to the right of } a_2(s') \text{ is such that } \lim_{t \rightarrow \infty} x(s', t) = a_3(s'), \text{ for every } 0 \leq s' \leq s\}$. It follows easily that $0 < s_0 < 1$ and that $\lim_{t \rightarrow \infty} x(s_0, t) \notin \{a_3(s_0), a_5(s_0)\}$. Hence $\lim_{t \rightarrow \infty} x(s_0, t) = a_4(s_0)$, i.e., $H(s_0)$ represents the saddle connection $\{2[1,4], 4[3,5]\}$. By using g' instead of g where g' is analytic and represents class $\{2[1,3], 4[1,5]\}$, we analogously prove that there is an analytic function $\in G_2$ representing the saddle connection $\{2[1,3], 4[2,5]\}$. This completes the proof of Theorem 3.3.

References

- [1] J. K. Hale, Theory of Functional Differential Equations, Springer-Verlag, 1977.
- [2] J. K. Hale, Asymptotic behavior of an integro-differential equation, preprint.
- [3] H. Whitney, Analytic extensions of differentiable functions defined on closed sets, Trans. Amer. Math. Soc., 36(1934), 369-387.

DATE
FILMED
-8