AD-A119	005	AIR FOR AN ANAL MAY 82	RCE INS	T OF TE	CH WRIG Radiat	HT-PATT ION DAT	ERSON /	FB OH	LINS,	F	/6 4/1 0,(U)		18
UNCLASS	Algons		I NK7 82	-301									
									END DATE FILMED IO ,82 DTIC				

REPORT DOCUMENT	ATION PAGE	BEFORE COMPLETING FORM
AFIT/CI/NR/82-30T	2. GOVT ACCESSION NO	3. RECIPIENT'S CATALOG NUMBER
. TITLE (and Subtitie)	VIV-1117005	5. TYPE OF REPORT & PERIOD COVERED
An Analysis of Solar Radiation	Data for Fort	THESIS/DASSERTAT/JON
coronado	U	6. PERFORMING ORG. REPORT NUMBER
· AUTHOR(+)		B. CONTRACT OR GRANT NUMBER(a)
John Robert Conley (Capt)		
PERFORMING ORGANIZATION NAME AND	ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
AFIT STUDENT AT: Colorado Si	tate University	
CONTROLLING OFFICE NAME AND ADDRE	ESS	12. REPORT DATE
AFIT/NR	· · ·	May_1982
WPAFB OH 45433		13. NUMBER OF PAGES 37
4. MONITORING AGENCY NAME & ADDRESS	If different from Controlling Office)	15. SECURITY CLASS. (of this report)
		UNCLASS
		154. DECLASSIFICATION/DOWNGRADING
		1 SCHEDULE
6. DISTRIBUTION STATEMENT (of this Report APPROVED FOR PUBLIC RELEASE; 7. DISTRIBUTION STATEMENT (of the abatrac) DISTRIBUTION UNLIMITED t entered in Block 20, 11 different fro	an Report)
5. DISTRIBUTION STATEMENT (of this Report APPROVED FOR PUBLIC RELEASE; I 7. DISTRIBUTION STATEMENT (of the abatrac) DISTRIBUTION UNLIMITED t entered in Block 20, 11 different fro	m Report)
5. DISTRIBUTION STATEMENT (of this Report APPROVED FOR PUBLIC RELEASE; I 7. DISTRIBUTION STATEMENT (of the abatrac 9. SUPPLEMENTARY NOTES) DISTRIBUTION UNLIMITED t entered in Black 20, if different fro	m Report)
APPROVED FOR PUBLIC RELEASE; I DISTRIBUTION STATEMENT (of the abatrac DISTRIBUTION STATEMENT (of the abatrac S. SUPPLEMENTARY NOTES) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17	en Report)
APPROVED FOR PUBLIC RELEASE; I O DISTRIBUTION STATEMENT (of the abatrac Supplementary notes APPROVED FOR PUBLIC RELEASE:) DISTRIBUTION UNLIMITED t entered in Block 20, 11 different fro IAW AFR 190-17	ETAN E. WOLAVER Dean for Research and Professional Development
5. DISTRIBUTION STATEMENT (of this Report APPROVED FOR PUBLIC RELEASE; I 7. DISTRIBUTION STATEMENT (of the abatrac 8. SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE:	DISTRIBUTION UNLIMITED t entered in Block 20, 11 different fro IAW AFR 190-17 30 AUG 1982	ETAX E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
APPROVED FOR PUBLIC RELEASE; I DISTRIBUTION STATEMENT (of the abatrac D. DISTRIBUTION STATEMENT (of the abatrac S. SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE:) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by block number)	TWN E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
APPROVED FOR PUBLIC RELEASE; I OISTRIBUTION STATEMENT (of the abatrac OISTRIBUTION STATEMENT (of the abatrac SUPPLEMENTARY NOTES NPPROVED FOR PUBLIC RELEASE: KEY WORDS (Continue on reverse aide if nec) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 30 AUG 1982 esseary and identify by block number)	TReport)
APPROVED FOR PUBLIC RELEASE; I DISTRIBUTION STATEMENT (of the abetrac DISTRIBUTION STATEMENT (of the abetrac SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE: KEY WORDS (Continue on reverse aide 11 nec) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 30 AUG 1982 eccentry and identify by block number)	TAR Report)
ABSTRACT (Continue on reverse side if nece) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> esseary and identify by block number)	TAR Report)
ABSTRACT (Continue on reverse side if nece) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by block number))	TReport) TYM E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
ABSTRACT (Continue on reverse side if nece ATTACHED) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by block number) >	The Report) ETAN E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
ABSTRACT (Continue on reverse side if nece ABSTRACT (Continue on reverse side if nece	DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by block number) presery and identify by block number)	an Report) ETAN E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
APPROVED FOR PUBLIC RELEASE; I DISTRIBUTION STATEMENT (of the abatrac DISTRIBUTION STATEMENT (of the abatrac S. SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE: KEY WORDS (Continue on reverse aide if nece ABSTRACT (Continue on reverse aide if nece ABSTRACT (Continue on reverse aide if nece) DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by block number) >	SCHEDULE SCHEDU
APPROVED FOR PUBLIC RELEASE; I DISTRIBUTION STATEMENT (of the abetrac S. DISTRIBUTION STATEMENT (of the abetrac S. SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE: . KEY WORDS (Continue on reverse aide if nece . ABSTRACT (Continue on reverse aide if nece . TTACHED) DISTRIBUTION UNLIMITED t entered in Black 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essary and identify by black number) }	TYNN E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB
APPROVED FOR PUBLIC RELEASE; I S. DISTRIBUTION STATEMENT (of the abstrac D. DISTRIBUTION STATEMENT (of the abstrac D. SUPPLEMENTARY NOTES APPROVED FOR PUBLIC RELEASE: ABSTRACT (Continue on reverse side if nece ABSTRACT (Continue on reverse side if nece ABSTRACT (Continue on reverse side if nece ABSTRACT (Continue on reverse side if nece	DISTRIBUTION UNLIMITED t entered in Block 20, if different fro IAW AFR 190-17 <u>30 AUG 1982</u> essery and identify by block number) > Preary and identify by block number)	an Report) ETMN E. WOLAVER Dean for Research and Professional Development AFIT, Wright-Patterson AFB

ABSTRACT

82~ 30

Work with solar radiation reaching the earth's surface frequently requires a knowledge of the extraterrestrial solar radiation. A method of calculating extraterrestrial solar radiation over various time scales (i.e. instantaneous, hourly, daily, etc.) with known errors is detailed.

Solar radiation data for Fort Collins, Colorado, for the years 1977-1980 are examined to determine extended periods of low surface solar radiation. During the 58 days from 5 January to 5 March 1978 the normalized direct component is shown to have been less than 0.15 for three separate periods totaling 29 days.

From this same data set correlation equations between the hemispheric/extraterrestrial and diffuse/extraterrestrial ratios are developed, allowing prediction of the diffuse component, given the hemispheric, for this high altitude, dry environment.

Primary sources: Liu, B. Y. H., R. C. Jordan, 1960: The interrelationship and characteristic distribution of direct, diffuse, and total radiation. Solar Energy, IV,1-9.

Orgill, J.F., K. G. T. Hollands, 1976: Correlation equation for hourly diffuse radiation on a horizontal surface. Solar Energy, 19, 325-329.

Thompson, T. M., 1981: Pyrheliometer observations as an indication of the climatological persistance of clouds. Technical Memo NOAA ERL ARL-97. Air Resources Laboratory, Silver Spring, MD, Feb., 78pp. Author: John Robert Conley Title: An Analysis of Solar Radiation Data for Fort Collins, Colorado Rank: Captain Branch: U.S. Air Force Date: 1982 Number of Pages: 37 Degree Awarded: Master of Science

Institution: Colorado State University

-



AFIT/CI/NR/82-30T

AFIT RESEARCH ASSESSMENT

The purpose of this questionnaire is to ascertain the value and/or contribution of research accomplished by students or faculty of the Air Force Institute of Technology (ATC). It would be greatly appreciated if you would complete the following questionnaire and return it to:

AFIT/NR

Wright-Patterson AFB OH 45433

RESEARCH TITLE: An Analysis of Solar Radiation Data for Fort Collins, Colorado

AUTHOR: <u>Capt John Robert Conley</u> RESEARCH ASSESSMENT QUESTIONS:

1. Did this research contribute to a current Air Force project?

() a. YES

() b. NO

2. Do you believe this research topic is significant enough that it would have been researched (or contracted) by your organization or another agency if AFIT had not?

() a. YES

() b. NO

3. The benefits of AFIT research can often be expressed by the equivalent value that your agency achieved/received by virtue of AFIT performing the research. Can you estimate what this research would have cost if it had been accomplished under contract or if it had been done in-house in terms of manpower and/or dollars?

() a. MAN-YEARS

() b. \$____

4. Often it is not possible to attach equivalent dollar values to research, although the results of the research may, in fact, be important. Whether or not you were able to establish an equivalent value for this research (3. above), what is your estimate of its significance?

() a. HIGHLY () b. SIGNIFICANT () c. SLIGHTLY () d. OF NO SIGNIFICANT SIGNIFICANT SIGNIFICANCE

5. AFIT welcomes any further comments you may have on the above questions, or any additional details concerning the current application, future potential, or other value of this research. Please use the bottom part of this questionnaire for your statement(s).

NAME

GRADE

LOCATION

POSITION

ORGANIZATION

STATEMENT(s):

AN ANALYSIS OF SOLAR RADIATION DATA FOR FORT COLLINS, COLORADO

Submitted by John R. Conley Department of Atmospheric Science

In partial fulfillment of the requirements for the Degree of Master of Science Colorado State University Fort Collins, Colorado Summer, 1982

ABSTRACT

Work with solar radiation reaching the earth's surface frequently requires a knowledge of the extraterrestrial solar radiation. A method of calculating extraterrestrial solar radiation over various time scales (i.e. instantaneous, hourly, daily, etc.) with known errors is detailed.

Solar radiation data for Fort Collins, Colorado, for the years 1977-1980 are examined to determine extended periods of low surface solar radiation. During the 58 days from 5 January to 5 March 1978 the normalized direct component is shown to have been less than 0.15 for three separate periods totaling 29 days.

From this same data set correlation equations between the hemispheric/extraterrestrial and diffuse/extraterrestrial ratios are developed, allowing prediction of the diffuse component, given the hemispheric, for this high altitude, dry environment.

> John R. Conley Atmospheric Science Department Colorado State University Fort Collins, Colorado 80523 Summer, 1982

iii

ACKNOWLEDGEMENTS

The author would like to thank Mr. John Kleist for his aid and guidance in computer programming, Ms. Odilia Panella for her help in preparing the manuscript, and Dr. Stephen K. Cox for his many helpful suggestions. My special thanks go to Dr. Thomas B. McKee, my mentor, for helping to keep this research on track.

TABLE OF CONTENTS

N-30

	Pa	ge
ABSTR	RACT	ii
ACKNO	DWLEDGEMENTS	iv
LIST	OF FIGURES	vi
LIST	OF TABLES	ii
LIST	OF SYMBOLS	ii
I.	INTRODUCTION	1
11.	CALCULATION OF EXTRATERRESTRIAL SOLAR RADIATION	3
	A. The Method	4
	B. The Computational Procedure	7
	C. Accuracy of the Method	8
III.	ANALYSIS OF FORT COLLINS SOLAR RADIATION DATA 1	0
	A. Data Handling	0
	B. Results of Analysis	2
	C. Relationships Between Solar Radiation Components 2	1
IV.	SUMMARY AND CONCLUSIONS	7
	REFERENCES	9
	APPENDIX	0

v

LIST OF FIGURES

Figure	Pag	e
1.	Time series plots of average daylight hours cloudiness, average daily direct/direct extraterrestrial ratios, and average daily hemispheric/extraterrestrial ratios for 5 January to 5 March 1978 2	20
2.	Correlation equations between daily hemispheric/ extraterrestrial and diffuse/extraterrestrial ratios for means, 80%, and 20% confidence levels 2	24
3.	Comparison of the hemispheric/extraterrestrial to diffuse/extraterrestrial correlations of the present study with those of Liu and Jordan (1960) for Blue Hill, MA and Orgill and Hollands (1976) for Toronto Airport, Canada 2	26

.....

LIST OF TABLES

Table		Page
1.	Frequency distribution of the daily Diffuse/ Hemispheric ratio	13
2.	Frequency distribution of the daily Diffuse/ Extraterrestrial ratio	14
3.	Frequency distribution of the daily Hemispheric/ Extraterrestrial ratio	15
4.	Frequency distribution of the Direct/ Direct extraterrestrial ratio	16

LIST OF SYMBOLS

7 1

ANGVEL	Angular velocity of the earth.
DIF	diffuse component of the incoming solar radiation.
DIF/ETR	dimensionless ratio of the diffuse component and the extraterrestrial solar radiation.
DIF/HEM	dimensionless ratio of the diffuse and hemispheric components of incoming solar radiation.
DIR	direct component of the incoming solar radiaion.
DIRETR	normal component of the extraterrestrial solar radiation.
DIR/DIRETR	dimensionless ratio of the direct component and the nor- mal component of the extraterrestrial solar radiation.
ET	number of days from an established base date.
ETR	extraterrestrial solar radiation.
GMT	Greenwich mean time.
h	local hour angle.
HEM	hemispheric component of the incoming solar radiation.
hr	hour.
HEM/ETR	dimensionless ratio of the hemispheric component and the extraterrestrial solar radiation.
m	meter.
NOON	local noon in Greenwich mean time.
Q	instantaneous extraterrestrial solar radiation.
Q ₁	extraterrestrial solar radiation integrated over some time interval.
R	distance between the sun and the earth.

viii

Ř	annual mean distance between the sun and the earth.
S	solar constant.
Т	equation of time.
TIME	mid-point of an integration interval.
W	watt.
Z	solar zenith angle.
α	time parameter for calculating the relative sun~earth distance.
δ	solar declination.
ρ	time parameter for calculating solar declination and the equation of time.
φ	latitude.
ψ	longitude.

ix

CHAPTER I

INTRODUCTION

The emphasis in recent years concerning alternate sources of energy has resulted in a large body of research concerning the solar radiation received at the earth's surface. This surface solar radiation, termed insolation or hemispheric radiation, is comprised of two components; the direct and diffuse. The hemispheric radiation, variously called the total or total hemispheric, is the total irradiance received from the upward hemisphere. The direct, variously termed direct beam or simply beam, component is that irradiance received directly from the sun (usually within a field of view of about 5°). The direct irradiance multiplied by the cosine of the solar zenith angle gives the contribution to the hemispheric irradiance from the direct component. The remainder of the hemispheric irradiance, which comes from all angles in the upward hemisphere not contained in the direct measurement, is called the diffuse component. One final definition is required. The extraterrestrial solar radiation is that irradiance available at the top of the atmosphere on a surface normal to the local vertical. The value of this extraterrestrial irradiance is affected by variations in the sun-earth geometry. These effects include an annual variation of about $\pm 3.3\%$ in the sun-earth distance due to the ellipticity of the earth's orbit, the inclination of the earth's axis, and the rotation of the earth about

its axis. These effects are predictable and a method of calculating the extraterrestrial solar radiation is detailed in Chapter II.

Liu and Jordan (1960) pioneered research in determining the relationship between the hemispheric and the diffuse component. Their work was based primarily on data measured at Blue Hill, Massachusetts. Since this initial study, many authors have investigated these relationships for various other locations. Most notable among these studies are those by Ruth and Chant (1975) for selected Canadian locations, Orgill and Hollands (1976) for Toronto, Canada, and Collares-Pereira and Rabl (1978) for locations in New Mexico, Texas, California, and Massachusetts. Chapter III of this paper will deal with the diffuse-hemispheric relationship based on data measured at Fort Collins, Colorado -- a high altitude, dry environment.

CHAPTER II

CALCULATION OF EXTRATERRESTRIAL SOLAR RADIATION

Normalization of the hemispheric irradiance and its components is accomplished by forming the dimensionless ratio of the hemispheric and/or a given component to an appropriate value of the extraterrestrial solar radiation (ETR). It should be pointed out, however, that the direct component should be normalized by that irradiance available at the top of the atmosphere on a surface normal to the sun's rays. For the purposes of this study, this modified ETR is termed the direct ETR (DIRETR). The purpose of this normalization procedure is to eliminate the dependence of the surface values on solar position. This is especially important whenever comparisons are to be made for different times (i.e. one hour to another, one day to another, etc.) or for different locations.

Sellers (1965) gives procedures for calculating instantaneous and daily extraterrestrial solar radiation values and a graphical method for determining daily and monthly values. Guttman and Mathews (1979) outline a method for calculating values for variable time periods of an hour up to one day. The remainder of this chapter will deal with a method of calculating extraterrestrial solar radiation over various time spans and with known errors.

A. The Method

Instantaneous extraterrestrial solar radiation can be calculated from the relationship (Sellers, 1965):

$$Q = S(\bar{R}/R)^2 \cos Z$$
 [2.1]

where,

Martin Laboration

Q is the extraterrestrial solar radiation,

S is the solar constant, taken to be 1377 Watts per

square meter (U.S. Department of Commerce, 1978),

 (\bar{R}/R) is the ratio of the mean to actual sun-earth distance,

cos Z is the cosine of the solar zenith angle.

Determination of the relative sun-earth distance parameter $((\bar{R}/R)^2)$ and the solar position using Fourier Series was first proposed by Spencer (1971). Thompson (1981) devised more accurate Fourier Series representations for these parameters. The Fourier Series presented in this paper are based on the formulations of Thompson (1981). They have been modified, however, to facilitate computer programming. All series formulations were based on data obtained from the <u>American Ephemeris and Nautical Almanac</u> for 1965. The Fourier Series formulation for calculating the relative sun-earth distance parameter is detailed in section II A.1. The solar position is determined from the geometric formula:

 $\cos Z = \sin \phi \sin \delta + \cos \phi \cos \delta \cos h \qquad [2.2]$ where,

φ is the latitude of the observation,
δ is the solar declination, + for north latitudes,
- for south latitudes,
h is the local hour angle of the sun.

The Fourier Series formulation for calculating solar declination is detailed in section II A.2. The local hour angle is related to Greenwich mean time (GMT) as follows:

$$h = \frac{2\pi}{24} (GMT - 12 + T) - \psi$$
 [2.3]

where,

 ψ is the longitude of the observation in radians, + for west longitude, - for east longitude,

T is the equation of time.

The equation of time corrects for the difference between true solar time and mean time which is in common use. The Fourier Series used to calculate the equation of time is detailed in section II A.3.

The Fourier series formulations are approximations to the exact values. Data was obtained from the <u>American Ephemeris</u> <u>and Nautical</u> <u>Almanac</u> for the years 1974, 1979, 1980 and 1981 and compared to values calculated using these series representations. The errors thus obtained are given at the end of the section detailing each series.

1. Relative sun-earth distance

The Fourier series representation for the relative sun-earth distance parameter is:

> $(\bar{R}/R)^2 = 1.000145 + 0.033382 \cos \alpha + 0.001952 \sin \alpha$ + 0.000698 cos (2 α) [2.4]

The time parameter (α) is given by:

$$\alpha = 0.0172017221 [ET + (NOON/24)]$$
 [2.5]

where,

ET is the number of days from a base date of 0 January 1974

(i.e. 1 January 1974 is day one),

NOON is local noon (in Greenwich mean time) for the longitude of the point for which the calculation is being made, the constant (0.0172017221) is 2π divided by the annual period for the earth to return to the same distance from the sun (365.2649 days).

The maximum error in the calculated square of the relative sunearth distance using this formula is $\pm 0.011\%$.

2. Solar declination

The Fourier series representation of the sun's declination is: $\delta = 0.00666 - 0.400866 \cos \rho + 0.06523 \sin \rho$ $- 0.00667 \cos (2\rho) + 0.00057 \sin (2\rho)$ $- 0.00274 \cos (3\rho) + 0.00124 \sin (3\rho)$ [2.6]

The time parameter (ρ) is given by:

$$\rho = 0.0172027912 [ET + (TIME/24)]$$
 [2.7]

where,

TIME is the mid-point of the integration period (in Greenwich mean time),

the constant is derived from an annual period of

365.2422 days.

The maximum error in the calculated declination using this formula is ± 1 minute.

3. Equation of time

The Fourier series for calculating the equation of time is: $T = 0.0098 \cos \rho - 0.1227 \sin \rho - 0.0510 \cos (2\rho)$ $- 0.1581 \sin (2\rho) - 0.0056 \sin (3\rho)$ $- 0.0021 \cos (4\rho) - 0.0030 \sin (4\rho)$ [2.8]

where the time parameter (ρ) is calculated by equation [2.7].

and the second second

The maximum absolute error in the calculated equation of time using this formula is ± 12 seconds.

B. The Computational Procedure

Integration of equation [2.1] over some time interval for which the declination and relative sun-earth distance can be assumed constant yields:

$$Q_1 = S(\overline{R}/R)^2 [\sin \phi \sin \delta (h_2 - h_1) + \cos \phi \cos \delta (\sin h_2 - \sin h_1)] / ANGVEL [2.9]$$

where,

ANGVEL is the earth's angular velocity in radians per hour. Because it changes so slowly the relative sun-earth distance was calculated for noon of each day and assumed constant for that day. Thompson (1981) points out that calculation of the solar declination once every four hours gives satisfactory results; however, for this study values of both declination and the equation of time were calculated at the midpoint of each integration interval and assumed constant for that interval. A one hour time interval was chosen for use in this study, except that when h_1 was less than the sunrise hour angle it was set equal to the sunrise hour angle and, similarly, when h_2 was greater than the sunset hour angle is was set equal to the sunset hour angle.

The appendix is a program listing (FORTRAN V) of this computational scheme with the time interval (TIMINC) set to one hour. The program will output daily total extraterrestrial solar radiation for the years and location specified. Values other than daily totals (i.e. instantaneous, hourly, etc.) can be obtained by changing the beginning (BEGGMT)

and ending (ENDGMT) times. For mean monthly values a simple averaging loop should be added.

C. Accuracy of the Method

Given the errors inherent in calculating the relative sun-earth distance, declination, and equation of time, the maximum error in calculated extraterrestrial solar radiation has been determined. The errors were maximized with the declination set at 23° and the hour angles (which are a function of the equation of time) set at 90° for the instantaneous error determination and at 0° and 15° for the hourly determination. Hourly errors were then summed for 15 hourly calculations to arrive at the daily errors. It should be noted that the improved performance of the daily calculation over the hourly calculation is due to the fact that the equation of time correction is symmetrical about true solar noon. For this reason whenever the calculation is performed over an interval which is also symmetrical about true solar noon, the equation of time correction is cancelled and the error associated with it makes no contribution to the calculated extraterrestrial solar radiation. It should be emphasized that the stated errors are due solely to the computational procedure and do not account for any changes in solar output of long term changes in sun-earth geometry.

1. Instantaneous values

Maximum absolute error 1.3 Wm⁻² Maximum percent error 0.10%

2. Hourly values

Maximum absolute error 4 W hr $m^{-2}hr^{-1}$ Maximum percent error 0.32%

3. Daily values

Maximum absolute error 5 W hr m⁻²day⁻¹

Maximum percent error 0.05%

CHAPTER III

ANALYSIS OF FORT COLLINS SOLAR RADIATION DATA

Both the direct component (DIR) and hemispheric irradiance (HEM) have been measured at the Atmospheric Science Department of Colorado State University since the spring of 1975. The hemispheric irradiance is measured by an Eppley Precision Pyranometer in the spectral interval $0.3 \ \mu m$ to $3.0 \ \mu m$, while an Eppley Pyrheliometer measures the direct component in the same wave length band. Cox and McKee (1978) and Cox and McKee (1980) give a complete description of the facility and the instrumentation, as well as hourly, daily, and monthly values of the direct component and hemispheric irradiance. Because of the completeness of the data, the years 1977 through 1980 were chosen as the data base for this study. The diffuse component (DIF) was calculated from the relation:

$$DIF = HEM - (DIR)(\cos Z).$$
[3.1]

The method described in Chapter II for determining solar position was used in calculating cos Z.

A. Data Handling

Quality of solar radiation data can be affected by the calibration of the pryanometer or pyrheliometer, mechanical or electronic problems with the recording system, misalignment of the pyrheliometer, and/or dirty or snow covered optical surfaces. Therefore, the following decision criteria were applied to the data prior to further analysis and resulted in a total data set of 707 days.

1. Missing data

Data for an entire day was disregarded if for one or more hours either the direct or hemispheric values were missing. A total of 506 days fell into this category.

2. Suspicious data

If the calculated diffuse component was less than zero the entire day's data was disregarded. This condition results from the direct component being too large or the hemispheric being too small, either of which indicate instrument problems. Only nine days fell into this category.

3. Problems at sunrise/sunset

Near the times of sunrise and sunset it is possible for the solar zenith angle to exceed $\pm 90^{\circ}$, in which case the cosine of the solar zenith angle will be negative and the resulting calculated diffuse component will be greater than the hemispheric (assuming the direct is non-zero). For these events the hourly diffuse component was set equal to the hemispheric value.

4. Excessively large values of diffuse

All hours for which the inequality

$$DIF \leq 0.5(S \times \cos Z)$$
 [3.1]

was not true were eliminated (Collares-Pereira and Rabl, 1978). This condition results from the direct component being too small with respect to the hemispheric, the most probable cause being misalignment of the

pyrheliometer. A total of 239 days were eliminated for not satisfying the above inequality.

B. Results of Analysis

1. Frequency distributions

Table 1 gives the number of days, by month for which the diffuse component to hemispheric (DIF/HEM) ratio fell within the various intervals. Table 2 and Table 3 are similar, but for the diffuse to extraterrestrial (DIF/ETR) and hemispheric to extraterrestrial (HEM/ETR) ratios. respectively. Inspection of Table 1 reveals a maximum occurrence of days (23%) in the 0.10-0.19 range. These represent days of relatively clear conditions. A secondary peak (11%) is observed in the range 0.90-1.00, representing overcast conditions when the daily direct component is equal to or very nearly zero. From Table 2 we find that on 75% of the days the DIF/ETR ratio falls in the 0.10-0.29 range. Table 3 indicates that on 29% of the days the HEM/ETR ratio falls in the 0.60-0.69 range, and no evidence of a secondary maximum exists. Table 4 gives the frequency distribution of the direct to direct extraterrestrial ratio (DIR/DIRETR). Most notable here is the relatively uniform distribution in the range of 0.00 to 0.59, with the difference in the maximum and minimum occurrences being only about 9.8 percent.

2. An extended period of low surface solar radiation

The Front Range of the Colorado Rocky Mountains has the reputation of being generally sunny and mild and, therefore, an excellent environment for solar energy applications. The question arises as to how well deserved this reputation may be. In an attempt to at least partially answer this question, the data was scanned to determine the number of

								•					
Interval	Jan	Feb	Mar	Apr	May	Jun	[ทก	Aug	Sep	0ct	Nov	Dec	Total
0.00-0.09	1	4	Ч	н	1	0	0	2	1	13	11	9	41
0.10-0.19	7	12	17	7	6	19	22	11	21	24	12	7	163
0.20-0.29	5	5	13	7	11	13	14	٢	11	٢	4	7	105
0.30-0.39	2	9	7	13	19	6	18	7	2	m	2	9	97
0.40-0.49	2	4	11	2	11	4	13	7	4	9	4	4	78
0.50-0.59	ę	4	8	7	10	2	9	Ч	2	2	9	m	54
0.60-0.69	0	2	m	ε	٢	9	m	Υ	Ч	4	4	2	38
0.70-0.79	2	ŝ	m	2	4	2	0	9	1	2	0	m	28
0.80-0.89	0	2	ε	ε	m	2	9	Н	0	2	ŝ	ŝ	28
0.90-1.00	اف	비	1 1	9	위	ام	~	~	1	4	∞	-1	<u> </u>
Total	23	53	80	54	85	63	84	47	44	70	56	48	707

Table 2. Frequency distribution of the daily Diffuse/Extraterrestrial ratio. Entries are the number of days.

					ļ	!							
Interval	Jan	Feb	Mar	Apr	May	Jun	<u>In</u>	Aug	Sep	lt IS	Nov	Dec	Total
0.00-0.09	1	11	9	2	2	თ	2	ß	12	24	18	10	102
0.10-0.19	10	16	26	18	34	33	46	26	21	21	14	13	278
0.20-0.29	٢	16	33	29	37	13	34	14	10	23	16	17	249
0.30-0.39	5	10	14	2	12	7	2	2	7	~	80	7	75
0.40-0.50	이	0		0	0	7	0	<u> </u>	0	0	미	-1	۳ ا
Total	23	53	80	54	85	63	84	47	44	70	56	48	707

distribution of the daily Hemispheric/Extraterrestria	itries are the number of days.
ncy di	Entr
Freque	ratio.
Table 3.	

Interval	Jan	Feb	Mar	Apr	May	Jun	<u>Jul</u>	Aug	Sep	oct	Nov	Dec	Total
0.00-0.09	0	-1	0	н	1	0	0	0	0	0	1	0	4
0.10-0.19	2	Ś	٦	2	9	с	2	2	0	0	2	Ч	24
0.20-0.29	2	4	7	4	m	ς	4	5	1	m	5	4	45
0.30-0.39	2	2	7	7	ω	4	ς	7	1	S	с	5	52
0.40-0.49	ŝ	2	11	2	13	8	14	7	2	S	7	9	83
0.50-0.59	5	5	80	10	13	8	18	11	4	11	10	10	113
0.60-0.69	4	13	22	16	25	14	24	10	15	21	6T	19	202
0.70-0.79	പ	17	23	12	16	23	19	10	20	25	ნ	ĸ	182
0.80-0.89	0	9	-1	이	이	0	0	0	-1	0	0	이	2
Total	23	53	80	54	85	63	84	47	44	70	56	48	707

	days.
ب	of
:t/Direc	number
)irec	the
ily C	are
of the dai	Entries
Frequency distribution o	extraterrestrial ratio.
Table 4.	

					1								
Interval	Jan	Feb	Mar	Apr	May	Jun	Inf	Aug	Sep	0ct	Nov	Dec	Total
0.00-C.09	7	14	17	11	16	10	8	7	2	10	11	12	125
0.10-0.19	m	4	6	4	15	ω	٢	9		4	7	5	73
0.20-0.29	4	ß	11	10	13	4	18	12	വ	10	7	Q	105
0.30-0.39	2	7	12	12	20	6	17	9	9	2	£	10	108
0.40-0.49	4	8	11	10	12	18	16	7	11	12	ნ	ъ	123
0.50-0.59	7	13	16	9	8	13	18	7	18	29	16	10	156
0.60-0.69	-1	~	4	-1]	-1	-1	이	~	-1	mj	-1	0	1
Total	23	53	80	54	85	63	84	47	44	70	56	48	707

times when the daily hemispheric to extraterrestrial ratio was less than 0.35 for a period of three or more consecutive days. The criteria (HEM/ETR) < 0.35 was chosen because Orgill and Hollands (1976) point out that values in this interval represent extremely cloudy days with over 90% of the total insolation being diffuse. Six such cases were found. They were 27-29 April 1977, 16-18 August 1977, 6-9 February 1978, 1-3 May 1978, 6-8 May 1980. Because of the duration of the February 1978 event, the fact that this is the only period which was found in the primary heating season, and having noted that the HEM/ETR values for 11 and 12 February were also below the threshold, a closer examination of this period was made.

a. The meteorological situation. A high pressure cell was centered over southwestern Colorado on 5 February with a stationary front lying just east of Fort Collins. Fort Collins experienced light (less than 5 knots) south easterly flow throughout the entire day and by 1900 LST began observing the classic upslope conditions (ceiling at 100 feet and two miles visibility in foq). On 6 February the front moved west over the Rocky Mountains. Ceilings were generally at 500 feet, visibility three to four miles in fog, and winds remained light out of the south through southeast. A cold front, having moved off the Pacific Ocean, merged with the stationary front on the 8th and produced, on the 7th and 8th of February, ceilings 100 to 300 feet, visibility ½ to 5 mile in fog and occasional light snow, and winds calm to light southeasterly. On the morning of the 9th, a second frontal system moved on shore and stagnated over central Nevada and central California and remained in this location through the 10th as the low pressure cell centered over central Nevada intensified. The system began a slow eastward movement on 11 February which continued until around midnight

when it merged with the stationary front over the Colorado Rockies. During this entire period Fort Collins continued to experience up-slope conditions, with ceilings generally 300 to 800 feet (except 5,000 to 7,000 on the 10th), visibilities from less than one up to five miles and light southeasterly flow. The active portion of the cold front then dropped south into east central New Mexico, and by 0800 LST the stationary front had dissipated and the surface winds became light out of the northeast, although the ceiling remained at between 200 and 500 feet throughout most of the daylight hours. By early morning on the 13th the clearing trend was well established.

b. <u>Radiative observations</u>. The hemispheric/extraterrestrial ratio and direct/direct extraterrestrial for each day during this period is given below.

DAY	HEM/ETR	DIR/DIRETR
6	0.25	0.00
7	0.21	0.00
8	0.11	0.00
9	0.29	0.00
10	0.43	0.09
11	0.15	0.00
12	0.23	0.11

In reviewing the weather records for this February 1978 case, it appeared that there were other periods of extreme cloudiness that had not been detected by the proposed threshold. Time series plots of average cloudiness during daylight hours, the DIR/DIRETR ratios, and the HEM/ETR ratios for the period 5 January to 5 March were made, and are given in Figure 1. Three periods stand out in the cloudiness and DIR/ DIRETR plots -- 12-20 January, 5-16 February, and 24 February through 3 March. Average cloudiness during the 5-16 February event was 9.6/10, while the climatological mean for February is only 5/10. The average DIR/DIRETR for the period was only 0.04. Temperature deviations

Figure 1. Time series plots of average daylight hours cloudiness, average daily direct/direct extraterrestrial ratios, and the average daily hemispheric/extraterrestrial ratios for 5 January to 5 March 1978. Points indicated by an M are missing, those indicated by an E are estimated.



(observed average minus climatological mean) were:

Maximum	(°F)	-13.8
Minimum	(°F)	+2.3

The synoptic situation during the January and February/March cases proved to be practically identical to that detailed for 5-12 February. The results of further investigation are as follows:

	12-20	January	<u>24 Febru</u>	<u>ary - 3 March</u>
	Mean	<u>Observed</u>	Mean	Observed
Cloudiness	4.5/10	8.9/10	5/10	9.9/10
DIR/DIRETR		0.03		0.04
Maximum Temperature (°F)	40.6	33.0	44.8	36.5
Minimum Temperature (°F)	14.6	10.0	18.0	20.2

We therefore have three events covering 29 days in a total period of only 58 days where the average daylight hours cloudiness exceeded 88% cloud cover, the average daily DIR/DIRETR ratio was only 0.04, and temperatures where highly variable with maximums from 7.6 to 13.8 degrees below the means and minimums from 2.3 degrees above to 4.6 degrees below mean values.

A four year data base is insufficient to establish a real solar radiation climatology. In general, however, it would seem that the reputation of the Front Range is well deserved, though occasional extended periods (three days or more) of low insolation may be expected. The January-March 1978 event is certainly an anomally and should not be considered representative of a normal year.

C. Relationships Between Solar Radiation Components

Subsequent to the initial work by Liu and Jordan (1960) relating the diffuse and hemispheric irradiance, many authors have investigated this relationship for various other locations. Among these Ruth and Chant (1975), Orgill and Hollands (1976), and Collares-Pereira and Rabl (1978) found correlations which deviated significantly from those of Liu and Jordan (1960). The remainder of this chapter will deal with the diffuse-hemispheric irradiance correlations determined from the 1977 through 1980 data base for Fort Collins, Colorado.

1. Method of correlation

The daily DIF/ETR and HEM/ETR ratios were calculated. For each HEM/ETR interval of 0.10 the corresponding values of DIF/ETR were averaged and these mean values were plotted against the value of HEM/ETR for the mid-point of that interval. The only exception to this approach was for the HEM/ETR interval 0.00-0.09. Only four days fell into this interval with values of 0.07-0.09. Therefore, the mean DIF/ETR value (0.08) was plotted against the mean HEM/ETR value (0.08). A least squares polynomial fit was then obtained for these means.

The cumulative density function of DIF/ETR values within each HEM/ETR interval was also obtained, and the 20 and 80 percent levels determined. The values of each of these levels were again plotted against the value of HEM/ETR for the mid-point of that interval (except that values for the lowest interval were again plotted against a HEM/ETR value of 0.08), and a least squares ploynomial fit obtained for each of these levels.

2. Correlation equations

The interval 0.35 \leq (HEM/ETR) \leq 0.80 included approximately 86.3 percent of the total number of days. The following correlations were obtained:

Mean	-0.348 + 3.93(HEM/ETR)	
	-7.51(HEM/ETR) ² + 4.14(HEM/ETR) ³	[3.2]
20% level	-0.129 + 2.72(HEM/ETR)	
	-5.81(HEM/ETR) ² + 3.42(HEM/ETR) ³	[3.3]
80% level	0.243 + 0.386(HEM/ETR)	

There were no points in the interval (HEM/ETR) > 0.81.

The remaining days fell in the interval $0.00 \leq (\text{HEM/ETR}) < 0.35$. The correlations obtained were:

Mean
$$0.010 - 0.863(HEM/ETR)$$

+ $0.200(HEM/ETR)^2 - 1.13(HEM/ETR)^3$ [3.5]
20% level $0.0913 - 0.724(HEM/ETR)$
+ $7.62(HEM/ETR)^2 - 12.0(HEM/ETR)^3$ [3.6]
80% level $0.0006 + 1.34(HEM/ETR)$
- $1.20(HEM/ETR)^2 + 0.273(HEM/ETR)^3$ [3.7]

It interesting to note that while 32.4 percent of the values in the Orgill and Hollands (1976) study (based on four years of data for Toronto Airport, Canada) fell in this interval, which again represents extremely cloudy conditions, only 13.7 percent of Fort Collins days were included. Caution should be used in any direct comparison, however, since their work was based on hourly values and the present study deals only with daily totals.

Figure 2 shows plots of the above correlation equations. A comparison of the correlation equations for the mean values in the present study with those developed by Liu and Jordan (1960), based on ten years of data for Blue Hill, Massachusetts, and by Orgill and Hollands (1976),

See See





based on four years of data for Toronto Airport, Canada, is given in Figure 3. The correlations for the three different regions are very similar for HEM/ETR values less than 0.15. This is no doubt due to the fact that in this range the direct component is zero (or very nearly so) and, therefore, the total insolation is very nearly all diffuse. In the HEM/ETR range of 0.15 to around 0.45, the present study gives DIF/ETR values that lie about midway between the other two correlations, while above HEM/ETR values of 0.60 the present study and that of Liu and Jordan (1960) agree quite well.

1.1.1



Comparison of the hemispheric/extraterrestrial to diffuse/extraterrestrial correlations of the present study with those of Liu and Jordan (1960) for Blue Hill, MA and Orgill and Hollands (1976) for Toronto Airport, Canada.

CHAPTER IV

SUMMARY AND CONCLUSIONS

An investigation of the solar radiation environment of Fort Collins, Colorado, was performed. A method of calculating the extraterrestrial solar radiation over various time scales and with known errors was accomplished. Uncertainty of the extraterrestrial values resulting from this calculation are 1.3 Wm^{-2} in calculating instantaneous values, $4 \text{ W hr m}^{-2} \text{ hr}^{-1}$ for hourly values, and $5 \text{ W hr m}^{-2} \text{ day}^{-1}$ for daily total calculations. These errors are certainly less than those to be expected in the measurement of surface solar radiation. The computer program developed to perform the ETR calculation is given in the Appendix and will produce daily totals for one year for about \$1.20 on a Control Data Corporation CYBER system. (Note: This is not intended to be an endorsement of the CDC CYBER and is given only as an example of expected cost.)

Equations correlating the diffuse and hemispheric components were developed. Since establishment of confidence intervals for these correlations would entail assumptions regarding the statistical normalicy of the data, correlations for the 20 percent level and 80 percent level obtained from cumulative density functions of the actual data were determined. This was done in an attempt to provide the maximum information with the fewest assumptions. Using these correlations the mean daily diffuse component of the insolation can be predicted, given the value of the hemispheric component. These present correlations were also compared to those found by Liu and Jordan (1960) and Orgill and Hollands (1976) and were found to agree well with the results of Liu and Jordan (1960) except in the HEM/ETR range of 0.20 to 0.60 where the present work gives values of DIF/ETR ranging from 0.01 to 0.04 higher. Except in the very lowest values ((HEM/ETR) \leq 0.20) the correlations of Orgill and Hollands (1976) consistently over estimate the DIF/ETR values of the present study by as much as 0.05. While this study was based on data taken only at Fort Collins, Colorado, and the correlations have not been applied to an independent data set for other locations, the author believes the correlations given should prove useful for other climatologically similar, high altitude sites.

Finally, an investigation into extended periods of low insolation was made. For the four year data set six periods of three days or more were found where the diffuse component would account for more than 90% of the total insolation received. Additionally, during a 58 day period three cases totaling 29 days were found in which the direct component to extraterrestrial direct component ratio did not exceed 0.15. The four year data set is certainly too short a time period to establish a climatology. However, the results of this study tend to lend creedence to the reputation of the Colorado Front Range as being very suitable for solar energy applications.

REFERENCES

- Collares-Pereira, M., A. Rabl, 1979: The average distribution of solar radiation -- correlations between diffuse and hemispherical and between daily and hourly insolation values. <u>Solar Energy</u>, <u>22</u>, 155-164.
- Cox, S. K., T. B. McKee, 1978: Fort Collins solar radiation data --May 1975 through December 1977. Climatology Report 78-1, Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado.
- Cox, S. K., T. B. McKee, 1980: Fort Collins solar radiation data --January 1978 through December 1979. Climatology Report 80-1, Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado.
- Liu, B. Y. H., R. C. Jordan, 1960: The interrelationship and characteristic distribution of direct, diffuse and total radiation. <u>Solar</u> <u>Energy</u>, <u>IV</u>, 1-9.
- Orgill, J. F., K. G. T. Hollands, 1976: Correlation equation for hourly diffuse radiation on a horizontal surface. <u>Solar Energy</u>, <u>19</u>, 325-329.
- Ruth, D. W., R. E. Chant, 1976: The relationship of diffuse radiation to total radiation in Canada. Solar Energy, 18, 153-154.
- Sellers, W. D., 1965: <u>Physical Climatology</u>. The University of Chicago Press, Chicago, Illincis.
- Spencer, J. W., 1971: Fourier Series representation of the sun. Search, 2, 172.
- Thompson, T. M., 1981: Pyrheliometer observations as an indication of the climatological persistance of clouds. Technical Memo NOAA ERL ARL-97. Air Resources Laboratory, Silver Spring, MD, Feb., 78 pp.
- U. S. Department of Commerce, 1978: <u>SOLMET, Volume 1 -- Users Manual</u>, Hourly Solar Radiation -- Surface Meteorological Observations. NOAA, Environmental Data and Information Service, National Climatic Center, Asheville, North Carolina.

APPENDIX

This Appendix is a listing of a program to compute daily values of extraterrestrial solar radiation (HEMETR), the normal component of the extraterrestrial solar radiation (DIRETR), and the length of the day (DAYLEN).

.

```
PROCRAII EXTRA
С
      COMPUTES EXTRATERRESTRIAL SOLAR RADATRION ON BOTH A HORIZONTAL
С
      PLANE (HEMETR) AND A PLANE NORMAL TO THE SUN'S RAYS (DIRETR) BY
С
      T.M. THOMPSON'S (1981) METHOD (MODIFIED BY J.R. CONLEY(1982))
С
      FOR DETERMINING SOLAR POSITION. (ETR IS IN WATT HOURS PER
С
      SQUARE METER). ALSO COMPUTES DAYLENGTH (DAYLEN).
С
      DAYLEN UNITS ARE HUNDREDTHS OF HOURS (IE. 932 IS 9.32 HOURS).
С
С
      INTEGER ALLSOL
      DIMENSION ALLSOL(3,31,12), NDAY(12)
      COMMON TIME1, TIME2, SINLAT, COSLAT, XLONG, DECL, EQT, DIST,
             BEGGMT, ENDGMT, TIMINC, SOLC, ANGVEL, HEMETR, DIRETR,
     1
             SUNRIS, SUNSET, DAYLEN, IYR, IMO, IDAY, IET
     2
      DATA NDAY/31,28,31,30,31,30,31,31,30,31,30,31/
      OPEN(UNIT = 7, FILE = 'RADAT')
С
С
      INITIALIZE ARRAY ALLSOL WHICH WILL CONTAIN VALUES OF
      EXTRATERRESTRIAL SOLAR RADIATRION AND DAYLENGTH.
С
С
С
С
      DO 1000 I = 1,3
      DO 1000 J = 1,31
      DO 1000 K = 1, 12
 1000 \text{ ALLSOL}(I, J, K) = -9.00
С
С
      PROVIDE LATITUDE, LONGITUDE, AND STATION NUMBER
С
      (EXAMPLE: 40 DEG 35 MIN NORTH = 40.5833 DEG, 40 DEG SOUTH =
С
      -40.0000, 90 DEG WEST = 90.0000 DEC, 90 DEG EAST =
С
      -90.0000 DEG).
С
С
      XLAT = 40.5333
      XLONG = 105.1333
      ISTA = 53006
С
С
      CONVERT LATITUDE AND LONGITUDE TO RADIAN MEASURE
С
С
      PI = 3.14159
      CONV = PI/180.
      XLAT = XLAT*CONV
      SINLAT = SIN(XLAT)
      COSLAT = COS(XLAT)
      XLONG = XLONG*CONV
С
С
      SET SOLAR CONSTANT (WATTS PER SQUARE METER) AND EARTH'S ANGULAR
      VELOCITY (RADIANS PER HOUR)
С
С
С
      SOLC = 1377.
      ANGVEL = 15.*CONV
С
```

С SET THE TIME PERIOD (IN DECIMAL CHT HOURS) OVER WHICH THE С INTEGRATION OF ETR IS TO BE PERFORMED (EX: 1010GMT = 10.1667) С AND THE YEAR DESIRED. С С BEGG11T = 11.00ENDGMT = 05.00IYR = 78С С С С SET THE INTERVAL (IN DECIMAL HOURS) OVER WHICH INDIVIDUAL С INTECRATIONS OF ETR ARE TO BE PERFORMED. С NOTE: THIS IS THE INTEGRATION INTERVAL (NOT THE PERIOD С FOR WHICH A FINAL VALUE IS DESIRED) AND IS NORMALLY С SET TO ONE HOUR. С С TIMINC = 1.0С С С DO 1500 IMO = 1,12ND = NDAY(IMO) $IF(MOD(IYR,4) \cdot EQ \cdot 0 \cdot AND \cdot IMO \cdot EQ \cdot 2)MD = 29$ DO 1500 IDAY = 1, NDС С THE ADDITION OF 0.5 TO VALUES BELOW IS DONE TO ROUND С THEM UP BEFORE FORCING THEM TO BECOME INTEGERS. IN С THE CASE OF DAYLEN THE !WLTIPLICATION FACTOR OF 100 С FORCES DAYLEN TO BE AN INTEGER. С С ALLSOL(1, IDAY, IMO) = ETR() + 0.5ALLSOL(2, IDAY, IMO) = HEMETR + 0.5ALLSOL(3, IDAY, INO) = DAYLEN * 100 + 0.51500 CONTINUE WRITE(7,6000) ISTA, IYR DO 1600 J = 1,31WRITE(7,6010) $J_{((ALLSOL(I,J,K),K = 1,12),I = 1,3)}$ 1600 CONTINUE 5000 CONTINUE 6000 FORMAT(///' STATION NO.', IE, 5X, ' YEAR 19', 12//6X, 'DAY', 6X, 1 'JAN', 7X, 'FEB', 7X, 'MAR', 7X, 'APR', 7X, 'MAY', 7X, 'JUN', 7X, 2 'JUL', 7X, 'AUG', 7X, 'SEP', 7X, 'OCT', 7X, 'NOV', 7X, 'DEC') 6010 FORMAT(18,12110,2(/8X,12110)) END

FUNCTION ETR() С THIS FUNCTION OPERATES AS THE TIMING CONTROL FOR COMPUTATION OF EXTERRESTRIAL SOLAR RADIATRION AND TOTALS DAILY VALUES. С CONMON TIME1, TIME2, SINLAT, COSLAT, XLONG, DECL, EQT, DIST, BEGGHT, ENDGMT, TIMINC, SOLC, ANGVEL, HEHETR, DIRETR, l 2 SUNRIS, SUNSET, DAYLEN, IYR, IMO, IDAY, IET С С CALL SUBROUTINE TO CALCULATE ELAPSED TIME SINCE BASE DATE С С CALL ELPSTM С С С CALL SUBROUTINE TO CALCULATE SUN-EARTH DISTANCE PARAMETER С С CALL DISTMN C . С С TIME1 = BECGMT HEMETR = 0.0ETRDIR = 0.0SUNRIS = 0.0SUNSET = 0.0IF(ENDGMT .LT. BEGGMT)ENDGMT = ENDGMT + 24. 10 CONTINUE IF(TIME1.GT. ENDCMT) GO TO 20 TIME2 = TIME1+TIMINC HEMETR = ETR2() + HEMETRETRDIR = DIRETR + ETRDIRTIME1 = TIME2GO TO 10 20 CONTINUE ETR = ETRDIRRETURN END

SUBROUTINE ELPSTM COMPUTES NUMBER OF DAYS SINCE THE BASE DATE O JAN 1974 FOR С С USE IN TIME PARAMETERS RHO AND ALPHA. COMMON TIME1, TIME2, SINLAT, COSLAT, XLONG, DECL, EQT, DIST, BEGGMT, ENDGNT, TIMINC, SOLC, ANCVEL, HEMETR, DIRETR, 1 SUNRIS, SUNSET, DAYLEN, IYR, IMO, IDAY, IET 2 С С С INITIALIZE 0 JAN 1974 TO NUMBER OF DAYS SINCE 1 JAN 1901 IBASE = 26663С COMPUTE NUMBER OF DAYS FROM 1 JAN 1901 TO DATE DESIRED С LEAPYR = 2 $IF((IYR/4*4) \cdot EQ \cdot IYR) LEAPYR = 1$ IDYYR = IMO*275/9 + IDAY - 30 $IF(IMO \cdot GT \cdot 2)IDYYR = IDYYR-LEAPYR$ IDYCT = (IYR-1)*1461/4 + IDYYRС С С COMPUTE ELAPSED TIME IN DAYS FROM BASE DATE TO DESIRED DATE IET = IDYCT-IBASERETURN END

SUBROUTINE DISTMN COMPUTES THE SQUARE OF THE RATIO OF THE MEAN TO ACTUAL С С SUN-EARTH DISTANCE COMMON TIME1, TIME2, SINLAT, COSLAT, XLONG, DECL, EQT, DIST, BEGGMT, ENDCMT, TIMINC, SOLC, ANGVEL, HEMETR, DIRETR, 1 2 SUNRIS, SUNSET, DAYLEN, IYR, IMO, IDAY, IET С С С С COMPUTE TIME PARAMETER ALPHA FOR LOCAL NOON OF GIVEN DAY XNOON = 12.+XLONG/ANGVEL ALPHA ≈ .0172017221*(IET+(XNOON/24.)) SINAL = SIN(ALPHA)COSAL = COS(ALPHA)COS2AL = 2.*COSAL*COSAL-1. DIST = 1.000145 + 0.033382*COSAL + 0.001952*SINAL + 1 0.000698*COS2AL RETURN END

```
FUNCTION ETR2()
С
      COMPUTES EXTERRESTIAL SOLAR RADIATION
С
      COMMON TIME1, TIME2, SINLAT, COSLAT, XLONG, DECL, EQT, DIST,
             BEGGMT, ENDGMT, TIMINC, SOLC, ANGVEL, HEMETR, DIRETR,
     1
     2
             SUNRIS, SUNSET, DAYLEN, IYR, IMO, IDAY, IET
      OBTAIN MIDPOINT OF CURRENT TIME INTERVAL FOR USE IN CALCULATING
С
С
      RHO
      TIME = (TIME1 + TIME2)/2.
С
С
      COMPUTE TIME PARAMETER RHO
      RHO = 0.0172027912*(IET + (TIME/24.))
С
С
      COMPUTE TRIGONOMETRIC VARIABLES
      SINRHO = SIN(RHO)
      COSRHO = COS(RHO)
      SIN2RH = 2.*SINRHO*COSRHO
      COS2RH = 2.*COSRHO*COSRHO - 1.
      SIN3RH = 3.*SINRHO-4.*SINRHO*SINRHO
      COS3RH = 4.*COSRHO*COSRHO*COSRHO - 3.*COSRHO
      SIN4RH = 4.*SINRHO*COSRHO - 8.*SINRHO*SINRHO*SINRHO*COSRHO
      COS4RH = 8.*COSRHO*COSRHO*COSRHO*COSRHO - 8.*COSRHO*COSRHO +1.
С
С
С
      COMPUTE DECLINATION AND EQUATION OF TIME
      DECL = 0.00666 - 0.400866 \times COSRHO + 0.06523 \times SINRHO - 0.00667 \times
            COS2RH + 0.00057*SIN2RH - 0.00274*COS3RH + 0.00124*
     1
            SIN3RH
     2
      EQT = 0.0098*COSRHO - 0.1227*SINRHO - 0.0510*COS2RH - 0.1581*
           SIN2RH - 0.0056*SIN3RH - 0.0021*C0S4RH - 0.0030*SIN4RH
     1
С
С
      SINDEC = SIN(DECL)
      COSDEC = COS(DECL)
      HR1 = ANGVEL*(TIME1 - 12. + EQT) - XLONG
      HR2 = ANGVEL*(TIME2 - 12. + EQT) - XLONG
      HRO = ACOS(-(SINLAT*SINDEC)/(COSLAT*COSDEC))
      IF(HR1 .GT. HRO) GO TO 10
      IF(HR2 .LT. -HRO) GO TO 10
      IF(HR1 \cdot LT \cdot -HRO) HR1 = -HRO
      IF(HR2 \cdot GT \cdot HRO) HR2 = HRO
      ETR2 = 0.0
      DIRETR = 0.0
      IF(HR1 .GE. HR2) RETURN
      ETR2 = SOLC*DIST*(SINLAT*SINDEC*(HR2-HR1)+
          COSLAT*COSDEC*(SIN(HR2)-SIN(HR1)))/ANGVEL
     1
      DIRETR = SOLC*DIST*((HR2 - HR1)/ANGVEL)
С
      CALCULATE SUNRISE AND SUNSET HOUR ANGLES (DEFINITIONS: SUNRISE -
      CENTER OF SOLAR DISK IS ON THE HORIZON WITH SUN RISING; SUNSET -
С
      CENTER OF SOLAR DISK IS ON THE HORIZON WITH SUN GOING DOWN).
С
      IF (HR1 .EQ. -HR0) SUNRIS = HR1
      IF (HR2 .EQ. HR0) SUNSET = HR2
С
      CALCULATE DAY LENGTH FROM SUNRISE AND SUNSET HOUR ANGLES, ADDING
С
      CORRECTION FOR ATMOSPHERIC REFRACTION (-34 MINUTES OF ARC) AND
```

C THE SUN'S SEMIDIAMETER (-16 MINUTES OF ARC). DAYLEN = (SUNSET - SUNRIS)/ANGVEL + 0.16 RETURN 10 CONTINUE ETR2 = 0.0 DIRETR = 0.0 RETURN END

