



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

ć

MRC Technical Summary Report #2512

A VERY SMALL REMARK ON SMOOTH DENJOY MAPS OF THE CIRCLE

AD A 129185

Glen Richard Hall

Mathematics Research Center University of Wisconsin—Madison 610 Walnut Street Madison, Wisconsin 53706

April 1983

(Received March 30, 1983)



DTIC FILE COPY

Sponsored by

U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 Approved for public release Distribution unlimited

anđ

National Science Foundation Washington, DC 20550

88 06 07 069

UNIVERSITY OF WISCONSIN - MADISON MATHEMATICS RESEARCH CENTER

A VERY SMALL REMARK ON SMOOTH DENJOY MAPS OF THE CIRCLE

Glen Richard Hall

Technical Summary Report #2512

April 1983

ABSTRACT

In this note we show that if f is a smooth homeomorphism of the circle with one 'non degenerate' critical point such that f has an irrational rotation number which is poorly approximable by rationals (e.g., a guadratic irrational) and f is a Denjoy map, then the length of the associated wandering interval cannot tend to zero too rapidly.

AMS (MOS) Subject Classifications: 58F99, 54H20 Key Words: Denjoy Maيs, wandering intervals Work Unit Number 1 - Applied Analysis

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and National Science Foundation Postdoctoral Fellowship.

SIGNIFICANCE AND EXPLANATION

Maps of a circle into itself, when considered as discrete dynamical systems, have been observed to contain some of the qualitative behavior of complicated experimental situations. Much of this structure can be understood for homeomorphisms of the circle by considering the invariant called the rotation number. In fact, the homeomorphisms of the circle with irrational rotation number can be classified up to topological conjugacy, provided the map and its inverse are sufficiently smooth. Counter examples to this Sub 1classification scheme exist which are C_1^{i} diffeomorphisms or C_1^{i} homeomorphisms with non differentiable inverse. Whether C_1^{i} can be replaced with analytic in the second examples is an open question. $-f_{i} \in A = f_{i}h_{i}rS$

In this report we show that if certain assumptions are made on the rotation number and on the derivative of a homeomorphism from the circle onto itself, then even if the map does not fit into the classification scheme we can still obtain estimates on the behavior of its orbits under iteration.

Accession For NTIS GRA&I DTIC TAB Unarmounded \square Justification By_ Distribution/ Availabil (ty Godes Avail Midy +r Dist Appelel

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

- a ~

A VERY SMALL REMARK ON SMOOTH DENJOY MAPS OF THE CIRCLE

Glen Richard Hall

1. <u>Introduction</u>: Let T = R/Z be the circle with unit circumference. We say a homeomorphism f : T + T is a Denjoy map if f has no periodic and no dense orbits. A theorem of Denjoy [2] states that if f is a C^2 diffeomorphism then f is not a Denjoy map. He also constructs C^1 diffeomorphisms which are Denjoy maps (see Herman [4]). Recently, examples of homeomorphisms f : T + T which are C^{∞} Denjoy maps have been constructed ([3]). These maps fail to be diffeomorphisms because they have points where the derivative of f is zero. However, the question of the existence of analytic homeomorphisms which are Denjoy maps remains open.

It turns out that a homeomorphism $f: T \to T$ is a Denjoy map if and only if it has no periodic orbits and there exists a "wandering" interval $I \subseteq T$ such that $\{f^{n}(I)\}_{n=\infty}^{\infty}$ forms a disjoint family of intervals. An interesting property of the examples of [3] is that they can be constructed so that length $f^{n}(I) + 0$ arbitrarily rapidly. (This requires an addition to [3] which states that "case B" will not occur. This was shown to the author by J. C. Yoccoz (personal communication).)

In this note we consider homeomorphisms f : T + T with only a "nondegenerate" critical point (e.g., analytic homeomorphisms) and rotation number which is very poorly approximable by rationals (e.g., quadratic irrationals). If f is a Denjoy map with $I \subseteq T$ a wandering interval as described above, then we show length $f^{n}(I)$ cannot tend to zero <u>arbitrarily</u> rapidly.

As we will note in section 3, recent computer studies (Shenker, [5]) indicate that the estimates of this note might be significantly improved, yielding a corresponding improvement of the theorem.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and National Science Foundation Postdoctoral fellowship.

2. <u>Definitions and Notations</u>: Fix a map f_0 : $\mathbb{R} + \mathbb{R}$ satisfying the following conditions

- 1) f_0 is increasing, C^3 and $f_0(0) \in [0,1)$,
- 2) $\Psi_x \in \mathbf{R}, f_0(x+1) = f_0(x) + 1, (i.e. f_0)$

is the lift of a homeomorphism of the circle),

- 3) there exists precisely one point $x_0 \in [0,1)$ such
 - that $f_0^*(x_0) = 0$ (hence $f_0^*(x_0) = 0$),
- 4) $f_0^{+++}(x_0) \neq 0$,
- 5) $f_n^w(x) = 0$ for only finitely many $x \in [0,1)$.

We let $f(\cdot,\phi)$: R + R denote the one parameter family of maps given by

$$f(x,\phi) = f_0(x) + \phi$$

Notation: Fix an interval $J \subseteq [0,1)$, $x_0 \in \text{interior } J$. Then let

$$M_{1} = \sup_{\substack{x \in [0,1)}} \frac{\partial f}{\partial x} (x,\phi) ,$$

$$M_{2} = \sup_{\substack{x \in [0,1)}} |\frac{\partial^{2} f}{\partial x^{2}} (x,\phi)| ,$$

$$M_{3} = \inf_{\substack{x \in J}} |\frac{\partial^{3} f}{\partial x^{3}} (x,\phi)| ,$$

$$M_{4} = \inf_{\substack{x \in [0,1) \sim J}} \frac{\partial f}{\partial x} (x,\phi) .$$

We may choose J small so that $M_3 > 0$. (Clearly these constants are independent of ϕ .) We define $f^n(x,\phi) = f(f^{n-1}(x,\phi),\phi)$ for $n \in \mathbb{Z}$, i.e., $f^n(x,\phi)$ is the n^{th} iterate of $f(\cdot,\phi)$ with ϕ fixed. Definition: The rotation number of $f(\cdot,\phi)$ denoted $\rho(\phi)$ is given by

$$\frac{1}{1}$$

$$\rho(\phi) = \lim_{n \to \infty} \frac{f^n(\mathbf{x}, \phi)}{n}$$

-2-

<u>Remarks</u>: 1) The rotation number $\rho(\phi)$ exists and is independent of the x in the definition. Moreover, $\rho(\phi)$ is a continuous increasing function of ϕ .

2) The function $\rho(\phi)$ is strictly increasing when its value is irrational, i.e., if $\rho(\overline{\phi}) \notin Q$ and $\phi \neq \overline{\phi}$ then $\rho(\phi) \neq \rho(\overline{\phi})$ (see Herman [4] for proofs of these remarks).

Notation: For each $\beta \in \mathbb{R}$, let $A_{\beta} = \{\phi : \rho(\phi) = \beta\}$. Then for every β , A_{β} is a closed interval, which is a single point when β is irrational.

Finally, we fix $\alpha_0 \in [0,1]$, α_0 an irrational such that there exists a constant $M_5 > 0$ so that if

$$a_0 = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{2}}}, a_1 \in \mathbb{Z}$$

Then $|a_i| < M_5$ for all i, (e.g., any quadratic irrational will satisfy this condition). We let $\frac{p_n}{q_n}$ denote the nth convergent of α_0 , i.e.

$$\frac{p_{n}}{n} = \frac{1}{a_{1} + \frac{1}{a_{2} + \frac{1}{2}}}, + \frac{1}{a_{n}}$$

Let $\phi_n \in \mathbb{R}$ be the unique number such that

$$\rho(\phi_0) = \alpha_0$$

3. <u>Statement of the theorem</u>: Suppose $f(\cdot, \phi_0)$ is a Denjoy map, i.e. $f(\cdot, \phi_0)$ is not continuously conjugate to rigid rotation by its rotation number a_0 . Then there exists a nontrivial (i.e., non empty, non singleton)

interval $I \subseteq \{0,1\}$ such that $\{f^n(x) + m : x \in I\}_{n,m\in\mathbb{Z}}$ form a family of disjoint intevals. Let $\mu_n = \text{length}(f^n(I))$, then $\mu_n + 0$ as $n + \pm \infty$ (recall f is the lift of a circle map).

Theorem: There exists a constant $M_0 > 1$ such that

(3.1)
$$\mu_{\pm q_n} > 1/M_0^{9n}$$

infinitely often.

That is, the length of the wandering interval cannot decrease too fast. The outline of the proof is as follows: First we derive an estimate on the length of the intervals A_{p_n/q_n} . Next we show that the orbit of the critical point must return to a neighborhood of the critical point in each set of q_n iterates, moreover, the size of this neighborhood is controlled by the size of μ_{q_n} . Hence, if μ_{q_n} is very small, a small change in ϕ will make a relatively large change in the rotation number, not leaving "enough room" for the A_{p_n/q_n} intervals. <u>Remark</u>: It is important to note that the estimates we use on the derivatives of high iterates of $f(\cdot,\phi)$ are very crude (see lemma 1 below). Recent numerical studies (Shenker, [5]) indicate <u>much</u> better estimates may hold, and

if so, much better bounds on μ_{α} would result.

4. <u>The lemmas</u>: In this section we give the necessary lemmas. Lemma 1: For all $x \in \mathbb{R}$, $\phi \in \mathbb{R}$ and $n \in \mathbb{Z}^+$

(4.1)
$$\frac{\partial (f^n)}{\partial x} (x,\phi) \leq M_1^n ,$$

(4.2)
$$\left|\frac{\partial^2(\underline{f}^n)}{\partial x^2}(x,\phi)\right| \le M_2 \cdot M_1^{2n-2}/(M_1^{-1})$$

(4.3)
$$\frac{\partial (\underline{r}^n)}{\partial \phi} (\mathbf{x}, \phi) \leq M_1^{n+1} / (M_1^{-1}) \quad .$$

-4-

Proof: These estimates follow immediately from the chain rule and (2.1),

(2.2) (see remark, end of section 3). // Lemma 2: For any n > 0, $3q_n$ length(A_{p_n/q_n}) > $1/(\frac{4M_2 \cdot M_1}{(M_1-1)})$.

<u>Proof</u>: Fix n > 0 and let $[\phi_1, \phi_2] = A_{p_n/q_n}$. Then $\forall x \in R, f^n(x, \phi_1) \leq x + p_n$,

and there exists $x_1 \in (0,1)$ with

$$f^{q}(x,\phi_{1}) = x_{1} + p_{n}$$

Now, $\frac{\partial f^{n}_{n}}{\partial x}(x_{0},\phi) = 0$, so by (4.2) for $\delta = 1/(\frac{2M_{2} \cdot M_{1}}{(M_{1}-1)})$ we have for any ϕ ,

$$\frac{\partial f^n}{\partial x}(x,\phi) \leq \frac{1}{2} \text{ whenever } |x_n - x| \leq \delta$$

But then it follows that for any ϕ ,

$$\max_{x \in \{0,1\}} (x - f^{n}(x,\phi)) - \min_{x \in \{0,1\}} (x - f^{n}(x,\phi))$$

$$\approx \{0,1\}$$

$$2q_{n}^{-1}$$

$$> 1/(\frac{4M_{2} \cdot M_{1}}{M_{1}^{-1}}) .$$

By (4.3) we then have that there exist x_2 , $x_3 \in \mathbb{R}$ such that

$$f^{n}(x_{2},\phi) \leq x_{2} + p_{n}$$
 and $f^{n}(x_{3},\phi) \geq x_{3} + p_{n}$

whenever $0 < \phi - \phi_1 < \frac{1}{\frac{q+1}{2q-1}}$ and the proof is complete. // $\left(\frac{4M_1 \cdot M_1}{(M_1 - 1)^2}\right)$

-5-

The following lemmas are standard results on the orbit structure for maps with irrational rotation number.

Lemma 3: The sets $\{f^i(x,\phi_0) + m : i, m \in \mathbb{Z}, 0 \le i \le q_n\}$ and $\{f^{-i}(x,\phi_0) + m : i, m \in \mathbb{Z}, 1 \le i \le q_n\}$ alternate on R (i.e. between any two points of the first set there is a point of the second set and vice versa) for any fixed $x \in \mathbb{R}$ and q_n .

Lemma 4: For each n > 0, x_1 , $x_2 \in \mathbb{R}$, if for some $q_1 > i, j > 0$, m_1 , $m_2 \in \mathbb{Z}$, $f^i(x_1, \phi_0) + m_1 < f^j(x_1, \phi_0) + m_2$ then there exists k and m_3 with $0 < k < q_{n+1}$ with

$$f^{i}(x_{1},\phi_{0}) + m_{1} \leq f^{k}(x_{2},\phi_{0}) + m_{3} \leq f^{j}(x_{1},\phi_{0}) + m_{2}$$

The proofs of these lemmas follow from, for example, Theorem 4.1, page 409 of Coddington, Levinson [1] and II. 7.1 of Herman [4], respectively.

Lemma 5: For n sufficiently large, the set $\{f^i(y,\phi_0)-[f^i(y,\phi_0)]: i = 0,...,n\}$ (where [*] denotes greatest integer) contains more than two points in J for any y $\in [0,1)$.

<u>Proof</u>: It suffices to show that the orbit of any point, say x_0 , contains the critical point in its closure. But if $\{f^i(x_0,\phi_0) - [f^i(x_0,\phi_0)] : i > 0\}$ does not contain x_0 then by altering f in a neighborhood of x_0 we can make a C^3 diffeomorphism without altering the orbit of x_0 , (see Figure 1) which contradicts Denjoy's theorem and the proof is complete. //



Figure 1

<u>Remark</u>: In fact the orbit of x_0 must approach x_0 from "both sides" as was shown by J. C. Yoccoz (personal communication).

5. <u>Proof of the theorem</u>. Without loss of generality we may assume $\mu_0 = \max \mu_i$. Now fix n > 0 and for y, $\hat{y} \in I$ let $y_i = f^i(y, \phi_0) - [f^i(y, \phi_0)]$, ies $\hat{y}_i = f^i(\hat{y}, \phi_0) - [f^i(\hat{y}, \phi_0)]$ where [•] denotes the greatest integer. Fix i_0 with $0 \le i_0 \le q_n$ and $|y_{i_0} - x_0| = \min\{|y_i - x_0| : i = 0, \dots, q_n - 1\}$. By condition (5) in the definition of f_0 there exist, say r, points in [0,1) such that $\frac{\partial^2 f}{\partial x^2}(x, \phi_0) = 0$. Hence by lemma 3 we see that taking n sufficiently large, so that the conclusion of lemma 5 holds,

$$\frac{\partial (f^{n})}{\partial x} (y, \phi_{0}) = \frac{\pi}{\pi} \frac{\partial f}{\partial x} (y_{j}, \phi_{0})$$

$$> \frac{\pi}{\pi} \frac{\partial f}{\partial x} (\hat{v}_{-j}, \phi_{0}) \cdot \mu_{4}^{r-1} \cdot \frac{\partial f}{\partial x} (y_{i_{0}}, \phi_{0})$$

-7-

This follows from lemma (3) since for each i, $0 \le i \le q_n$ there corresponds j, $0 > j > -q_n$ such that $\frac{\partial f}{\partial x}(y_j,\phi_0) \le \frac{\partial f}{\partial x}(y_i,\phi_0)$ except for $i = i_0$ and the r - 1 choices of $i \ne i_0$ where y_i is adjacent to a point with $\frac{\partial^2 f}{\partial x^2} = 0$. Hence, by an appropriate choice of $y \in I$ we obtain

$$\frac{\partial f^{n}_{n}}{\partial x} (y, \phi_{0}) > \frac{\mu_{0}}{\mu_{-q_{n}}} M_{4}^{r-1} \frac{\partial f}{\partial x} (y_{1_{0}}, \phi_{1})$$
$$> M_{4}^{r-1} \frac{\partial f}{\partial x} (y_{1_{0}}, \phi_{1}) .$$

Noting that

$$\frac{\partial f}{\partial x} (y_{i_0}, \phi_1) > 6M_3 \cdot (y_{i_0} - x_0)^2$$

we see that

$$\frac{\partial f^{n}}{\partial x}(y,\phi_{0}) > 6M_{4}^{r-1} \cdot M_{3}(y_{i_{0}} - x_{0})^{2}$$

for all y @ I. Hence, for some y @ I

(5.1)
$$\mu_{q_{n}} > 6\mu_{0}M_{4}^{r-1} \cdot M_{3}(y_{i_{0}} - x_{0})^{2}, \text{ i.e.,}$$
$$(\mu_{q_{n}}/(\mu_{0} \cdot 6M_{4}^{r-1} \cdot M_{3}))^{1/2} > |y_{i_{0}} - x_{0}|$$

Suppose $\mu_{q_n} < 1/c^n$ for all n, for some c > 1. Then, by (5.1), for each n there exists $y_n \in I$ such that if $\delta_n = \min_{\substack{i=0,\ldots,q_n=1 \\ i=0,\ldots,q_n=1}} |f^i(y_n,\phi_0) - f^i(y_n,\phi_0)| - x_0|$ then

$$\delta_n < c_2/c_1^{4n}$$

where $c_1 = c^{1/2}$ and $c_2 = (6 \cdot \mu_0 M_4^{r-1} M_3)^{-1/2}$. Fix m such that $\delta_{m+1} < \delta_m$. Then by lemma 4 there exists t, $0 < t < q_{m+2}$ with

$$|f^{t}(x_{0},\phi_{0}) - [f^{t}(x_{0},\phi_{0})] - x_{0}| < \delta_{m}$$

But since $1 \leq \frac{\partial f^n}{\partial \phi}(x,\phi)$ for all x, ϕ, n we see that there exists ε such that $f^t(x_0,\phi_0+\varepsilon) - [f^t(x_0,\phi_0+\varepsilon)] = x_0$ and $|\varepsilon| < \delta_m$. But then either $A_{p_{m+3}/q_{m+3}} \subseteq [\phi_0 - |\varepsilon|, \phi_0 + |\varepsilon|]$ or $A_{p_{m+4}/q_{m+4}} \subseteq [\phi_0 - |\varepsilon|, \phi_0 + |\varepsilon|]$ and hence by lemma 2

$$\delta_{m} > |\epsilon| > (M_{1}-1)^{2}/(4M_{2} \cdot M_{1}^{3q_{m+4}}) > (M_{1}-1)^{2}/(4M_{2} \cdot M_{1}^{3(M_{4}+1)^{4}q_{m}})$$

i.e. $c_2/c_1^m \ge (M_1-1)^2/(4M_2 \cdot M_1^m)$. Hence, we must have $3(M_4+1)^4 q_m$. Hence, we must have $3(M_4+1)^4$ $c_1 \le M_1^m$,

which completes the proof of (3.1) for μ . To show $\mu_{-\alpha_n}$ also satisfies $9(M_4+1)^R$ the theorem for $M_0 = M_1$ + 1 is similar. //

6. Example: Let
$$f_0(x) = x + \frac{1}{2\pi} \sin(2\pi x)$$
. Then the family $f(x,\phi) = x + \phi + \frac{1}{2\pi} \sin(2\pi x)$ satisfies the hypotheses, and if we let $\alpha_0 = \frac{\sqrt{5} - 1}{2}$ then we obtain $M_0 = 2^{2304} + 1$.

-9-

REFERENCES

- E. A. Coddington, N. Levinson, <u>Theory of Ordinary Differential Equations</u>, McGraw-Hill Book Company, New York, 1955.
- A. Denjoy, Sur les courbes definies par les equation differentiables a la surface du tore. J. de Math. Pure et Appliquees, 11 (1932), 33-375.
- G. R. Hall, A C Denjoy counter example, Ergodic Theory and Dynamical Systems, 1 (1981), 261-272.
- M. R. Herman, Sur la conjugaison differentiable des diffeomorphismes du cercle a des rotation, Publ. I.H.E.S., 49 (1979), 5-234.
- S. J. Shenker, Scaling Behavior in a map of a circle onto itself: Empirical results, (preprint).

GRH/jvs

#2512		BEFORE COMPLETING FORM
¥0510	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
# & J I Z		
TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
A Very Small Remark on Smooth Denjoy Maps of the Circle		Summary Report - no specifi
		reporting period
of the cifcle		6. PERFORMING ORG. REPORT NUMBER
AUTHOR(.)		8. CONTRACT OR GRANT NUMBER(+)
Glen Richard Hall		DAAC 20 80 C 0041
gien kichard mari		DAAG27-80-0-0041
PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK
Mathematics Research Center, Univ	versity of	Work Unit Number 1 -
610 Walnut Street	Wisconsin	Applied Analysis
Madison, Wisconsin 53706		
U. S. Army Research Office		April 1983
P. O. Box 12211		13. NUMBER OF PAGES
Research Triangle Park, North Caro	lina 27709	10
14. MONITORING AGENCY NAME & ADDRESS(I dittore	t from Controlling Office)	15. SECURITY CLASS. (of this report)
		UNCLASSIFIED
		15. DECLASSIFICATION/DOWNGRADING
		<u> </u>
7. DISTRIBUTION STATEMENT (of the abstract entered	in Block 20, if different fro	6 41
		м кероп)
8. SUPPLEMENTARY NOTES		m Kepok)
8. SUPPLEMENTARY NOTES 9. KEY WORDS (Continue on reverse eide if necessary ar	nd identify by block number)	
 SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde il necessary and Denjoy Maps, wandering intervals 	nd identify by block number)	m Kepok)
 SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse side if necessary and Denjoy Maps, wandering intervals ABSTRACT (Continue on reverse side if necessary and set of the s	nd identify by block number) d identify by block number)	
 SUPPLEMENTARY NOTES KEY WORDS (Continue on reverse elde il necessary and Denjoy Maps, wandering intervals ASSTRACT (Continue on reverse elde il necessary and In this note we show that if with one 'non degenerate' critical rotation number which is poorly apirrational) and f is a Denjoy matering interval cannot tend to 	d identify by block number) d identify by block number) f is a smooth point such that oproximable by ra ap, then the leng zero too rapidly	homeomorphism of the circle f has an irrational tionals (e.g., a quadratic th of the associated

D