AD-A136 470 UNCLASSIFIE	(LOWERSE MATHEMAT	IMAX THEOREI MICONTINU. TICAL STUDI TR-418 NOOO	.(U) STA ES IN TH	ANFORD HE SOCI	UNIV CA	J F MI	FOR	LSC 1	/•	
						END DATE FUND 2 84 DTIC				
1										
				_	_					



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS 1963 A

A136470

THE MINMAX THEOREM FOR U.S.C.-L.S.C. PAYOFF FUNCTIONS

by

Jean François Mertens



COPY NSPECTED

8

PREPARED UNDER

NATIONAL SCIENCE FOUNDATION GRANTS SES 8201373

AND THE

CENTER FOR RESEARCH ON ORGANIZATIONAL EFFICIENCY STANFORD UNIVERSITY CONTRACT ONR-N00014-79-C-0685

THE ECONOMICS SERIES

INSTITUTE FOR MATHEMATICAL STUDIES IN THE SOCIAL SCIENCES Fourth Floor, Encina Hall Stanford University Stanford, California 94305

St. Sec. 2 de

at the second second

THE MINMAX THEOREM FOR U.S.C.-L.S.C. PAYOFF FUNCTIONS*

. .

by

Jean-François Mertens**

Introduction

1.

Our aim is to get a "general minimax theorem" whose assumptions and conclusions are phrased only in terms of the data of the problem, i.e. the pair of pure strategy sets S and T and the payoff function on $S \times T$. For the assumptions, this means that we want to avoid any assumption of the type "there exists a topology (or a measurable structure) on S and (or) T such that..." For the conclusions, we are led to require that players have ε -optimal strategies with finite support, both because those are the easiest to describe in intrinsic terms, and because in any game where the value would not exist in strategies with finite support, all known general minmax theorems implicitly select as "value" either the sup inf or the inf sup by in effect restricting either player I or player II arbitrarily to strategies with finite support - so that the resulting "value" is completely arbitrary and misleading.

Those points are discussed in more detail in Section 3, after having proved a first theorem in Section 2.

*This research was supported in part by the National Science Foundation Grant SES 8201373 and in part by the Office of Naval Research Contract ONR-N00014-79-C-0685 at the Institute for Mathematical Studies in the Social Sciences, Stanford University.

**CORE, 34 Voie du Roman Pays, B-1348 Louvain La Neuve, Belgium

2. A First Theorem

<u>Minmax Theorem</u>: A two person zero sum game with compact strategy spaces has a value and each player has ϵ -optimal strategies with finite support if each player's payoff is, for any fixed strategy of his opponent, an uppersemicontinuous (u.s.c.) function of his strategy, and is bounded either from above or from below.

[In this Section 2, "compact" means compact and Hausdorff.]

<u>Proof</u>: Let f denote the first player's payoff function, S his strategy space, and T his opponent's strategy space. For any pair of mixed strategies σ and τ , we will denote by $f(\sigma,\tau)$ the expectation of f under the corresponding product measure, whenever this expectation is unambiguously defined (via Fubini's theorem).

A function f on a compact space K is u.s.c. if and only if f: $K + R \cup \{-\infty\}$ and $\forall \alpha \in R, \{x \mid f(x) \ge \alpha\}$ is closed, or equivalently, if and only if f is the pointwise limit of a decreasing net of continuous real valued functions on K. f is lowersemicontinous (l.s.c.) if and only if -f is u.s.c.. Remark that if f is l.s.c., then f is bounded from below, so each player guarantees himself a finite payoff with any pure strategy.

The proof starts by proving particular cases of the statement; the first of them is standard and is just recalled for the reader's convenience.

-2-

(A) Case where S or T -say T- is finite (Von Neumann).

<u>Proof</u>: Let $T = \{t_1, \ldots, t_n\}$ and let C denote the closed convex set $\{x \in \mathbb{R}^n | \exists \sigma \text{ probability on } S \pmod{\# \operatorname{Supp}(\sigma)} \leq n \operatorname{(Caratheodory)}\}$: $\forall i, \int f(s,t_i) d\sigma(s) \geq x_i \}$. Let $v = \max \min x_i$; then $\forall \varepsilon > 0$, $x \in \mathbb{C} \quad i$ $[C - (v + \varepsilon)] \cap \mathbb{R}^n_+ = \phi$, so that for a separating hyperplane $(\lambda_1, \ldots, \lambda_n)$ we have $\sup_{\substack{\langle \lambda, y \rangle < 0, \ \lambda_i \geq 0, \ \lambda_i \geq 0, \ \lambda_i \leq v = 1}} \langle \lambda_i = 1; y \in \mathbb{C}_{-}(v + \varepsilon)$ player II has a strategy λ such that $\forall x \in \mathbb{C}, \langle \lambda, x \rangle \leq v + \varepsilon$. When $\varepsilon \neq 0$, we get by compactness a strategy λ : $\langle \lambda, x \rangle \leq v, \ \forall x \in \mathbb{C}$.

(B) <u>Corollary</u>: Let T be an arbitrary set, f(s,t) u.s.c. on S for each $t \in T$. Then max $\inf_{\sigma \tau} f(\sigma,\tau) = \inf_{\tau} \max_{\sigma \tau} f(\sigma,\tau)$, where σ ranges over all regular Borel probabilities on S and τ over all probabilities with finite support on T.

<u>Proof</u>: For any finite subset \tilde{T} of T, let $\sum_{\varepsilon, T}$ denote the set of σ 's satisfying $\inf_{t \in T} f(\sigma, t) \ge \inf_{T} \max_{\sigma} f(\sigma, \tau) - \varepsilon$. By (A), the $\sum_{\varepsilon, T}$ form a decreasing net of non-empty compact sets, so have a non-empty intersection. Any σ_0 in the intersection yields $\inf_{T} f(\sigma_0, \tau) = \inf_{T} \max_{\sigma} f(\sigma, \tau)$.

(C) Case where one of the strategy spaces -say T- is metrisable.

(i) Claim: f is Borel measurable.

Let $\phi_1, \phi_2, \phi_3, \dots$ be a dense sequence of continuous functions on T. Denote by S_i the closed set $\{s \mid \forall t \in T, f(s,t) \ge \phi_i(t)\}$, and let $\psi_i(s,t) = \phi_i(t)$ for $s \in S_i$, = - ∞ otherwise. Obviously ψ_i is Borel measurable, and therefore also $f = \sup \psi_i$.

-3-

(ii) Denote by \sum (resp. \overline{T}) the space of regular Borel probabilities on S (resp. T), and by \sum_{f} (resp. \overline{T}_{f}) the probabilities with finite support.

f being Borel measurable and bounded either from above or from below, $f(\sigma,\tau)$ is well defined on $\sum \times \mathbf{T}$.

By (B), there exist $\bar{v} \in R$ and $\sigma_0 \in \Sigma$ such that

$$f(\sigma_0, t) \geq \vec{v} = \inf_{\tau \in \mathbf{I}_f} \sup_{\tau \in \mathbf{I}_f} f(\sigma, \tau) \qquad \forall t \in \mathbf{I}$$

and there exist $\underline{v} \in \mathbb{R}$ and $\tau_0 \in \overline{\Gamma}$ such that

$$f(s,\tau_0) \leq \underline{v} = \sup \inf_{\sigma \in \mathbf{f}_f} f(\sigma,\tau) \qquad \forall s \in S$$

On the one hand, one has always $\underline{v} = \sup \inf \underline{s} = \overline{v}$, on the other hand one gets $f(\sigma_0, \tau_0) = \int f(\sigma_0, t) d\tau_0(t) \ge \overline{v} \ge \underline{v} \ge \int f(s, \tau_0) d\sigma_0(s)$ $= f(\sigma_0, \tau_0)$, and thus the desired equality $\underline{v} = \overline{v}$.

(D) General Case

(i) Construction of a countable set of best replies.

Let $\overline{v} = \inf_{T} \sup_{f} f(\sigma, \tau)$, and let \overline{T}_{n} denote those $\tau \in \overline{T}_{f}$ with $\overline{T}_{f} \sum_{f} \sum_{f} f$ with $\overline{T}_{f} \sum_{f} \sum_{f} f$ with $\overline{T}_{f} \sum_{f} \sum_{f} f$ with $\overline{T}_{f} \sum_{f} f$ wi

$$\forall \phi \in F$$
, let $\circ_{\phi,k,n} = \{\tau \in \overline{T}_n \mid f \phi \, d\tau > \overline{v} - \frac{1}{k}\}$.

4

The $0_{\phi,k,n}$ form, for each fixed k and n, an open covering of the compact space \overline{I}_n . Let $\Phi_{k,n}$ denote a finite subset of F such that $\bigcup_{\phi \in \Phi_{k,n}} 0_{\phi,k,n} = \overline{I}_n$.

Then $\phi = \bigcup \phi_{k,n}$ is a countable subset of F, such that $\bar{v} = \inf_{\tau \in T_{p}} \sup \int \phi d\tau.$

(ii) Reduction to Case (C).

If ϕ_i enumerates ϕ , let $d(t_1, t_2) = \sum_i 2^{-i} |\phi_i(t_1) - \phi_i(t_2)| / \|\phi_i\|$. d defines a metrisable quotient space \tilde{T} of T, such that, if ψ denotes the quotient mapping, any $\phi \in \phi$ can be written as $g \circ \psi$, for some $g \in C(\tilde{T})$ - where, for some $s \in S$, $(g \circ \psi)(t) \leq f(s, t)$ $\forall t \in T$. Let Ψ denote the set of all $g \in C(\tilde{T})$ having this property.

Define f on $S \times T$ by $\tilde{f}(s,\tilde{t}) = \sup \{g(\tilde{t}) \mid g \in C(\tilde{T}), (g \circ \psi)(.) \leq f(s,.)\}$. Then, from (i), $\tilde{v} = \inf \sup_{\tau \in \overline{T}_{f}} \sup_{\phi \in \Phi} \int d\tau \leq \inf_{\tau \in \overline{T}_{f}} \sup_{\tau \in \overline{T}_{f}} \int (g \circ \psi)(t) d\tau(t) \leq \inf_{\tau \in \overline{T}_{f}} \sup_{s \in S} \int \tilde{f}(s,\tilde{t}) d\tilde{\tau}(\tilde{t})$. Obviously \tilde{f} is l.s.c. on \tilde{T} for each $s \in S$, and is the largest such function satisfying $\tilde{f}(s,\psi(t)) \leq f(s,t)$.

Let $h(s,\tilde{t}) = \inf \{f(s,t) \mid t \in \psi^{-1}(\tilde{t})\}$: if we show that $h(s,\tilde{t})$ is l.s.c. on \tilde{T} for each $s \in S$, it will follow that $\tilde{f} = h$, and therefore that \tilde{f} is also u.s.c. on S for each $\tilde{t} \in \tilde{T}$ - as an infimum of u.s.c. functions.

Since \tilde{T} is metrisable, denote by \tilde{t}_i a sequence converging to \tilde{t}_{∞} . Choose $t_i \in T$ such that $\Psi(t_i) = \tilde{t}_i$ and $f(s,t_i) \leq h(s,\tilde{t}_i) + 1/i$, and let t_{∞} be a limit point of t_i : we have $\Psi(t_{\infty}) = \tilde{t}_{\infty}$, and
$$\begin{split} h(s,\tilde{t}_{\infty}) &\leq f(s,t_{\infty}) \leq \lim_{i \to \infty} \inf f(s,t_{i}) \leq \lim_{i \to \infty} \inf h(s,\tilde{t}_{i}) - \text{hence the result.} \\ (\text{iii}) \quad \text{Conclusion.} \end{split}$$

 \tilde{f} on $S \times \tilde{T}$ satisfies all assumptions of (C), and from (ii) we have $\tilde{f}(s,\psi(t)) \leq f(s,t)$ and $\bar{v} \leq \inf_{\tilde{\tau} \in \widetilde{T}_{f}} \sup_{s \in S} \int \tilde{f}(s,\tilde{t}) d\tilde{\tau}(\tilde{t})$.

From (C), we know therefore that

$$\overline{v} \leq \sup_{\sigma \in \sum_{f}} \inf_{\tilde{t} \in \tilde{T}} \int \tilde{f}(s, \tilde{t}) d\sigma(s) \leq \sup_{\sigma \in \sum_{f}} \inf_{t \in T} f(\sigma, t)$$

which completes the proof, the reverse inequality being obvious.

3. Comments on the Present Result

 (i) The compactness assumption on both sides is really needed, as the following example due to H. Kuhn shows (private communication through R.J. Aumann, example originates from Kuhn's "Lecture Notes in Game Theory"
 [~1949], unpublished).

Player I picks a number x in [0,1], and player II chooses a continuous function ψ from [0,1] to itself with $\int \psi(t) dt = 1/2$. The payoff is $\psi(x)$. Then player I can guarantee 1/2 by choosing x uniformly distributed and player II by taking $\psi(t) = 1/2 \ \forall t$. But for any strategy of I with finite support, player II can choose an appropriate ψ that vanishes on the support - so player I cannot guarantee more than zero using strategies with finite support.

Remark here that if player II's strategy space is endowed with the uniform topology, we have a compact metric strategy space for player I, and a complete separable metric for II, and the payoff is uniformly bounded and a uniformly continuous function of player II's strategy -

-6-

in particular it is jointly continuous on $S \times T$: even with much stronger assumptions everywhere else, just the failure of the compactness assumption on one side makes the theorem break down.

(ii) Two directions of extension seem conceivable:

- The first would rely on some idea that our whole line of proof, of reducing oneself to some case where Fubini's theorem could be applied, is artificial, and that Fubini's theorem is irrelevant for this problem. In that case, one ought to be able to remove from the assumptions the last trace of Fubini's theorem - i.e. that the payoff function be uniformly bounded either from below or from above, and one might try to extend this to some "intrinsic" setting, i.e. essentially drop the assumption that the strategy spaces are Hausdorff, and just ask, using Alexander's subbase theorem, that the family of sets $S_{t,a} = \{s \in S \mid f(s,t) \geq a\}$ ($t \in T$, $a \in R$) has the finite intersection property, and similarly for Player II.

- The other direction would on the contrary look at Fubini's theorem as being basic, and consider that the "good" minmax theorem is the corollary sub (B) in the proof (from which all other minmax statements are easily obtained as corollaries - cfr. Section 4). So one would try to show that, for any payoff function satisfying the assumptions of the theorem, Fubini's theorem applies for any product of regular Borel measures. Then the theorem would become an immediate corollary of the statement sub (B). Furthermore, this would be a much more flexible tool, in conjunction with the methods mentioned in Section 4, using the stability properties of the set of functions that are measurable for

-7-

any product of tight probabilities (pointwise limits, algebraic operations, composition, etc.).

The best result we know in this direction requires however, f(s,t) to be continuous in each variable separately (and implies then, whenever S and T are Hausdorff spaces, that f is measurable for any tight probability on $S \times T$ - cfr. Bourbaki([1959] § 2, ex. 26). It is by the way the basic idea of a fundamental lemma underlying this theorem (that points in the closure of some set in C(K) are already in the closure of some countable subset of it) that we have used in part (D)(i) of the proof.

(iii) Motivations for this type of theorem are multiple.

(a) There is first the aesthetic motivation of obtaining the minmax theorem for mixed strategies under such assumptions that, by just adding the quasi-concavity-quasi-convexity assumption, one obtains Sion's assumptions for the existence of pure strategy solutions (adding a boundedness assumption to Sion's does not weaken his theorem).

(b) There is also the "effectiveness" point of view, that only discrete random variables can be effectively generated in finite time. (Any such discrete r.v. can be generated with a fair coin, by generating by successive tosses the successive bits of some random number $x (0 \le x < 1)$, stopping as soon as it is clear that, for some n, $\sum_{i \le n} p_i \le x < \sum_{i \le n} p_i$ and deciding then in favor of the n-th outcome: clearly a decision will be reached a.s. in finite time, and the n-th outcome will have probability p_n . This procedure can even be used with a biased coin, with unknown bias, by counting a pair of

-8-

successive tosses as one unit, giving a bit of "1" if first heads, then tails, a bit of "0" if first tails, then heads, and being inconclusive otherwise, in which case one should look to the next pair.)

It follows then that, for an operational concept of value, the players should be able to construct ε -optimal strategies where, in each information set, they are restricted to discrete mixtures.

In particular, for general minmax theorems, for normal form games, which may represent games with a single information set, one is led to the requirement of ε -optimal discrete mixtures. Under our assumptions, the players have ε -optimal finite mixtures, and the guaranteed payoff varies continuously with the probability vector used, so in any realistic model the player can play ε -optimally, even in bounded time.

(c) More important is the "safety" consideration, that, even neglecting the above, the player should always consider that his opponent might have an infinity of information sets at his disposal, or some analog device such as a continuous roulette wheel, and thus would be able to generate continuous random variables. In such a case, the evaluation of the expected payoff (even if players agree on some compact topologies on S and T, and to use only integration theory for regular Borel measures on compact spaces) would depend on the order of integration - except if one knows in addition that Fubini's theorem holds for the payoff function. For instance, players may think that the corollary sub (B) in the proof is the good minmax theorem; given a payoff function that would satisfy those assumptions for both players, player I might

-9-

find an optimal strategy that "guarantees" him +1, and similarly, player II applying the same corollary for him might find an optimal strategy that "guarantees" to him that the payoff will not exceed -1. Obviously, realizing this, both players can only conclude that the only thing they really can guarantee, is what they can guarantee with mixtures with finite support. Since "-1" is always \leq to "+1" this argument is perfectly general. (By our theorem, one could, in this example, have a strict inequality only for unbounded payoff functions; but still, the theorem is needed for that, and anyway the argument remains, since it depends only on the weak inequality $-1 \leq +1$.)

And things can be much worse, because nothing compels the player to agree on some integration theory - or even on some topology for the strategy spaces. You could even happen to be playing against hubins and Savage [1965], using finitely additive randomizations. In such cases, only finite mixtures remain safe, unambiguous, and devoid of arbitrariness (i.e. only such a solution depends only on the data of the problem, i.e. the pure strategy sets -as sets- and the payoff function -as a real valued function on the product of the pure strategy sets).

(d) The proof shows that the result is much easier to get for compact metric spaces. Is it really worthwhile to make a substantially bigger effort in order to get rid of the metrisability assumption?

A first answer would be that this simplifies the statement of the theorem (one can drop "metric"), and that any effort in a proof is worthwhile if it leads to a simpler (and more powerful) theorem.

-10-

A much more important reason is, however, that in any game where a continuous variable can be observed before some action is taken, the strategy spaces are non-metrisable. Certainly one does not want to exclude such models from game theory. Besides the obvious cases (observation of a price, or a quantity, or a continuous random variable), this would also exclude any differential game (time being a continuous variable) and any game with a continuum of players (just to define its characteristic function, one needs the value of a zero sum game between two opposing coalitions, where the strategy space of each coalition, as an uncountable product, cannot be metrisable). Those problems are amply documented in the literature; for instance, R.J. Aumann [1964] suggests the use of behavioral strategies to get around this type of problem; in other contexts (differential games, games with a continuum of players, etc.) various other restrictions are imposed on the strategy sets.

Besides the inconvenience of requiring additional structure (a topology, or a measurable structure), as was discussed in the previous point, such procedures have the unpleasant feature of restricting the strategy space -for instance, an arbitrary pure strategy is not necessarily a behavioral strategy, since it may lack the measurability requirement. For a minmax theorem, it is of course a net gain if one can show that some player's (ε)-optimal strategies have in fact some additional regularity property; however, it is disturbing when it is only shown the the strategy is safe against some subclass of the opponent's strategies. Indeed, the opponent should ideally not even be assumed to follow a strategy, -he is just playing- and one would like the minmax theorem

-11-

to have essentially the same force as in the perfect information case, i.e. that any "play" consistent with your ε -optimal strategy yields a payoff $\geq v - \varepsilon$.

4. Other Minmax Theorems

A) Measure Theory (non-Hausdorff compact spaces)

For any compact space S, denote by C the convex cone of bounded l.s.c. functions on S, and let $E = C \setminus C$. Denote by P the set of monotonic real valued sublinear (i.e. positively homogeneous of degree l and subadditive $(p(x + y) \leq p(x) + p(y))$ functions on C. P is ordered in the usual way $(p_1 \leq p_2 \text{ iff } p_1(f) \leq p_2(f) \forall f \in C)$.

Definition: M(S) is the set of minimal elements of P.

Lemma 1: $\forall p \in P, \exists \mu \in M(S): \mu \leq p$.

<u>Proof</u>: $\forall p \in P$, $\forall f \in C$ let $\alpha \leq 0$ be such that $f \geq \alpha$. Then $\forall q \in P$ s.t. $q \leq p$: $p(f) \geq q(f) \geq q(\alpha) \geq -q(-\alpha) \geq -p(-\alpha)$, so that the set of possible values q(f) is a bounded interval. Zorn's lemma then yields a minimal such q.

Lemma 2: Any $\mu \in M(S)$ is a positive linear functional on E, satisfying

 $\mu(f) = \inf \{\mu(g) \mid g \in C, g \ge f\} \quad \forall f \in E .$

<u>Proof</u>: Let $\tilde{\mu}$: $E \neq R$ be defined by the above formula. Clearly $\tilde{\mu}$ is real valued (if $f \geq a$, then $\tilde{\mu}(f) \geq \tilde{\mu}(a) \geq \mu(a) > -\infty$). Further, $\tilde{\mu}$ is obviously monotonic and sublinear, coinciding with μ on C. The Hahn-Banach theorem says that any such $\tilde{\mu}$ is the supremum of a family of linear functionals. Any linear functional $\zeta \leq \tilde{\mu}$ is positive, because ≤ 0 on the negative functions. If there was a second such ζ , say $\tilde{\zeta}$, one would have $\zeta(f) \neq \tilde{\zeta}(f)$ for some $f \in C$. Then either ζ or $\tilde{\zeta}$, restricted to C, would be some element of P $\leq u$, and $\neq u$, thus μ would not be minimal. Therefore this ζ is unique, and thus coincides with $\tilde{\mu}$: $\tilde{\mu}$ is a positive linear functional on E.

Lemma 3: M(S) is the set of positive linear functionals μ on E such that

 $\forall f \in C , \mu(-f) = \inf \{\mu(g) \mid g \in C, g \ge -f\} .$

<u>Proof</u>: One direction is given by Lemma 2. In the other direction, we have clearly $\mu \in P$, thus (Lemma 1) $\exists \nu \in M(S)$: $\nu(h) \leq \nu(h) \forall h \in C$. Thus $\nu(-f) \geq \mu(-f) = \inf \{\mu(g) \mid g \in C, g \geq -f\}$ $\geq \inf \{\nu(g) \mid g \in C, g \geq -f\} = \nu(-f)$ (by Lemma 2), from which the equality of μ and ν .

Lemma 4: M(S) is the set of regular Borel measures on S, i.e. the set of positive, bounded, countably additive measures on the Borel sets of S satisfying $\mu(A) = \sup \{\mu(F) \mid F \subseteq A, F \text{ closed}\}$ = inf $\{\mu(0) \mid 0 \supseteq A, 0 \text{ open}\}$ for any μ -measurable set A.

<u>Proof</u>: Follows from a standard Daniell-type extension procedure. We sketch just a typical sequence of steps:

Sal.

• Denote by LSC the set of l.s.c. functions with values in $R \cup \{+\infty\}$, and USC = - LSC. Let $\mu_1(f) = \sup \{\mu(g) \mid -g \in C, g \leq f\} \forall f \in LSC$

Let $\mu^{*}(f) = \inf \{\mu_{1}(g) \mid g \in LSC, g \geq f\}$ for any \overline{R} -valued f.

• Lemma 2 implies $\mu^* = \mu$ on E, and $\mu^* = \mu_1$ on LSC.

• Dini's theorem implies that, if f $\ensuremath{_\alpha} \in \mbox{LSC}$ is filtering increasing, then

$$\mu^*(\lim f_{\alpha}) = \lim \mu^*(f_{\alpha}) .$$

• It follows that $\mu^*(f + g) = \mu^*(f) + \mu^*(g)$ on LSC.

• The sublinearity of μ^* follows (avoiding $(+\infty) + (-\infty)$ right-hand members, and using the convention $(0) \cdot (\infty) = 0$).

• Finally one gets: if h_n is any increasing sequence, with lim $\mu^*(h_n) > -\infty$, then $\mu^*(\lim h_n) = \lim \mu^*(h_n)$. (It is sufficient to consider $\mu^*(h_n)$ finite; choose $f_n \in LSC$, $f_n \ge h_n$, $\mu^*(f_n) \le \mu^*(h_n) + \varepsilon \cdot 2^{-n}$. Let $\psi_n = \sup f_1$. ψ_n is an increasing sequence in LSC such that $\psi_n \ge h_n$. $\psi_n = \max(\psi_{n-1}, f_n)$ implies $\mu^*(\psi_n) + \mu^*(h_{n-1}) \le \mu^*(\psi_n) + \mu^*(\min(f_n, \psi_{n-1})) = \mu^*(\psi_n + \min(f_n, \psi_{n-1}))$ $= \mu^*(f_n + \psi_{n-1}) = \mu^*(f_n) + \mu^*(\psi_{n-1})$, thus $\mu^*(\psi_n) - \mu^*(h_n)$ $\le [\mu^*(f_n) - \mu^*(h_n)] + [\mu^*(\psi_{n-1}) - \mu^*(h_{n-1})]$ and therefore $\mu^*(\psi_n) - \mu^*(h_n) \le \sum_{\substack{i \le n}} [\mu^*(f_i) - \mu^*(h_i)] \le \varepsilon \sum_{\substack{i \le n}} 2^{-i} \le \varepsilon$, so that lim $\mu^*(h_n) \le \mu^*(\liminf h_n) \le \mu^*(\liminf \psi_n) = \lim \mu^*(\psi_n) \le \lim \mu^*(h_n) + \varepsilon$).

• It follows that μ^* , restricted to indicator functions, is an outer measure.

-14-

• Let $L = \{f \mid \mu^{\#}(f) > -\infty, \mu^{\#}(-f) > -\infty, \mu^{\#}(f) + \mu^{\#}(-f) \leq 0\}$. By the sublinearity of $\mu^{\#}$, L is a vector space and $\mu^{\#}$ is a linear functional on L. Further $E \subseteq L$ since $\mu^{\#}$ coincides with μ on E, and in addition $f \in LSC$, $\mu^{\#}(f) < \infty$ implies $f \in L$.

• Since for any open sets **0** and U, $I_0 \in L$, and $I_{O\cap U} \in L$, we have also $I_{O\setminus U} \in L$, so we get $\mu^*(0) = \mu^*(0 \cap U) + \mu^*(0 \setminus U)$. Thus, for any set A, since $\mu^*(A) = \inf \{\mu^*(0) \mid A \subseteq 0\}$, we get $\mu^*(A) \ge \mu^*(A \cap U) + \mu^*(A \setminus U)$, so that any open set U is μ^* -measurable and thus all Borel sets are μ^* -measurable.

• The formula $\mu(A) = \inf \{\mu(0) \mid A \subseteq 0, 0 \text{ open}\}$ for any μ^* -measurable set A implies $\mu(A) = \sup \{\mu(F) \mid F \subseteq A, F \text{ closed}\}$, and thus the μ^* -measurable sets are just the completion of the Borel sets, and μ is a regular Borel measure.

• Finally, $\int f d\mu = \mu^*(f)$ first for the convex cone spanned by the constants and the indicators of open sets, by linearity, next for all $f \in LSC$, by monotone convergence, and therefore for all $f \in L$ and in particular all $f \in E$. It also follows that $L = L_1(\mu)$.

• The regular Borel measure μ is unique, because its value on the open sets is determined.

 \cdot Conversely, clearly the integral for any regular Borel measure is a functional in M(S).

The essential results of this section are summarized in the following:

Proposition and Denote by M(S) the set of positive, regular Borel measures on S. Then:

• $\forall p \in P \exists \mu \in M(S): \mu \leq p \text{ on } C \text{ (Lemma 1).}$

M(S) is a convex cone (and a complete lattice) (immediate from Lemma 4).
If the sum of 2 positive linear functionals on E is in M(S), each one is also (immediate from Lemma 3).

. Define the "weak*-topology" on M(S) as the coarsest topology for which $\int f d\mu$ is lowersemicontinuous $\forall f \in C$, and thus for all l.s.c. f. Then addition and scalar multiplication are continuous, points are closed and all sets $\{\mu \mid \mu(1) = \lambda\}$ and $\{\mu \mid \mu(1) \leq \lambda\}$ ($\lambda > 0$) are closed and compact.

<u>Proof</u>: The continuity of addition and scalar multiplication is immediate. { μ } is closed because, if $\tilde{\mu}$ is in the closure of μ , one has $\tilde{\mu}(f) \leq \mu(f) \quad \forall f \in C$, thus $\tilde{\mu} = \mu$ by the minimality of μ .

Since all constants belong to C, there only remains to prove the compactness of $\{\mu \mid \mu(1) \leq \lambda\}$. For any ultrafilter on this set, let ϕ denote its pointwise limit in the set of positive linear functionals on E. By Lemma 1, $\exists \nu \in M(S)$ with $\nu \leq \phi$ on C: ν is a limit point in M(S).

Note that the above applies whenever C is a lattice and convex cone of bounded functions on some set S, containing the constants and such that $f_i \in C$, $f_i \leq f_{i+1} \leq 0$ implies $\lim f_i \in C$, and $\lim f_i > -1$ implies $\exists i: f_i > -1$. The Borel sets are then interpreted as the σ -field generated by C, and the regularity is with respect to the

-16-

 $\{f \leq 0\}$ ($f \in C$) as closed sets. (Dini's theorem should be used only for sequences then.)

In particular, everything applies as soon as S is countably compact (any sequence has a cluster point). Compactness only yields the additional τ -smoothness property: $\mu(\bigcup_{\alpha} 0_{\alpha}) = \sup_{\alpha} \mu(0_{\alpha})$ for any increasing net of open sets 0_{α} .

In the particular case where S is Hausdorff, i.e., $\forall f. g \in C$ s.t. $f > -g \exists h: -g \leq h \leq f$, h continuous, the above results yield Riesz' theorem.

We now obtain also, using standard techniques, a Fubini theorem (the same theorem and proof obviously holds for arbitrary products, finite or infinite; the minmax theorem is however, concerned only with products of two factors. The contribution of the theorem is obviously to get the measure on the Borel σ -field of the product, instead of the product σ -field):

<u>Proposition 2</u>: Let K_1 and K_2 denote two compact spaces, $K = K_1 \times K_2$. Let $\mu_1 \in M(K_1)$, $\mu_2 \in M(K_2)$. Then

(a) There exists a unique $\mu \in M(K)$, denoted $\mu_1 \otimes \mu_2$, such that $\mu(F_1 \times F_2) = \mu_1(F_1)\mu_2(F_2) \forall F_i$ closed in K_i .

(b) For any lower semicontinuous f, there exists a sequence of functions $\psi_i = \varepsilon_i \prod_{\substack{i = 1 \ F_1^i \times F_2^i}} with \varepsilon_i > 0$, F_j^i closed in K_j , and a constant function $\psi_0 \leq 0$, such that

$$\sum_{i} \psi_{i} \leq f \text{ and } \int f d\mu = \sum \int \psi_{i} d\mu .$$

(c) For any μ -quasi-integrable f one has that

- for μ_1 - almost every x, f(x,.) is μ_2 -quasi-integrable.

- $\int f(x,y)d\mu_2(y)$ is μ_1 -quasi-integrable (and l.s.c. if f is) - $\int [\int f(x,y)d\mu_2(y)]d\mu_1(x) = \int f d\mu$.

(An extended real valued function f called μ -quasi-integrable if it is μ -measurable and either $\int f^+ d\mu \neq \infty$ or $\int f^- d\mu \neq \infty$).

<u>Proof</u>: Denote by $\tilde{\mu}$ the product measure on the product of the Borel σ -fields; let $\tilde{\mu}$ be a positive linear functional on all bounded functions that extends $\int (.) d\tilde{\mu}$, and let (Lemma 1) $\tilde{\mu} \in M(K)$ minorize $\tilde{\mu}$ on C. Then we have

$$\tilde{\mu}(0_1 \times 0_2) \leq \tilde{\mu}(0_1 \times 0_2) = \tilde{\mu}(0_1 \times 0_2) = \mu_1(0_1)\mu_2(0_2)$$

for all 0 open in K_i, and

$$\overline{\overline{\mu}}(\mathbf{F}_{1} \times \mathbf{F}_{2}) \geq \overline{\mu}(\mathbf{F}_{1} \times \mathbf{F}_{2}) = \widetilde{\mu}(\mathbf{F}_{1} \times \mathbf{F}_{2}) = \mu_{1}(\mathbf{F}_{1})\mu_{2}(\mathbf{F}_{2})$$

for all \mathbf{F}_{1} closed in K_{1} .

Using the regularity of μ_i , it follows that, for any μ_i measurable sets A_i , $\overline{\mu}(A_1 \times A_2) = \mu_1(A_1)\mu_2(A_2)$.

Thus the product measure $\tilde{\mu}$ can be extended to a regular Borel measure. By the regularity of μ_1 and μ_2 , any Borel measure μ satisfying $\mu(F_1 \times F_2) = \mu_1(F_1)\mu_2(F_2)$ will satisfy $\mu(A_1 \times A_2) \ge \mu_1(A_1)\mu_2(A_2)$ for any Borel sets A_1 , and thus going to complements, the inverse inequality: it will be an extension of $\tilde{\mu}$. Since any open set 0 is the limit of the increasing net of sets 0_{α} , where the sets 0_{α} are all finite unions of products $0_1 \times 0_2 \subseteq 0$, and thus $\tilde{\mu}$ -measurable, the regularity of μ will imply that μ is uniquely determined on all open sets, and thus on all Borel sets.

This proves statement (a), and (b) is obtained by pursuing a bit the same argument (representing the $\begin{array}{c} \mathbf{0}\\ \alpha\end{array}$ as finite disjoint unions of products of Borel sets, and using there the regularity of the μ_i 's, going from indicator functions of open sets to l.s.c. functions f, one obtains a finite sequence ψ_i with $\sum_i \psi_i \leq f$ and $\sum_i \psi_i d\mu$ as close as required to $\int f d\mu$. Iterating this conclusion with the l.s.c. function $f - \sum_i \psi_i$ yields (b)).

(c) is proved by first noting its truth when $f = I_{C}$, next, a using the regularity of all measures, that it remains therefore true for indicators of open sets, and hence for all lowersemicontinuous f. The general statement follows then because, for any quasi-integrable f, $\int fd\mu = \inf \{ \int gd\mu \mid g \mid 1.s.c. \geq f \} = \sup \{ \int hd\mu \mid h \mid u.s.c. \leq f \};$ this was proved in Lemma 4.

B) A Basic Tool

We return now to the corollary sub (B) of our minmax theorem, dropping the Hausdorff assumption on S. Remark that, for any τ with finite support, $f(s,\tau)$ is u.s.c. in s; it therefore achieves its maximum, and max $f(s,\tau) = \max_{\sigma \in M(S)} f(\sigma,\tau)$. Let $\bar{v} = \inf_{\sigma \in M(S)} \max_{\tau = \sigma} f(\sigma,\tau)$ (τ ranging over probabilities with finite support). By part (A) of the proof there exists, for any finite subset \tilde{T} of T, a probability σ with finite support on S such that $\min_{\tau \in T} f(\sigma,\tau) \geq \bar{v}$. Since f(.,t) is u.s.c., and σ is a positive $t\in \tilde{T}$ linear functional on E, there exists by Lemma 1, $\tilde{\sigma} \in M(S)$ with $\tilde{\sigma}(g) \geq \sigma(g)$ for all u.s.c. g. In particular, $\tilde{\sigma}(1) = \sigma(1) = 1$, and $\min_{\tau} f(\tilde{\sigma}, t) \geq \bar{v}$. Let $\tilde{\lambda}_{\tilde{T}} = \{\sigma \in M(S) \mid \sigma(1) = 1 \text{ and } f(\tilde{\sigma}, t) \geq \bar{v}$ $t\in \tilde{T}$. \sum_{T} is obviously closed in M(S), thus, by Proposition 1, the sets T \sum_{T} form a decreasing net of nonempty, closed, compact sets, and therefore have a nonempty intersection. Any σ in the intersection fills the bill.

Observe that it follows from this that we can phrase the assumptions of that corollary in a fully intrinsic way, using Alexander's subbase theorem:

<u>Theorem 2</u>: Denote by S and T two sets, f: $S \times T \rightarrow R \cup \{-\infty\}$. For any $\alpha \in \mathbb{R}$ and $t \in T$, let $S_{t,\alpha} = \{s \in S \mid f(s,t) \geq \alpha\}$. If the family of sets $S_{t,\alpha}$ has the finite intersection property, and if M(S) denotes the set of regular Borel probabilities on S endowed with the coarsest topology for which the functions f(.,t) $(t \in T)$ are u.s.c., then

$$\begin{array}{ll} \max & \inf f(\sigma, t) = \inf \max f(s, \tau) \\ \sigma \in \mathbb{M}(S) \ t \in \mathbb{T} & \tau \in \mathbb{T}_{f} \ s \in S \end{array}$$

Remarks:

1) The finite intersection property can be rephrased as asking that any pointwise limit of pure strategies, i.e. of functions f(s,.), be dominated by some pure strategy, i.e. by some function $f(s_0,.)$.

2) This property is substantially weaker than the usual compactness assumption. For instance, if T is a single point, it just asks that f(s) attains its maximum.

3) Using the remark after Proposition 1, it is sufficient to ask that the class of sets $S_{t,\alpha}$ be countably compact, i.e. that any countable family of such sets has the finite intersection property. In that case M(S) becomes the set of probabilities on the σ -field

-20-

generated by the functions f(.,t), regular w.r.t. the countable intersections of finite unions of sets S_t , (in particular, if S with its coarsest topology is countably compact, M(S) is just the set of regular Borel probabilities on S).

This amounts to ask that, for any countable subset \tilde{T} of T, any pointwise limit of pure strategies s_n be dominated on \tilde{T} by some pure strategy s_0 .

4) Given the above measure theory, including Fubini, essentially our whole proof of Section 2 would go through without the Hausdorff assumption, and thus give the corresponding "intrinsic" result. Indeed, the use of continuous functions in part "D" has nothing essential; lowersemicontinuous ones would do as well, giving a reduction to the case where T is compact with countable base, in which case the proof sub (C) works just as well. The only troublesome point lies in the last sentence sub (ii) of case (D) where the regularity of the topology ("every point has a basis of closed neighborhoods") seems to be used in an essential way.

Anyways, it goes through if one assumes that one of the spaces S and T either has a countable basis or is Hausdorff.

C) Other Techniques from the Literature

Most results of the literature (except of course Sion's) are obtained, or are at least obtainable, by applying to some particular case of Theorem 2 one of the two following generalization techniques:

1) The first one goes back to Wald. It states that, if for any $\varepsilon > 0$ it is possible to find subsets S_{ε} of S and T_{ε} of T such that, by some independent argument, the game restricted to S_{ε} and T_{ε} has a value, and such that $\forall s \in S$, $\exists s' \in S_{\varepsilon}$: $f(s,t) \leq f(s',t) + \varepsilon \quad \forall t \in T_{\varepsilon}$, and similarly $\forall t \in T$, $\exists t' \in T_{\varepsilon}$: $\forall s \in S_{\varepsilon}$ $f(s,t) \geq f(s,t') - \varepsilon$, then the given game has a value.

2) The other idea is based on Karlin, and amounts essentially to using the monotone convergence theorem, or Fatou's lemma.

For instance, after applying Theorem 2 to some game, and having found the value v and player I's optimal strategy μ , one looks for some class C of functions $\phi(s)$ such that $\int \phi(s) d\mu \geq v$. Say C contains all bounded measurable functions that are minorized by some f(.,t) ($t \in T$) and their convex combinations, and also the limit of any decreasing sequence ϕ_n of such functions (one could still add all functions ψ which are, for any regular Borel probability μ , in the equivalence class of some $\phi \in C$)^{$\underline{1}'$}.

Then any game $\tilde{f}(s,t)$ such that $\tilde{f} \leq f$ and $\forall t, \tilde{f}(.,t) \in C$ would have the same value and optimal strategy.

Usually f is constructed from \tilde{f} by taking the smallest u.s.c. function majorizing it, but sometimes the argument has to be applied both ways.

To illustrate, we give a typical application: S and T are compact metric, f is a bounded measurable (i.e. measurable for any product measure) function on $S \times T$ such that, if $E = \{(s,t) \mid f$ is not continuous in s or in t at (s,t)}, then $\forall s$, $\#\{t \mid (s,t) \in E\} \leq 1$ and $\forall t$, $\#\{s \mid (s,t) \in E\} \leq 1$. (Remark that measurability would follow for instance if we had, denoting by \tilde{E} the (F_{σ}) set of points of discontinuity of f, $\forall s \{t \mid (s,t) \in \tilde{E}\}$ and $\forall t \{s \mid (s,t) \in \tilde{E}\}$ are at most countable.) Such a game has a value.

Indeed, let $\phi_1(s_0, t_0) = \lim \sup f(s, t_0), \phi_2(s_0, t_0) = \lim \inf f(s_0, t).$ $s + s_0$ $s + s_0$ Consider an optimal strategy σ of player I in the game with payoff $f \lor \phi_1$ (Theorem 2). $\forall t_0 \in T$, by considering an appropriate sequence t_i converging to t_0 , one can have $\lim_{i \to \infty} (f \lor \phi_1)(s, t_i) = [\phi_2 \land (f \lor \phi_1)]$ (s, t_0) . Thus, by the above argument, say in the form of Fatou's lemma, σ is still an optimal strategy of player I for $[\phi_2 \land (f \lor \phi_1)]$, guaranteeing the same value \lor_1 , and player II's ε -optimal strategy with finite support τ_{ε} guarantees the same \lor_1 for $(f \lor \phi_1)$.

Inverting the roles of the players, we get a value v_2 for both $f \wedge \phi_2$ and $[\phi_1 \vee (f \wedge \phi_2)]$, an optimal strategy τ for II and an ϵ -optimal strategy with finite support σ_{ϵ} for I. But $\phi_1 \vee (f \wedge \phi_2) \ge \phi_2 \wedge (f \vee \phi_1)$, so, by applying Fubini's theorem to $\sigma \odot \tau$ and those payoff functions, we get $v_2 \ge v_1$. Since $f \vee \phi_1 \ge f \ge f \wedge \phi_2$, we have $v_2 = v_1 = v$, and σ_{ϵ} and τ_{ϵ} guarantee v in those 3 games, so in f.

Karlin's "general game of timing of class II" (Karlin, [1959], Chap. V, Ex. 20), falls in this category.

Similarly, assume S and T are compact, and f(s,t) is bounded and Borel-measurable on $S \times T$. Assume $\exists S^0 \subseteq S$ and $T^0 \subseteq T$ such that f(s,.) is l.s.c. on T for $s \in S^0$, f(.,t) is u.s.c. on S for $t \in T^0$, and such that $\forall s \in S \exists s_n \in S^0$ with $\liminf_{n \to \infty} f(s,t)$ $\geq f(s,t) \forall t \in T$, and that $\forall t \in T \exists t_n \in T^0$ with lim sup $f(s,t_n) \leq f(s,t) \forall s \in S$. Then this game has a value, $n \to \infty$ and both players have ε -optimal strategies with finite support carried by S^0 and T^0 respectively.

-23-

Indeed, Theorem 2 yields a value \overline{v} for the game on $S \times T^0$, and by Fatou's lemma, player I's optimal strategy is still safe against all t \notin T. Similarly one can apply Theorem 2 and next Fatou's lemme to the game on $S^0 \times T$, with value \underline{v} . Fubini's theorem applied to f and the product of the optimal strategies yields then $\underline{v} = \overline{v}$, hence the result.

C. Waternaux [1983]'s "auxiliary game" is of this type.

Most other classical examples (like all examples in Karlin [1950], Restrepo's "general silent duel" or Karlin's "two machinegun duel") can be shown to have a value by the same technique, applied in a more or less similar (and often easier) way.

Footnotes

1/

If this is done, the resulting class C is identical to the set of all functions ϕ , such that $\forall n$, $\phi v(-n)$ is in the closed convex hull of the set of bounded measurable function \geq some f(.,t), in the space of all bounded universally measurable functions in duality with the space of regular Borel measures on S. This shows that our technique is really equivalent to Karlin's apparently more powerful closure methods (Karlin, [1950]).

The only point to show is that the class \tilde{C} of bounded functions in C is closed, it is thus sufficient to show closedness in $L_{\omega}(\mu) \sigma(L_{\omega}(\mu), L_{1}(\mu))$. By the Krein-Smulian theorem on weak*closed convex sets, it is sufficient to show that its intersection with any ball of L_{ω} is $\tau(L_{\omega}, L_{1})$ -closed. Since the Mackey topology coincides on balls of L_{ω} with the topology of convergence in measure (this result of Grothendieck follows easily from Dunford-Pettis' equiintegrability criterion for weak compactness in L_{1}), it follows from Egorov's theorem that it is sufficient to show that the limit of any uniformly bounded a.e. convergent sequence (g_{n}) in \tilde{C} is in \tilde{C} .

There is obviously no loss in assuming further that each g_n is larger than some convex combination of functions f(.,t). Then lim sup g_n is obviously in \tilde{C} .

-25-

References

For general reference in mathematics, the reader may want to consult Halmos [1950], Kelley [1955] and Kelley and Namioka [1963]; and on minmax theorems, Parthasarathy and Raghavan [1971].

- Aumann, R.J. [1964], "Mixed and Behavioural Strategies in Infinite Extensive Games," in Dresher, Shapley and Tucker (eds.), pp. 627-650.
- Bourbaki, N. [1959], <u>Eléments de Mathematique</u>, Livre VI: Intégration, Ch. VI: <u>Intégration vectorielle</u>, Hermann, Paris.
- Dresher, M., A.W. Tucker and P. Wolfe (eds.) [1957], <u>Contributions to</u> <u>the Theory of Games III (Annals of Mathematical Studies</u>, Vol. 39), Princeton University Press, Princeton, New Jersey.

Dresher, M., L.S. Shapley and A.W. Tucker (eds.) [1964], <u>Advances in</u> <u>Game Theory</u> (<u>Annals of Mathematical Studies</u>, Vol. 52) Princeton University Press, Princeton, New Jersey.

- Dubins, L. and L.J. Savage [1965], <u>How to Gamble if You Must</u>, MacGraw Hill, New York.
- Grothendieck, A. [1952], "Critères de compacité dans les espaces fonctionnels généraux," American Journal_of Mathematics 74, pp. 168-186.

Grothendieck, A. [1953], "Sur les applications linéaires faiblement compactes d'espaces de type C(K)," <u>Canadian Journal of</u> <u>Mathematics</u> 5, pp. 129 - 173.

Halmos, P.R. [1950], Measure Theory, Van Nostrand, New York.

- Karlin, S. [1950], "Operator Treatment of the Minmax Principle, I," in H.W. Kuhn and A.W. Tucker (eds.) pp. 133 - 154.
- Karlin, S. [1959], <u>Mathematical Methods and Theory in Games</u>, Programming and Economics, Vol. II, Addison-Wesley.

Kelley, J.L. [1955], General Topology, Van Nostrand, New York.

- Kelley, J.L., I. Namioka, et al. [1963], <u>Linear Topological Spaces</u>, Van Nostrand, New York.
- Kneser, H. [1952], "Sur un théorème fondamental de la théorie des jeux," Comptes Rendus de l'Académie des Sciences 234, pp. 2418-2420.

-.'0-

- Kuhn, H.W. and A.W. Tucker (eds.) [1950], <u>Contributions to the Theory</u> of <u>Games, I</u> (<u>Ann. Math. Studies</u>, Vol. 24), Princeton University Press, Princeton, New Jersey.
- Nikaido, H. [1954], "On von Neumann's Minimax Theorem," <u>Pacific Journal</u> of <u>Mathematics</u> 4, pp. 65-72.
- Parthasarathy, T. and T.E.S. Raghavan [1971], <u>Some Topics in Two-Person</u> <u>Games</u>, American Elsevier Publishing Company, New York.
- Restrepo, R. [1957], "Tactical Problems Involving Several Actions," in Dresher, Tucker, and Wolfe (eds.), pp. 313-337.
- Sion, M. [1958], "On General Minimax Theorems," <u>Pacific Journal of</u> Mathematics 8, pp. 171-176.
- Sion, M. and P. Wolfe [1957], "On a Game without a Value," in Dresher, Tucker and Wolfe (eds.), pp. 299-305.
- Teh Tjoe-Tie [1963], "Minmax Theorems on Conditionally Compact Sets," Annals of Mathematical Statistics 34, pp. 1536-1540.
- Wald, A. [1945], "Generalization of a Theorem by von Neumann Concerning Zero-Sum Two-Person Game," Annals of Mathematics 46, pp. 281-286.
- Wald, A. [1950], "Note on Zero-Sum Two-Person Games," <u>Annals of</u> Mathematics 52, pp. 739-742.
- Waternaux, C. [1983], "Minmax and Maxmin of Repeated Games Without A Recursive Structure," CORE Discussion Paper 8313, Université Catholique de Louvain.
- Yanovskaya, E.B. [1964], "Minimax Theorems for Games on the Unit Square," Theory of Probability and its Applications 9, pp. 500-502.

REPORTS IN THIS SERIES

. Bis

- The Structure and Stability of Competitive Dynamical Systems," by David Cass and Karl Shell.
- wopulistic Competition and the Capital Market," by J. E. Stiglitz.
 - The Corporation Tax." by J. E. Stightz.
- Messuring Returns to Scale in the Aggregate and the Scale Effect of Public Goods," by David A. Starrett
- Monopoly, Quality, and Regulation," by Michael Spence.
- A Note on the Budget Constraint in a Model of Borrowing," by Duncan K. Foley and Martin F. Hellwig.
- contitives, Risk, and Information. Notes Towards a Theory of Hierarchy," by Joseph E. Stightz. * = 2 = 3 = 5 = 5 = 5
- **taymetronic Expansions** of the Dutributions of Estimates in Simultaneous Equations for Alternative Parameter quencies, "by T. W. Anderson.
- ar of Lanear Functional Relationships. Approximate Distributions and Connections with Simultaneous in Econometrics," by T.W. Anderson. 3
 - Monopoly and the Rate of Extraction of Exhaustible Resources," by Joseph E. Suglitiz. <u>\$</u> 8
- Equilibrium as Competitive Insurance Markets: An Essay on the Economics of Imperfect Information." by dischael Rothschäfd and Joseph Stightz.
- Sirong Consistency of Least Squares Estimates in Normal Linear Regression," by T. W. Anderson and John B. Taylor Incentive Schemes under Differential Information Structures. An Application to Trade Policy." by Pariha Daugupta and Joseph Strights. ΞË
 - The incidence and Efficiency Effects of Taxes on income from Capital," by John B. Shoven. 173.
- Distribution of a Maximum Likelihood Estimate of a Slope Coefficient: The LIML Estimate for Known Covanance Vatrix," by T. W. Anderson and Takamitsu Sava.
 - "A Comment on the Test of Overidentifying Restrictions," by Joseph B. Kadane and T. W. Anderson. 175.
- 176.
- An Aymptotic Expansion of the Distribution of the Maximum Likelihood Estimate of the Slope Coefficient in a Linear Functional Relationship." by T. W. Anderson.
 - Some Experimental Results on the Statistical Properties of Least Squares Estimates in Control Problems, "hy F W. Anderson and John B. Taylor. 17.
 - "A Note on "Fuffilled Expectations" Equilibria." by David M. Kreps.
- Uncertainty and the Rate of Extraction under Alternative Institutional Arrangements." by Partha Dasgupta and losesh E. Stabita. R R
 - Budget Displacement Effects of Inflationary Finance," by Jerry Green and E. Sheshinski.
 - Towards a Marxust Theory of Money," by Duncan K. Foley.

**

- "The Existence of Futures Markets, Noisy Rational Expectations and Informational Externalities," by Sanford Grossman On the Efficiency of Competitive Stock Markets where Traders have Diverse Information." by Sanford Grossman
- A Bayessan Approach to the Production of Information and Learning by Dwing." hy Santord J. Grossinan, Richard E. Ribbarom and Leonard J. Mirman. "A Bidding Model of Perfect Competition," by Robert Wilson
 - Deequilibrium Allocations and Recontracting," by Jean-Michel Grandment, Guy Laroque and Yves Younes **38** 12 18
 - "Agreeing to Disagree," by Robert J. Aumann.
- The Maximum Lifelihood and the Numlinear Three Stage Least Squares Estimator in the General Numlinear Sumultaneous Equation Model." by Takesh Amemya

 - The Modified Second Round Estimator in the General Qualitative Response Model," by Takedu Amemica 822
 - Some Theorems in the Linear Probability Model," by Takeshi Amenuya.
- The Bilinear Complementarity Problem and Competitive Equilibria of Linear Economic Models," by Robert Wilson
 - Noncooperative Equilibrium Concepts for Obsorpedy Theory," by L. A. Gerard-Varet

 - Inflation and Costs of Price Adjustment." by Eytan Sheshimtki and Yoram Weiss.
 - Power and Laxes in a Multi-Commodity Economy," by R. J. Aumann and M. Kurz
- Disjortion of Preferences, Income Distribution and the Case for a funear fraction Eax." by Mordeon Kurz
 - "Demand for Fixed Factors, Inflation and Adjustment Costs." by Extan Shedinski, and Yotum Weise Search Strategies for Nonrenewahle Resource Deposits," hy Richard J. Gilhert
- Begaues and Rapolfs. A Model of Monopolistically Competitive Price Dispersions," by Steve Salop and Joweph Stight

 - The Design of Tax Structure Direct Versus Indirect Extation by A B Atkinson and J E Stiglitz Warket Altimations of Limition Choice in a Model with Free Mobility," by David Starrett
 - "Efficiency in the Optimum Supply of Public Goods," by Lowrence J. Luo, Extan Shedninski and Docpli E. Stightz

 - Rick Sharing, Sharecropping and Uncertain Labor Markets," hy David M. G. Newherry
 - On Non-Walrasan Equilibria." by Frank Habn
- A Note on Elasticity of Substitution Functions," by Lawrence J. Law
- "Duantity Constraints as Substitutes for Inoperative Markets The Case of the Credit Markets" by Morde an Kurz
 - Incremental Consumer's Surplus and Hedomic Price Adjustment," by Robert D. Willig

2

"Optimal Depletion of an Uncertain Stock," by Richard Gilbert.

- Some Minimum Chi-Square Estimators and Comparisons of Normal and LogisticaModels in Qualitative Response Analysis, "by Kunno Morimune 202
- "A Characterization of the Optimality of Equilibrium in Incomplete Markets," by Sanford J. Grossman
- "Inflation and Taxes in a Growing Economy with Debi and Equity Finance," by M. Feldistern. J. Green and E. Sheahinaki
 - "The Specification and Estimation of a Multivariate Logit Model," by Takeshi Amemiya.
 - Prices and Queues as Screening Devices in Competitive Markets," by Joseph E. Stightz.
- Conditions for Strong Consistency of Least Squares Estimates in Linear Models," by T. W. Anderwin and John B. Taylor
 - "Utilitarianism and Horizontal Equity The Case for Random Taxation," by Joseph E. Stightz.
 - Simple Formulae for Optimal Income Taxation and the Measurement of Inequality." by Joseph F. Stightz
- Temporal Resolution of Uncertainty and Dynamic Choice Behavior," by David M. Kreps and Evan L. Porteus
- The Estimation of Nonlinear Labor Supply Functions with Taxes from a Truncated Sample," by Michael Hurd
 - "The Welfare Implications of the Unemployment Rate," by Michael Hurd.
- "Keynesian Economics and General Equilibrium Theory: Reflections on Some Current Debates." by Frank Mahn
 - "The Core of an Exchange Economy with Differential Information," by Robert Wilson.
 - "A Competitive Model of Exchange." by Robert Wilson.
- Intermediate Preferences and the Majorsty Rule." by Jean-Michel Grandmunt
- "The Fixed Price Equilibria" Some Results in Local Comparative Statics," by Guy Lanoque
- On Stockholder Unanimity in Making Production and Financial Decisions," by Saniord J. Guossman. and Joseph E. Stiglitz
- "Selection of Regressors," by Takeshi Amemiya.
- "A Note on A Random Coefficients Model." by Takeshi Amemiya
 - "A Note on a Heteroscedastic Model," by Takeshi Amemiya
- "Welfare Measurement for Local Public Finance," by David Starrett
- "Unemployment Equilibrium with Rational Expectations," by W. P. Heller and R. M. Starr
- "A Theory of Competitive Equilibrium in Stock Market Economies," by Sanford J. Grossman and Oliver D. Hart.
- "An Application of Stem's Methods to the Problem of Single Period Control of Repression Models," by ALAD Zaman
- Second Best Welfare Economics in the Mixed Economy ." by David Starrett
- "The Logic of the Fix-Price Method." by Jean-Michel Grandmont
- Tables of the Distribution of the Maximum Likethbood Estimate of the Slope Coefficient and Approximations." by F. W. Anderson and Takamitsu Sawa
 - "Further Results on the Informational Efficiency of Competitive Stock Markets," by Sanford Grossman

 - "The Estimation of a Simultaneous-Equation Tobit Model." by Takeshi Amerinya
- The Estimation of a Simultaneous-Equation Generalized Probit Model." by Takeshi Amemiya
- The Consistency of the Maximum Likelihood Estimator in a Disequilibrium Model," by T. America and G. Son
- "Numerical Realisation of the Exact and Approximate Distribution Earctions of the Two Stage Least Squares Estimate" by E. W. Anderson and Edvantisu Sawa 22222
 - 'Risk Measurement of Public Projects," by Robert Wilson

 - On the Capitalization Hypothesis in Cloved Communities," by David Starrett
- A vite on the Uniqueness of the Representation of Commodity Augmenting Technical Change." In Lawrence J. Law
 - The Property Rights Doctrine and Demand Revelation under Incomplete Information," by Kenneth J. Arrow
- Optimal Capital Gains Lavation Under Limited Information," by Jerry R. Green and Eytan Shedunski.
- Straightforward Individual Incentive Compatibility in Large Economies," by Peter J. Hammond
 - On the Rate of Convergence of the Core 7 by Robert J. Aumann

 - "Unsatisfactory Equilibria," by Frank Hahn
- "Existence Conditions for Aggregate Demand Functions. The Case of a Single Index," by Lawrence J. Law
- Textence Conditions for Aggregate Demand Functions The Case of Multiple Indexes," by Lawrence 3, Lau
 - "A Note on Exact Index Numbers," by Lawrence J. Lau
- "Uncar Repression Using Both Temporally Aggregated and Temporally Disaggregated Data," by Obeng Hyao-
- "A Complete Class Theorem for the Control Problem and Eurifier Results on Admissibility and Inadmissibility," 154 Aud Zaman "The Exitence of Economic Equilibria. Continuity and Mixed Strategies," by Parilia Dasgupta and Eric Maskin 展主任主要主要主要主要工作的基的工作

"The Existence of Efficient and Incentive Compatible Equilibria with Public Gowds," 55. Theology Groves and

"A Representation Theorem for Preference for Elexibility 27 by David W. Kreps

"Altruism as an Outcome of Social Interaction," by Mordecai Kurz-

Measure-Based Values of Market Games," hy Sergin Hart

On the Impossibility of Informationally Efficient Markets," by Sanford J. Grossman and Joseph E. Stightz

"Efficient Collective Choice with Compensation," by Theodore Groves

John O. Ledyard

¥ 3.

i

REPORTS IN THIS SERIES

- "Vallees for Guartes Without Sadepus ments: Sume Dirikultees With Current Cuncepts." by Alond Roch "Mertangtes and the Valuation of Redundant Arosts." by J. Michael Harrison and David M. Kreps
- e Modelling and Money Income Councility Detection." by Cheng Huuci Autom
- **een Errec en a Dynamik Simultaneous Equations Model without Stationary Undurhance.**" hy Cherk Hydor
 - ent of Deadweight Loss Revisited," by W. F. Diewert
- The Electricity of Derived Net Supply and a Generalized Lr. Charelier Principle." hy W. E. Diewert
- Theome Distributions and Distortion of Preferences the ₹ (commodity Case," by Mordevia kurz "The a²⁴ Codes Massis Squared Errors of the Maximum Listetikhood and the Mormium Listeti Chis Square Aximutur," by Talechi Americu.

 - aral Yon Neumana-Murgenstern and Indused Preferences," by Duvid M. Kreps and Evan L. Porteus
 - TabeOver Buds and the Theory of the Corporation." by Stanford Grossman and Oliver D. Harl
 - mercul Values of Some Key Parameters in Econometric Models," by T. W. Anderwin, Kimio Mortmune, and Lakamitou Sawa Ē
 - Any approve. Expansions of the Distributions of Estimation is a Linear Euricitonal Relationship when the Sample Size is Large." his Wastic Kantonico "Two Representations of Information Structures and their Comparisonss." by Jerry Green and Nancy Stokey

 - Twis. Goods and Poner," by R. J. Aumana, M. Kurz and A. Neyman.
- "As Assemble: Approach to the Efficiency of NewCoopergine Fquidhform in Economics with a Continuum of Traders" by A MacCooff Tables of the Tsue Obstantion Distance of the Annuel Information Maximum Lifetihioad Estimator when the Coostunice Maxim in Revense" by T. A Matheward and Educative Size Example Distance Maximum Lifetihioad Estimator when the Coostunice
- e Modeling of Canadian Money and Income Data," by Cheng Huuri
- Te Can't Deagnee Forever," by Judia D. Geznadioptics and Heraklis Polemarchakis
 - of Excens Demand Functions." by Herklis M. Polemarchakis. 'On the Bayessan Selection of Nucl Equilibrium." by Akira Formola
- ibream Econometrics in Simultaneous Equations Systems," by C. Gourierous, J. J. Lattont and A. Montort
- Durkiny Approx. has to Marton. cauma: Theory." by W. F. Diewert
- *A Time Series Analysis of the frequent of Canadian Wage and Price Cuntrols." by Cheng Huan and Olivia tau tais of *A Stranger Theory of laritations." by Mondocan Kurz.
 - zation of Vector Meanure Games in pNA." by Yair Lauman Č.
- "On the Machined of Francisco and the Phorenica of Local Public Condo," by David A Statrett "An Optimization Problem Arange in Economics: Approximate Solutions, Linearity, and a Lux of Large Vumberi," by Sergia Mar "Subdom, Emon Strandsman, Nanok Estimation of Cheffmann, of Coefficientis in a Smultaneous Equation System," by Yusum-ri "Subdom, Emon Strandsman, Nanok Estimations of Cheffmann, of Coefficientis in a Smultaneous Equation System," by Yusum-ri
 - Optimal & Voluntary Income Distribution," by K. J. Arrow.
- totic Valuers of Maxed Games," by Abraham Neyman
- Time Series Modelling and Caseal Ordering of Canadian Minicy, Income and Interest Rate," his Cheng History
- Als Analysis of Poster in Evchange Evonomen." by Martin J. Oxburne. Estimations of the Reciprocal of a Normal Mean." by Acid Zaman
- <mark>theprives the Maximum Latchhood Prim</mark>ate m Linear Punchinal Relativiships for Alternative Parameter Sequence. " by Kimis Bernamar and Norto Russionio.
- Calculations of Broartste Normal Integrals by the Use of Incomplete Negative-Order Moments." by Kei Laken, hi and Akimichi Lakenura
- De Burteneme of a Semple with Breary Type Orections in Levin Guideling (Deuration). By Korneth J. Arrow, Four Pourh Inda Million Sobel.
- The few Stage Least Abualure Deviations Estimation." by Ealerthi Amemira "Three Franss on Capital Markets." by David M. Kreps

 - Safrance Norwon Programs." by Michael J. P. Mapili
- Electoral Outcomes and Social Log-Litelibord Maxima," by Peter Coughlin and Shmuel Nitzan
 - Notes on Social Choice and Volume," by Peter Coughlin
- mag Expectations and Hart's Conditions for Equilibrium in a Securities Medel." by Peter 1. Hammond
- Directional and Local Flectorate Competitions with Probabilistic Voting." by Peter Coughlin and Shmuel Nitzan
- Asymptotic Expansions of the Distributions of the Test Statistics (Decidentifying Restrictions in a System of Simultancous Generation," by Kensionae, Moremane, and Fuskida 医牙浆液素医素浆素原因的 化乙酸 尤指花花用的复数形式的 医角色的的的 医乳 美国家的复数 医皮 医 医里耳氏管 带带带带带带
 - tecomplete Murkets and the Observability of Risk Preference Properties." In H. H. Polemarchakos and L. Seiden

13.7

- Asymptotic Exploration of the Distributions of AcTass Extimutions when the Disturbance are Small." In Number Aumiumo Kumu Morimume, and Yomhindin Fouriscie
 - Arbitrage and Equiptorium in Economics with Infinitely Mains Commodition." by David M. Kreps
 - loyment Equilibrium in an Economy with Linked Prices." by Mordecar Kurz ļ
- Parsin (Spinnaul Namb Equilibria are Competitive in a Repeated Economy." by Mordecal Kurz and Sericia Hari
 - Mentification." by Cheng Huuu
- "An Introduction to Two-Person Zaro Sum Repeated Games with Incomplete Information." by Sylvain Sector
- "Estimation of Dynamic Models with Error Components," by T. W. Anderson and Cheng Histor
- "On Robust Estimation in Certainty Equivalence Control." by Anders H. Westlund and Hans Stenlund
 - "On Industry Equilibrium Under Uncertainty," by J. Dieze and E. Sheshinski.
- "Cost Benefit Analysis and Project Evaluation from the Viewpoint of Productive Efficiency," by W. F. Diewert
- "On the Cham-Store Paradox and Predation Reputation for Toughnes," by D. W. Kreps and Robert Wilson "On the Number of Commodities Required to Represent a Market Games," Sergiu Hart
 - "Evaluation of the Distribution Function of the Limited Information Maximum Likelihood Atimator " hy T. W. Anderson, Naoto Kunitomo, and Takanitsu Sawa

"Asymptotic Bas of the Least Squares Estimator for Wultivariate Autoregressive Models" by 1. Yamamotic and N. Kurutomo

Rational Cooperation in the Emitely Repeated Prisoner's Discours." by D. Kreps. P. Wilgrom, J. Roberts, and R. Wilson

The Asymptotic Normality of Two Staye Least Absolute Deviations Estimators " by F. Powell

"Partially Generalized Last Squares and Two Stage Erast Squares and Two Stage Least Squares Estimations," by T. Amemica

Non-Zero-Sun Two-Person Repeated (sames with Incomplete Information," by S. Hart

"Semi-Values of Political Economic Games," by A. Neyman

"Lipwhitz Properties of Solution in Mathematical Programming," by B. Cornet and G. Laroque

"Team Theory and Deceminalized Resource Allocation An Example " by K. J. Arrow

Multiple Hypothesis Testing," by N. E. Savin

333322228

Texting for Unit Roots [7] by G. B. A. and Fvans and N. F. Savin

Asymptotic Efficiency and Higher Order Efficiency of the Limited Information Maximum Likelihood Extimator in Large Econometric Models," by N. Kumitomo

"Capital Accumulation and the Characteristics of Private Interprintizational Transfers." by M. Kurz

"Optimum Pricing Policy Under Stochastic Inflation." by E. Sheshinski and Y. Wens-

"The Computative Dynamics of Efficient Programs of Capital Accumulation and Resource Depletion." hy W. F. Dewert and T. R. Lewis.

"The Individual Freedom Allowed by the Value Restriction Condition." hy Herve Raynaud

"Incomplete Resource Allocation Mechanisms." hy P. J. Hammond

"A Theory of Auctions and Competitive Bidding." by Paul Milgrom and Robert Weber

'Facing an Uncertain Future," by W. M. Gorman

"Relatively Recursive Rational Choice," by Alain Lewis

Recurrave Rational Choice," by Alam Lewis

- 2
- "A Comparison of the Logit Model and Normal Discriminant Analysis when the Independent Vanables are Roury ^{TI} by Takeah Amemya and James L. Prowell

- "Efficiency of Resource Allocation by Uninformed Demand," by Theodore Groves and Sergui Hart 22
- "A Comparison of the Box Cox Maximum Likelihood Estimator and the Nonlinear Two Stage Least Squares Estimator," by Takehi Amenuya and James L. Powell.
 - Comparison of the Dematics of the TSLS and LIMLK Fatimators for Simultaneous Equations." by J. W. Anderson, Notice Kunitoma, and Takanutsu Sawa. 323
 - "Admissibility of the Bayes Procedure Corresponding to the Unitorn Prior Distribution for the Control Problem in Four Dimensions but Not in Five," by Charles Stein and Asad Zaman 324

 - "Some Recent Developments on the Distributions of Single-Equation Estimators," by T. W. Anderson
 - On Inflation." by Frank Hahn. 326.327.
- Two Papers on Majority Rule "Continuity Properties of Majority Rule with Intermediate Preferences." by Peter and KumPan Lin. and. "Thertonal Outcomes with Probabilistic Volting and Nanh Social Welfare Maxum." by Peter Gouphin and Shmedi Nitzer.
 - "On the Endogenous Formation of Coalitions," by Sergiu Hart and Mordecai Kurd
- "Controlushisty, Pecuniary Externalities and Optimal Taxation." by David Starrett.
 - Nonlinear Regression Models," by Takeshi Amemiya.
- Paradoxical Results from Inada's Conditions for Majonty Rule." by Herve Raynaud.
- "On Weffare Economics with Incomplete Information and the Social Value of Public Information." by Peter J Hammond

 - "Equilibrium Policy Proposals with Abstentions." by Peter J. Coughtin. "Infinite Excessive and Invariant Measures." by Michael I. Takiar
- The Life-Cycle Hypotheus and the Effects of Social Scourity and Private Pensions on Family Savings." by Mordeua Kurz

 - "Optimal Retirement Age," by Mordecai Kurz.
 - Bayeman Incentive Compatible Behefs," by Claude d'Asprement and Louis-Andre Gerard-Varet.
- The Social Costs of Monopuly and Regulation A Game Theoretic Analysis," by William P. Rugeraun Qualitative Response Models A Survey." by Takeshi Amemiya
 - Sequential Equilibria," by David M. Kreps and Robert Wilson

 - Enhancing of Semigroups," by Michael 1. Taksar
- Formulation and Estimation of Dynamic Models Using Panel Data," by T. W. Anderson and Cheng Haav
 - Ex-Post Optimality as a Dynamically Consistent Objective for Collective Choice Under Undertainty " by Peter Hammend "Three Lectures in Monetary Theory," by Frank H. Hahn **雅松说明的现在分词的形式的现在分词**

 - Socially Optimal Investment Rules in the Presence of Incomplete Markets and Other Second Berl Distortions." by Tank Müne and Divid A. Starrett.
 - Approximate Punikation of Mixed Strategies." by Robert Aumann, Yitzhak Karznelovn, Risy Radner, Robert W Coenthal, and Benjaman Wens 346
 - "Conditions for Transitivity of Majority Rule with Algorithmic Interpretations," by Herve Raynaud
 - How Restrictive Actually are the Value Restriction Conditions," hy Herve Raynaud
- Cournot Duopoly in the Style of Fulfilled Expectations Equilibrium." by William Norshek and Hugh Somernichein
- "Law of Large Numbers for Random Sets and Alboration Processes." by Zvi Arititein and Sergiu Hari
- "Risk Perception in Psychology and Economics," by Kenneth J. Arrow
- Shrunken Predictors for Autoregressive Models," by Taku Yamanisito
- Predation, Reputation, and Entry Deterrence." by Paul Milgrom and John Roberts
- Social and Private Production Objectives in the Sequence Economy." by David Starrett

"Least Absolute Deviations Estimation for Censored and Truncated Regression Models." by James Powell.

Reports in this Series

- 376. "Necessary and Sufficient Conditions for Single-Peakedness Along a Linearly Ordered Set of Policy Alternatives" by P.J. Coughlin and M.J. Hinich.
- 377. "The Role of Reputation in a Regeated Agency Problem Involving Information Transmission" by W. P. Rogerson.
- 378. "Unemployment Equilibrium with Stochastic Rationing of Supplies" by Ho-mou Wu.
- 379. "Optimal Price and Income Regulation Under Uncertainty in the Model with One Producer" by M. I. Taksar.
- 380. "On the NTU Value" by Robert J. Aumann.
- 381. "Best Invariant Estimation of a Direction Parameter with Application to Linear Functional Relationships and Factor Analysis" by T. W. Anderson, C. Stein and A. Zaman.
- 382. "Informational Equilibrium" by Robert Kast.
- 383. "Cooperative Oligopoly Equilibrium" by Mordecai Kurz.
- 384. "Reputation and Product Quality" by William P. Rogerson.
- 385. "Auditing: Perspectives from Multiperson Decision Theory" By Robert Wilson.
- 386. "Capacity Pricing" by Oren, Smith and Wilson.
- 387. "Consequentialism and Rationality in Dynamic Choice Under Uncertainty" by P.J. Hammond.
- 388. "The Structure of Wage Contracts in Repeated Agency Models" by W. P. Rogerson.
- 389. "1982 Abraham Wald Memorial Lectures, Estimating Linear Statistical Relationships by T.W. Anderson.
- 390. "Aggregates, Activities and Overheads" by W.M. Gorman.
- 391. "Double Auctions" by Robert Wilson.
- 392. "Efficiency and Fairness in the Design of Bilateral Contracts" by S. Honkapohja.
- 393. "Diagonality of Cost Allocation Prices" by L.J. Mirman and A. Neyman
- 394. "General Asset Markets, Private Capital Formation, and the Existence of Temporary Walrasian Equilibrium" by P.J. Hammond
- 395. "Asymptotic Normality of the Censored and Truncated Least Absolute Deviations Estimators" by J.L. Powell
- 396. "Dominance-Solvability and Cournot Stability" by Herve Moulin
- 397. "Managerial Incentives, Investment and Aggregate Implications" by B. Holmstrom and L. Weiss

4

Reports in this Series

- 398 "Generalizations of the Censored and Truncated Least Absolute Deviations Estimators" by J.L. Powell.
- 399. "Behavior Under Uncertainty and its Implications for Policy" by K.J. Arrow.
- 400. "Third-Order Efficiency of the Extended Maximum Likelihood Estimators in a Simultaneous Equation System" by K. Takeuchi and K. Morimune.
- 401. "Short-Run Analysis of Fiscal Policy in a Simple Perfect Foresight Model" by K. Judd.
- 402. "Estimation of Failure Rate From A Complete Record of Failures and a Partial Record of Non-Failures" by K. Suzuki.
- 403. "Applications of Semi-Regenerative Theory to Computations of Stationary Distributions of Markov Chains" by W.K. Grassmann and M.I. Taksar.
- 404. "On the Optimality of Individual Behavior in First Come Last Served Queues With Preemption and Balking" by Refael Hassin.
- 405. "Entry with Exit: An Extensive Form Treatment of Predation with Financial Constraints" by J.P. Benoit.
- 406. "Search Among Queues" by A. Glazer and R. Hassin
- 407. "The Space of Polynomials in Measures is Internal" by J. Reichers and Y. Tauman.
- 408. "Planning Under Incomplete Information and the Ratchet Effect" by X. Freixas, R. Guesnerie and J. Tirole.
- 409. "A Theory of Dynamic Oligopoly, I: Overview and Quantity Competition with Large Fixed Costs" by Eric Maskin and Jean Tirole.
- 410. "Approximate Measures of the Social Welfare Benefits of Labor Projects" by Peter Hammond
- 411. "Transversality Conditions for Some Infinite Horizon Discrete Time Optimization Problems" by Ivar Ekeland and Jose A. Scheinkman.
- 412. "Asymptotic Efficiency of the Partial Likelihood Estimator in the Proportional Hazard Model" by Aaron K. Hahn.
- 413. "A Monte Carlo Comparison of Estimators for Censored Regression Models" by darry J. Paarsch.
- 414. "Instrumental-Variable Estimation of an Error-Components Model" by Takeshi Amemiya and Thomas E. MaCurdy.
- 415. "An Axiomatization of the Non-Transferable Utility Value" by Robert J. Aumann.
- 416. "Diffusion Approximation in Arrow's Model of Exhaustable Resources" by Darrell Duffie and Michael Taksar.

<u>Reports in this Series</u>

417. "The Shapley Value in the Non Differentiable Case" by Jean François Mertens.

The seal

1

418. "The Minimax Theorem for U.S.C.-L.S.C. Payoff Functions" by Jean François Mertens.

