

AD-A158 535

COMPOUND POISSON APPROXIMATIONS FOR SUMS OF RANDOM
VARIABLES(U) GEORGIA INST OF TECH ATLANTA SCHOOL OF
INDUSTRIAL AND SYSTEMS. R F SERFOZO MAY 85

1/1


UNCLASSIFIED

AFOSR-TR-85-0623 AFOSR-84-0367

F/G 12/1

NL

					END								
					FILMED								
					DTIC								




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AFOSR-TR- 85-0623

AD-A158 555

(Georgia Institute of Technology)

U.S. Department of Defense



INDUSTRIAL and SYSTEMS ENGINEERING REPORTS SERIES

DTIC FILE COPY

DTIC
ELECTE
AUG 29 1985
S D

G

FOR INFORMATION WRITE:

REPORT SERIES LIBRARIAN
SCHOOL OF INDUSTRIAL & SYSTEMS
ENGINEERING
GEORGIA INSTITUTE OF TECHNOLOGY
ATLANTA, GEORGIA 30332

Approved for public release;
distribution unlimited.

85 8 21 010

COMPOUND POISSON APPROXIMATIONS FOR
SUMS OF RANDOM VARIABLES

by

Richard F. Serfozo

AIR FORCE SYSTEMS SCIENTIFIC ESTABLISHMENT (AFSC)
RESEARCH
TECHNICAL INFORMATION DIVISION
Chief, Technical Information Division

DTIC
ELECTE
AUG 29 1985
S D
G

Submitted for publication to
Annals of Probability
May, 1985

DISTRIBUTION STATEMENT
Approved for
Distribution

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

AD-A158 555

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY ---			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution Unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 85 - 0623		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION AFOSR		
6a. NAME OF PERFORMING ORGANIZATION Georgia Institute of Tech.		6b. OFFICE SYMBOL (If applicable)		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, D.C. 20332-6448	
6c. ADDRESS (City, State and ZIP Code) Atlanta Georgia, 30332		8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR			
8b. OFFICE SYMBOL (If applicable) NM		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0367			
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, D.C. 20332-6448		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.
11. TITLE (Include Security Classification) Compound Poisson Approximations for Sums of Random Variables		61102F	2304	A5	
12. PERSONAL AUTHOR(S) Richard F. Serfozo					
13a. TYPE OF REPORT Reprints		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) May 1985	
15. PAGE COUNT 14					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	Poisson Approximations, Markovian occurrences		
XXXXXXXXXXXX					
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
We show that a sum of dependent random variables is approximately compound Poisson when the variables are rarely nonzero and, given they are nonzero, their conditional distributions are nearly identical. We give several upper bounds on the total-variation distance between the distribution of such a sum and a compound Poisson distribution. Included is an example for Markovian occurrences of a rare event. Our bounds are consistent with those that are known for Poisson approximations for sums of uniformly small random variables.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☒ DTIC USERS ☐			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Brian W. Woodruff, Maj, USAF			22b. TELEPHONE NUMBER (Include Area Code) (202)767-5027		22c. OFFICE SYMBOL NM

COMPOUND POISSON APPROXIMATIONS FOR
SUMS OF RANDOM VARIABLES

(COMPOUND POISSON APPROXIMATIONS)

By Richard F. Serfozo
Georgia Institute of Technology

SUMMARY

We show that a sum of dependent random variables is approximately compound Poisson when the variables are rarely nonzero and, given they are nonzero, their conditional distributions are nearly identical. We give several upper bounds on the total-variation distance between the distribution of such a sum and a compound Poisson distribution. Included is an example for Markovian occurrences of a rare event. Our bounds are consistent with those that are known for Poisson approximations for sums of uniformly small random variables.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A/1	



Footnotes for Page 1

AMS 1980 subject classifications. Primary 60E15, 60F99; secondary 60J10.

Key words and phrases. Compound Poisson distribution, total variation distance, sums of dependent variables, rare Markovian events.

This research was supported in part of AFOSR Grant 84-0367.

1. Introduction

There have been a number of studies on Poisson approximations for sums of uniformly small random variables. Of paramount interest is the total-variation distance between a sum of random variables and a Poisson variable. The total-variation distance between two probability measures (distributions) F and G on some measurable space is defined by

$$(1.1) \quad d(F,G) = \sup_B |F(B) - G(B)|,$$

where the supremum is over all measurable sets ($2d(F,G)$ is the total variation of the signed measure $F-G$). The total-variation distance between random elements X and Y with the respective distributions F and G is $d(X,Y) = d(F,G)$.

Building on the works of Hodges and Le Cam (1960), Le Cam (1960), Franken (1964) and Freedman (1974), Serfling (1975) proved this result: If $X_1, \dots, X_n$ are non-negative integer-valued random variables adapted to the increasing σ -fields $\{F_i\}_{i=0}^n$, then

$$(1.2) \quad d\left(\sum_{i=1}^n X_i, N\right) \leq \sum_{i=1}^n [E^2(p_i) + E|p_i - Ep_i| + P(X_i > 2)],$$

where $p_i = P(X_i = 1 | F_{i-1})$ and N is Poisson with mean $\sum_{i=1}^n Ep_i$. Comparable bounds for other Poisson approximations appear in Barbour and Eagleson (1983), Brown (1983), Chen (1975), Kabanov et al. (1983), Kerstan (1964), Valkeila (1982) and their references. Such bounds are useful for proving limit theorems for random variables and point processes as well.

In this paper, we present analogues of (1.2) for compound Poisson approximations for sums. We consider sums of random elements that take

values in a measurable group S : the group operation, addition, is measurable. If X is a random element of S with the compound Poisson distribution $H(B) = \sum_{n=0}^{\infty} F^{n*}(B) \alpha^n e^{-\alpha} / n!$, then we say X is $CP(\alpha, F)$. If X has the distribution $EH(\cdot)$, where α or F are random, then we say X is mixed $CP(\alpha, F)$.

Here is our main result. Let $X_1, \dots, X_n$ be random elements of S adapted to the increasing σ -fields $\{F_i\}_{i=0}^n$, and define

$$p_i = P(X_i \neq 0 | F_{i-1}), \quad F_i(B) = P(X_i \in B | F_{i-1}, X_i \neq 0).$$

Let F be a distribution on S with $F(\{0\}) = 0$, and define, by (1.1), the random distance $d_i = d(F_i, F)$ (F_i is random but F is not).

Theorem 1. If Z is mixed $CP(\sum_{i=1}^n p_i, F)$, then

$$(1.3) \quad d(\sum_{i=1}^n X_i, Z) \leq E \left[\sum_{i=1}^n (d_i + p_i^2) \right].$$

If Z is $CP(\sum_{i=1}^n \alpha_i, F)$ where $\alpha_i = E p_i$, then

$$(1.4) \quad d(\sum_{i=1}^n X_i, Z) \leq E \left[\sum_{i=1}^n (d_i + |p_i - \alpha_i| + \alpha_i^2) \right].$$

If Z is $CP(\alpha, F)$, then

$$(1.5) \quad d(\sum_{i=1}^n X_i, Z) \leq E \left[\sum_{i=1}^n (d_i + p_i^2) + \left| \sum_{i=1}^n p_i - \alpha \right| \right].$$

This result says, roughly, that $\sum_{i=1}^n X_i$ is approximately compound Poisson when the X_i 's are rarely nonzero (the p_i 's are small), and given that the X_i 's are nonzero, their conditional distributions $F_1, \dots, F_n$

are nearly identical. Note that (1.5) with $\alpha = \sum_{i=1}^n \alpha_i$ is different from (1.4); in some cases the bound in (1.4) is smaller than that in (1.5) but in other cases the reverse is true.

For the degenerate distribution F on $\mathbb{R}$ with unit mass at 1, Theorem 1 yields bounds for Poisson or mixed Poisson approximations for sums. In this case, (1.4) is the same as (1.2), and (1.5) is consistent with the inequalities of Brown (1983) and Kabanov et al. (1983), which were established by martingale techniques.

Brown (1983) also obtains compound Poisson approximations for certain discrete variables via Poisson approximations. This approach, however, does not apply to the general case. We prove our results by rather direct arguments based on judicious conditioning and the use of (1.1) as a random distance for random distributions. Our approach also brings to light the key role of the F_i 's.

From its proof, one can easily see that Theorem 1 is also true when the number of variables n in the sum is a stopping time of $\{F_i\}$. For instance, Theorem 1 applies to sums of the form $\sum_{i=1}^{N(t)} X_i$, where $N(t) = \sum_i I(\tau_i \leq t)$ and $\tau_1 < \tau_2 < \dots$ are stopping times of the increasing σ -fields $\{F(t)\}$ and $F_i = F(\tau_i)$, respectively. Theorem 1 also holds when F and α are random; the Z in (1.4) and (1.5) would then be mixed compound Poisson.

The rest of this paper is organized as follows. Section 2 gives some basics on the total-variation distance, Section 3 consists of the proof of Theorem 1, and Section 4 gives an example for Markovian occurrences of an event.

2. Basic Inequalities for Distances

Let X and Y be random elements of some measurable space. A well-known coupling inequality is

$$(2.1) \quad d(X, Y) \leq P(X \neq Y).$$

The X, Y in the probability are the random elements -- with an arbitrary dependency -- defined on a common probability space. Inequality (2.1) follows because $P(X \in B) \leq P(X \neq Y) + P(Y \in B)$.

It is natural for us to analyze $d(X, Y)$ in terms of conditional probabilities. Accordingly, we sometimes refer to X as having a distribution $EF(\cdot)$ where F is a random distribution. Typically, $F(B) = P(X \in B | F)$, or F could be defined as a measurable function of random elements.

Lemma 2.1. Suppose X and Y have the respective distributions $EF(\cdot)$ and $EG(\cdot)$, where F and G are random distributions. Then

$$(2.2) \quad d(X, Y) \leq E[d(F, G)].$$

In case $F(B) = P(X \in B | F)$ and $G(B) = P(Y \in B | G)$, for some σ -fields F and G , then

$$(2.3) \quad d(X, Y) \leq E[d(F, G)] \leq E[P(X \neq Y | F, G)].$$

Proof. Expression (2.2) follows since

$$d(X, Y) = \sup_B |EF(B) - EG(B)| \leq \sup_B E|F(B) - G(B)| = E[d(F, G)].$$

Expression (2.3) follows from (2.2) and a random version of (2.1).

Remark. Keep in mind that F, G in the expectation in (2.2) are the random distributions on a common probability space and their dependency is arbitrary. A similar comment applies to the X, Y, F, G in the probability in (2.3).

Distances involving functions of random elements, such as sums or maxima, can generally be represented as $D = d(h(X), h(Y))$, where $X = (X_1, \dots, X_n)$, $Y = (Y_1, \dots, Y_n)$ and h is a measurable function from the range space of X and Y to some other measurable space. Here are some bounds on this distance.

Lemma 2.2. (i) $D \leq d(X, Y)$. (ii) $D \leq \sum_{i=1}^n P(X_i \neq Y_i)$.

(iii) If $X_1, \dots, X_n$ are independent and $Y_1, \dots, Y_n$ are independent, then $D \leq \sum_{i=1}^n d(X_i, Y_i)$.

(iv) If $X_1, \dots, X_n$ are adapted to the increasing σ -fields $\{F_i\}_{i=0}^n$ and $Y_1, \dots, Y_n$ are adapted to the increasing σ -fields $\{G_i\}_{i=0}^n$, and $F_i(B) = P(X_i \in B | F_{i-1})$, $G_i(B) = P(Y_i \in B | G_{i-1})$, then

$$(2.4) \quad D \leq E \left[\sum_{i=1}^n d(F_i, G_i) \right] \leq E \left[\sum_{i=1}^n P(X_i \neq Y_i | F_{i-1}, G_{i-1}) \right].$$

Proof. Statement (i) is true since

$$\begin{aligned} D &= \sup_B |P(h(X) \in B) - P(h(Y) \in B)| \\ &= \sup_B |P(X \in h^{-1}(B)) - P(Y \in h^{-1}(B))| \leq d(X, Y). \end{aligned}$$

Statement (ii) is true since by (i) and (2.1) we have

$$D \leq P(X \neq Y) = P\left(\bigcup_{i=1}^n \{X_i \neq Y_i\}\right) \leq \sum_{i=1}^n P(X_i \neq Y_i).$$

Now consider (iii) when $n=2$. From (i), the triangle inequality for d , and the independence, we have

$$\begin{aligned} D &\leq d((X_1, X_2), (Y_1, Y_2)) \leq d((X_1, X_2), (Y_1, X_2)) + d((Y_1, X_2), (Y_1, Y_2)) \\ &\leq d(X_1, Y_1) + d(X_2, Y_2). \end{aligned}$$

Using this inequality and induction yields (iii) for general n .

Under the hypotheses of (iv), it follows by successive conditioning that $P(X \in B_1 \times \dots \times B_n) = E[F_1(B_1) \dots F_n(B_n)]$, and a similar statement holds for Y . Then using (i), (2.2) and (iii) we have

$$D \leq d(x, y) \leq E[d(F_1 \dots F_n, G_1 \dots G_n)] \leq E \sum_{i=1}^n d(F_i, G_i).$$

The second inequality in (2.4) follows from (2.3).

The next two results deal with compound Poisson distributions.

Lemma 2.3. If X is $CP(\alpha, F)$ and Y is $CP(\beta, G)$, with $F(\{0\}) = 0$ and $G(\{0\}) = 0$, then $d(X, Y) \leq |\alpha - \beta| + (\alpha \wedge \beta)d(F, G)$.

Proof. First consider the case in which $\alpha < \beta$. Clearly Y is equal in distribution to $Y_1 + Y_2$, where Y_1, Y_2 are independent $CP(\beta - \alpha, G)$ and $CP(\alpha, G)$, respectively. Note that the distributions of X and Y_2 can be written as $EF^{N^*}(\cdot)$ and $EG^{N^*}(\cdot)$, respectively, where N is a Poisson random variable with mean α . Then applying the triangle inequality, (2.2), (2.1) and Lemma 2.2 (iii) in the form $d(F^{N^*}, G^{N^*}) \leq nd(F, G)$, we have

$$\begin{aligned} d(X, Y) &\leq d(X, Y_2) + d(Y_2, Y_1 + Y_2) \leq Ed(F^{N^*}, G^{N^*}) + P(Y_1 \neq 0) \\ &\leq ENd(F, G) + 1 - e^{-(\beta - \alpha)} \leq \alpha d(F, G) + \beta - \alpha. \end{aligned}$$

This proves the assertion when $\alpha < \beta$, and a similar proof applies when $\alpha > \beta$.

Lemma 2.4. Suppose X is a random element of S and let

$$(2.5) \quad p = P(X \neq 0) \quad \text{and} \quad F(B) = P(X \in B | X \neq 0).$$

If Z is $CP(p, F)$, then $d(X, Z) \leq p^2$.

Proof. It suffices, by (2.1), to construct X, Z on a common probability space such that $P(X \neq Z) \leq p^2$. To this end, let N, U and $Y_1, \dots, Y_n$ be independent random elements on a common probability space such that N is

a Poisson random variable with mean p , each Y_i has the distribution F , and $P(U = 0) = (1 - p)e^p = 1 - P(U = 1)$. Define

$$X = Y_1(1 - I(U = 0, N = 0)) \quad \text{and} \quad Z = \sum_{i=1}^N Y_i.$$

An easy check shows that X satisfies (2.5), and Z is clearly $CP(p, F)$.

Furthermore,

$$\begin{aligned} P(X \neq Z) &= P(X \neq Z, N = 0) + P(X \neq Z, N > 2) \\ &= P(U = 1)P(N = 0) + P(N > 2) = p(1 - e^{-p}) < p^2. \end{aligned}$$

This completes the proof.

We end this section by comparing two random elements that have certain conditional distributions that are equal.

Lemma 2.5. Let X and Y be random elements. If there is a measurable set A such that $P(X \in B | X \in A) = P(Y \in B | Y \in A)$ for each measurable B , then

$$(2.6) \quad d(X, Y) \leq |P(X \in A) - P(Y \in A)|.$$

Proof. Let U , V and W be independent random elements on a probability space. Assume that U is uniform on $(0, 1)$ and that V and W take values in A and A^c , respectively, and their distributions are $P(V \in B) = P(X \in B | X \in A)$ and $P(W \in B) = P(X \in B | X \in A^c)$. Let p and q denote the respective probabilities in (2.6), and define $X = VI(U < p) + WI(U > p)$ and $Y = VI(U < q) + WI(U > q)$. Clearly X and Y satisfy the hypotheses and, moreover, $P(X \neq Y) = P(p \wedge q < U < p \vee q) = |p - q|$. Thus the assertion follows by applying (2.1).

3. Proof of Theorem 1

In addition to the notation of Theorem 1, we let $G_p(\cdot) = pF(\cdot) + (1-p)\delta_0(\cdot)$, where δ_0 is the Dirac measure with unit mass at 0, and

we let Y be a random element with distribution $E(G_{p_1} * \dots * G_{p_n}(\cdot))$ (recall that p_i is random).

To prove (1.3), consider the inequality

$$(3.1) \quad d\left(\sum_{i=1}^n X_i, Z\right) < d\left(\sum_{i=1}^n X_i, Y\right) + d(Y, Z).$$

By the use of successive conditioning, it is clear that

$$P\left(\sum_{i=1}^n X_i \in B\right) = E[F'_1 * \dots * F'_n(B)], \quad \text{where } F'_i(B) = P(X_i \in B | F_{i-1}).$$

Note that $F'_i(\cdot) = p_i F_i(\cdot) + (1-p_i)\delta_0(\cdot)$, and so $d(F'_i, G_{p_i}) = d(F_i, F) = d_i$.

Then applying (2.2) and Lemma 2.2 (iii), we have

$$(3.2) \quad d\left(\sum_{i=1}^n X_i, Y\right) < E[d(F'_1 * \dots * F'_n, G_{p_1} * \dots * G_{p_n})] < E\left(\sum_{i=1}^n d_i\right).$$

Similarly, using $P(Z \in B) = E[H_{p_1} * \dots * H_{p_n}(B)]$, where the distribution H_p is $CP(p, F)$, and applying Lemmas 2.1, 2.2 (iii) and 2.4, we have

$$(3.3) \quad \begin{aligned} d(Y, Z) &< E[d(G_{p_1} * \dots * G_{p_n}, H_{p_1} * \dots * H_{p_n})] \\ &< E\left[\sum_{i=1}^n d(G_{p_i}, H_{p_i})\right] < E\left(\sum_{i=1}^n p_i^2\right). \end{aligned}$$

Then combining (3.1) - (3.3) yields the assertion (1.3).

Now consider the assertion (1.4). Here Z is $CP(\sum_{i=1}^n \alpha_i, F)$. Let

$U_1, \dots, U_n$ be independent random elements with the respective distributions $G_{\alpha_1}, \dots, G_{\alpha_n}$. Then by applications of (3.2), Lemmas 2.1, 2.2 (iii) and 2.5 (with $A = S \setminus \{0\}$), we have

$$\begin{aligned}
d\left(\sum_{i=1}^n X_i, Z\right) &< d\left(\sum_{i=1}^n X_i, Y\right) + d\left(Y, \sum_{i=1}^n U_i\right) + d\left(\sum_{i=1}^n U_i, Z\right) \\
&< E\left(\sum_{i=1}^n d_i\right) + E\left[d(G_{p_1} * \dots * G_{p_n}, G_{\alpha_1} * \dots * G_{\alpha_n})\right] \\
&\quad + d(G_{\alpha_1} * \dots * G_{\alpha_n}, H_{\alpha_1} * \dots * H_{\alpha_n}) \\
&< E\left[\sum_{i=1}^n d_i + |p_i - \alpha_i| + \alpha_i^2\right].
\end{aligned}$$

Finally, to prove (1.5), consider the inequality

$$(3.4) \quad d\left(\sum_{i=1}^n X_i, Z\right) < d\left(\sum_{i=1}^n X_i, Z'\right) + d(Z', Z),$$

where Z is $CP(\alpha, F)$ and Z' is mixed $CP(\sum_{i=1}^n p_i, F)$. By Lemmas 2.1 and 2.3 we

have $d(Z', Z) < E\left|\sum_{i=1}^n p_i - \alpha\right|$. Applying this and (1.3) to (3.4) yields

(1.5).

4. A Compound Poisson Approximation for Markovian Occurrences of an Event

Suppose that $Y_0, Y_1, \dots$ is a Markov chain with states 0 and 1 that represent the non-occurrence and occurrence, respectively, of a certain event E . Let $\epsilon = P(Y_1 = 1 | Y_0 = 0)$ and $p = P(Y_1 = 1 | Y_0 = 1)$, and assume that ϵ and p are not zero or one. The stationary distribution of this Markov chain is

$$\pi(0) = (1 - p)/(1 - p + \epsilon), \quad \pi(1) = \epsilon/(1 - p + \epsilon).$$

Consequently, when ϵ is small, then the event E is rare.

Consider the sum $N_n = \sum_{i=1}^n Y_i$, which is the number of occurrences of the event E in time n . We assume, for simplicity, that the Markov chain

is stationary. Isham (1980) and Böker and Serfozo (1983) showed that if ϵ varies with n such that $\epsilon \rightarrow 0$ and $n\epsilon \rightarrow \alpha > 0$ as $n \rightarrow \infty$, then N_n converges in distribution to a random variable Z that is $CP(\alpha, F)$ with $F(\{k\}) = p^{k-1}(1-p)$, $k \geq 1$. A bound on the rate of this convergence is given in the following result. Brown (1983) obtained a variation of this bound by another approach.

Theorem 4.1.

$$(4.1) \quad d(N_n, Z) \leq |n\epsilon - \alpha| + \epsilon(1 + p + \epsilon n(2 - p))/(1 - p + \epsilon).$$

Proof. Define the random variables

$$X_i = \sum_{k=1}^{\infty} k(1 - Y_{i-1})Y_i \cdots Y_{i+k-1}(1 - Y_{i+k}), \quad i=1, \dots, n,$$

$$X'_1 = \sum_{k=1}^{\infty} kY_1Y_2 \cdots Y_k(1 - Y_{k+1}).$$

When the Markov chain begins a sojourn in state 1 at time i (a success run of the event E), then X_i records the length of that sojourn.

Clearly

$$\begin{aligned} p_i &:= P(X_i > 1 | Y_0, \dots, Y_{i-1}) \\ &= \sum_{k=1}^{\infty} (1 - Y_{i-1}) \epsilon p^{k-1}(1-p) = \epsilon(1 - Y_{i-1}), \end{aligned}$$

$$F_i(k) := P(X_i \leq k | Y_0, \dots, Y_{i-1}, X_i > 1) = F(k).$$

Let $T_n = \sum_{i=1}^n X_i$ and $T'_n = T_n + X'_1$, and consider

$$(4.2) \quad d(N_n, Z) \leq d(N_n, T'_n) + d(T'_n, T_n) + d(T_n, Z).$$

Clearly

$$(4.3) \quad d(N_n, T'_n) \leq P(N_n \neq T'_n) = P(Y_n = 1, Y_{n+1} = 1) = \pi(1)p,$$

$$(4.4) \quad d(T'_n, T_n) \leq P(X'_1 \neq 0) = P(Y_1 = 1) = \pi(1),$$

and by (1.5)

$$\begin{aligned} (4.5) \quad d(T_n, Z) &\leq \sum_{i=1}^n E p_i^2 + E \left| \sum_{i=1}^n p_i - \alpha \right| \\ &= n\varepsilon^2 \pi(0) + E \left| \varepsilon \sum_{i=1}^n (1 - Y_{i-1}) - \alpha \right| \\ &\leq n\varepsilon^2 \pi(0) + \varepsilon n \pi(1) + |n\varepsilon - \alpha|. \end{aligned}$$

Combining (4.2) - (4.5) yields (4.1).

Remark. Note that the preceding proof applies (1.5) to the auxiliary sum T_n instead of to the original sum N_n . One could also apply (1.4) to T_n , but this would yield (4.1) with $2 - p$ replaced by $(2 - p)^2$, which is worse.

REFERENCES

- [1] Barbour, A. D. and Eagleson, G. K. (1983). Poisson approximation for some statistics based on exchangeable trials. Adv. Appl. Probability 15, 583-600.
- [2] Boker, F. and Serfozo, R. F. (1983). Ordered thinnings of point processes and random measures. Stochastic Process. Appl. 15, 113-132.
- [3] Brown, T. C. (1983). Some Poisson approximations using compensators. Ann. Probability 11, 726-744.
- [4] Chen, L.H.Y. (1975). An approximation theorem for convolutions of probability measures. Ann. Probability 3, 992-999.
- [5] Franken, P. (1964). Approximation der Verteilungen von Summen unabhängiger nichtnegativer ganzzahler Zufallsgrossen durch Poissonsche Verteilungen. Math. Nachr. 27, 303-340.
- [6] Freedman, D. (1974). The Poisson approximation for dependent events. Ann. Probability 2, 256-269.
- [7] Hodges, J. L. and Le Cam, L. (1960). The Poisson approximation to the Poisson binomial distribution. Ann. Math. Statist. 31, 737-740.
- [8] Isham, V. (1980). Dependent thinning of point processes. J. Appl. Probability 17, 987-995.
- [9] Kabanov, Y. M., Liptser, R. S. and A. N. Shirayev (1983). Weak and strong convergence of the distributions of counting processes. Theor. Probability Appl. 28, 303-336.
- [10] Kerstan, J. (1964). Verallgemeinerung eines Satzes von Prochorow und Le Cam. Z. Wahrscheinlichkeits-theorie verw. Gebide 2, 173-179.
- [11] Le Cam, L. (1960). An approximation theorem for the Poisson binomial distribution. Pacific J. Math. 10, 1181-1197.
- [12] Serfling, R. J. (1975). A general Poisson approximation theorem. Ann. Probability 3, 726-731.
- [13] Valkeila, E. (1982). A general Poisson approximation theorem. Stochastics 7, 159-171.

School of Industrial and Systems Engineering
Georgia Institute of Technology
Atlanta, GA 30332

END

FILMED

10-85

DTIC