

AD-A160 302

ON THE ASYMPTOTIC PROPERTIES OF A KERNEL-TYPE QUANTILE
ESTIMATOR FROM CEN. (U) SOUTH CAROLINA UNIV COLUMBIA
DEPT OF MATHEMATICS AND STATISTI... Y O LIO ET AL.

1/1

UNCLASSIFIED

MAY 85 TR-104 AFOSR-TR-85-0065


F/G 12/1

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Department of Mathematics
and Statistics
The University of South Carolina
Columbia, South Carolina 29208

AFOSR-TR- 85-0865

②

AD-A160 302


ON THE ASYMPTOTIC PROPERTIES OF A KERNEL-TYPE
QUANTILE ESTIMATOR FROM CENSORED SAMPLES *

by

Y. L. Lio, W. J. Padgett, and K. F. Yu**

University of South Carolina
Statistics Technical Report No. 104
62G05-13

DTIC
SELECTED
OCT 15 1985
E



DTIC FILE COPY

85 10 11 119

ON THE ASYMPTOTIC PROPERTIES OF A KERNEL-TYPE
QUANTILE ESTIMATOR FROM CENSORED SAMPLES *

by

Y. L. Lio, W. J. Padgett, and K. F. Yu **

University of South Carolina
Statistics Technical Report No. 104
62G05-13

May, 1985

Department of Mathematics and Statistics
University of South Carolina
Columbia, South Carolina 29208

DTIC
ELECTE
OCT 15 1985
S D
E

AIR FORCE
OFFICE OF SCIENTIFIC RESEARCH
AND ARMY RESEARCH OFFICE

* Research supported by the U.S. Air Force Office of Scientific Research and Army Research Office under grant AFOSR-84-0156.

** Partially supported by a University of South Carolina Research and Productive Scholarship Grant.

Abstract

Some asymptotic results for a kernel type estimator of the quantile function from right-censored data are obtained. The estimator is defined by $Q_n(p) = h_n^{-1} \int_0^1 \hat{Q}_n(t) K((t-p)/h_n) dt$, which is smoother than the usual product-limit quantile function $\hat{Q}_n(p) = \inf\{t: \hat{F}_n(t) \geq p\}$, where $\hat{F}_n$ denotes the product-limit estimator of the lifetime distribution F_0 . Under the random censorship model and general conditions on h_n, K , and F_0 , the asymptotic normality of $Q_n(p)$ is proven. In addition, an approximation to Q_n is shown to be asymptotically uniformly equivalent to Q_n in mean square.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Key Words: Random right-censorship; Kernel estimation; Product-limit quantile function; Asymptotic normality; Mean-square convergence.

1. Introduction

In reliability and medical studies, it is often of interest to estimate various quantiles of the unknown lifetime distribution. In particular, the median lifetime and extreme quantiles are of interest to the experimenter in such studies. In many life testing and medical follow-up experiments, however, arbitrarily right-censored data arise, and it is important to be able to estimate the quantiles of interest based on the censored data. For such data, some kernel-type quantile estimators are considered in this paper which give smoother estimates than the usual product-limit quantile function. $\rightarrow$

For any probability distribution function G , denote the quantile function by $Q(p) \equiv G^{-1}(p) = \inf\{x: G(x) \geq p\}$, $0 \leq p \leq 1$. For a random (uncensored) sample $Y_1, \dots, Y_n$ from G , the sample quantile function $G_n^{-1}(p) = \inf\{x: G_n(x) \geq p\}$, $0 \leq p \leq 1$, has been used to estimate $Q(p)$, where G_n denotes the sample distribution function. Csörgő (1983) gave many of the known results concerning $G_n^{-1}(p)$. Also, Falk (1984) studied the relative deficiency of the sample quantile with respect to kernel-type estimators, and Falk (1985) obtained asymptotic normality for kernel estimators. Yang (1985) has obtained some convergence properties of kernel estimators of $Q(p)$ and gave some simulation results comparing kernel-type estimators with other estimators. For arbitrarily right-censored data, Sander (1975) proposed estimation of $Q(p)$ by the quantile function of the product-limit estimator, and she and Cheng (1981) derived some asymptotic properties of that estimator. Also, Csörgő (1983) presented strong approximation results for that estimator.

Recently, Padgett (1985) studied a smoothed nonparametric estimator of

$Q(p)$ from arbitrarily right-censored data based on the kernel method. It was shown that his estimator, mentioned briefly by Parzen (1979, p. 119), was strongly consistent, and a small Monte Carlo study was performed to compare the estimator with the product-limit estimator. In addition, a simple approximation to this kernel estimator was shown to be almost surely asymptotically equivalent to it.

The purpose of this paper is to further study the asymptotic properties of the estimators proposed by Padgett (1985). In particular, the asymptotic normality will be proven, and the asymptotic equivalence in mean square of the estimator and its approximation will be shown under general conditions on the kernel function, bandwidth sequence, lifetime distribution, and censoring mechanism.

2. Notation and Preliminaries

Let $X_1^0, X_2^0, \dots, X_n^0$ denote the true survival times of n items or individuals which are censored on the right by a sequence $U_1, U_2, \dots, U_n$, which in general may be either constants or random variables. It is assumed that the X_i^0 's are nonnegative independent identically distributed random variables with common unknown distribution function F_0 and unknown quantile function

$$Q^0(p) \equiv \xi_p^0 = \inf\{t: F_0(t) \geq p\}, \quad 0 \leq p \leq 1.$$

The observed right-censored data are denoted by the pairs (X_i, Δ_i) , $i=1, \dots, n$, where

$$X_i = \min\{X_i^0, U_i\}, \quad \Delta_i = \begin{cases} 1 & \text{if } X_i^0 \leq U_i \\ 0 & \text{if } X_i^0 > U_i \end{cases}.$$

For the asymptotic results of this paper, the random right-censorship model will be assumed, that is, $U_1, \dots, U_n$ constitute a random sample from a distribution H (usually unknown) and are independent of $X_1^0, \dots, X_n^0$. The distribution function of each X_i , $i=1, \dots, n$, is then $F = 1 - (1-F_0)(1-H)$.

A popular estimator of the survival function $S_0(t) = 1 - F_0(t)$ based on (X_i, Δ_i) , $i=1, \dots, n$, is the product-limit estimator, proposed by Kaplan and Meier (1958) as the "nonparametric maximum likelihood estimator." Let (Z_i, Δ_i') , $i=1, \dots, n$, denote the ordered X_i 's along with their corresponding Δ_i 's. The product-limit estimator of $S_0(t)$, shown to be "self-consistent" by Efron (1967), is defined by

$$\hat{P}_n(t) = \begin{cases} 1, & 0 \leq t \leq Z_1, \\ \prod_{i=1}^{k-1} \left(\frac{n-i}{n-i+1} \right)^{\Delta_i'}, & Z_{k-1} < t \leq Z_k, \quad k=2, \dots, n \\ 0, & t > Z_n. \end{cases}$$

Denote the product-limit estimator of $F_0(t)$ by $\hat{F}_n(t) = 1 - \hat{P}_n(t)$, and let s_j denote the jump of $\hat{P}_n$ at Z_j , that is

$$s_j = \begin{cases} 1 - \hat{P}_n(Z_2), & j = 1 \\ \hat{P}_n(Z_j) - \hat{P}_n(Z_{j+1}), & j = 2, \dots, n-1 \\ \hat{P}_n(Z_n), & j = n. \end{cases}$$

Note that $s_j = 0$ if and only if $\Delta_j' = 0$, $j < n$, i.e. whenever Z_j is a censored observation. Also, denote $S_i \equiv \hat{F}_n(Z_{i+1}) = \sum_{j=1}^i s_j$, $i=1, \dots, n$, with

$S_0 \equiv 0$, $Z_0 \equiv 0$, and $Z_{n+1} \equiv Z_n + \epsilon$, for some positive constant ϵ .

It is natural to estimate ξ_p^0 by the product-limit quantile function $\hat{Q}_n(p) \equiv \hat{\xi}_p^0 = \inf\{t: \hat{F}_n(t) \geq p\}$. Sander (1975) and Cheng (1981) obtained

asymptotic normality results for $\hat{\xi}_p^0$. Padgett (1985) smoothed $\hat{Q}_n$ by the kernel method to obtain the estimator

$$\begin{aligned} Q_n(p) &= h_n^{-1} \int_0^1 \hat{Q}_n(t) K((t-p)/h_n) dt \\ &= h_n^{-1} \sum_{i=1}^n Z_i \int_{S_{i-1}}^{S_i} K((t-p)/h_n) dt, \end{aligned} \quad (2.1)$$

where K is an appropriate kernel function and $\{h_n\}$ is a bandwidth sequence. Also, a simpler kernel-type estimator which is an approximation to (2.1) was defined by

$$Q_n^*(p) = h_n^{-1} \sum_{i=1}^n Z_i S_i K((S_i - p)/h_n). \quad (2.2)$$

Note that only the uncensored observations actually appear in the sums of (2.1) and (2.2).

In the next section, the asymptotic normality of $Q_n(p)$ and $Q_n^*(p)$ will be obtained. The following general conditions on the kernel function, the bandwidth sequence, and the lifetime and censoring distributions will be assumed:

$$(h.1) \quad h_n \rightarrow 0 \text{ as } n \rightarrow \infty;$$

$$(K.1) \quad K(x) \text{ is a bounded probability density function which has finite support, i.e. } K(x) = 0 \text{ for } |x| > c \text{ for some } c > 0;$$

$$(K.2) \quad K \text{ is symmetric about zero;}$$

$$(K.3) \quad K \text{ satisfies a Lipschitz condition, i.e. there exist a constant } \Gamma \text{ such that for all } x, y,$$

$$|K(x) - K(y)| \leq \Gamma |x - y|;$$

$$(F.1) \quad F_0 \text{ is continuous with density function } f_0; \text{ and}$$

$$(F.2) \quad H(T_{F_0}) < 1, \text{ where } T_{F_0} = \sup\{t: F_0(t) < 1\}.$$

It should be noted that these conditions are not prohibitive and (F.1) and (F.2) are similar to conditions required by Cheng (1981). Also, (F.2) insures that observations will be available from the entire support of F_0 , a common condition in random right-censorship models.

3. The Main Results

In this section, the main results are summarized in Theorems 3.1 and 3.2. The proofs will be presented in the next section. Theorem 3.1 gives conditions for the asymptotic normality of $Q_n(p)$. The asymptotic uniform mean-squared equivalence of Q_n and Q_n^* will be shown in Theorem 3.2.

Theorem 3.1. Assume, in addition to conditions (h.1), (K.1) - (K.3), (F.1), and (F.2), that the derivative f'_0 is continuous at ξ_p^0 and $f_0(\xi_p^0) > 0$. Suppose $\{h_n\}$ is such that $n^{1/4}h_n \rightarrow 0$ as $n \rightarrow \infty$. Then for $0 < p < T$, where $T < 1$, as $n \rightarrow \infty$, $\sqrt{n}[Q_n(p) - Q^0(p)] \rightarrow Z$ in distribution, where Z is a normally distributed random variable with mean zero and variance

$$\sigma_p^2 = (1-p)^2 \int_0^{\xi_p^0} [1-F(u)]^{-2} \frac{dF_0^*(u)}{f_0^2(\xi_p^0)}$$

with $1-F(u) = [1-F_0(u)][1-H(u)]$ and $F_0^*(u) = P(X_1 \leq u, \Delta_1 = 1)$, the subdistribution function of the uncensored observations.

Note that an example of a bandwidth sequence that satisfies the conditions of Theorem 3.1 is $h_n = cn^{-\delta}$ with $\delta > 1/4$.

The next theorem gives some conditions for which Q_n^* and Q_n are asymptotically uniformly equivalent in mean square.

Theorem 3.2. Suppose F_0 and H are continuous and (h.1), (K.1), and (K.3) hold. Assume $E(X_1^{2q}) < \infty$ for some $q > 1$, where $X_1 = \min\{X_1^0, U_1\}$. Let η be such that $[1-F_0(\eta)][1-H(\eta)] > 0$ and let $T^* = F_0(\eta)$. Then for all $T \in [0, T^*)$, $\lim_{n \rightarrow \infty} E[\sup_{0 \leq p \leq T} |Q_n^*(p) - Q_n(p)|^2] = 0$, provided $n^{1/2} h_n^4 \rightarrow \infty$ as $n \rightarrow \infty$.

4. Proofs of Theorems

The following two lemmas will be needed in the proof of Theorem 3.1.

In this section, $\{K(s, t): 0 \leq s \leq 1, t \geq 0\}$ will denote the generalized Kiefer process as stated by Csörgő (1983, Ch. 8).

Lemma 1. For $0 < p < T$, where $T < 1$, and $\delta < \min\{T - p, p\}$, as $n \rightarrow \infty$

$$\sup_{|h| < \delta} |n^{-1/2} [K(p+h, n) - K(p, n)]| \rightarrow 0 \text{ in probability.}$$

The proof of Lemma 1 follows the same argument as the proof of Theorem 8.2.1 of Csörgő (1983).

Lemma 2. Suppose the derivative f'_0 is continuous at ξ_p^0 and $f_0(\xi_p^0) > 0$. Under assumptions (h.1), (K.1), (K.2), (F.1) and (F.2), for $0 < p < 1$,

$$\left| \int_0^1 [q_n(t) - q_n(p)] h_n^{-1} K\left(\frac{t-p}{h_n}\right) dt \right| \rightarrow 0 \text{ in probability}$$

as $n \rightarrow \infty$, where $q_n(t) = n^{1/2} [\hat{Q}_n(t) - Q^0(t)]$ denotes the product-limit quantile process.

Proof. For any given $\delta > 0$, there exists N such that when $n \geq N$,

$$\begin{aligned}
& \left| \int_0^1 [q_n(t) - q_n(p)] h_n^{-1} K\left(\frac{t-p}{h_n}\right) dt \right| \\
&= \left| \int_{A(\delta)} [q_n(t) - q_n(p)] h_n^{-1} K\left(\frac{t-p}{h_n}\right) dt \right| \\
&\leq \sup_{t \in A(\delta)} |q_n(t) - q_n(p)|, \tag{4.1}
\end{aligned}$$

where $A(\delta) = [p-\delta, p+\delta]$. By the conditions on f_o , for δ sufficiently small, $f_o(Q^o(t)) > 0$ for all $t \in A(\delta)$. Hence, the right-hand-side of (4.1) is less than or equal to

$$\begin{aligned}
& \sup_{t \in A(\delta)} |\beta_n(t) - \beta_n(p)| \cdot \left| \frac{1}{f_o(Q^o(t))} \right| \\
&+ \sup_{t \in A(\delta)} \left| \beta_n(p) \left[\frac{1}{f_o(Q^o(t))} - \frac{1}{f_o(Q^o(p))} \right] \right|,
\end{aligned}$$

where $\beta_n(t) = f_o(Q^o(t))q_n(t)$.

Let

$$a = \sup_{t \in A(\delta)} \left| \frac{1}{f_o(Q^o(t))} \right| \quad \text{and} \quad b = \sup_{t \in A(\delta)} \left| \frac{1}{f_o(Q^o(t))} - \frac{1}{f_o(Q^o(p))} \right|.$$

From Corollary 1 of Cheng (1981), since f_o is continuous at ξ_p^o , $\beta_n(p) \rightarrow Z$ in distribution as $n \rightarrow \infty$, where Z is a normally distributed random variable

with mean zero and variance $\sigma^2 = (1-p)^2 \int_0^{\xi_p^o} [1-F(u)]^{-2} dF_o^*(u)$,

with $1-F(u) = [1-F_o(u)][1-H(u)]$ and $F_o^*(u) = P(X_1 \leq u, \Delta_1 = 1)$. Therefore,

$$\sup_{t \in A(\delta)} \left| \beta_n(p) \left[\frac{1}{f_o(Q^o(t))} - \frac{1}{f_o(Q^o(p))} \right] \right| \leq b |\beta_n(p)| \tag{4.2}$$

and for given $\epsilon > 0$

$$\lim_{n \rightarrow \infty} \sup P(|\tilde{\beta}_n(p)| > \epsilon/b) \leq P(|Z| > \epsilon/b) \leq b\sigma^2/\epsilon^2. \quad (4.3)$$

Now,

$$\begin{aligned} \sup_{t \in A(\delta)} |\tilde{\beta}_n(t) - \tilde{\beta}_n(p)| \cdot \left| \frac{1}{f_0(Q^0(t))} \right| &\leq a \sup_{t \in A(\delta)} |\tilde{\beta}_n(t) - \tilde{\beta}_n(p)| \\ &\leq a \{ \sup_{t \in A(\delta)} |\tilde{\beta}_n(t) - n^{-1/2} K(t,n)| \\ &\quad + \sup_{t \in A(\delta)} |\tilde{\beta}_n(p) - n^{-1/2} K(p,n)| \\ &\quad + \sup_{t \in A(\delta)} |n^{-1/2} [K(p,n) - K(t,n)]| \}. \end{aligned} \quad (4.4)$$

For small enough δ , $p + \delta < T < 1$ and $p - \delta \geq 0$, so that by Corollary 8.3.3 of Csörgő (1983) as $n \rightarrow \infty$

$$\sup_{t \in A(\delta)} |\tilde{\beta}_n(t) - n^{-1/2} K(t,n)| \rightarrow 0 \text{ in probability}$$

and

$$\sup_{t \in A(\delta)} |\tilde{\beta}_n(p) - n^{-1/2} K(p,n)| \rightarrow 0 \text{ in probability.}$$

By Lemma 8.2.1 of Csörgő (1983) (or Berkes and Philipp, 1977) with $0 = t_0 < t_1 = n$, letting $B(s) \equiv n^{-1/2} K(s,n)$, $0 \leq s \leq 1$,

$$\sup_{t \in A(\delta)} |B(p) - B(t)| = \sup_{|h| < \delta} |B(p+h) - B(p)|,$$

which by Lemma 1 converges to zero in probability for sufficiently small δ .

Therefore, (4.4) converges to zero in probability as $n \rightarrow \infty$.

Finally, since b depends on δ , letting b become arbitrarily small gives from (4.2) and (4.4) that

$$\sup_{t \in A(\delta)} |\beta_n(p) [\frac{1}{f_0(Q^0(t))} - \frac{1}{f_0(Q^0(p))}]| \rightarrow 0 \text{ in probability.}$$

Thus, the result follows. ///

Proof of Theorem 3.1. Analogous to the beginning of the proof of Theorem 1 of Yang (1985), write

$$\begin{aligned} \sqrt{n} [Q_n(p) - Q^0(p)] &= \int_0^1 [q_n(t) - q_n(p)] h_n^{-1} K(\frac{t-p}{h_n}) dt \\ &+ n^{1/2} [\int_0^1 Q^0(t) h_n^{-1} K(\frac{t-p}{h_n}) dt - Q^0(p)] + q_n(p), \end{aligned} \quad (4.5)$$

where $q_n(t) = n^{1/2} [\hat{Q}_n(t) - Q^0(t)]$ as in Lemma 2.

From Lemma 2, the first term on the right hand side of (4.5) is $o_p(1)$, which means that it converges to zero in probability as $n \rightarrow \infty$. Similar to Yang's (1985) equation (10),

$$\begin{aligned} n^{1/2} [\int_0^1 Q^0(t) h_n^{-1} K(\frac{t-p}{h_n}) dt - Q^0(p)] \\ = o(n^{1/2} h_n^2) + n^{1/2} h_n^2 Q^{0''}(p) \int_{-\infty}^{\infty} \frac{t^2}{2} K(t) dt. \end{aligned} \quad (4.6)$$

With the assumption that $n^{1/4} h_n \rightarrow 0$ as $n \rightarrow \infty$, (4.6) is also $o_p(1)$.

Therefore, by Corollary 1 of Cheng (1981), the conclusion of the theorem follows. ///

Proof of Theorem 3.2. For $0 \leq p \leq T$, write

$$Q_n^*(p) - Q_n(p) = h_n^{-1} \sum_{i=1}^n Z_i [s_i K(\frac{S_i - p}{h_n}) - \int_{S_{i-1}}^{S_i} K(\frac{t-p}{h_n}) dt].$$

When $s_1 > 0$, that is, Z_1 is uncensored, let S_i^* be an interior point of the interval (S_{i-1}, S_i) with probability one so that

$$s_1 K((S_1^* - p)/h_n) = \int_{S_{i-1}}^{S_i} K((t-p)/h_n) dt \quad \text{almost surely.}$$

Then by condition (K.3), letting I_A denote the indicator function of the set A,

$$\begin{aligned} |Q_n^*(p) - Q_n(p)| I_{[0,T]}(p) &\leq h_n^{-1} \sum_{i=1}^n s_i Z_i |K((S_i^* - p)/h_n) \\ &\quad - K((S_i^* - p)/h_n)| I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p) \\ &\leq \Gamma h_n^{-2} \sum_{i=1}^n Z_i s_i |S_i - S_i^*| I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p) \\ &\leq \Gamma h_n^{-2} \sum_{i=1}^n Z_i s_i^2 I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p) \quad \text{almost surely.} \end{aligned}$$

So

$$|Q_n^*(p) - Q_n(p)|^2 I_{[0,T]}(p) \leq \Gamma^2 h_n^{-4} (\sum_{i=1}^n Z_i s_i^2 I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p))^2. \quad (4.7)$$

Now,

$$\begin{aligned} &(\sum_{i=1}^n Z_i s_i^2 I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p))^2 \\ &\leq \sum_{i=1}^n Z_i^2 s_i^3 I_{[0,T]}(p) I_{[S_i^* - ch_n, 1]}(p) \\ &\leq \sum_{i=1}^{n+1} Z_{i-1}^2 (|\hat{F}_n(Z_i) - F_0(Z_i)| + |F_0(Z_i^-) - \hat{F}_n(Z_i^-)|)^3 I_{[0,T]}(p) I_{[S_{i-2} - ch_n, 1]}(p), \quad (4.8) \end{aligned}$$

where $g(x^-)$ denotes the limit from the left at x of the function g .

There is an N such that for all $n \geq N$, $T + ch_n < T^*$ and

$T_N < F_0(\eta - \epsilon)$ where $T_N \equiv T + ch_N$. For all such n 's

$$\begin{aligned} & \sum Z_{i-1}^2 (|\hat{F}_n(Z_1) - F_0(Z_1)| + |F_0(Z_1^-) - \hat{F}_n(Z_1^-)|)^3 I_{[0,T]}^{(p)} I_{[S_{1-2}^{-ch_n}, 1]}^{(p)} \\ & \leq 8 \sum Z_{i-1}^2 \sup_{0 \leq x \leq Q_n(T_N)} |\hat{F}_n(x) - F_0(x)|^3, \end{aligned} \quad (4.9)$$

which is independent of p . Notice also that

$$\sup_{0 \leq x \leq \hat{Q}_n(T_N) + \epsilon} |\hat{F}_n(x) - F_0(x)|^3 \leq \sup_{0 \leq x \leq \eta} |\hat{F}_n(x) - F_0(x)|^3 + I_{[\hat{Q}_n(T_N) > \eta - \epsilon]}.$$

From the exponential bound in Theorem 2 of Földes and Rejtő (1981), for any $r \geq 1$, $\{n^{1/2} \sup_{0 \leq x \leq \eta} |\hat{F}_n(x) - F_0(x)|\}^r$, $n \geq 1$ is uniformly integrable. Also,

$$\begin{aligned} & E(n^{3/2} I_{[\hat{Q}_n(T_N) > \eta - \epsilon]})^r \\ & = n^{3r/2} P[\hat{Q}_n(T_N) > \eta - \epsilon] \\ & = n^{3r/2} P[\hat{Q}_n(T_N) - F_0^{-1}(T_N) > \eta - F_0^{-1}(T_N) - \epsilon]. \end{aligned}$$

Letting $\gamma = \eta - F_0^{-1}(T_N) - \epsilon$, where ϵ is chosen so that $\gamma > 0$,

$$\begin{aligned} & P[\hat{Q}_n(T_N) - F_0^{-1}(T_N) > \gamma] \\ & \leq P[T_N > \hat{F}_n(\eta - \epsilon)] \\ & = P[T_N - F_0(\eta - \epsilon) > \hat{F}_n(\eta - \epsilon) - F_0(\eta - \epsilon)] \\ & \leq P[|\hat{F}_n(\eta - \epsilon) - F_0(\eta - \epsilon)| > F_0(\eta - \epsilon) - T_N]. \end{aligned}$$

By the same exponential bound as in Theorem 2 of Földes and Réjtö (1981),
as $n \rightarrow \infty$

$$E(n^{3/2} I_{[\hat{Q}_n(T_N) > \eta - \epsilon]})^r \rightarrow 0.$$

Therefore, $\{(n^{1/2} \sup_{0 \leq x \leq \hat{Q}_n(T_N) + \epsilon} |\hat{F}_n(x) - F_0(x)|)^r, n \geq 1\}$ is uniformly integrable.

By hypothesis, $E(X_1^{2q}) < \infty$ for some $q > 1$, so $\{(n^{-1} \sum_{i=1}^n Z_i^2)^q, n \geq 1\}$ is uniformly integrable. Thus,

for $\frac{1}{s} + \frac{1}{q} = 1$,

$$\begin{aligned} & E(n^{3/2} \sup_{0 \leq x \leq \hat{Q}_n(T_N) + \epsilon} |\hat{F}_n(x) - F_0(x)|^3 n^{-1} \sum_{i=1}^n Z_i^2) \\ & \leq (E(n^{3/2} \sup_{0 \leq x \leq \hat{Q}_n(T_N) + \epsilon} |\hat{F}_n(x) - F_0(x)|^3)^s)^{1/s} E((n^{-1} \sum_{i=1}^n Z_i^2)^q)^{1/q} \\ & = o(1). \end{aligned} \tag{4.10}$$

Therefore, from (4.7) - (4.10), by the hypothesis $n^{1/2} h_n^4 \rightarrow \infty$,

$$E[\sup_{0 \leq p \leq T} |Q_n^*(p) - Q_n(p)|^2] = o(1),$$

completing the proof. ///

References

- Berkes, I. and Philipp, W. (1977). An almost sure invariance principle for the empirical distribution function of mixing random variables. Z. Wahrsch. Verw. Gebiete 41, 115-137.
- Cheng, K. F. (1981). On almost sure representations for quantiles of the product limit estimator with applications. Research Report No. 87, SUNY Buffalo, Dept. of Statistics.
- Csörgő, M. (1983). Quantile Processes with Statistical Applications. CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, PA.
- Efron, B. (1967). The two-sample problem with censored data. Proc. Fifth Berkeley Sympos. 4, 831-853.
- Falk, M. (1984). Relative deficiency of kernel type estimators of quantiles. Ann. Statist. 12, 261-268.
- Falk, M. (1985). Asymptotic normality of the kernel quantile estimator. Ann. Statist. 13, 428-433.
- Földes, A. and Rejtő, L. (1981). A LIL type result for the product limit estimator. Z. Wahr. Verw. Gebiete 56, 75-86.
- Kaplan, E. L. and Meier, P. (1958). Nonparametric estimation from incomplete observations. J. Amer. Statist. Assoc. 53, 457-481.
- Padgett, W. J. (1985). A kernel type estimator of a quantile function from right-censored data. Statistics Technical Report No. 101R, University of South Carolina, Dept. of Math. and Statist.
- Parzen, E. (1979). Nonparametric statistical data modeling. J. Amer. Statist. Assoc. 74, 105-121.
- Sander, J. (1975). The weak convergence of quantiles of the product limit estimator. Technical Report 5, Stanford University, Dept. of Statistics.
- Yang, S. S. (1985). A smooth nonparametric estimator of a quantile function. Technical Report, Kansas State University, Dept. of Statistics.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS AD-A160302	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Stat. Tech. Rep. No. 104 (62G05-13)		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 5- 305	
6a. NAME OF PERFORMING ORGANIZATION Department of Mathematics and Statistics	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) University of South Carolina Columbia SC 29208		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR, ARO	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0156	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 61102F	TASK NO. A5
		PROJECT NO. 2304	WORK UNIT NO.
11. TITLE (Include Security Classification) ON THE ASYMPTOTIC PROPERTIES OF A KERNEL TYPE QUANTILE ESTIMATOR FROM CENSORED SAMPLES			
12. PERSONAL AUTHOR(S) Y. L. Lio, W. J. Padgett, and K. F. Yu			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day) 1985 May	15. PAGE COUNT 13
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		Random right-censorship; Kernel estimation; Product-limit quantile function; Asymptotic normality; Mean-square con- vergence.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
Some asymptotic results for a kernel type estimator of the quantile function from right-censored data are obtained. The estimator is defined by $Q_n(p) = h_n^{-1} \int_0^1 \hat{Q}_n(t) K((t-p)/h_n) dt$, which is smoother than the usual product-limit quantile function $\hat{Q}_n(p) = \inf\{t: \hat{F}_n(t) \geq p\}$, where $\hat{F}_n$ denotes the product-limit estimator of the lifetime distribution F_0. Under the random censorship model and general conditions on h_n, K, and F_0, the asymptotic normality of $Q_n(p)$ is proven. In addition, as approximation to Q_n is shown to be asymptotically uniformly equivalent to Q_n in mean square.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL CPT Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5027	22c. OFFICE SYMBOL NM

END

FILMED

11-85

DTIC