

AD-A168 460

DEMONSTRATION OF A JACOBI POLYNOMIAL REPRESENTATION OF
A KROMECKER DELTA. (U) JOHNS HOPKINS UNIV LAUREL MD
APPLIED PHYSICS LAB R A FARRELL ET AL. SEP 85

1/1

UNCLASSIFIED

JHU/APL-TG-1330 N00024-85-C-5301


F/G 12/1

NL

END

FILED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963-A

12

JHU APL

TG 1350

SEPTEMBER 1985

AD-A160 460


DEMONSTRATION OF A JACOBI POLYNOMIAL REPRESENTATION OF A KRONECKER DELTA FUNCTION

RICHARD A. FARRELL

ERNEST P. GRAY

ROBERT W. HART

OTIC FILE COPY



Approved for public release; distribution is unlimited

85 10 22 042

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

PLEASE FOLD BACK IF NOT NEEDED
FOR BIBLIOGRAPHIC PURPOSES

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE None			
4 PERFORMING ORGANIZATION NUMBER(S) JHU API TG 1350		5 MONITORING ORGANIZATION REPORT NUMBER(S) JHU API TG 1350	
6a NAME OF PERFORMING ORGANIZATION The Johns Hopkins University Applied Physics Laboratory	6b OFFICE SYMBOL (If Applicable) RTP	7a NAME OF MONITORING ORGANIZATION Naval Plant Representative Office	
6c ADDRESS (City, State, and ZIP Code) Johns Hopkins Road Laurel, Maryland 20707		7b ADDRESS (City, State, and ZIP Code) Johns Hopkins Road Laurel, Maryland 20707	
8a NAME OF FUNDING SPONSORING ORGANIZATION Naval Sea Systems Command	8b OFFICE SYMBOL (If Applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00024-85-C-5301	
8c ADDRESS (City, State, and ZIP Code) Washington, DC 20362		10 SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO PROJECT NO TASK NO WORK UNIT ACCESSION NO X83J	
11 TITLE (Include Security Classification) Demonstration of a Jacobi Polynomial Representation of a Kronecker Delta Function (U)			
12 PERSONAL AUTHOR(S) Farrell, Richard A.; Gray, Ernest P.; Hart, Robert W.			
13a TYPE OF REPORT Technical Memorandum	13b TIME COVERED FROM TO	14 DATE OF REPORT (Year, Month, Day) 1985, September	15 PAGE COUNT 14
16 SUPPLEMENTARY NOTATION			
17 LOCATOR CODES FIELD GROUP SUB GROUP		18 SUBJECT TERMS Kronecker delta function Jacobi polynomials fluid shells	
19 ABSTRACT (Include in reverse if necessary and identify by block number) A formula derived by Noble has been generalized to obtain an expansion of the Kronecker delta function as an infinite series involving the products of Jacobi polynomials.			
20 ABSTRACT SECURITY CLASSIFICATION Unclassified		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a TELEPHONE (Include Area Code) (301) 953-5000		22b OFFICE SYMBOL NPRO	

UNCLASSIFIED

JHU/APL

TG 1350

SEPTEMBER 1985

Technical Memorandum

**DEMONSTRATION OF A
JACOBI POLYNOMIAL REPRESENTATION
OF A KRONECKER DELTA FUNCTION**

RICHARD A. FARRELL

ERNEST P. GRAY

ROBERT W. HART

THE JOHNS HOPKINS UNIVERSITY ■ APPLIED PHYSICS LABORATORY

Johns Hopkins Road, Laurel, Maryland 20707

Operating under Contract N60024-85-C-5301 with the Department of the Navy

Approved for public release; distribution is unlimited

ABSTRACT

A formula derived by Noble has been generalized to obtain an expansion of the Kronecker delta function as an infinite series involving the products of two Jacobi polynomials.

INTRODUCTION

In our investigations of the inviscid, incompressible flow of rotating fluid shells confined between concentric, spherical, rigid, co-rotating boundaries, we encountered the need for a proof of a Kronecker delta function representation in terms of a series of products of two Jacobi polynomials. In particular, we required a proof that

$$\delta_{l,0} = \frac{(\mu + 1/2)(1 - \Lambda)^{\mu + 1/2}}{2^{\mu + 1/2}} F_{\mu + l, \mu - l}(\Lambda), \quad (1)$$

where

$$F_{\mu + l, \mu - l}(\Lambda) = \sum_{k=0}^{\infty} \frac{(k + \mu/2 + l/2 + 1/4)}{(k + \mu + l + 1/2)(k + \mu + 1/2)} \times P_k^{(\mu + l + 1/2, -1)}(\Lambda) P_k^{(\mu - l + 1/2, -1)}(\Lambda), \quad 0 \leq |\Lambda| \leq 1. \quad (2)$$

For $(\alpha, \beta) > -1$, the $P_n^{(\alpha, \beta)}(\Lambda)$ are the Jacobi polynomials which are orthogonal over the interval $(-1, +1)$ with the integration weight factor $w(x) = (1-x)^\alpha(1+x)^\beta$. They are normalized by the relation $P_n^{(\alpha, \beta)}(1) = \Gamma(\alpha + n + 1)/[\Gamma(n+1)\Gamma(\alpha+1)]$, with $\Gamma(z)$ the gamma (factorial) function. For $(\alpha, \beta) \leq -1$, the Jacobi polynomials¹ $P_n^{(\alpha, \beta)}(\Lambda)$ are defined in terms of the polynomials for $(\alpha, \beta) > -1$ through repeated applications of the contiguous relations^{1,2}

$$(2n + \alpha + \beta) P_n^{(\alpha - 1, \beta)}(\Lambda) = (n + \alpha + \beta) P_n^{(\alpha, \beta)}(\Lambda) - (n + \beta) P_{n-1}^{(\alpha, \beta)}(\Lambda) \quad (3a)$$

and

$$(2n + \alpha + \beta) P_n^{(\alpha, \beta - 1)}(\Lambda) = (n + \alpha + \beta) P_n^{(\alpha, \beta)}(\Lambda) + (n + \alpha) P_{n-1}^{(\alpha, \beta)}(\Lambda) \quad (3b)$$

with $P_0^{(\alpha, \beta)}(\Lambda) = 1$ and $P_{-1}^{(\alpha, \beta)}(\Lambda) = 0$.

Certain sums involving products of two Jacobi polynomials have been previously evaluated (see, for example, Refs. 3-6). Our method for evaluating the sum in Eq. 2 employs mathematical procedures analogous to those which Noble⁵ used to show that⁶

¹W. Magnus, F. Oberhettinger, and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer-Verlag, New York (1966).

²I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series, and Products*, Alan Jeffrey, ed., Academic Press, New York (1965).

³H. I. Manocha and B. I. Sharma, "Some Formulae for Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 459-462 (1966).

⁴H. I. Manocha and B. I. Sharma, "Generating Functions of Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 431-433 (1966).

⁵B. Noble, "Some Dual Series Equations Involving Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 363-371 (1963).

⁶E. R. Hansen, *A Table of Series and Products*, Prentice-Hall, New York (1975).

$$2^{c-1}(1-y)^{-a}(1+x)^{-b}(x-y)^{a+b-c-1}\Delta(x,y) \\ = \Gamma(a+b-c) \sum_{k=0}^{\infty} \frac{k!\Gamma(k+c+1)(2k+c+1)}{\Gamma(k+a+1)\Gamma(k+b+1)} P_k^{(c-b,b)}(x) P_k^{(a,c-a)}(y), \quad (4)$$

where

$$\Delta(x,y) = \begin{cases} 1 & \text{if } -1 \leq y < x \leq 1 \\ 0 & \text{if } -1 \leq x \leq y \leq 1. \end{cases} \quad (5)$$

PROOF

The proof is done in two stages: first the $l = 0$ case is treated, and then the $l \neq 0$ case is analyzed. Both stages use the expansion of a function of two variables in an infinite series of Jacobi polynomials of one variable and coefficients that depend on the other variable. In particular, the coefficients in the expansion are obtained by representing the Jacobi polynomial as a finite series, evaluating the resulting elementary integrals, and re-summing the resultant series.

In proving Eq. 1, it is convenient to use the relationship

$$P_n^{(\alpha, -1)}(x) = \left(\frac{\alpha+n}{2n} \right) (1+x) P_{n-1}^{(\alpha,1)}(x) \quad \text{for } n \geq 1, \quad (6)$$

in order to write Eq. 2 as

$$F_{\mu+l, \mu-l} = \frac{1}{(2\mu+1)} P_l^{(\mu-l+1/2, -1)}(\Lambda) \\ + \frac{(1+\Lambda)^2}{8} \sum_{k=1}^{\infty} \frac{(2k+\mu+l+1/2)}{k(k+l)} P_{k-1}^{(\mu+l+1/2, 1)}(\Lambda) P_{k+l-1}^{(\mu-l+1/2, 1)}(\Lambda), \quad (7)$$

where the first term is the $k=0$ term of the sum appearing in Eq. 2.

CASE OF $l = 0$

For $l=0$, we expand

$$\begin{aligned} H^{(0)}(x,y) &\equiv \frac{2^{\mu+5/2}}{(1+x)(1+y)} \int_{-1}^w (1-t)^{-(\mu+3/2)} dt \\ &= \sum_{k=0}^{\infty} C_k^{(0)}(y) P_k^{(\mu+1/2,1)}(x), \end{aligned} \quad (8a)$$

where

$$w = \begin{cases} x & \text{if } y > x \\ y & \text{if } y \leq x. \end{cases} \quad (8b)$$

Multiplying Eq. 8a by $(1+x)(1-x)^{\mu+1/2} P_m^{(\mu+1/2,1)}(x)$ and integrating from $x = -1$ to 1 yields

$$\begin{aligned} &\frac{(m+1)2^{\mu+5/2}C_m^{(0)}(y)}{(m+\mu+3/2)(2m+\mu+5/2)} = \frac{2^{\mu+5/2}}{(1+y)} \\ &\times \left\{ \int_{-1}^y dx(1-x)^{\mu+1/2} P_m^{(\mu+1/2,1)}(x) \int_{-1}^x dt(1-t)^{-(\mu+3/2)} \right. \\ &+ \left[\int_{-1}^1 dx(1-x)^{\mu+1/2} P_m^{(\mu+1/2,1)}(x) \right] \left[\int_{-1}^y (1-t)^{-(\mu+3/2)} dt \right] \Big\} \\ &= (1+y)^{-1}(\mu+1/2)^{-1} \left[-4 \int_{-1}^1 dx(1-x)^{\mu+1/2} P_m^{(\mu+1/2,1)}(x) \right. \\ &\quad + 2^{\mu+5/2} \int_{-1}^y dx P_m^{(\mu+1/2,1)}(x) \\ &\quad \left. + 2^{\mu+5/2}(1-y)^{-(\mu+1/2)} \int_y^1 dx(1-x)^{\mu+1/2} P_m^{(\mu+1/2,1)}(x) \right]. \end{aligned} \quad (9)$$

The integrals are readily evaluated by expanding the Jacobi polynomial as¹

$$P_m^{(\alpha,\beta)}(x) = \sum_{r=0}^m \gamma(m,\alpha,\beta;r) (1-x)^r, \quad (10a)$$

with

$$\gamma(m, \alpha, \beta; r) \equiv \Gamma(\alpha + m + 1) \Gamma(\alpha + \beta + m + r + 1) (-1/2)^r / [r! (m - r)! \Gamma(\alpha + r + 1) \Gamma(\alpha + \beta + m + 1)]; \quad (10b)$$

integrating each term; and re-summing the series. In particular,

$$\begin{aligned} & \frac{(m+1)2^{\mu+5/2}}{(m+\mu+3/2)(2m+\mu+5/2)} C_m^{(0)}(y) = (1+y)^{-1} (\mu+1/2)^{-1} \\ & \times \sum_{r=0}^m \gamma(m, \mu+1/2, 1; r) \left\{ \frac{-2^{r+\mu+7/2}}{(r+\mu+3/2)} - \frac{2^{\mu+5/2}}{(r+1)} [(1-y)^{r+1} - 2^{r+1}] \right. \\ & \left. + 2^{\mu+5/2} \frac{(1-y)^{r+1}}{(r+\mu+3/2)} \right\} = (1+y)^{-1} (\mu+1/2)^{-1} \sum_{r=0}^m \gamma(m, \mu+1/2, 1; r) \\ & \times \left[2^{r+\mu+7/2} - 2^{\mu+5/2} (1-y)^{r+1} \right] \frac{(\mu+1/2)}{(r+1)(r+\mu+3/2)} \\ & = -2^{\mu+7/2} (1+y)^{-1} (m+\mu+3/2)^{-2} \left[P_{m+1}^{(\mu+1/2, -1)}(-1) - P_{m+1}^{(\mu+1/2, -1)}(y) \right] \\ & = \frac{2^{\mu+5/2}}{(m+1)(m+\mu+3/2)} P_m^{(\mu+1/2, 1)}(y), \quad (11) \end{aligned}$$

where the fact that

$$\gamma(m+1, \mu+1/2, -1; r+1) =$$

$$-(m+\mu+3/2)^2 \gamma(m, \mu+1/2, 1; r) / [2(r+\mu+3/2)(r+1)]$$

is used to arrive at the next to the last line, and Eq. 6 is used in the final step. Inserting this result in Eq. 8a yields

$$H^{(0)}(x,y) = \frac{2^{\mu+5/2}}{(1+y)(1+x)} \left[\frac{(1-w)^{-(\mu+1/2)} - 2^{-(\mu+1/2)}}{(\mu+1/2)} \right]$$

$$= \sum_{k=0}^{\infty} \frac{(2k+\mu+5/2)}{(k+1)^2} P_k^{(\mu+1/2,1)}(y) P_k^{(\mu+1/2,1)}(x) \quad (12a)$$

or

$$\frac{1}{(2\mu+1)} \left(\frac{2}{1-w} \right)^{\mu+1/2} = \frac{1}{(2\mu+1)} + \frac{(1+y)(1+x)}{8}$$

$$\times \sum_{k=1}^{\infty} \frac{(2k+\mu+1/2)}{k^2} P_{k-1}^{(\mu+1/2,1)}(y) P_{k-1}^{(\mu+1/2,1)}(x), \quad (12b)$$

with w defined by Eq. 8b. Since $P_0^{(\alpha,\beta)}(\Lambda) = 1$, comparing Eq. 7 with $l = 0$ to Eq. 12b with $x=y=\Lambda$ yields

$$F_{\mu,\mu} = \frac{1}{(2\mu+1)} \left(\frac{2}{1-\Lambda} \right)^{\mu+1/2}, \quad (13)$$

which demonstrates that Eq. 1 holds for $l = 0$.

CASE $l \neq 0$

For $l \neq 0$, we consider

$$H^{(l)}(x,y) \equiv \begin{cases} 0 & \text{for } 1 \geq y \geq x \geq -1, \\ \frac{\Gamma(\mu+5/2)}{\Gamma(\mu-l+5/2)l!} \frac{\partial^{l-1}}{\partial y^{l-1}} [(1-y)^{-(\mu+3/2)}(x-y)^l] & \text{for } 1 \geq x > y \geq -1. \end{cases} \quad (14)$$

Application of Leibnitz's rule for differentiating a product yields

$$H^{(l)}(x,y) = \sum_{i=0}^{l-1} S_i \frac{(x-y)^{i+1}}{(1+x)} U(x,y), \quad (15)$$

$$\text{with } S_l \equiv \frac{(\mu+3/2)\Gamma(\mu+l+3/2)\Gamma(l)(-1)^{l-t-1}(1-y)^{-(\mu+3/2+t)}}{\Gamma(\mu-l+5/2)\Gamma(t+2)\Gamma(t+1)\Gamma(l-t)},$$

and $U(x,y) = 1$ if $1 > x > y \geq -1$, and $U(x,y) = 0$ otherwise. The function $H^{(l)}(x,y)$ is expanded in terms of Jacobi polynomials

$$H^{(l)}(x,y) = \sum_{k=0}^{\infty} C_k^{(l)}(y) P_k^{(\mu-l+1/2,1)}(x), \quad (16)$$

with the coefficients determined from

$$\begin{aligned} I(m,l) &\equiv \frac{(m+1)2^{\mu-l+5/2}}{(m+\mu-l+3/2)(2m+\mu-l+5/2)} C_m^{(l)}(y) \\ &= \int_{-1}^1 H^{(l)}(x,y) (1-x)^{\mu-l+1/2} (1+x) P_m^{(\mu-l+1/2,1)}(x) dx \\ &= \sum_{t=0}^{l-1} S_t \sum_{r=0}^m \gamma(m, \mu-l+1/2, 1; r) \int_y^1 dx (1-x)^{\mu-l+1/2+r} (x-y)^{t+1}. \end{aligned} \quad (17)$$

The integral is evaluated by changing the variable from x to $z \equiv (x-y)/(1-y)$, which leads to

$$\begin{aligned} I(m,l) &= \sum_{r=0}^m \gamma(m, \mu-l+1/2, 1; r) (1-y)^{\mu-l+1/2+r+2} \Gamma(\mu-l+r+3/2) \\ &\quad \times \sum_{t=0}^{l-1} \left[\frac{\Gamma(t+2) S_t (1-y)^t}{\Gamma(\mu-l+r+t+7/2)} \right]. \end{aligned} \quad (18)$$

Substituting for S_t , the inner summation appearing in Eq. 18 reduces to

$$\begin{aligned} \sigma(r,l) &= \frac{(\mu+3/2)(-1)^{l-1}}{\Gamma(\mu-l+5/2)} \sum_{t=0}^{l-1} \left\{ \frac{\Gamma(\mu+3/2+t)}{\Gamma[\mu+3/2+t-(l-2-r)]} \right\} \\ &\quad \times \left[\frac{(l-1)!}{t!(l-1-t)!} \right] (-1)^t. \end{aligned} \quad (19)$$

For $l \geq r+2$, we note that

$$\sigma(r, l) = \frac{(\mu + 3/2)(-1)^{r+1}}{\Gamma(\mu - l + 5/2)} \frac{\partial^{l-2-r} [z^{\mu+1/2} (1-z)^{l-1}]}{\partial z^{l-2-r}} \Big|_{z=1} = 0 \text{ for } r \geq 0. \quad (20)$$

The derivative is zero at $z = 1$ because $(l-2-r) < l-1$. The fact that the derivative reproduces the series in Eq. 19 can be readily verified by writing the factor $(1-z)^{l-1}$ as a polynomial in z , differentiating each term, and then setting $z=1$. For $l < r+2$, we note that

$$\begin{aligned} \sigma(r, l) &= \frac{(-1)^{l-1}(\mu + 3/2)}{\Gamma(\mu - l + 5/2)} \int_0^1 ds_1 \int_0^{s_1} ds_2 \dots \int_0^{s_{j-1}} ds_j s_j^{\mu+1/2} (1-s_j)^{l-1} \Big|_{j=r+2-l} \\ &= \frac{(-1)^{l-1} \Gamma(\mu + 5/2) \Gamma(r+1)}{\Gamma(\mu - l + 5/2) \Gamma(r+2-l) \Gamma(\mu + 5/2 + r)}. \end{aligned} \quad (21)$$

The multiple integral in Eq. 21 is readily evaluated by recalling that integration by parts leads to

$$\int_0^1 ds_1 \int_0^{s_1} ds_2 \dots \int_0^{s_{j-1}} ds_j f(s_j) = \int_0^1 \frac{(1-s_1)^{j-1}}{(j-1)!} f(s_1) ds_1. \quad (22)$$

The fact that the multiple integral reproduces the series in Eq. 19 can be readily verified by writing $(1-s_j)^{l-1}$ as a polynomial in s_j and integrating each term separately. Inserting Eqs. 20 and 21 into Eq. 18 yields

$$I(m, l) = \begin{cases} \frac{(-1)^{l-1} \Gamma(\mu + 5/2)}{\Gamma(\mu - l + 5/2)} \sum_{r=l-1}^m \gamma(m, \mu - l + 1/2, 1; r) (1-y)^{r-l+1} \\ \times \frac{\Gamma(\mu - l + r + 3/2) r!}{\Gamma(\mu + 5/2 + r) (r+1-l)!} & \text{for } m \geq l-1 \\ 0 & \text{otherwise.} \end{cases} \quad (23)$$

For $m \geq l-1$, using the definition of $\gamma(m, \mu - l + 1/2, 1; r)$ and introducing the indices $r' = r - (l-1)$ and $m' = m - (l-1)$ leads to

$$\begin{aligned}
 I(m, l) &= \frac{\Gamma(\mu + 5/2) (\frac{1}{2})^{l-1}}{(m' + \mu + \frac{1}{2}) \Gamma(\mu + 5/2 - l)} \\
 &\times \sum_{r'=0}^{m'} \frac{[(y-1)/2]^{r'} \Gamma(m' + \mu + l + \frac{1}{2} + r')}{(m' - r')! r'! \Gamma(\mu + l + 3/2 + r')} \\
 &= \frac{\Gamma(\mu + 5/2) (\frac{1}{2})^{l-1}}{(m + \mu + 3/2) (m + \mu - l + 3/2) \Gamma(\mu - l + 5/2)} P_{m-(l-1)}^{(\mu+l+\frac{1}{2}, -1)}(y), \quad (24)
 \end{aligned}$$

where the definition of $\gamma(m-l+1, \mu+l+\frac{1}{2}, -1; r)$ was employed to obtain the last line. Inserting Eqs. 23 and 24 into Eqs. 17 and 16 yields

$$\begin{aligned}
 H^{(l)}(x, y) &= \frac{2^{-(\mu+3/2)} \Gamma(\mu + 5/2)}{\Gamma(\mu + 5/2 - l)} \sum_{m=l-1}^{\infty} \frac{(2m + \mu - l + 5/2)}{(m+1)(m + \mu + 3/2)} \\
 &\times P_{m-(l-1)}^{(\mu+\frac{1}{2}+l, -1)}(y) P_m^{(\mu-l+\frac{1}{2}, 1)}(x). \quad (25)
 \end{aligned}$$

Changing the summation index to $m' = m - (l-1)$, writing the $m' = 0$ term separately, and using Eq. 6 in both the $m' = 0$ and the $m' \neq 0$ terms lead to

$$\begin{aligned}
 H^{(l)}(x, y) &= \frac{2^{-(\mu-\frac{1}{2})} \Gamma(\mu + 5/2)}{(1+x) \Gamma(\mu + 5/2 - l)} \left[\frac{P_l^{(\mu-l+\frac{1}{2}, -1)}(x)}{(2\mu+1)} + \frac{(1+x)(1+y)}{8} \right. \\
 &\times \sum_{m'=1}^{\infty} \frac{(2m' + l + \mu + \frac{1}{2})}{m' (m' + l)} P_{m'-1}^{(\mu+l+\frac{1}{2}, 1)}(y) P_{m'+l-1}^{(\mu-l+\frac{1}{2}, 1)}(x) \left. \right]. \quad (26)
 \end{aligned}$$

Setting $x=y=\Lambda$ in Eq. 26 and comparing with Eq. 7, one finds

$$H^{(l)}(\Lambda, \Lambda) = \frac{2^{-(\mu-\frac{1}{2})} \Gamma(\mu + 5/2)}{(1+\Lambda) \Gamma(\mu + 5/2 - l)} F_{\mu+l, \mu-l} = 0, \quad l \neq 0, \quad (27)$$

where the vanishing of $H^{(l)}(\Lambda, \Lambda)$ follows directly from the form of Eq. 15. This completes the proof of Eq. 1.

REFERENCES

- ¹W. Magnus, F. Oberhettinger, and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer-Verlag, New York (1966).
- ²I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series, and Products*, Alan Jeffrey, ed., Academic Press, New York (1965).
- ³H. L. Manocha and B. L. Sharma, "Some Formulae for Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 459-462 (1966).
- ⁴H. L. Manocha and B. L. Sharma, "Generating Functions of Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 431-433 (1966).
- ⁵B. Noble, "Some Dual Series Equations Involving Jacobi Polynomials," in *Proc. Camb. Phil. Soc.*, pp. 363-371 (1963).
- ⁶E. R. Hansen, *A Table of Series and Products*, Prentice-Hall, New York (1975).

INITIAL DISTRIBUTION EXTERNAL TO THE APPLIED PHYSICS LABORATORY*

The work reported in TG 1350 was done under Navy Contract N00024-85-C-5301. This work is related to Task X831 which is supported by Naval Sea Systems Command.

ORGANIZATION	LOCATION	ATTENTION	No. of Copies
DEPARTMENT OF DEFENSE			
DTIC	Alexandria, VA		12
DEPARTMENT OF THE NAVY			
NAVSEA/SSCOM	Washington, DC	SEA 9961	2
NAVAIR/SSCOM	Washington, DC	AIR 7226	2
NAVPAC	Laurel, MD		1
ONR	Arlington, VA	Code 412	1
CONTRACTOR			
Lockheed Palo Alto Research Lab	Palo Alto, CA	E. R. Hansen	1
UNIVERSITIES			
Catholic Univ. of America, Dept. of Physics	Washington, DC	P. Meier	1
Royal College of Science and Technology, Dept. of Mathematics	Glasgow, Scotland	B. Noble	1
The Johns Hopkins Univ., Dept. of Earth and Planetary Science	Baltimore, MD	O. M. Phillips	1
		P. L. Olson	1
		R. R. Long	1

Requests for copies of this report from DoD activities and contractors should be directed to DTIC, Cameron Station, Alexandria, Virginia 22314 using DTIC Form 1 and, if necessary, DTIC Form 55.

*Initial distribution of this document within the Applied Physics Laboratory has been made in accordance with a list on file in the APL Technical Publications Group.

END

FILMED

12-85

DTIC