

45 50 32 32 34 36 40 40 1.0 2.5 2.0 1.8 1.000 1.25 1.6 ż

المريحة ا

ALLEY STRATES COLUMN STRATES AND STRATES

1

10" . 14" . Rat . 14" . 1

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

The Art and the transments

<u>د ا</u>

The Johns Hopkins University

l Z

Solarian on the Might rate Discourt, partian for Parabally Speeda

0

്





Solution to the Algebraic Riccati Equation for Parabolic Systems

Laurence R. Riddle and Howard L. Weinert Report JHU/EE-86/21



This work supported by the Office of Naval Research under Contract N00014-85-K-0255.



Solution to the Algebraic Riccati Equation for Parabolic Systems

Laurence R. Riddle and Howard L. Weinert

Department of Electrical Engineering

The Johns Hopkins University

Baltimore, MD 21218

ABSTRACT

This paper presents an analytical solution to the operator algebraic Riccati equation (ARE) for selfadjoint parabolic systems. The solution to the operator ARE is important in the design of the steady-state, on-line filter for estimating the system's states. This analytical solution is derived by considering the operator analog of Potter's method of using the Hamiltonian system's eigenvectors and eigenvalues to solve a finite-dimensional ARE. As an example of using this analytical solution, the steady-state filtering error covariance for the 2-D heat equation is studied.

This work supported by the Office of Naval Research under Contract N00014-85-K-0255.



Diet.

1. Introduction

In this paper, we derive an analytical solution to the operator algebraic Riccati equation (ARE) on a Hilbert space H:

$$AP + PA' + Q = PRP \tag{1.1}$$

when -A is a strongly positive (coercive) selfadjoint operator that generates a continuous semigroup, Q is positive-definite, bounded and commutes with A, and R is bounded and nonnegative-definite. We also assume that the domain of A^2 is dense in H, and is contained in the domain of A, and that the positive square root of Q commutes with A. In the context of distributed parameter filtering and control, one needs a solution to Eq. (1.1) that is bounded, nonnegative, selfadjoint and that maps the domain of A into itself. Gibson [3] has shown that under the above assumptions, Eq. (1.1) has a unique solution with these properties.

An example where one would solve Eq.(1.1) with the above assumptions on A, Q and R is the design of a steady-state filter to estimate a process governed by the heat equation:

$$\frac{\partial}{\partial t}u(\mathbf{x},t) = \nabla^2 u(\mathbf{x},t) + \epsilon(\mathbf{x},t) , \mathbf{x} \in \Omega \subset \mathbf{R}^3, t \ge 0$$

$$\frac{\partial}{\partial n}u(\mathbf{x},t) = 0, \mathbf{x} \in \partial\Omega$$
(1.2)

where $\frac{\partial}{\partial n}$ denotes the outward normal derivative, and the observations are

$$y_j(t) = \int C_j(\mathbf{x})u(\mathbf{x},t)d\mathbf{x} + w_j(t) , \ j = 1, \cdots, N$$

In Eq (1.2), $\epsilon(\mathbf{x},t)$ is a random process that is white in space and time with

constant intensity Q, $w_j(t)$ is a scalar white noise process with intensity R^{-1} , and $A = \nabla^2$. Several applications using this model to describe the temperature distribution of heated metals have been reported [1],[9]. Numerical examples of calculating the steady-state filtering error covariance P for this model are given in [11].

2. Solving the ARE

In this section we will solve Eq. (1.1) by using the operator analog to the Potter method [10] of solving the ARE in finite dimensions. The Potter method of solving the ARE is summarized as follows: let the Hamiltonian matrix L be defined as

$$L = \begin{bmatrix} A & Q \\ R & -A \end{bmatrix}$$

and diagonalize L via

$$LM = M\Lambda$$

where

$$\Lambda = diag(\lambda_1, \cdots, \lambda_{2n}), \lambda_1, \cdots, \lambda_n > 0$$

and

$$M = \begin{bmatrix} M_{11} & M_{12} \\ \\ M_{21} & M_{22} \end{bmatrix}$$

Then

$$P = M_{11} M_{21}^{-1}$$

is a nonnegative symmetric solution to the ARE , assuming $M_{\rm 21}$ is invertible.

Using this approach in an operator setting, we will consider first a formal solution to the operator ARE. Let Q = I and consider the following operator equation:

$$\begin{bmatrix} A & I \\ R & -A \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} = \begin{bmatrix} m_1 \\ m_2 \end{bmatrix}$$
(2.1)

where m_{1}, m_{2} and λ are operators. It is easy to verify that the choices

$$m_1 = I$$
, $m_2 = -A + (A^2 + R)^{4}$

$$\lambda = (A^2 + R)^{\frac{1}{2}}$$

satisfy Eq. (2.1) and thus

$$P_{1} = (-A + (A^{2} + R)^{\frac{1}{2}})^{-1}$$
(2.2)

satisfies Eq. (1.1) in a formal sense. If $Q \neq I$ then

$$P = S(-A + (A^{2} + R)^{\frac{1}{2}})^{-1}S$$
(2.3)

satisfies Eq. (1.1), where $S^2 = Q$ and S is positive-definite.

To justify the formal solution given by Eq. (2.3) we need to show that P is bounded, nonnegative, selfadjoint, and maps the domain of A into itself. Since -A > 0 it follows that A^2+R has a unique positive square root that is selfadjoint and that the domain of A^2 is dense in the domain of $(A^2+R)^{\frac{1}{2}}$ [7, p. 281]. Therefore $-A + (A^2+R)^{\frac{1}{2}}$ is strongly positive and selfadjoint with dense domain:

 $D(-A + (A^2 + R)^{\frac{1}{2}}) = D(-A) \cap D((A^2 + R)^{\frac{1}{2}}) \subset D(A^2) \subset D(A)$ (2.4) where $D(\cdot)$ denotes domain; so that P_1 as defined in Eq. (2.2) is bounded [4, p. 209]. Furthermore, P_1 is selfadjoint and nonnegative because $(-A + (A^2 + R)^{\frac{1}{2}})$ is [7, p. 272]. We will prove that P_1 maps the domain of A into the domain of The range of P_1 is contained in the domain of A by Eq. (2.4), and so it is sufficient to show that the domain of P_1 contains the domain of A. Since P_1 is bounded and necessarily closed (being selfadjoint), it follows [7, p. 269] that the domain of P_1 is H, and hence contains the domain of A.

- 5 -

Since we have assumed that the operator S commutes with A, if $x \in D(A)$ then $Sx \in D(A)$, hence P in Eq. (2.3) also maps D(A) into D(A). We have thus shown that the operator defined in Eq. (2.3) is the correct solution to Eq. (1.1) needed for filtering and control applications.

3. Numerical Considerations and Example

The analytical approach to solving the operator ARE presented in this chapter provides a computationally faster way of implementing the optimal gain for filtering and control applications. In our approach, finite-dimensional approximations to the operators in Eq (1.1) are done after the analytical solution has been obtained, resulting in a computational complexity of $O(6n^3)$, where *n* is the size of the matrix that approximates the operator *A*, whereas in other techniques [2] one approximates Eq (1.1) by a matrix ARE and then solves this equation using algorithms developed for the finite-dimensional case which require approximately $O(75n^3)$ operations [8]. We remark that a similar approach (with the same computational complexity $O(6n^3)$) to the finite-dimensional ARE has been considered [5],[6].

Another important difference between the analytical approach used in this chapter and those using a high order finite-dimensional approximation to Eq.

(1.1), is that the implementation of Eq. (2.3) calls for approximations of A and A^2 separately, whereas in approximating Eq. (1.1) one is implicitly using \overline{A}^2 , where \overline{A} is a finite-dimensional approximation to A, as an approximation to A^2 . The fact that \overline{A}^2 may not be a good approximation to A^2 suggests that our approach may be more accurate.

The steady-state filtering error covariance for estimating the temperature profile of a heated square aluminum slab was calculated using the results obtained in the previous section. These calculations were based on the following model for the variation in temperature u(x,y,t) above an assumed known ambient temperature T_0 :

$$\frac{\partial}{\partial t}u(x,y,t) = \alpha \nabla^2 u(x,y,t) + \epsilon(x,y,t) \quad 0 \le x , y \le L$$

with boundary conditions

$$u(0,y,t) = u(L,y,t) = u(x,0,t) = u(x,L,t) = 0$$

where

ウバイ ファイト・キャー しょうけい ビング イオ・エット ノイト・

$$L = 1meter$$
, $\alpha = 5 \times 10^{-5}meter^2/second$

and the observations are

$$y(t) = u(.5,.5,t) + w(t)$$

The input $\epsilon(x,y,t)$ is assumed to be white in space and time with intensity

$$Q = 1(degreeC)^2 / (meter^2)$$
 second

and the observation noise w(t) is white with intensity

$$R^{-1} = 10^{-2} (degreeC)^2 / second$$

In Eq. (2.3), a 10×10 finite-difference approximation to the operators A and

 A^2 was used. Figure 1 shows a cross-section of the variances of the temperature variations u(x,.5,t) and the filtering error $\tilde{u}(x,.5,t)$. It is evident from Figure 1 that the unobserved temperatures more that 10 cm away from the sensor are not being estimated very effectively. This result places some doubt on the possibility of estimating, with few sensors, the unobserved temperature distribution of heated metals in the presence of spatially white process noise.

References

- M. B. Ajinkya, M. Köhne, M. F. Mäder, and W. H. Ray, 'The experimental implementation of a distributed parameter filter,' *Automatica*, vol 11, pp. 571-577, 1975.
- H.T. Banks and K. Kunisch, 'The linear regulator problem for parabolic systems,' SIAM J. Contr. Opt., vol. 22, pp 684-695, 1984.
- J.S. Gibson, 'The Riccati integral equations for optimal control problems on Hilbert spaces,' SIAM J. Contr. Opt. vol. 17, pp 537-565, 1979.
- [4] K. E. Gustafson, Introduction to partial differential equations and Hilbert space methods, Wiley, New York, 1980.
- [5] M. Hamidi, 'A method of optimal actuator and sensor placement', Proc. Fourth IFAC Symp. Contr. Dist. Parm. Syst., Los Angeles, 1986.
- [6] F. Incertis, 'An extension on a new formulation of the algebraic Riccati equation problem', *IEEE Trans. Auto. Contr.*, vol AC-28, pp. 235-237, 1983.

- [7] T. Kato, Perturbation theory for linear operators, Springer-Verlag, New York, 1966.
- [8] A.J. Laub, 'A Schur method for solving algebraic Riccati equations,' IEEE Trans. Auto. Contr., vol. AC-24, pp. 913-922,1979.

- [9] G. K. Lausterer, W. H. Ray and H. R. Martens, 'The real time application of distributed parameter state estimation theory to a two dimensional heated ingot,' pp. 451-467 of Proc. Second IFAC Symp. Contr. Dist. Parm. Syst., S. P. Banks, A. J. Pritchard Eds., Peragamon Press, Oxford, 1978.
- [10] J.E. Potter, 'Matrix quadratic solutions,' SIAM J. Appl. Math., vol. 14, pp 496-501, 1966.
- [11] K. Watanabe, M. Iwasaki, 'A fast computational approach in optimal distributed parameter state estimation,' J. Dynamical Systems, Measurement and Contr., vol. 105, pp. 1-10, 1983.



	REPORT DOCUM	ENTATION PAG	E		
REPORT SECURITY CLASSIFICATION		16. RESTRICTIVE N	ARKINGS		
UNCLASSIFIED		3 DISTRIBUTION/AVAILABULITY OF REPORT			
28. SECURITY CLASSIFICATION AND HORITY					
20. DECLASSIFICATION/DOWNGRADING SCHEDULE		UNLIMITE	D		
PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
1HU/EE - 86/21					
NAME OF PERFORMING ORGANIZATION 66. OFFICE SYMBOL		78. NAME OF MONITORING ORGANIZATION			
HE JOHNS HOPKINS UNIVERSITY		OFFICE OF NAVAL RESEARCH			
c. ADDRESS (City, State and ZIP Code)		7b. ADDRESS (City, State and ZIP Code)			
CHARLES AND 34TH STREETS		800 N. QUINCY ST.			
BALTIMORE, MARYLAND 21218		ARLINGTON,	VA 22217		
Be. NAME OF FUNDING/SPONSORING ORGANIZATION (11 applicable)		9. PROCUREMENT	NSTRUMENT IC	ENTIFICATION N	JMBER
		N00014-85-K-0255			
c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NOS.			
		PROGRAM ÉLEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT
		-	_		
TITLE (Include Security Classification) SOL II	TION TO THE ALG	EBRAIC		NK661-019	
TCCATT FOUNTION FOR DADADOLT	<u> CVCTENC (INCT</u>				
ICCATI EQUATION FOR PARABOLI	C SYSTEMS (UNCLA	ASIFIED)	l	<u></u>	
RICCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS) RIDDLE, L.R. AND WEINERT, H.L	C SYSTEMS (UNCLA	ABSIFIED)	1		
RICCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS RIDDLE, L.R. AND WEINERT, H.L IDDLE, L.R. AND WEINERT, H.L NTERIM INTERIM SUPPLEMENTARY NOTATION	C SYSTEMS (UNCLA COVERED /1/85 to 10/17/8	14. DATE OF REPOR	ат (Yr. Mo., Day , 1986	15. PAGE C 9	OUNT
ICCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS IDDLE, L.R. AND WEINERT, H.L TYPE OF REPORT NTERIM COSATI CODES SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB.GR. ASSTRACT (Continue on reverse if necessary on This paper presents an anal ARE) for selfadjoint parabol: n the design of the steady-st his analytical solution is defined the design of the steady-st	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 0PERATOR RICO PARAMETER FIL d dentify by block number systems. The cate, on-line fi erived by consider	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON Tr, h to the operation solution to the left for estimated	, 1986 PARABOLIC NTROL. tor algebra he operator mating the rator analo	n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter	ount STRIBUTED equation ortant ates. 's method
ICCATI EQUATION FOR PARABOLI .PERSONAL AUTHORIS IDDLE, L.R. AND WEINERT, H.L TYPE OF REPORT NTERIM SUPPLEMENTARY NOTATION COSATI CODES SUPPLEMENTARY NOTATION COSATI CODES SUPPLEMENTARY NOTATION ABSTRACT (Continue on reverse if necessary and the design of the steady-steady and the design of the steady-steady steady steady and the design of the steady steady steady steady and the design of the steady steady steady steady and the design of the steady stead	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FIL d identify by block number ic systems. The cate, on-line fil erived by consider this analytical	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TO to the operation to the liter for estimates of the operation is and eigenval	AT (Yr. Mo. Day , 1986 PARABOLIC NTROL. tor algebra he operator mating the rator analo lues to so	15. PAGE C 9 h/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite	ount STRIBUTED equation ortant ates. 's method ≻dimension
AGSTRACT (Continue on reverse of necessary on This paper presents an anal ARE) for selfadjoint parabol: n the design of the steady-st bis analytical solution is defined and the steady st his an example of using ovariance for the 2-D heat exponential of the steady states of the stea	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (C OPERATOR RICC PARAMETER FII d dentify by block number systems. The cate, on-line fi erived by consider em's eigenvector of this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON Try to the operation solution to the let for estimate is and eigenvalue of solution, the colution, the	AT (Yr. Mo. Day , 1986 PARABOLIC NTROL. tor algebra he operator mating the rator analo lues to so he steady-s	n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	ounT , STRIBUTED equation ortant ates. 's method ing error
ABSTRACT (Continue on reverse if necessary and ARE) for selfadjoint parabol: n the design of the steady-st his analytical solution is defined and the Hamiltonian system ovariance for the 2-D heat economic PARABOLICATION FOR PARABOLI SUPLEMENTARY NOTATION	18. SUBJECT TERMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (C OPERATOR RICC PARAMETER FIL d identify by block number ic systems. The cate, on-line fi erived by consider is eigenvector this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TRAINE AND CONT I to the operation solution to the liter for estimates and eigenvalues and e	recuary and ident PARABOLIC NTROL. tor algebra he operator mating the rator analo lues to so he steady-s	15. PAGE C 9 n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	ount STRIBUTED equation ortant ates. 's method ⊢dimension ing error
ALEST ACT (Continue on reverse of necessary and ARE) for selfadjoint parabol: ARE) for selfadjoint parabol: ARE AS an example of using tovariance for the 2-D heat eco	18. SUBJECT TERMS (UNCLA OVERED (1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FII d dentify by block number bytical solution ic systems. The cate, on-line fi erived by conside em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON in to the operation solution to the liter for estimates and eigenvalues and	AT (Yr. Mo. Day , 1986 PARABOLIC PARABOLIC NTROL. tor algebra he operator mating the rator analo lues to so he steady-s	15. PAGE C 9 h/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	oun⊤ STRIBUTED equation ortant ates. 's method ⊢dimension ing error
ACCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS) RIDDLE, L.R. AND WEINERT, H.L. INTERIM COSATI CODES FIGLO GROUP SUB.GR. AGSTRACT (Continue on reverse () necessary on This paper presents an anal ARE) for selfadjoint parabol: In the design of the steady-st This analytical solution is defined and the steady-st this analytical solution is defined and the steady-st Cosation of the steady-st ARE. As an example of using tovariance for the 2-D heat economic State of the steady-state of the steady state of the steady-state of the steady state of the stea	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICO PARAMETER FIL d dentify by block number by tical solution ic systems. The cate, on-line file rived by consider em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON Tr, to the operation solution to the lering the operation is and eigenvalue is solution, the ed.	AT (Yr., Mo., Day , 1986 PARABOLIC NTROL. tor algebra tor algebra he operator mating the rator analo lues to so he steady-s	n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	ounT ortant ates. ≻dimension ing error
RICCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS RIDDLE, L.R. AND WEINERT, H.L ATYPE OF REPORT INTERIM COSATI CODES FIELD GROUP COSATI CODES FIELD GROUP COSATI CODES FIELD GROUP SUB.GR. ABSTRACT (Continue on reverse if necessary and This paper presents an anal ARE) for selfadjoint parabol: In the design of the steady-st This analytical solution is defined and the steady-st This analytical solution is defined and the steady start I ARE. As an example of using covariance for the 2-D heat eco	18. SUBJECT TERMS (UNCLA OVERED (1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FII d dentify by block number lytical solution ic systems. The cate, on-line fi erived by consider em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TO to the operation to the liter for estimates and eigenvalues and eigenvalu	AT (Yr. Mo. Day , 1986 PARABOLIC NTROL. tor algebra tor algebra he operator mating the rator analo lues to so he steady-s	15. PAGE C 9 h/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	oun⊤ STRIBUTED equation ortant ates. 's method ≻dimension ing error
AGETRICUTION/AVAILABILITY OF ABSTRACE	C SYSTEMS (UNCLA COVERED (1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FII d dentify by block number systems. The cate, on-line first erived by consider em's eigenvector this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TO to the operation solution to the lering the operation is and eigenvalue of solution, the ed.	AT (Yr., Mo., Day , 1986 PARABOLIC PARABOLIC NTROL. tor algebra tor algebra he operator nating the rator analo lues to so he steady-s	15. PAGE C 9 h/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	oun⊤ STRIBUTED equation ortant ates. 's method ≻dimension ing error
ABSTRACT (Continue on reverse if necessary and This paper presents an anal ARE) for selfadjoint parabol: n the design of the steady-si his analytical solution is defined this analytical solution is defined this analytical solution is defined the design of the steady-si his analytical solution is defined the design of the 2-D heat equilated the d	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FIL d identify by block number lytical solution ic systems. The cate, on-line fi erived by conside em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TRADICTION OF THE Solution to the liter for estimation of the liter for estimation of the liter for estimation of the liter for estimation of the liter for estimation of the liter for estimation of the liter for estimation of the liter for estimation of th	AT (Yr., Mo., Day , 1986 PARABOLIC PARABOLIC NTROL. tor algebra he operator nating the rator analo lues to so he steady-s	15. PAGE C 9 n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	ounT STRIBUTED equation ortant ates. 's method ≻dimension ing error
AGE TRACT CONTINUE ON REVERENCE IN THE AND WEINERT, H.L. COSATI CODES FIELD GROUP SUB. GR. AGESTRACT CONTINUE ON REVERENCE IN RECEIPANT ON AGESTRACT CONTINUE ON REVERENCE IN A REASE ON AGESTRACT CONTINUE ON REVERENCE IN A REASE ON AGESTRACT CONTINUE ON REVERENCE IN A REASE ON A REASE OF RESPONSIBLE INDIVIDUAL	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FII d identify by block number bytical solution ic systems. The cate, on-line fi erived by conside em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON Try to the operation solution to the let for estimate is and eigenvalue of solution, the ed.	AT (Yr., Mo., Day , 1986 PARABOLIC PARABOLIC NTROL. tor algebra tor algebra he operator nating the rator analo lues to so he steady-s	15. PAGE C 9 n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter	OUNT STRIBUTED equation ortant ates. 's method ing error ing error
ALCATI EQUATION FOR PARABOLI PERSONAL AUTHORIS RIDDLE, L.R. AND WEINERT, H.L DATYPE OF REPORT INTERIM COSATI CODES FIELD GROUP SUB GR COSATI CODES FIELD GROUP SUB GR C	C SYSTEMS (UNCLA COVERED /1/85 TO 10/17/8 18. SUBJECT TERMS (OPERATOR RICC PARAMETER FIL d identify by block number lytical solution ic systems. The cate, on-line fi erived by conside em's eigenvector g this analytical quation is studi	Continue on reverse if ne Continue on reverse if ne CATI EQUATION, TERING AND CON TR to the operation solution to the liter for estimate solution to the liter for estimate solution, the ed.	AT (Yr., Mo., Day , 1986 PARABOLIC PARABOLIC NTROL. tor algebra he operator nating the rator analo lues to so he steady-s	15. PAGE C 9 n/y by block number SYSTEMS, DI aic Riccati r ARE is imp system's st og of Potter lve a finite state filter ICATION 22c. OFFICE SYM	ounT STRIBUTED equation ortant ates. 's method ≻dimension ing error BOL

 Γ,

بالمرجع من

1.1

1

......

 \overline{x}

