



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

| | PORT DOCUME | | | | |
|--|---|---|--|---|--|
| AD-A 161 009 | | ID. RESTRICTIVE M | ARKINGS | | |
| | FIC | 1.0.57818117108/A | | | |
| | ECTE | APPROVED F | OR PUBLI | C RELEASE | |
| DECLASSIFICATION/DOWNERACE OF MEDUC | 2 4 1097 | DISTRIBUTI | ON UNLIM | ITED | |
| PERPORMING ORGANIZATION REPORTIUNSE | A(8) | MONITORING OR | GANIZATION R | EPORT NUMBER(S) | |
| | D 70 | A | FOSR - TD | - 87-0 | 619 |
| A NAME OF PERFORMING ORGANIZATION | If applicable | TE NAME OF MONIT | ORING ORGAN | ZATION | |
| ARIZONA STATE UNIVERSIT | | AFOSR | | | |
| C. ADDRESS (City. Siste and ZIP Cate) | | 76. ADORESS (City. S | State and ZIP Co | de i | |
| Tempe, Az 85287 | | Bidg 410 Bol⊥ing AF | B, DC 20 | 332-6448 | |
| NAME OF FUNDING/BRONSORING | OFFICE SYMEGI | | | ENTIFICATION MUS | |
| AFOSR | NE | AFOSR-84-0290 | | | |
| . ADORESE (City. State and EIP Code) | | 19. SOURCE OF FUNDING NOR | | | |
| Bldg 410 | | PROGRAM | PROJECT | TASK | WORK UN |
| Bolling AFB, DC 20332 | | ELEMENT NO. | NO. | NO. | NO. |
| 1. TITLE Include Security Classification The or at | ical Studied | 61102F | 2305 | C1 | |
| of Experiments and Applicati | ons of Su | DDIC osecond | Pho too | onau ctivity | |
| 2.PERSONALAUTHORUS) Dr. Robert O. Grondin | | | | | |
| SA TYPE OF REPORT 138. TIME COV | ERED | 14. DATE OF REPOP | T (Yr. Mo., De | I IS PAGE CO | |
| | | | | | •••• |
| Annual FROM 01Se | <u>p85 to31Aug8</u> | 87 | April | 10 | |
| Annual FROM <u>Q1Se</u> C. SUPPLEMENTARY NOTATION | <u>p85</u> ro <u>31Aug8</u> | 87 | April | 10 | |
| Annual FROM OISE | <u>p85</u> 7031Aug8 | 87 | April | 10 | |
| Annual FROM OISE C. SUPPLEMENTARY NOTATION 7. COBATI CODES 11 MELO GROUP SUB. GR. | D85 TO 31Aug8 | 87 | April | afy by block numbers | |
| Annual FROM Olse C. SUPPLEMENTARY NOTATION 7. COBATI CODES MELD GROUP SUB. GR. | D85 TO 31Aug8 | 87 | April | afy by block number: | |
| Annual PROM Olse COBATI CODES MELO GROUP SUB GR. MARETRACT (Continue on reverse if receivery and id | D85 TO 31Aug8 | 87 | April | aty by block number: | |
| Annual FROM OISE COBATI CODES THE CODES T | D85 TO 31Aug8 | 87 | April cemery and iden o year p | 10 Bry by block number, rogram whose | e goal |
| Annual FROM OISE COLATI CODES THE O GROUP SUB GR. MANDTRACT (Continue on reverse if recensory and id The reporting period covers the development and use of m as probes of hot carried tra | D85 TO 31Aug8 SUBJECT TERMS (Co In S/7 by Neck number the second y odels of fem psport in se | ear of a tw tosecond ph | April cemery and iden o year p otocondu | 10 ary by block number, rogram whose ctive experi- program bas | e goal iments |
| Annual FROM OISE COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENTARY COMPLEMENTARY NOTATION COMPLEMENTARY COM | p85 to 31Aug8 subject terms (contraction of fer isodels of fer isport in second to the second to | ear of a tw tosecond ph miconductor experiment | April o year p otocondu s. The s are be | 10 Rfy by block number, rogram whose ctive experi program has ing carried | e goal iments out |
| Annual FROM OISE COMATICODES COMATICODES MANY CODES MANY CODES | p85 to 31Aug8 Subject terms (Control of the second y odels of fer insport in second structures) ears. These ed by Dr. Ge | 87 Tear of a two tosecond ph miconductor experiment rald Mourou | April o year p otocondu s. The s are be at the | 10 ary by block number, rogram whose ctive experi program has ing carried University of | e goal iments out of |
| Annual FROM OISE COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENTARY COMPLEMENTARY NOTATION COMPLEMENTARY COMPLEMENTAR | p85 to 31Aug86 subject teams (Co budges of fer nsport in se ears. These ed by Dr. Ge k. There ar | ear of a tw tosecond ph miconductor experiment rald Mourou e several m | April o year p otocondu s. The s are be at the ain comp | 10 ary by block number, rogram whose ctive experi program has ing carried University consts to the eration of e | e goal iments out of |
| Annual FROM OISE COLATICODES COLATICODES MANDTRACT CONTINUE OF SUR GR. MANDTRACT CONTINUE | p85 to 31Aug86 SUBJECT TERMS (Co build by block number the second y odels of fem nsport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of | April o year p otocondu s. The s are be at the ain comp the gen the inci | nogram whose ctive experi program has ing carried University conets to the eration of e dence of a f | e goal iments out of electro femto- |
| Annual FROM <u>Olse</u> COMATICODES MARTRACT (Continue on rearro if recempt and in The reporting period covers the development and use of m as probes of hot carried tra been extended for two more y in a companion effort direct Rochester, Rochester New Yor modeling of such experiments hole pairs inside a semicond second optical pulse. Then | p85 to 31Aug8 subject terms (Control of the second y odels of fer insport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the one must mod | 87 Tear of a two tosecond ph miconductor experiment rald Mourou e several m first model result of tel the proc | April o year p otocondu s. The s are be at the ain comp the gen the inci esses by | 10 ary by block sumber, ctive experi program has ing carried University to onets to the eration of e dence of a f which the t | e goal iments out of electro femto- resulti |
| Annual PROM <u>Olse</u> COMATICODES MLD CROUP SUB GR MANDTRACT /Continue on rearry // recempt and use The reporting period covers the development and use of m as probes of hot carried tra been extended for two more y in a companion effort direct Rochester, Rochester New Yor modeling of such experiments hole pairs inside a semicond second optical pulse. Then current transient is develop | p85 to 31Aug86 subject terms ice the second y odels of fem nsport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the one must mod ed. Lastly, | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver | April o year p otocondu s. The s are be at the ain comp the gene the inci esses by sion of | 10 afy by block number, rogram whose ctive experi program has ing carried University con onets to the eration of e dence of a f which the r the current | e goal iments out of electro femto- resulti |
| Annual FROM <u>Olse</u> COLATI CODES MADDIALS COLATI CODES MADDIALS COLATI CODES MADDIALS COLATI CODES MADDIALS COLATI CODES MADDIALS COLATI CODES MADDIALS SUB GROUP SUB GROUP | p85 to 31Aug86 subject terms (ca buddy by block number the second y odels of fer nsport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the one must mod ed. Lastly, e that is di | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver | April o year p otocondu s. The s are be at the ain comp the inci esses by sion of ured in | afy by block number, rogram whose ctive experi program has ing carried University of onets to the eration of e dence of a f which the n the current the experime | e goal iments out of electro femto- resulti ents of |
| Annual PROM <u>Olse</u> COMATICODES MADDERACT (Continue on reverse if recemery and in the reporting period covers the development and use of m as probes of hot carried tra been extended for two more y in a companion effort direct Rochester, Rochester New Yor modeling of such experiments hole pairs inside a semicond second optical pulse. Then current transient is develop transient into a voltage wav interest. During the first | p85 to 31Aug8 subject terms (Co budgets of fer insport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the one must mod ed. Lastly, e that is di year the fir | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t | April o year p otocondu s. The s are be at the ain comp the gen the inci- esses by sion of ured in hese nec | nogram whose ctive experi program has ing carried University of onets to the eration of e dence of a f which the n the current the experime essary steps | e goal iments out of electro fesulti ents of swere |
| Annual PROM <u>Olse</u> COLATICODES MLS <u>GROUP</u> SUB GR MANDTRACT (Continue on rearry of access) and a Continue on rearry (I access) and a Continue on rearry (I access) and a Continue on rearry (I a | p85 ro31Aug8 SUBJECT TERMS (Co SUBJECT TERMS (Co | 87 manue on muere if ne tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t models wer | April o year p otocondu s. The s are be at the ain comp the inci- esses by sion of ured in hese nec- e refine pment of | nogram whose ctive experi- program has ing carried University of onets to the eration of e dence of a f which the n the current the current the experime essary steps d and used to an experime | e goal iments out of electro femto- results ents of s were to student whi |
| Annual CONTRACT CODES CONTRACT CODES Contract Codes Contract Codes Contract Codes Contract Codes Contract Codes Contract Contract on Norre 1/ Accessory and the Contract on Norre 1/ Accessory and the | p85 ro3lAug8 subject reams (Co build by black number the second y odels of fer nsport in se ears. These ed by Dr. Ge k. There ar . ()ne must uctor as the one must mod ed. Lastly, e that is di year the fir d year these e usedful in | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t models wer the develo | April o year p otocondu s. The s are be at the at the at the gene the inci- esses by sion of ured in hese nec- e refine pment of | rogram whose ctive experi program has ing carried University of onets to the eration of e dence of a f which the n the current the current the experime essary steps d and used f an experime | e goal iments out of electro cesult: ents of s were to stude ent what |
| Annual PROM <u>Olse</u> Supplementany Notation Magnetic Continue on Norre if Account on the Magnetic Continue on Norre if Account of Acco | DB5 TO 31Aug8 SUBJECT TERMS (Ca SUBJECT TERM | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t models wer the develo 21. AGSTRACT SEC UNCLA | April o year p otocondu s. The s are be at the ain comp the inci- esses by sion of ured in hese nec- e refine pment of SSIFIED | rogram whose ctive experi- program has ing carried University of onets to the eration of e dence of a f which the n the current the experime essary steps d and used to an experime | e goal iments out of electro femto- resulti ents of s were to stud |
| Annual PROM <u>Olse</u> COLATI CODES MADDIALS COLATI CODES MADDIALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIONALS COLATIO | D85 TO 31Aug8 SUBJECT TERMS (Ca SUBJECT TERM | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t models wer the develo 21. AGSTRACT SEC UNCLA | April o year p otocondu s. The s are be at the ain comp the inci- esses by sion of ured in hese nec- e refine pment of SSIFIED | ary by North Aumber, rogram whose ctive experi program has ing carried University of onets to the eration of e dence of a f which the n the current the experime essary steps d and used to an experime | e goal iments out of electro femto- resulti ents of s were to stud ent whi |
| Annual PROM <u>Olse</u> COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENTARY NOTATION COMPLEMENT ANY NOTATION COMPLEMENT ODES COMPLEMENT ODES COMPLEMENT ODES COMPLEMENT OF SUBJECT ON COMPLEMENT OF ADDATES COMPLEMENT OF SUCH EXPERIMENTS COMPLEMENT OF SUCH EXPERIMENTS COMPLETED COMPLEMENT OF ADDATES COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETED COMPLETE | D85 TO 31Aug8 SUBJECT TERMS (Construction the second y odels of fem nsport in second ears. These ed by Dr. Geck. There ar . ()ne must uctor as the one must mod ed. Lastly, e that is di year the fir d year these e usedful in D71C USERS | ear of a tw tosecond ph miconductor experiment rald Mourou e several m first model result of el the proc the conver rectly meas st two of t models wer the develo 21. AGGTRACT SEC UNCLA | April o year p otocondu s. The s are be at the ain comp the gen the inci esses by sion of ured in hese nece pment of UNITY CLASSI SSIFIED | rogram whose ctive experi program has ing carried University of onets to the eration of e dence of a f which the n the current the current the experime essary steps d and used to an experime PICATION | e goal iments out of electro femto- resulti ents of swere to stud ent whi |

AFOSR-TR- 87-0619

Theoretical Studies of Experiments and Applications of Subpicosecond Photoconductivity

Annual Technical Report September 1, 1985 to August 31, 1986

Award Number AFOSR-84-0290 A

Submitted to Dr. Gerald L. Witt Air Force Office of Scientific Research Air Force Systems Command, USAF Bolling AFB, DC 20332

Submitted by Dr. Robert O. Grondin Center for Solid State Electronics Research Arizona State University Tempe, AZ 85287

> Approved for public release; distribution unlimited.



Ø

D

Accession For

MTIS GRAMI

Unannounged

Justification

tritution/

Availability Codes

Aveil and/or Special

DTIC TAB

87

pist

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC) NOTICE OF TRANSMITTAL TO DTIC This technical report has been reviewed and is approved for public release IAW AFR 190-12. Disclosulation se unimited. MATTEL was KENPER Chief, Technical Information Division

Summary Abstract

The reporting period covers the second year of a two year program whose goal is the development and use of models of femtosecond photoconductive experiments as probes of hot carrier transport in semiconductors. The program has been extended for two more years. These experiments are being carried out in a companion effort directed by Dr. Gerard Mourou at the University of Rochester, Rochester New York. There are several main components to the modeling of such experiments. One must first model the generation of electron-hole pairs inside a semiconductor as the result of the incidence of a femtosecond optical pulse. Then one must model the processes by which the resulting current transient is developed. Lastly, the conversion of the current transient into a voltage wave transmitted down a transmission line must be understood. It is this voltage wave that is directly measured in the experiments of interest. During the first year the first two of these necessary steps were completed. During the second year these models were refined and used to study some assumptions which may be useful in the development of an experiment which can be quantitatively analyzed.

Statement of Work

For approximately 15 years there has been interest in the possibility of constructing transistors with extremely fast switching properties by using novel features associated with charge transport in semiconductors over distances of submicron length. While there has been a large effort at theoretically analyzing these possibilities, the time and space scales are sufficiently short that meaningful experimental efforts have been precluded. Subpicosecond optical techniques may overcome this problem. The primary goal of this program is the development and use of theoretical tools for the analysis of one key example of such optical experiments.

The experiment of interest is being performed in a companion effort headed by G. Mourou of the University of Rochester, Rochester New York. In it a laser pulse is used to activate a photoconductive transient in a gap in a microstrip line on GaAs. A second pulse then is used to electro-optically sample the voltage wave triggered in this line as a result of the photoconductive transient. Both portions of the experiment are done on a subpicosecond scale thus making it the first direct probe of subpicosecond current transients in semiconductors. It differs from the vast majority of optical experiments in that information directly pertinent to the carrier momentum is collected whereas spectroscopic techniques tend to probe only carrier energy.

There are three main steps which must be performed in the development of a model of this experiment. First, we must accurately model the processes by which the laser pulse is converted into electron-hole pairs inside the gap. Secondly we must model the ensuing transport transients and their conversion into a current transient seen at the gap terminals. Lastly, we then must model the process by which this current transient is represented as a voltage wave traveling down a microstrip line as it is this wave that is sampled in electro-optic experiments of this sort. Once the models are developed they then are to be used in conjunction with actual experiments in probes of our understanding of carrier transport on the subpicosecond scale.

Status of Research

During the first portion of the first year our efforts centered on the . transformation of optical pulse data into a set of electron-hole pair generation events which are distributed over the transmission line gap in both space and time. As the transport model to be used in the second step is a Monte Carlo model, we chose to use Monte Carlo techniques for this portion as well. The result was an ability to simulate spatial, temporal and spectrally varying electron-hole pair generation events. Attention was then turned to the second problem. Here the difficulty is that while there has been a great deal of effort at developing good Monte Carlo models for electron transport in GaAs, holes have been neglected. They however cannot be neglected here. The development of a good hole model was the other main effort of the first year of the program. The hole model which was developed includes both light and heavy holes. The hole masses used are .45m for heavy holes and .082m for light holes¹. The scattering processes include interband and intraband transitions through the mechanisms of acoustic phonon, optical phonon and ionized impurity scattering². The drift velocity curves for holes show a peak velocity of 7 x 10^6 cm/s at 80 kV/cm.

The main task for the second year was the extension of these models into a thorough simulation of the experiment of interest. We began by exploring variations on the seminal Monte Carlo study of Ruch³. Ruch showed that if one suddenly stepped the electric field in a semiconductor that the electrons present would overshoot their expected final velocity value during a transient of several picoseconds duration. The optical experiments considered here however may significantly differ from the study of Ruch in that here we start with an existing field and instead create a collection of electron-hole pairs. Furthermore while these pairs will have an initial average momentum of zero they may be quite energetic if the laser wavelength corresponds to above-gap excitation. In fact we expect to see a transient in which the carrier gas is losing energy while simultaneously gaining a nonzero average momentum. We used the Monte Carlo tools to study the effects of such excitation. We found that when the electrons were excited into the bottom of the central valley that overshoots quite similar to those of a typical Ruch-like velocity overshoot study were expected. When higher photon energies were used to place electrons above the threshold for intervalley transfer changes were seen as these electrons rapidly transfer to the higher energy satellite valleys. Velocity overshoots then were seen only for relatively large electric fields and even then a smaller overshoot is expected. We also varied the temperature and found that cooling to 77 K should increase the amount of overshoot but did not dramatically increase the duration of the transient in time.

The next step was the coupling of the electron and hole transport models with Poisson's equation. The difficulty faced here is the treatment of carriers which leave the region simulated during the course of the simulation. We used a periodic boundary condition in which every time a carrier left one end of the region, it was reinjected from an equilibrium distribution at the other end. We developed a more complete methodology for treating this problem but found that for the relatively simple case considered here that the more powerful technique is not needed. The essential concept is to couple this reinjection problem with the already assumed boundary condition for the field by insuring that the induced current needed to maintain the field boundary condition through induced charges at some interface be consisent with the current associated with the net flow of carriers in and out of the region of simulation.

Once the boundary condition issue was resolved the crucial question which we asked was whether or not the fields and carrier densities would be significantly inhomogenous during the early transient. Such homogeneity was assumed by Ruch and would greatly simplify the interpretation of any experiment. We found that at least in a one-dimensional model homogeneity is expected. We then coupled the photoconductive gap simulated by Monte Carlo techniques to the transmission line and compared our results with those of experiment. Qualitative agreement was obtained.

There are several main scientific questions which we hope to answer. The first is the nature and existence of velocity overshoots of carriers suddenly introduced to a high electric field in a semiconductor. All transport calculations that allow for this possibility predict it yet no unambiguous experimental observation of such an overshoot has been made. Our goal is to use these techniques in an effort at performing such an experiment. We have made good progress in establishing an effective experimental protocol. High quality material must be used and effective contacts to it must be made. It is important that a low excitation experiment be carried out in which difficulties associated with substantial temporal variations in the voltage across the photoconductive gap, strong carrier-carrier interactions and the generation of nonequilibrium phonons are avoided. The sensitivity of the electro-optical sampling technique should allow this to be possible.

Even in this protocol we face a difficulty. The desired transport transient is convolved with the capacitive response of the gap. We will use an extension of Auston's model⁴ of these experiments as a basis for understanding how the deconvolution is to be performed. Here the Monte Carlo models will play an important role. We will simulate an experiment and then use our proposed data analysis techniques on the waveforms generated by the simulation. The answers obtained will be compared with the stored details of the actual transient response of the gap during the simulation. Such a test is useful as deconvolution is an ill-posed problem.

A second important question involves the role of intervalley scattering in such experiments. The intervalley coupling coefficients are extremely important parameters and yet are difficult to either measure or compute. It may be possible to probe these processes by choosing laser energies that introduce carriers into the conduction band at the level of the satellite valley minima.

A third set of questions involve the role of carrier-carrier scattering. Optical experiments offer one vehicle of studying these interactions without the possible competing effects of increased doping density and associated ionized impurity scattering.

We also will be investigating the utility of such techniques as a basis for high-speed device characterizations. The first question here will be the development of an effective equivalent circuit model for the photoconductive response of the gap. This is closely related to the development of methods of extracting transient transport data from the photoconductive experiment. The second main question is the use of such techniques as a vehicle for determining the S-parameters of high frequency transistors. Here we will first use analytical models in which both the transient time-domain response and the sinusoidal steady-state S-parameters are expressable in closed form. The second feature will be the use of a SPICE based, time-domain GaAs MESFET circuit model for which the Sparameters are known. We will simulate the transient experiments and investigate various techniques for extracting S-parameters from the time series data.

Professional Personnel

Christopher Caruso is a graduate research assistant working on this program. His undergraduate degree is in Electrical Engineering from General Motors Institute, Flint MI. He presently is a graduate student in Electrical Engineering at Arizona State University and anticipates the completion of his masters degree in November 1986.

Dr. Robert O. Grondin is an assistant professor of Electrical and Computer Engineering at Arizona State University and the principal investigator of the present effort. He received the BS, MS and Ph.D. degrees in Electrical Engineering from the University of Michigan. He was a post-doctoral fellow. at Colorado State University prior to his acceptance of his present faculty position in 1983. In 1985 he was named a Presidential Young Investigator by the National Science Foundation.

Interactions

As reported in the first year a strong interaction has developed between this effort and the companion experimental effort at the University of Rochester. All of the personnel described above have visited the experimental site and regular communication via telephone, letter and running into each other at conferences has developed. During the second year this form of interaction has continued to grow. In the upcoming years this will also be augmented by the URI on ultrafast electronics which has been established jointly at Rochester and Cornell.

Another strong interaction noted during the first year was one developed between the investigators on this project and others who are performing investigations into transient carrier transport at Arizona State University. This has greatly aided in the development of the Monte Carlo models. Several of the other investiagtors have now graduated and are located at the University of Illinois and at Scientific Research Associates in Glastonbury Connecticut. Both continue to be actively involved in this general activity.

References

- 1. K. Brennan and K. Hess, Physical Review B 29, 5581 (1984)
- 2. C. Jacoboni and L. Reggiani, Reviews of Modern Physics, 55, 645 (1983)
- 3. J. Ruch, IEEE Trans. Electron Devices, ED-19, 652 (1972)
- 4. D. Auston, IEEE J. Quantum Electronics, QE-19, 639 (1983)

