



ECURITY CLASSIFIC			011		
		VENTATION	PAGE		
UNCLASSIE AD-A1	83 188	16 RESTRICTIVE NONE	MARKINGS		
SECURITY CLAS	ECIE	3 DISTRIBUTION	AVAILABILITY O	OF REPORT	
DECLASSIFICATION / DOWNGRADING SCHEE	JG 1 2 1987	UNLIMITE	D		
PERFORMING ORGANIZATION REPORTIUM	BER(S)	5 MONITORING	ORGANIZATION R	REPORT NUMBE	R(S)
NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b OFFICE SYMBOL - (If applicable)	7a NAME OF M Chemistry Office of	ONITORING ORGA Division the Chief (NIZATION	esearch
ADDRESS (City, State, and ZIP Code) Ashington, DC 20375-5000		76. ADDRESS (Cr Arlington	ty, State, and ZIP , VA 22217-5	Code) 5000	
NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9 PROCUREMEN	T INSTRUMENT ID	DENTIFICATION	NUMBER
ADDRESS (City, State, and ZIP Code)		10. SOURCE OF I	UNDING NUMBER	RS	
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
		61153N	RR013-01-40		61-2804-0-7
		USING DIOLE	Baser Mbser		
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball	la* and H.H. Nels	on			
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a TYPE OF REPORT chnical	La* and H.H. Nels COVERED	on 14 DATE OF REPC 1987 Ju	DRT (Year, Month,	Day) 15 PAC	
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a type of REPORT chnical SUPPLEMENTARY NOTATION IRC Postdoctoral fellow	La* and H.H. Nels COVERED LO/86 TO 10/87	on 14 DATE OF REPC 1987 Ju	DRT (Year, Month, ly 30	Day) 15 PAC 2	SE COUNT 1
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a type of REPORT chnical SUPPLEMENTARY NOTATION IRC Postdoctoral fellow COSATI CODES	La* and H.H. Nels COVERED LO/86 TO 10/87 18 SUBJECT TERMS (ON 14 DATE OF REPO 1987 Ju Continue on revers	ORT (Year, Month, ly 30 the if necessary and	Day) 15 PAC 2: d identify by b	SE COUNT 1 lock number)
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a TYPE OF REPORT connical SUPPLEMENTARY NOTATION IRC Postdoctoral fellow COSATI CODES FIELD GROUP SUB-GROUP	La* and H.H. Nels COVERED LO/86 TO_10/87 18 SUBJECT TERMS (on 14 DATE OF REPC 1987 Ju Continue on revers	PRT (Year, Month, ly 30 The if necessary and	Day) 15 PAC 2: d identify by bi	SE COUNT 1 lock number)
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a TYPE OF REPORT Schnical SUPPLEMENTARY NOTATION IRC Postdoctoral fellow COSATI CODES FIELD GROUP SUB-GROUP	La* and H.H. Nels COVERED LO/86TO10/87 18 SUBJECT TERMS (ON 14 DATE OF REPC 1987 Ju Continue on revers	DRT (Year, Month, ly 30 The if necessary and	Day) 15 PAC 2: d identify by b	SE COUNT 1 lock number)
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a TYPE OF REPORT Cchnical SUPPLEMENTARY NOTATION RC Postdoctoral fellow COSATI CODES FIELD GROUP SUB-GROUP ABSTRAT/(Continue on reverse if necessar	La* and H.H. Nels COVERED LO/86TO10/87 18 SUBJECT TERMS (18 SUBJECT TERMS (ny and identify by block (ON 14. DATE OF REPO 1987 Ju Continue on revers	DRT (Year, Month, ly 30 Se if necessary and	Day) 15 PAC 2: d identify by b	iE COUNT 1 lock number)
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball a TYPE OF REPORT Chnical SUPPLEMENTARY NOTATION RC Postdoctoral fellow COSATI CODES FIELD GROUP SUB-GROUP ABSTRUT/(Continue on reverse if necessar Absolute rate constants for reaction of BH ₃ ^T with O ₂ ^T and H The reactions were studied at ro	a* and H.H. Nels COVERED LO/86TO0/87 18 SUBJECT TERMS (18 SUBJECT TERMS (ry and identify by block (r the reaction of (BH) 20 are reported using boom temperature over	on 14. DATE OF REPO 1987 Ju Continue on revers number) Juliu (with CO, NO, a diode laser r a pressure rai	PRT (Year, Month, 1y 30 The if necessary and $C_2^TH_4$ and up to probe BH_3^{-7} nge of 6 to 620	Day) 15 PAC 2: d identify by b per limits fc disappearand 0 torr using	The count the count N_2 as
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Ball TYPE OF REPORT Chnical SUPPLEMENTARY NOTATION RC Postdoctoral fellow COSATI CODES FIELD GROUP SUB-GROUP ABSTRUT/(Continue on reverse if necessar Absolute rate constants for reaction of BH ₃ ⁻¹ with O ₂ ⁻¹ and H The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10 ⁻¹⁸ cm ³ - ¹ for BH ₃ ⁻¹ limit of 3.7 x 10 ⁻¹³ cm ³ - ¹ for BH ₃ ⁻¹	18 SUBJECT TERMS (18 SUBJECT TERMS (18 SUBJECT TERMS (19 and identify by block (19 and identify by block (10 are reported using 10 are reported using 10 to 620 to 10 to 10 to 620 to 10 to 1	Continue on reverse with CO, NO, a diode laser r a pressure rai + CO was found our of N_2^2 with ant was found our of N_2^2 with	PRT (Year, Month, 1y 30 The if necessary and $C_2^TH_4$ and up to probe BH_3^T nge of 6 to 620 and to be in the rate constants to approach the a pressure index	Day) 15 PAC 2: d identify by be per limits for disappearance 0 torr using e intermedia ranging from high press	The count 1 for the ce. N_2 as the number 1.5 for the ce. 1.5 for 1.5 for 2 for 1.5 for 2 for 1.5 for 2 for 1.5 for 2 for 1.5 for
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey BallTYPE OF REPORT chnical13b TIME FROM_1SUPPLEMENTARY NOTATION RC Postdoctoral fellow13b TIME FROM_1COSATI CODES FIELDGROUPSUB-GROUPSUB-GROUPABSTRUT/(Continue on reverse if necessar Absolute rate constants for reaction of BH3 with O2 and H2 The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10^{-13} cm3s^{-1}. For BH37 limit of 3.7 x 10^{-13} cm3s^{-1} at pr constant (p \geq 6 torr) for the rea reactions of BH3 + O2 and BH3 Upper limits of 5 x 10^{-15} cm3s^{-15}	La* and H.H. Nels COVERED LO/86TO	Continue on reverse 14. DATE OF REPORT 1987 Ju Continue on reverse for a pressure rate + CO was found for of N ₂ with ant was found for of N ₂ . The was found to than could be re- 1 - 1 respective	PRT (Year, Month, 1y 30 The if necessary and $C_2^TH_4$ and up to probe BH_3^T nge of 6 to 620 and to be in the rate constants to approach the pressure inde be 5.2 x 10 ⁻¹¹ neasured in the were placed o	Day) 15 PAC 2: d identify by b per limits fc disappearance 0 torr using e intermedia ranging from high press ependent rat cm ³ s ⁻¹ . Th ese experime-	E COUNT 1 Pock number) Pock number)
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Balla. TYPE OF REPORT13b TIME FROM_1SUPPLEMENTARY NOTATIONRC Postdoctoral fellowCOSATI CODESFIELDGROUPSUB-GROUPABSTRUT'(Continue on reverse if necessal Absolute rate constants for reaction of BH3' with O2' and H2 The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10 ⁻¹³ cm3s ⁻¹ . For BH3' limit of 3.7 x 40 ⁻¹³ cm3s ⁻¹ at pr constant ($p \ge 6$ torr) for the rea reactions of BH3 + O2 and BH3 Upper limits of 5 x 10 ⁻¹⁵ cm3s these reactions.	La* and H.H. Nels COVERED LO/86TO	Continue on reverse 14 DATE OF REPC 1987 Ju Continue on reverse 1987 Ju 1987 J	PRT (Year, Month, 1y 30 The if necessary and r_{2} r_{4} and up to probe BH ₃ nge of 6 to 620 and to be in the rate constants to approach th e pressure inde be 5.2 x 10 ⁻¹¹ neasured in th were placed o	Day) 15 PAC 2: d identify by b per limits for disappearand 0 torr using e intermedia ranging from high press ependent rat cm ³ s ⁻¹ . Th ese experime on the rate of	be COUNT 1 lock number) for the ce. N_2 as te N_2 as te $n \ 1.5$ sure e e ents. f
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Balla. TYPE OF REPORT13b TIME FROM_1chnical13b TIME FROM_1SUPPLEMENTARY NOTATIONIRC Postdoctoral fellowCOSATI CODESFIELDGROUPSUB-GROUPAbsolute rate constants for reaction of BH3 with O2 and H3The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10 ⁻¹⁸ (1)3s ⁻¹ . For BH3Imit of 3.7 x 10 ⁻¹³ cm3s ⁻¹ at pr constant (p \geq 6 torr) for the rea reactions of BH3 + O2 and BH3 Upper limits of 5 x 10 ⁻¹⁵ cm3s these reactions.CCDS TRIBUTION AVAILABILITY OF ABSTRACCDINCLASSIFIED UNLIMITED	La* and H.H. Nels COVERED LO/86 TO 10/87 18 SUBJECT TERMS (18 SUBJECT TERMS (1	Continue on reverse with CO, NO, a diode laser r a pressure rai + CO was found forr of N ₂ ? The was found to than could be r -1 respectivel 21 ABSTRACT SE UNCLASSIF	PRT (Year, Month, 1y 30 The if necessary and C_2H_4 and up to probe BH_3^{-1} inge of 6 to 620 and to be in the rate constants to approach the pressure inder be 5.2 x 10 ⁻¹¹ neasured in the were placed of CURITY CLASSIFIC LED	Day) 15 PAC 2: d identify by be per limits for disappearance 0 torr using e intermedia ranging from high press ependent rat cm ³ s ⁻¹ . Th ese experime on the rate of	be COUNT 1 lock number) for the ce. N_2 as te N_2 as te $n \ 1.5$ sure e e ents. f
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Balla. TYPE OF REPORT13b TIME FROM_1SUPPLEMENTARY NOTATION IRC Postdoctoral fellow13b TIME FROM_1COSATI CODESCOSATI CODESFIELDGROUPSUB-GROUPABSTRAT' (Continue on reverse if necessal Absolute rate constants for reaction of BH3' with O2' and H2 The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10^{-13} cm3s^{-1}. For BH3' limit of 3.7 x 10^{-13} cm3s^{-1} at pr constant (p \geq 6 torr) for the rea reactions of BH3 + O2 and BH3 Upper limits of 5 x 10^{-15} cm3s^{-16} these reactions.DIS'RIBUTION AVAILABILITY OF ABSTRACE UNCLASSIFIED/UNLIMITEDSAME AS SAME AS A NAVE OF RESPONSIBLE INDIVIDUAL R. MCDOnald	La* and H.H. Nels COVERED LO/86TO10/87 18 SUBJECT TERMS (18 SUBJECT T	Continue on revers 14. DATE OF REPC 1987 Ju Continue on revers with CO, NO, a diode laser r a pressure ran + CO was found forr of N ₂ with ant was found forr of N ₂ . The was found to than could be re- 1 respectivel 21 ABSTRACT SE UNCLASSIF 225 TELEPHONE (202 767-33	PRT (Year, Month, 1y 30 The if necessary and C_2H_4 and up to probe BH_3^{TT} nge of 6 to 620 and to be in the rate constants to approach the pressure inde be 5.2 x 10 ⁻¹¹ neasured in th were placed o CURITY CLASSIFIC IED Include Area Code 40	Day) 15 PAC 2 2 d identify by bi i i c i f oper limits for disappearand 0 torr using e intermedia ranging from high press ependent rat cm ³ s ⁻¹ . Th ese experime on the rate of ATION	SE COUNT L lock number) or the ce. N ₂ as te n 1.5 sure e e ents. f SYMBOL
PERSONAL AUTHOR(S) Pasternack, R. Jeffrey Balla TYPE OF REPORT13b TIME FROM_1SUPPLEMENTARY NOTATIONJRC Postdoctoral fellowCOSATI CODESFIELDGROUPSUB-GROUPAbsolute rate constants for reaction of BH3 with O2 and H2 The reactions were studied at ro a buffer gas. The rate constant pressure regime over the pressu to 47 x 10 ⁻¹⁵ cm3s ⁻¹ . For BH3 limit of 3.7 x 10 ⁻¹³ cm3s ⁻¹ at pr constant (p \geq 6 torr) for the rea reactions of BH3 + O2 and BH3 Upper limits of 5 x 10 ⁻¹⁵ cm3s these reactions.CCDISTRIBUTION AVAILABILITY OF ABSTRACE UNCLASSIFIED-UNLIMITEDSAME AS AME OF RESPONSIBLE INDIVIDUAL .R. McDonaldPORM 1473, 84 MAR83.	La* and H.H. Nels COVERED LO/86TO_10/87	Continue on revers	PRT (Year, Month, 1y 30 The if necessary and the if necessary and C_2H_4 and up to probe $BH_3^{(7)}$ nge of 6 to 620 and to be in the rate constants to approach the pressure inde be 5.2 x 10 ⁻¹¹ neasured in th were placed of CURITY CLASSIFIC LED Include Area Code 40 SECURITY	Day) 15 PAC 2: d identify by bi i i c i f oper limits for disappearance 0 torr using e intermedia ranging from high press ependent rat cm ³ s ⁻¹ . Th ese experime on the rate of ATION e) 22c OFFICE CLASSIFICATION	SE COUNT L lock number) or the ce. N ₂ as te n 1.5 sure e ents. f SYMBOL N OF THIS PAGE

Vorsila soccessa extended resterado reverso recerció prontro arrente antenno arrente antena presenta recerció n

OFFICE OF NAVAL RESEARCH

PROGRAM ELEMENT 61153N

TECHNICAL REPORT NO. 2

REACTIONS OF BH₃ WITH CO, NO, O₂, C₂H₄.

AND H₂O USING DIODE LASER ABSORPTION

by

L. PASTERNACK, R. JEFFREY BALLA AND H.H. NELSON

Prepared for publication

Contraction of the second s

in the

Journal of Physical Chemistry

CHEMISTRY DIVISION NAVAL RESEARCH LABORATORY WASHINGTON, DC 20375-5000

JULY 1987

Reproduction in whole or in part is permitted for

any purpose of the United States Government

This document has been approved for public release and sale, its distribution is unlimited

Submitted to J. Phys. Chem.

REACTIONS OF BH₃ WITH CO, NO, O₂, C_2H_4 , and H_2O USING DIODE LASER ABSORPTION

L. Pasternack, R. Jeffrey Balla[†] and H.H.Nelson

Chemistry Division Naval Research Laboratory Code 6111 Washington, D. C. 20375-5000

Accesio	For		
NTIS	CRA&I	r.	
DTIC	TAB		
Unation	21-16 d	[]	
JUSTIC	121601) 	·····	
By Distrib	ition]	••••	
Ĥ.	valiability	Codes	
Dist	esenti un Succi	djor al	
A-1			

† NRC/NRL Postdoctoral Research Associate. Current address: Morgantown Energy Technology Center, Morgantown, WV 26505.

DTIC

OPY

Abstract

Absolute rate constants for the reaction of BH₃ with CO, NO, C_2H_4 and upper limits for the reaction of BH₃ with O₂ and H₂O are reported using a diode laser to probe BH₃ disappearance. The reactions were studied at room temperature over a pressure range of 6 to 620 torr using N₂ as a buffer gas. The rate constant for reaction of BH₃ + CO was found to be in the intermediate pressure regime over the pressure range 10 to 620 torr of N₂ with rate constants ranging from 1.5 to 47 x 10⁻¹³ cm³s⁻¹. For BH₃ + NO, the rate constant was found to approach the high pressure limit of 3.7 x 10⁻¹³ cm³s⁻¹ at pressures above ≈ 200 torr of N₂. The pressure independent rate constant ($p \ge 6$ torr) for the reaction of BH₃ + C₂H₄ was found to be 5.2 x 10⁻¹¹ cm³s⁻¹. The reactions of BH₃ + O₂ and BH₃ + H₂O were slower than could be measured in these experiments. Upper limits of 5 x 10⁻¹⁵ cm³s⁻¹ and 6 x 10⁻¹⁵ cm³s⁻¹ respectively were placed on the rates of these reactions.

200000

たんがべきない

2222222226

200705601h

172202020

There has been a resurgence of interest in the reactions of boranes due to the need to develop boron-assisted propellants. In studying the combustion of these systems, it has become evident that there is a lack of knowledge about basic chemical processes involving small boron containing species.¹⁻³ Although several models for B/O/H and B/C/O/H combustion have been proposed^{2,4} there are still large uncertainties in the thermochemistry of key radicals in the models and very few rate constants have been measured.

In this paper we report rate constants from the measurement of BH_3 reactions with CO, NO, and C_2H_4 and upper limits for the rate constants for BH_3 reactions with O_2 and H_2O . There are no previous experimental measurements of the rates of these reactions. with the exception of $BH_3 + C_2H_4 \rightarrow H_2BC_2H_5$ which has been reported by Fehlner.⁵ Several other elementary reactions of BH_3 have previously been studied in order to determine initial products and measure rate constants.⁶ It has been found that, in general, BH_3 reactions proceed via adduct formation followed by rearrangement or elimination.³

たんしょう がた ういいいいいたい ういんりょう

We use a tunable diode laser as the probe source in a pump-probe experiment to study the kinetics of BH₃. The technique has been described previously.⁷ BH₃ is formed by excimer laser photolysis (193 nm) of B₂H₆ as described by Irion and Kompa⁸ and detected by IR absorption using a tunable diode laser at the frequency of a rotation line of the ν_2 vibration of BH₃.⁹ We measure the rate of disappearance of BH₃ with added reactant gases.

L

Experimental

The reaction cell is a cylindrical pyrex tube of 2m length and 4cm diameter with BaF₂ windows. The BH₃ radicals are produced by 193 nm photolysis of diborane using an ArF excimer laser (Lumonics, Model 860). The excimer beam is mildly focussed by a 2m focal length lens and turned 90° by a dichroic beam splitter (Acton Research). The radical concentration is probed using a tunable diode laser (Laser Analytics, Model LS-3) operating near 1140 cm⁻¹. The IR radiation is transmitted by the dichroic beam splitter and makes 3 passes through the reaction cell. This arrangement permits collinear propagation of the pump and probe lasers. After passing through the reaction cell, the IR radiation from the diode laser passes through a 0.5m monochromator which selects a single laser emission mode. The probe laser intensity is measured using a HgCdTe detector. A schematic of the apparatus is shown in Figure 1.

Service States

والديد ويديدونا الأقصاء متحرا المعمومون والموكرين يراي بالمتعودين والديد برخرش إيذري وووف

The tunable diode laser is a cryogenically cooled lead-salt diode that emits cw radiation in many monochromatic and near monochromatic modes. Tuning is accomplished by varying either the refrigerator temperature or the laser current. A reference cell or a one inch germanium etalon is placed in the beam for wavelength calibration. After the wavelength for IR absorption by the radical has been determined, the temperature and current to the diode are fixed and the monochromator is set to pass the wavelength of the absorption.

To collect kinetic data, the diode laser frequency is tuned to the absorption feature of interest. The digital oscilloscope (LeCroy model 9400) and excimer laser are triggered sequentially to obtain the transient absorption signal. The oscilloscope is used to digitize and average the signal from 50-500 experimental repetitions. The signal is then sent to an IBM PC-AT for processing. A typical plot of the transient absorption of BH₃ is shown in Figure 2 using the ν_2 , J=6 line at 1141.0245 cm^{-1.9} The intensity of the

absorption feature is less than 2% of the diode laser intensity to maintain linearity of signal with concentration.

and the stand of the stand of the stand

The positions of some of the rotational lines in the Q branch of the ν_2 vibrational band of ¹¹BH₃ (X¹A'₁) have been determined by Kawaguchi et al.⁹ They have identified the positions of lines with J=1 to 9 of the ^qQ(J,K=J) branch in the region from 1140.5 to 1142.8 cm⁻¹. In this region, we use NH₃ and N₂O as reference gases.¹⁰ Although there is overlap from the very strong ν_{18} band of B₂H₆ at 1172 cm⁻¹,¹¹ we are able to discriminate between the B₂H₆ and BH₃ lines because of the high resolution of the diode laser. We have also produced BH₃ from BH₃CO photolysis at 193 nm and there is less interference from BH₃CO than from B₂H₆ in this wavelength region. However, we observed a lower quantum yield for BH₃ production from BH₃CO than from B₂H₆. In addition, BH₃CO must be synthesized (following the method of Burg and Schlesinger¹²). Since our experiments required a relatively fast flow of materials to reduce product buildup, we primarily photolyzed B₂H₆ rather than BH₃CO.

Experiments are performed under pseudo-first-order conditions with the reactant gas present in large excess over BH₃. When no reactant gas is added, we observe a slow disappearance of the BH₃ absorption which we attribute to reaction with the precursor B_2H_6 and to BH₃ recombination. These rates are fast compared to diffusion and pump out rates. Under our experimental conditions with total flow rates of 2 to 20 l/min the disappearance of BH₃ with no added reactant does not depend on the residence time in the cell. Under these conditions, the BH₃ decays in the presence of reactant are firstorder and can be fit by a single exponential extending over 2 to 4 reaction lifetimes. The transient absorption signal is analyzed starting 15 - 20 μ s after the excimer laser pulse. This allows time for vibrational and rotational equilibration and for the diode laser frequency to stabilize after the rf pulse generated by the firing of the excimer laser

The pressure of the precursor B_2H_6 was typically 80 - 90 mtorr. At these pressures, less than 10% of the excimer laser photons were absorbed by the precursor over the 2m path thus generating a homogeneous concentration of BH_3 over the length of the cell. For the $BH_3 + C_2H_4$ reaction studies the pressure of the precursor was reduced to 10 to 20 mtorr to enable the reactant gas to be present in large excess over BH_3 to maintain pseudo-first-order conditions.

Metered flows of the precursor (B_2H_6) , reactants $(C_2H_4, CO, NO, O_2, and H_2O)$ and buffer gases $(N_2 \text{ and } SF_6)$ were mixed and flowed through the cell using Tylan mass flow controllers and mass flow meters. The following gases were used without further purification: B_2H_6 (Matheson, 1% in He and 30% in He), N_2 and O_2 (Air Products, Industrial Grade purity), and SF_6 , C_2H_4 , CO, and NO (Matheson, Commercial purity). Water was admitted to the cell both by using a bubbler with N_2 and by using premixed tanks of water saturated N_2 .

A COMPANY OF SALARY SALE OF SALARY

Results and Discussion

 $BH_3 + CO$ reactions. The reaction of $BH_3 + CO \rightarrow BH_3CO$ is expected to be exothermic by approximately 23 kcal/mole.¹ We measured the rate of reaction of BH_3 with CO at total pressures of 10 to 600 torr at room temperature. A typical time resolved plot of the transient absorption of BH_3 in the presence of CO is shown in Figure 2. The curve shown is the result of averaging 100 experimental repetitions. The first-order decay constant is obtained from the slope of the log of the absorption signal vs. time, using a weighted linear-least-squares fitting technique. Second-order rate constants at a given total pressure are obtained from a plot of first-order rate constants vs. reactant pressure, as shown in Figure 3. The intercept represents the sum of all BH₃ removal processes other than reaction with CO. These are typically at least an order of magnitude slower than the fastest measured first-order decay constant. The pressure dependence of this reaction was measured using N₂ as a buffer gas at total pressures of 10 to 600 torr. The results are shown in Figure 4 and listed in Table 1. The uncertainties listed in Table 1 ($\pm 2\sigma$) refer to the precision of the data as determined from the scatter in the second-order plots. In general, the high pressure data is the least precise, due to pressure broadening of the absorption. We also used SF₆ as a buffer gas and found it to be about a factor of 2 more efficient at promoting this reaction. However, SF₆ weakly absorbs 1140 cm⁻¹ radiation and therefore could not be used as a buffer gas at high pressures. From Figure 4, it is evident that the high pressure limit for the BH₃ + CO reaction is likely greater than 10^{-11} cm³s⁻¹ and is approached at greater than atmospheric pressure.

There are no prior direct measurements of the $BH_3 + CO + M \rightarrow BH_3CO + M$ reaction. By comparison to the $BH_3 + PF_3$ reaction measured by Fridmann and Fehlner,¹³ Bauer¹ concludes that the rate should be approximately 3 x 10⁻¹² cm³s⁻¹. However, these experiments were performed at 500K and 6 torr total pressure in helium.

 $BH_3 + NO$ reactions. We have measured the reaction of BH_3 with NO at room temperature at total pressures of 11 to 310 torr using N₂ as a buffer gas. Experiments were performed analagously to the $BH_3 + CO$ experiments. The second-order rate constants are listed in Table 2 and shown as a function of total reaction pressure in Figure 5. We approach a high pressure limit to the rate constant for the $BH_3 + NO$ reaction of approximately 3.7 x 10⁻¹³ cm³s⁻¹. The pressure at which the high pressure limit is reached is on the order of 200 torr of N₂. This is in contrast to the $BH_3 + CO$ reaction where the high pressure limit to the rate constant was found to occur at pressures greater than atmospheric.

There have been no previous measurements of the $BH_3 + NO$ reaction. Hoffmann and Engelhardt¹⁴ have observed a low concentration of the BH_3NO complex from $BH_3CO +$ NO reactions using mass spectroscopy. The different pressure dependence of the $BH_3 +$ CO and $BH_3 + NO$ reactions suggest that a mechanism other than complex stabilization might be involved. The thermochemistry of species containing boron and nitrogen is not well known so it is difficult to determine whether products from rearrangement or elimination channels are exothermic. Some possible exothermic channels, all of which involve extensive rearrangement, include:

$$BH_3 + NO \rightarrow [BH_3NO]^{\ddagger} \rightarrow H_3BN + O$$
 (1)

$$\rightarrow$$
 HNBOH + H (2)

L'OCTATION OF

$$\rightarrow$$
 H₂BN + OH (3)

- \rightarrow HBN + H₂O (4)
- \rightarrow HBO + NH₂ (5)

 $BH_3 + O_2$ reactions. The reaction of $BH_3 + O_2$ has several exothermic pathways, all of which involve extensive rearrangement, similar to the $BH_3 + NO$ reaction pathways. These include:

$$BH_{3} + O_{2} \rightarrow [BH_{3}O_{2}]^{\ddagger} \rightarrow HBOH + OH$$
(6)
$$\rightarrow HBO + H_{2}O$$
(7)
$$\rightarrow H_{2}BOH + O$$
(8)
$$\rightarrow B(OH)_{2} + H$$
(9)
$$\rightarrow HBO_{2} + H_{2}$$
(10)

Unlike $BH_3 + NO$, we see no apparent disappearance of BH_3 with pressures of 100 torr of O_2 at total pressures up to 400 torr with N_2 buffer gas. It is possible that spin constraints lead to large barriers in some of the $BH_3 + O_2$ channels which would not occur in $BH_3 + NO$ reactions.

The BH₃ signal does not decay exponentially in the presence of large amounts of O₂. This is most likely due to the formation of a significant amount of O³P atoms from 193 nm photolysis of O₂. The O³P may react with B₂H₆ to generate BH₃ from the reaction O³P + B₂H₆ \rightarrow BH₃O + BH₃. This reaction has been proposed as a step in ^Aiborane oxidation.¹⁵ From our results, we estimate that the rate of reaction of BH₃ + O₂ is less than 5 x 10⁻¹⁵ cm³s⁻¹. Although there are no prior measurements of this reaction, Shaub and Lin⁴ estimate the rate constant for reaction (6) to be 7 x 10⁻¹⁷ cm³s⁻¹ from their model of diborane oxidation.

 $BH_3 + C_2H_4$ reactions. We have measured the rate of reaction of $BH_3 + C_2H_4$ to be (5.2 \pm 1.0) x 10⁻¹¹ cm³s⁻¹ at room temperature. By monitoring the rate of disappearance of BH_3 in the presence of excess C_2H_4 at a series of ethylene pressures we determined second-order rate constants for this reaction using the same techniques as in the other BH_3 reactions presented in this report. The experiments were performed at 9 different total pressures between 6 and 90 torr using N₂ as a buffer gas. We observed no pressure dependence for this reaction. Our results are reported in Table 3.

The hydroboration reaction has been studied more extensively than the other reactions reported in this paper. Fehlner⁵ has measured the rate of production of ethylborane ($C_2H_5BH_2$) from this reaction at temperatures from 430 to 590K and 4.7 torr total pressure in helium to be in the range of 1.5 x 10^{-12} cm³s⁻¹ to 4.2 x 10^{-12} cm³s⁻¹, which is more than an order of magnitude slower than our room temperature results. If we assume that this represents the high pressure limit for the reaction and that the only pathway is to ethylborane formation, we can combine their results with ours to determine an activation energy of -4 kcal/mole for this reaction.

There have also been several theoretical studies of this reaction.¹⁶⁻¹⁹ The reaction to form ethylborane is predicted to proceed through a two step mechanism.^{17,19} The first step involves the formation of a π -complex held together by a three center CBC bond. The complex is calculated to be 10 kcal/mole more stable than the reactants.¹⁷ The second step involves the formation of a transition state via a donation-back-donation mechanism. In this step a transition state is postulated in which ethylene donates electrons through a carbon atom to the vacant orbital on boron at the same time as a hydrogen on BH₃ donates electrons back to the other carbon atom of ethylene in a four center arrangement. The calculations of Sundberg et al.¹⁷ show no activation barrier for this reaction. Nagase et al.¹⁹ predict an activation energy of ~4 kcal/mole.

8

1222.52

 $BH_3 + H_2O$ reactions. We observe no reaction between BH_3 and H_2O with 5 torr of water and total pressures up to 200 torr using N₂ as a buffer gas. Water was introduced into the reaction cell both by bubbling N₂ through water and by making mixtures of H₂O in N₂. We were able to confirm the presence and concentration of water in the cell by monitoring absorption in the ν_2 band of H₂O in the region 1120 - 1142 cm⁻¹.^{20,21} In addition, by monitoring the B₂H₆ absorption both with and without added water, we did not observe room temperature reaction between B₂H₆ and H₂O, although Weiss and Shapiro²² have observed diborane hydrolysis at 324K. We estimate that the rate constant for reaction of BH₃ + H₂O is less than 6 x 10⁻¹⁵ cm³s⁻¹. Shaub and Lin⁴ predict a room temperature rate constant of 7×10^{-17} cm³s⁻¹ for the BH₃+H₂O reaction from their model of diborance oxidation. However, Weiss and Shapiro²² have postulated that one step in the hydrolysis of diborane at 324K is BH₃ + H₂O \rightarrow [H₃BH₂O][‡] \rightarrow BH₂OH + H₂. Based on the work of Weiss and Shapiro, Bauer has derived a rate constant for the BH₃ + H₂O reaction of approximately 10⁻¹³ cm³ s⁻¹. The disagreement between our results for the BH₃ + H₂O reaction and the results derived from the experiments of Weiss and Shapiro may be indicative of a different mechanism for the diborane hydrolysis than that which they proposed. However, evidence for the existence of a borane-water complex comes from a kinetic study of the intermediates in hydroborate ion hydrolysis by Wang and Jolly.²³ In addition, Dill et al.²⁴ have calculated a configuration for this complex that is stable relative to BH₃ + H₂O by 6 kcal/mole. In this complex, BH₃ and H₂O are in a staggered conformation.

Conclusions

We have studied the reactions of BH₃ with CO, O₂, NO, H₂O and C₂H₄. It is difficult to justify our results in terms of the model for BH₃ reactions proposed by Fehlner.³ By this theory, BH₃ reactions are expected to proceed via reaction with Lewis bases to produce tetra-coordinate boron compounds. These reactions are expected to be fast. Consequently they involve early transition states which result from reactions controlled by early interactions. For BH₃ reactions, these interactions are due to the attractive force generated by the empty B 2p₂ orbital and the donor p_{σ} orbital.³ Although this type of reaction is expected to be faster for reactants containing lower nuclear charge, it is difficult to justify such large differences in rates and pressure dependences as we observe. Our results suggest that different mechanisms may be involved for BH₃ reactions with CO, O₂, and NO. Theoretical calculations on BH₃CO, BH₃NO, and BH₃O₂ complexes could help determine the mechanism. Experimental studies to determine reaction products would also be useful.

Acknowledgment

We thank Dr. J. E. Butler for providing the BH_3 spectroscopic information prior to publication. We also thank Dr. S. H. Bauer for providing reference 1 prior to publication and for his helpful discussions.

Total Pressure (torr)	k x 10 ¹³ (cm ³ s ⁻¹)
10	1.5 ± 0.1
10	1.7 <u>+</u> 0.1
20	2.7 + 0.4
30	4.3 ± 0.7
40	4.6 <u>+</u> 0.6
50	5.4 ± 1.0
60	6.6 ± 0.8
80	6.2 <u>+</u> 0.8
100	9.9 <u>+</u> 2.3
100	12.0 <u>+</u> 2.3
203	16.6 <u>+</u> 2.1
300	21.0 <u>+</u> 3.1
400	21.7 <u>+</u> 4.1
400	22.6 <u>+</u> 3.4
500	39.6 <u>+</u> 5.7
500	34.6 <u>+</u> 4.6
600	38.8 <u>+</u> 5.6
620	46.6 ± 4.3

20000000

Table 1. Room temperature rate constants for the reaction of $BH_3 + CO$ with N_2 buffer gas $(\pm 2\sigma)$.

Table 2. Room temperature rate constants for the reaction of $BH_3 + NO$ with N_2 buffer gas $(\pm 2\sigma)$.

k x 10 ¹³ (cm ³ s ⁻¹)		
$\frac{1}{2.0 \pm 0.2}$		
2.9 ± 0.3		
3.0 ± 0.6		
3.7 <u>+</u> 1.3		
3.7 <u>+</u> 0.6		

Table 3.	Room temperature rate constants for the reaction of
	$BH_3 + C_2H_4$ with N ₂ buffer gas ($\pm 2\sigma$).

Total Pressure (torr)	k x 10 ¹¹ (cm ³ s ⁻¹)
6	5.2 ± 1.8
12	5.7 <u>+</u> 1.7
20	5.1 ± 0.4
22	5.4 ± 0.8
45	4.3 ± 0.8
46	4.1 ± 0.6
47	3.5 ± 0.4
88	7.1 ± 2.7
89	6.0 <u>+</u> 2.1

DECKORSED CONDECCO

References

- 1. Bauer, S. H. In Boron Chemistry, Liebman, J. F. and Williams, R.; in press.
- 2. Brown, R. C.: Kolb, C. E.; Yetter, R. A.; Dryer, F. L.; Rabitz, H. R.; Aerodyne Report ARI-RR-580, 1987.
- 3. Fehlner, T. P.; Housecroft, C. E. In *Molecular Structure and Energetics*; Liebman, J. F.; Greenberg, A., Eds.; VCH: Deerfield Beach, 1986; Vol. 1, Chapter 6.
- 4. Shaub, W. M.; Lin, M. C. NBS Special Publication 1979, 561, 1249.
- 5. Fehlner, T. P. J. Am. Chem. Soc. 1971, 93, 6366.
- 6. Fehlner, T.P. Int. J. Chem. Kinetics 1975, 7, 633.
- 7. Balla, R. J.; Pasternack, L. J. Phys. Chem. 1987, 91, 74.
- 8. Irion, M. P.; Kompa, K. L. J. Photochem. 1986, 32, 139.
- 9. Kawaguchi, K.; Butler, J. E.; Yamada, C.; Bauer, S. H.; Minowa, T.; Kanamori, H.; Hirota, E. J. Chem. Phys., in press.

- 10. Guelachvili, G.; Rao, K. N. Handbook of Infrared Standards; Academic: New York, 1986; pp 120 and
- Duncan, J. L.; McKean, D. C.; Torto, I.; Nivellini, G. D. J. Molec. Spectrosc. 1981, 85, 16.
- 12. Burg, A. B.; Schlesinger, H. I. J. Am. Chem. Soc. 1937, 59, 780.
- 13. Fridmann, S. A.; Fehlner, T. P. J. Phys. Chem. 1971, 75, 2711.
- 14. Hoffmann, V. K.; Engelhardt, U. Z. Anorg. Allg. Chemie 1972, 389, 97.
- 15. Hand, C. W.; Derr, L. K. Inorg. Chem. 1973, 13, 339.
- 16. Clark, T.; Schleyer, P. v. R. J. Organomet. Chem. 1978, 156, 191.
- 17. Sundberg, K. R.; Graham, G. D.; Lipscomb, W. N. J. Am. Chem. Soc. 1979, 101, 2863.
- 18. Dewar, M. J. S.; McKee, M. L. Inorg. Chem. 1978, 17, 1075.
- 19. Nagase, S.; Ray, N. K.; Morokuma, K. J. Amer. Chem. Soc. 1980, 102, 4536
- 20. Guelachvili, G. J. Opt. Soc. Am. 1983, 73, 137.
- 21. Benedict, W. S.; Claassen, H. H.; Shaw, J. H. J. Res. Natl. Buo. Stand (1952) 49, 91
- 22. Weiss, H. G.; Shapiro, I. J. Am. Chem. Soc. 1953, 75, 1221.

- 23. Wang, F. T.; Jolly, W. L. Inorg. Chem. 1972, 11, 1933.
- 24. Dill, J.; Schleyer, P. v. R.; Pople, J. A. J. Am. Chem. Soc. 1975, 97, 3402.

Figures

- Fig. 1. Schematic diagram of the apparatus.
- Fig. 2. Plot of the transient absorption (in arbitrary units) of BH₃ for the reaction of BH₃ + CO. The plot is from the average of 100 decays at 298K. P(B₂H₆)=66 mtorr, P(CO)=3 torr, and P(total)=50 torr using N₂ is a buffer gas. The oscilloscope is triggered 10 μs prior to the excimer laser.
- Fig. 3. Plot of the observed BH₃ decay constant, k_1 , vs CO pressure for the reaction of BH₃ + CO at 298K. N₂ was used as a buffer gas. $P(B_2H_6) = 66$ mtorr and the total pressure was 50 torr. The slope gives a reaction rate of 5.4×10^{-12} cm³s⁻¹ and the intercept represents the rate for other BH₃ removal processes.
- Fig. 4. Pressure dependence of the bimolecular rate constant for the reaction of BH_3 with CO at 298K. The buffer gas used is N₂. The error bars represent $\pm 2\sigma$.
- Fig. 5. Pressure dependence of the bimolecular rate constant for the reaction of BH_3 with NO at 298K. The buffer gas used is N₂. The error bars represent $\pm 2\sigma$



Chier Chier

1.222222

and the second









TECHNICAL REPORT DISTRIBUTION LIST, GEN

the bar out the watt

.

......

1

	No.		No.
	Copies		Copies
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 1911	1 2
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

Harren Corri

