

AD-A186 290

BINDING OF PENTAMMINERUTHENIUM(II) RESIDUES TO THE
TRIS(BIPYRAZINE)RUTHENIUM(II) COMPLEX (U) YORK UNIV NORTH YORK
(ONTARIO) DEPT OF CHEMISTRY H E TOMA ET AL. AUG 87

UNCLASSIFIED

IR-18 SN00014-84-G-0201

F/G 7/3

1-1

NL

END
DATE
FILMED
12 87



MICROCOPY RESOLUTION TEST CHART
 NATIONAL BUREAU OF STANDARDS-1963-A

AD-A186 290

(12)

DTIC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-84-G-0201

Task No. 0051-865

Technical Report #18

DTIC
ELECTE
OCT 14 1987
S D

Binding of Pentaammineruthenium(II) Residues to the
Tris(bipyrazine)ruthenium(II) Cation

By

Henrique E. Toma, Pamela R. Auburn, Elaine S. Dodsworth,
M. Neal Golovin and A.B.P. Lever

in

Inorganic Chemistry

York University
Department of Chemistry
4700 Keele St., North York
Ontario, Canada M3J 1P3

Reproduction in whole, or in part, is permitted for any purpose of the United States Government

*This document has been approved for public release and sale; its distribution is unlimited

*This statement should also appear in Item 10 of the Document Control Data-DD form 1473. Copies of the form available from cognizant contract administrator

87 10 2 046

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY Unclassified		3 DISTRIBUTION/AVAILABILITY OF REPORT As it appears on the report	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Report # 18		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION A.B.P. Lever, York University Chemistry Department	6b OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c ADDRESS (City, State, and ZIP Code) 4700 Keele St., North York, Ontario M3J 1P3 Canada		7b. ADDRESS (City, State, and ZIP Code) Chemistry Division 800 N. Quincy Street Arlington, VA 22217 U.S.A.	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b OFFICE SYMBOL (if applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-84-G-0201	
8c. ADDRESS (City, State, and ZIP Code)		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Binding of Pentaammineruthenium(II) Residues to the Tris(bipyrazine)ruthenium(II) Cation			
12 PERSONAL AUTHOR(S) Toma, Henrique E.; Auburn, Pamela R.; Dodsworth, Elaine S.; Golovin, M. Neal; and Lever, Alfred B.P.			
13a TYPE OF REPORT Technical	13b TIME COVERED FROM Aug/86 TO Aug/87	14 DATE OF REPORT (Year, Month, Day) August 1987	15 PAGE COUNT 27
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		pentaammineruthenium, tris(bipyrazine)ruthenium cation, peripheral binding, electronic spectra, charge transfer spectra, electrochemistry, ruthenium ←	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) A series of $[(\text{NH}_3)_5\text{Ru}]_n^{\text{2n+}}(\text{bpz})_n^{\text{2n-}}$ ($n=1,2$) and $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n^{\text{2+2n}}]$ ($n = 1,3,6$, bpz = bipyrazine) complexes has been synthesized and characterized by means of electronic spectra, ^1H NMR, cyclic voltammetry and spectroelectrochemistry. The complexes are very stable and inert in aqueous solution. The peripheral pentaammineruthenium(II) groups in the tris(bipyrazine) series exhibit strong absorption bands in the visible, around 490 and 660 nm, ascribed to metal-to-ligand charge-transfer transitions. The attached ions can be reversibly oxidized, with formal potentials of approximately 0.65 V versus NHE, in comparison to 0.52 V for the $[(\text{NH}_3)_5\text{Ru}]_n^{\text{2n+}}(\text{bpz})_n^{\text{2n-}}$ complexes. No evidence of intervalence bands with $\epsilon > 100 \text{ M}^{-1} \text{ cm}^{-1}$ (#19 con'd over)			
20 DISTRIBUTION AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Robert K. Grasselli		22b TELEPHONE (Include Area Code)	22c OFFICE SYMBOL

#19 con'd

has been observed in the IR region of mixed valence species. The spectroscopic and electrochemical data are consistent with a very weak interaction between the peripheral pentaammineruthenium(II) groups.

#19 con'd

has been observed in the IR region of mixed valence species. The spectroscopic and electrochemical data are consistent with a very weak interaction between the peripheral pentaammineruthenium(II) groups.

cal

Introduction

Ruthenium(II) complexes of bipyrazine (bpz) are of special interest in coordination chemistry, because of their pronounced photochemical activity and of the existence of six peripheral nitrogen atoms available to bind Lewis acids and transition metal ions.²⁻¹⁰ Recently, we reported a detailed kinetic and spectroscopic study on a series of di-to-heptanuclear tris(bipyrazine)-ruthenium(II)-pentacyanoferrate(II) complexes in aqueous solution.¹¹ In this work, we extend the investigation to a series of pentaamineruthenium(II) derivatives, including the $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ species in solid state and in aqueous solution. Electrochemical and electronic spectroscopic data reveal that the several pentaamineruthenium(II) groups are essentially uncoupled to one another, and to the central ruthenium atom.

Experimental Section

The bipyrazine ligand was prepared by the pyrolysis of the bis(2-pyrazinecarboxylate)copper(II) complex at 270-290°C, under an argon atmosphere, as described in the literature.^{3,12} The solid was recrystallized from toluene, yielding pale yellow crystals, mp 186-190°C. [Lit. 190°C].¹³

The complexes $[\text{Ru}(\text{bpz})_3](\text{PF}_6)_2$ and $[\text{Ru}(\text{NH}_3)_5(\text{H}_2\text{O})](\text{PF}_6)_2$ were prepared by literature methods.^{2,14} $[\text{Ru}(\text{NH}_3)_5(\text{bpz})](\text{PF}_6)_2$ and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})](\text{PF}_6)_4$ were obtained by reacting 0.20 mmol of bpz with 0.20 or 0.40 mmol of $[\text{Ru}(\text{NH}_3)_5(\text{H}_2\text{O})](\text{PF}_6)_2$ in boiling acetone (25ml) under an argon atmosphere. After 30 min, the deep purple solution was evaporated to dryness at 55°C under a vigorous stream of argon. The black residue was kept under vacuum for one day, and washed with toluene. The compounds were recrystallized by dissolving in a small volume of 1:1 water-acetone solution, and allowing the solvent to evaporate very slowly under reduced pressure. Anal.

calcd for $[\text{Ru}(\text{NH}_3)_5(\text{bpz})](\text{PF}_6)_2$: C, 15.15; H, 3.33; N, 19.87. Found: C, 14.6; H, 3.2; N, 19.7. Calcd for $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})](\text{PF}_6)_4 \cdot \text{C}_3\text{H}_6\text{O}$: C, 11.31; H, 3.62; N, 16.78. Found: C, 11.0; H, 3.6; N, 15.8.

The polynuclear complexes $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ $n = 1, 3$ and 6 , were synthesized by reacting 0.025 mmol of $[\text{Ru}(\text{bpz})_3](\text{PF}_6)_2$ with 0.025, 0.075 or 0.150 mmol of $[\text{Ru}(\text{NH}_3)_5(\text{H}_2\text{O})](\text{PF}_6)_2$ in boiling acetone (25 ml) under an argon atmosphere. After 30 min, the violet-blue solution was evaporated to dryness, at 55°C , with a vigorous stream of argon. Recrystallization was

carried out as in the preceding examples. Anal. calcd. for

$[\text{Ru}(\text{bpz})_3\text{Ru}(\text{NH}_3)_5](\text{PF}_6)_4$: C, 21.48; H, 2.48; N, 17.74. Found: C, 21.6;

H, 3.0; N, 17.9. Calcd for $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3](\text{PF}_6)_8$: C, 12.56; H, 2.76;

N, 16.49. Found: C, 12.7; H, 2.9; N, 15.5. Calcd. for

$[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_6](\text{PF}_6)_{14}$: C, 7.74; H, 2.92; N, 15.79. Found: C, 8.1;

H, 3.0; N, 15.2.

The electronic spectra in the visible-UV region were recorded on a Hewlett-Packard 8451-A diode-array spectrophotometer. The absorption spectra in the near-infrared region were obtained with a Cary 17 spectrophotometer, with the samples dissolved in D_2O . Nuclear magnetic resonance spectra were recorded on a Varian T-60 spectrometer, at a probe temperature of 33°C . The measurements were made in D_2O , with use of 0.03-0.05 M solutions, after converting the hexafluorophosphate salts into the more soluble chloride form, with tetraphenylarsonium chloride. The chloride salts can also be obtained by dissolving the hexafluorophosphate compounds in acetone, and adding one or two drops of concentrated hydrochloric acid. The chloride salts precipitate almost

quantitatively in the presence of acetone and can be easily separated by filtration. *data were collected in the presence of 0.1 M methanol as internal solvent, and converted to Ru. This scale using 247 nm for CH_3*

Cyclic voltammetry was carried out with Princeton Applied Research Corporation (PARC) instrumentation, consisting of a model 173 potentiostat and

a model 175 universal programmer. A glassy carbon electrode was employed for the measurements in aqueous solution, with use of a conventional Luggin capillary arrangement with a Ag/AgCl (1 M KCl) reference electrode. A platinum wire was used as the auxiliary electrode. The measured potentials were converted to the NHE scale by adding 0.222 V. For the measurements in acetonitrile solutions, a platinum disk electrode was used versus a Ag/Ag⁺ (0.01 M silver nitrate in acetonitrile) reference electrode in the presence of 0.10 M tetraethylammonium perchlorate ($E^0 = 0.503$ vs NHE).¹⁵ For the spectroelectrochemical measurements, the PARC 173 potentiostat was used in parallel with the Cary 17 or HP 8451-A spectrophotometer. A three electrode system was designed for a rectangular quartz cell of 0.030 cm optical pathlength. A gold minigrid was used as a transparent working electrode, in the presence of a small Ag/AgCl reference electrode, and a platinum auxiliary electrode.

Results and Discussion

The electronic spectra of the $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ complexes are illustrated in Fig.1. The absorption bands at 230 and 286 nm also occur in free bpz and are ascribed to internal $\pi \rightarrow \pi_2^*$ and $\pi \rightarrow \pi_1^*$ transitions in the aromatic ligand respectively. The absorption band around 500 nm is assigned to the $\text{Ru}(\pi) \rightarrow \text{bpz}(\pi_1^*)$ charge-transfer (MLCT) transition, in comparison with the spectra of pentaammineruthenium(II) complexes of aromatic N-heterocycles.¹⁶ A second MLCT band associated with the $\text{Ru}(\pi) \rightarrow \text{bpz}(\pi_2^*)$ transition is expected in a cis-bipyrazine metal complex at a wavenumber approximately $8 \times 10^3 \text{ cm}^{-1}$ higher than that of the first one, i.e. around 360 nm. This band was not observed in either species and its absence may be evidence of a trans configuration for these species. The uv $\pi \rightarrow \pi^*$ transition (Table 1) in these species lie at the same energies as in the free (trans) ligand and differ in both position and separation from those obtained for the cis

bidentate species. A summary of the electronic transitions can be seen in Table I. While both the 1:1 and 2:1 species can theoretically exist in two isomeric forms depending upon which bpz nitrogen atoms are involved, the clarity of the NMR and UV/VIS spectra lead one to believe that only one isomer is present in each case. On the basis of minimising steric hindrance, binding only to N(4) is most likely (see labeling in Figure 4), and will be assumed henceforth. Note that neither the ^1H NMR nor the uv spectrum of the 1:1 adduct is consistent with the alternative formulation of this species with a bidentate bipyrazine ligand, i.e. as $[\text{Ru}(\text{NH}_3)_4(\text{bpz})]^{2+}$.

Protonation of the pentaammineruthenium(II)-bipyrazine complexes leads to a bathochromic shift of the $\pi\text{-}\pi^*$ and MLCT bands, with several isosbestic points in the visible and UV region, as shown in Fig. 1.A-B.

Quantitative measurements were carried out rapidly, using freshly prepared solutions, in order to minimize the acid catalysed aquation of the ruthenium(II)-ammines.¹⁷ In 0.01-6 M HCl solutions, a single protonation equilibrium is involved in the complexes. The corresponding $\text{pK}_a = 0.85 (\pm .05)$ for both the mononuclear and binuclear complexes was calculated from the linear plots of pH versus $\log[A - A_i]/[A_f - A]$, where A is absorbance and the subscripts i and f refer to the initial and final solutions, respectively. A slope of unity was observed, in agreement with the single protonation hypothesis.¹⁸ The similar pK_a 's of the mononuclear and binuclear complexes are consistent with protonation at the free N-atom of the coordinated pyrazine ring. This atom is expected to have its basicity increased by the backbonding interactions from the Ru(II) ion. In accordance, the pK_a values of the complexes are higher than that previously reported for the free bpz ligand ($\text{pK}_a = 0.45$).³ The increase in basicity of the peripheral nitrogen atom of the coordinated ring is very small, certainly far smaller than the increase observed in $[(\text{NH}_3)_5\text{Ru}(\text{Pyz})]^{2+}$ (Pyz = pyrazine) when it is protonated (from 0.6 to 2.5)¹⁹ or 2,6-dimethylpyrazine (from 1.94 to

3.70).¹⁸ Possibly the existence of the N=C-C=N framework created in the bipyrazine system, generates an alternate path for delocalisation such that the basicity of the far nitrogen is not greatly enhanced (vide infra).

Approximate values for the excited state pK_a 's were calculated based on the equation (1) excepting however, that such values are approximate, especially if there is excited state structural distortion, or solvent dipole induced distortion.^{19,20}

$$pK_a^* = pK_a(\text{gs}) + (\nu_1 - \nu_2)/476.5 \quad (1)$$

In the case of the mononuclear complex, $\nu_1 = 20,000 \text{ cm}^{-1}$, $\nu_2 = 17,580 \text{ cm}^{-1}$ and $pK_a^* = 5.9$. In the case of the binuclear complex, $\nu_1 = 19,700 \text{ cm}^{-1}$, $\nu_2 = 17,480 \text{ cm}^{-1}$ and $pK_a^* = 5.5$. The value of pK_a^* is some 2 pK units less than for the pyrazine complex but the difference, $pK_a - pK_a^*$ is actually rather larger, in both complexes, for the bipyrazine system than for the pyrazine system (5.05 and 4.65 vs. 3.9 units)¹⁹ but less than the difference (5.95 units) seen with the 2,6-dimethylpyrazine system.¹⁸ This discussion must be tempered with the suspicion that the unprotonated species has a trans bpz group, while, possibly, either of the two protonated species in question, may be cis.

Further protonation of the complexes can be observed in concentrated sulfuric acid; however, accurate measurements were precluded by the rapid decomposition of the complexes.

The electronic spectra of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ complexes with $n=1,3$ and 6 can be seen in Fig. 2A-C. The absorption bands around 230, 260, 300, 340 and 440 nm were also observed in the $[\text{Ru}(\text{bpz})_3]^{2+}$ cation. The last two bands were ascribed³ to the $\text{Ru}_c(\pi) \rightarrow \text{bpz}(\pi_2^*)$ and $\text{Ru}_c(\pi) \rightarrow \text{bpz}(\pi_1^*)$ MLCT transitions (subscript c refers to central).

Two additional bands appearing around 490 and 660 nm in the spectra of the polynuclear complexes exhibit intensities which increase with the number of attached $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups (Table I). The energy difference of $5,300 \text{ cm}^{-1}$ is

comparable to that observed for the MLCT bands of the $[\text{Ru}(\text{bpz})_3]^{2+}$ complex, and these new bands can be assigned to the $\text{Ru}_p(\pi) \rightarrow \text{bpz}(\pi_2^*)$ and $\text{Ru}_p(\pi) \rightarrow \text{bpz}(\pi_1^*)$ MLCT transitions from the peripheral (p) metal ions.

In Fig. 3 one can see two different aspects associated with the presence of free peripheral N-atoms in the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ complex. First, the complex can be protonated in concentrated HCl, with a set of four simultaneous isosbestic points, and $\text{pK}_a = -2.1$. This value has been evaluated from the Ho acidity functions for concentrated HCl solutions,²¹ and can be compared with the $\text{pK}_a = -2.2$ of the $[\text{Ru}(\text{bpz})_3]^{2+}$ complex^{3b}. In the presence of concentrated sulphuric acid, further changes are observed in the spectra which seem to indicate more drastic effects in the structure of the complexes. The possibility of decomposition reactions cannot be excluded in this case. The red shifted absorption can be assigned to a transition of the type $\text{Ru}(\text{II}) \rightarrow \pi^*(\text{bpzH}^+)$ as discussed previously.^{3b} The second aspect illustrated in Fig. 3.B refers to the binding¹¹ of $[\text{Fe}(\text{CN})_5]^{3-}$ groups in the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ complex. The successive attachment of 1 to 3 pentacyanoferrate(II) groups increase the intensity of the MLCT bands at 490 and 660 nm, with a small bathochromic shift (Table I). The $\text{Fe}(\pi) \rightarrow \text{bpz}(\pi_2^*)$ and $\text{Fe}(\pi) \rightarrow \text{bpz}(\pi_1^*)$ transitions in the series of $[\text{Ru}(\text{bpz})_3\{\text{Fe}(\text{CN})_5\}_n]^{2-3n}$ complexes ($n=1$ to 6) were observed near 500 and 700 nm.¹¹ Therefore, in the mixed polynuclear complexes, these bands should occur superimposed upon those from the peripheral ruthenium(II) ions. The spectrum of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_6]^{14+}$ complex is not modified by the presence of cyanoferrate(II) ions, excluding the possibility that the changes illustrated in Fig. 3.B arise from outer-sphere association. The assignments of the three charge transfer bands in the visible region, described here, are supported by resonance Raman data which will be discussed in detail elsewhere.²²

NMR Spectra ^1H NMR spectra of the $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ complexes are shown in Fig. 4.B-C, in comparison with the spectrum of the free ligand (Fig. 4.A). The NMR peaks at 9.46 ppm and 8.86 ppm in the free ligand have been assigned to the protons H3 and to the pair of protons H5 and H6, respectively. The NMR spectrum of the $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ complex exhibits an apparent doublet at 9.41 ppm, a singlet peak at 8.83 ppm and a doublet signal at 8.27 ppm, having relative intensities of 2:3:1, respectively. The binding of $[\text{Ru}(\text{NH}_3)_5]^{2+}$ ions to the N-heterocyclic ligands has been reported²³ to produce a shielding effect at the ortho protons (here denoted as H3 and H5), as a consequence of the paramagnetic anisotropy and backbonding interactions which predominate over the inductive effects from the Ru(II) ion. These effects have already been discussed in the literature.^{3,9,11, 23-25} The chemical shifts will depend on the extent of the shielding effects from π backbonding and of the deshielding, inductive effects from the metal ion. Previously,¹¹ the H6 protons shifted upfield, while the H5 protons shifted downfield, but to a lesser extent, upon coordination by the pentacyanoferrate(II) group. Therefore, the apparent doublet at 9.4 ppm in the monomeric pentaammineruthenium(II)-bipyrazine complex can be ascribed to the nonequivalent H3 and H3' protons. The singlet peak at 8.83 ppm is associated with the remote H6, H6' and H5' protons. The new doublet at 8.27 ppm is shifted upfield (shielding), being consistent with the H6 resonance signals. The NMR spectrum of the binuclear complex is shown in Fig.4C. In this case, the pair of H3, H3'; H5, H5' and H6, H6' protons are equivalent and the three resonance signals have the same intensity.

A potential problem in discussing the NMR spectra is the probable presence of structural isomers for certain values of n . Thus, for example, for $n = 2$, two pentaammineruthenium groups may bind to different bpz ligands, or conceivably, to the same bpz ligand. Moreover even where they bind to different

bpz ligands, two isomers exist differing in the relative positions of the pentammine groups. We discuss here the low resolution NMR considering the different types of proton present without considering their relative numbers, which will depend upon the isomer mix.

The methods used to prepare these compounds might be expected to yield a statistical mixture of products. Analytical data suggest but do not provide definitive proof of the unique n value of each species isolated. On the basis of the NMR data reported here ^{and} some preliminary high field NMR experiments, the following statements may be made:

- (i) Mixing of the $n = 1$ and $n = 3$, or $n = 0$ and $n = 6$ species causes no re-equilibration below 50° in water, the mixed (not averaged) spectra being obtained.
- (ii) The $n = 1$ species appears to be a single compound but the presence of some $n = 0$ cannot be ruled out. Higher values of n are absent.
- (iii) The $n = 3$ species spectrum is complex, indicative at least of several $n = 3$ isomers. If $n = 4, 5, 6$ are present, they can only be present in small amount.
- (iv) The $n = 6$ species is not contaminated with $n = 0-3$, but the presence of small amounts of $n = 4, 5$ cannot be excluded at this time. The evidence would indeed favour the isolation of compounds of unique n , as was apparently the case with the pentacyanoferrate(II) species.¹¹ The absence of a statistical mixture is likely a consequence of the kinetic factors controlling the binding of each ruthenium pentaammine residue.

The ^1H NMR spectra of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ complexes with $n=0, 3, 6$ are shown in Fig. 5.A-C. For the $[\text{Ru}(\text{bpz})_3]^{2+}$ complex, the resonance signals observed at 10.05, 8.85 and 8.17 ppm were previously assigned to the H3, H5 and H6 protons, respectively. The NMR spectrum of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ complex does not seem to differ appreciably from the spectrum of the $[\text{Ru}(\text{bpz})_3]^{2+}$ cation. Given the close similarity in the spectra

shown in Fig. 5A and B, and the small changes seen in the $\text{bpz}[\text{Ru}(\text{NH}_3)_5]_n$ system ($n=1,2$), with peripheral coordination, the assignment outlined in Table 2 is proposed. Any other assignment requires a substantial downfield shift in H6 or H6' which, again by analogy with the data in Fig. 4, seems unlikely. In the case of the heptanuclear complex, $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_6]^{14+}$ the resonance signals at 9.96 (singlet), 8.83 (doublet) and 7.65 ppm (doublet) can be ascribed to the H3, H3'; H5, H5' and H6, H6' protons, respectively.

Electrochemistry Cyclic voltammograms of the $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ complexes can be seen in Fig. 6. The voltammograms of the mononuclear complex follow very closely the several reversibility criteria based on the treatment of Nicholson and Shain,²⁶ with $E_0 = 0.52$ V versus NHE and a diffusion coefficient of $2.5 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$. The observed Ru(III)/Ru(II) potential is in the expected region.²⁷ In the case of the binuclear complex, only a single pair of cathodic and anodic waves is observed with a peak separation of 70 mV, and $E_0 = 0.53$ V. The voltammograms for the mononuclear and binuclear complexes normalised to constant [Ru] are shown in Fig. 6 and are very similar; however, the current profile for the binuclear species is broader than for the mononuclear species. As anticipated, the Faradaic current for the bis-ruthenium species, is twice as large as for the mono-ruthenium species. The voltammograms of the binuclear complex can be simulated by a sum of two mono-electronic reversible waves, separated by approximately 0.01 V.

Spectroelectrochemical oxidation of the binuclear species proceeds as shown in Fig. 7.A with a formal potential of 0.52 V, leading to complete decay of the two $\text{Ru(II)}_p \longrightarrow \pi^* \text{ bpz MLCT}$ bands in the visible region. The Nernst slope of 70 mV is slightly higher than that expected for a reversible mono-electronic

process (e.g. 59 mV) and is consistent with the occurrence of two successive steps having very similar E^0 . Measurements of the peripherally oxidised species in the near-ir region (800-1800 nm), in D_2O solutions, provided no evidence of low energy intervalence transfer (IT) transitions with $\epsilon > 100 \text{ M}^{-1} \text{ cm}^{-1}$.

Cyclic voltammograms of the $[\text{Ru}(\text{bpz})_3\text{Ru}(\text{NH}_3)_5]^{4+}$ complex in aqueous solution are shown in Fig. 8.A. The voltammograms are consistent with a reversible mono-electronic process, with $E^0 = 0.69 \text{ V}$ and $D = 2.3 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$. In the cases of the 1:3 and 1:6 polynuclear complexes, a pair of broad, quasi-reversible waves can be observed around 0.7 V in Fig. 8.B.C. The cathodic and anodic peaks are separated by almost 150 mV, having approximately 1:1 intensity ratio. The peak potentials do not vary with the potential scan rates in the range of 20-200 mV s^{-1} . It seems probable that the large separation between the cathodic and anodic waves arises from the superimposition of several parallel mono-electronic processes having very close E^0 values. A quantitative simulation can be made, as in Fig. 8. B, C, by assuming a set of three reversible waves for the 1:3 complex, with $E^0 = 0.67, 0.72$ and 0.77 V , ($D = 2.0 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$), and a set of six reversible waves for the 1:6 complex, with $E^0 = 0.62, 0.66, 0.70, 0.74, 0.78$ and 0.82 V ($D = 1.4 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$).

The spectroelectrochemistry of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ and $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_6]^{14+}$ complexes in aqueous solution is illustrated in Fig. 7.B,C. The oxidation process leads to a decay of the $\text{Ru}_p \rightarrow \text{bpz}(\pi_1^*, \pi_2^*)$ bands in the visible region, indicating that the peripheral $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups are totally oxidized within a narrow range of potentials around 0.7 V. This result confirms the occurrence of the several parallel steps in the cyclic voltammograms. The spectra can be regenerated in more than 90% and 70% yield for the 1:3 and 1:6 complexes, respectively, by reduction at a

potential just negative of 0.7V. As expected, the corresponding Nernst slopes of 100 and 130 mV differ from that of a reversible mono-electronic process, and are consistent with a series of parallel mono-electronic steps having very close E^0 values.

The electronic spectra of the $[\text{Ru(II)(bpz)}_3\{\text{Ru(III)(NH}_3)_5\}_n]^{2+3n}$ complexes resemble that of the $[\text{Ru(bpz)}_3]^{2+}$ cation, with an additional absorbance increase around 360 nm probably associated with LMCT charge-transfer transitions in the (N-heterocyclic) pentaammine-ruthenium(III) groups.^{28,29}

Spectroelectrochemical measurements in the near-IR region (800-1800 nm), in D_2O solutions, provided no evidence of intervalence bands with $\epsilon > 100 \text{ M}^{-1} \text{ cm}^{-1}$. The possibility that the IT band is associated with the residual absorption around 700-800 nm, observed spectroelectrochemically, cannot be excluded. However, considering that minor, but detectable amounts of decomposed oxidized products are always present in these systems, any assignment would be too speculative at the present stage.

Cyclic voltammograms in acetonitrile solutions are shown in Fig. 9. The quasi-reversible waves associated with the $[\text{Ru(NH}_3)_5] \text{ (III)/(II)}$ couples are similar to those measured in aqueous solution, except for an anodic shift of approximately 200 mv which can be ascribed to solvent effects.²⁷ The several waves observed at -0.5, -0.65 and -0.9 V versus NHE for the $[\text{Ru(bpz)}_3\text{Ru(NH}_3)_5]^{4+}$ complex are similar to those previously reported for the $[\text{Ru(bpz)}_3]^{2+}$ cation,³ and can be ascribed to the successive reduction of the bpz ligands. The electrochemical behavior becomes more complicated in the case of the 1:3 and 1:6 complexes, as shown in Fig. 9.B-C.

However this behaviour is, in part, due to severe adsorption problems.

Concluding Remarks

The polynuclear $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ complexes represent an interesting, cubic symmetry version of the linear array $[(\text{NH}_3)_5\text{Ru}(\text{pzRu}(\text{NH}_3)_4\text{pz})_n-\text{Ru}(\text{NH}_3)_5]^{4+2n}$ ($n=2,3,4$) complexes studied by von Kameke, Tom and Taube.³⁰ However, in contrast with the extended interactions observed in the linear systems, the peripheral $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups seem to be relatively isolated in the cubic complex. The energies of the MLCT bands and the electrochemical potentials for the $\text{Ru(III)}/\text{Ru(II)}$ couples vary only slightly with the number of $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups. No evidence of IT bands with $\epsilon > 100 \text{ M}^{-1} \text{ cm}^{-1}$ have been detected in the near IR region, in contrast with the linear pyrazine bridged systems and with Creutz and Taube's ion^{31,32} $[(\text{NH}_3)_5\text{Ru}-\text{pz}-\text{Ru}(\text{NH}_3)_5]^{5+}$. It is evident that up to six pentaamine-ruthenium(II) (or pentacyanoferrate(II)¹¹) groups may be coordinated by the $[\text{Ru}(\text{bpz})_3]^{2+}$ cation and that interaction between these groups is minimal. Surprisingly, the central $\text{Ru(II)}_c \longrightarrow \pi^* \text{ bpz}$ transition is also unaffected by such peripheral coordination, even when all the pentaamine groups are oxidised to Ru(III) . This is contrary to the situation where the bpz rings are protonated,^{3b} in which case there is an effect upon the $\text{Ru(II)}_c \longrightarrow \pi^* \text{ bpz}$ transition energy. Lavalley and Fleischer²³ noted that the MLCT transitions of $\text{LRu}(\text{NH}_3)_5$ species (L = a heterocyclic ligand) were linearly related to the redox potential $E(L/L^-)$ of the ligand L , where reduction occurs into the same orbital into which the electronic transition occurs. If we regard the unit $[\text{Ru}(\text{bpz})_3]^{2+}$ as a ligand towards $[\text{Ru}(\text{NH}_3)_5]^{2+}$, then we may add this datum point to the same graph, using the reduction potential for $[\text{Ru}(\text{bpz})_3]^{2+}$ instead of that for a simple ligand L . This extra datum point lies exceptionally well on the same line as the other ligands L but at very much more positive potential. The equation of this line for 11 points is:

$$\nu(\text{CT}) = 0.51 E(L/L^-) + 1.59 \text{ (in eV)} [R = 0.98]$$

It is surely of considerable interest that the electronic spectra of these species are almost independent of the degree of peripheral binding, that each ruthenium pentaammine (or pentacyanoferrate) fragment is independent of each other, and further that the central ruthenium to bpz MLCT transition is independent of the extent of peripheral binding, even when the $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups are oxidised to $[\text{Ru}(\text{NH}_3)_5]^{3+}$. Recall that this is not true for binding of protons.^{3b} Expressed in another way, the several $[\text{Ru}(\text{NH}_3)_5]^{2+}$ groups generate several MLCT excited states which do not couple to each other, nor to the central $[\text{Ru}(\text{III})_c(\text{bpz}^-)]$ excited MLCT state, at least on the time scale of absorption. Further, the $[\text{Ru}(\text{III})_p/\text{Ru}(\text{II})_p]$ couples appear identical, except for statistical effects, independent of the number of peripheral groups. Thus the ground state wavefunctions involving these groups are localized on each pyrazine ring. These results have considerable significance with respect to our understanding of the nature of charge transfer states, and their localisation and delocalisation and emission. Possibly each $\text{Ru}(\text{III})_p \longrightarrow \pi^*(\text{bpz})$ transition is located within its own pyrazine ring and the transition is orthogonal to the central $\text{Ru}(\text{II})_c \longrightarrow \pi^*(\text{bpz})$ transition which, possibly, is localised in the diimine framework. Note that such diimine localisation is consistent with the lack of a significant increase in the pK_a value of the nitrogen atom trans to the ruthenium atom in the complexes $\text{bpz}[\text{Ru}(\text{NH}_3)_5]_n$ ($n=1,2$).

Acknowledgement

Financial support from the FINEP and CNPq agencies in Brazil (HET), the Natural Science and Engineering Research Council (Ottawa), and the Office of Naval Research (Washington) are gratefully acknowledged.

References

1. (a) Sao Paolo; (b) York University. (c) Current address: Department of Chemistry, University College, Singleton Park, Swansea, Wales, SA2 8PP, Great Britain.
2. Crutchley, R.J.; Lever, A.B.P., J. Am. Chem. Soc., 1980, 102, 7128.
3. (a) Crutchley, R.J.; Lever, A.B.P., Inorg. Chem., 1982, 21, 2276.
(b) Crutchley, R.J.; Kress, N.; Lever, A.B.P., J. Am. Chem. Soc., 1983, 105, 1170.
4. Rillema, D.P.; Allen, G.; Meyer, T.J.; Conrad, D. Inorg. Chem. 1983, 22, 1617.
5. Crutchley, R.J., Lever, A.B.P.; Poggi, A., Inorg. Chem., 1983, 22, 2647.
6. Gonzales-Velasco, J.; Rubinstein, I.; Crutchley, R.J., Lever, A.B.P.; Bard, A.J., Inorg. Chem., 1983, 22, 822.
7. Dodsworth, E.S.; Lever, A.B.P., Chem. Phys. Lett., 1984, 112, 567.
8. Dodsworth, E.S.; Lever, A.B.P.; Eryavec, G; Crutchley, R.J., Inorg. Chem., 1985, 24, 1906.
9. Haga, M.-A., Dodsworth, E.S., Eryavec, G., Seymour, P.; Lever, A.B.P., Inorg. Chem., 1985, 24, 1901.
10. Balk, W.R.; Stufkens, D.J., Crutchley, R.J.; Lever, A.B.P., Inorg. Chim. Acta., 1982, 64, L49-50.
11. Toma, H.E.; Lever, A.B.P., Inorg. Chem., 1986, 25, 176.
12. Lafferty, J.J.; Case, F.H., J. Org. Chem., 1967, 32, 1591.
13. Lever, A.B.P.; Lewis, J.; Nyholm, R.S., J. Chem. Soc., 1964, 1187.
14. Kuehn, C.G.; Taube, H., J. Am. Chem. Soc., 1976, 98, 689.
15. Kratochvil, B; Lorah, E.; Garber, C., Anal. Chem., 1969, 41, 1793.
16. Ford, P.; Rudd, D.F.P.; Gaunder, R; Taube, H., J. Am. Chem. Soc., 1968, 90, 1187.
17. Ford, P.C.; Kuempel, J.R.; Taube, H., Inorg. Chem., 1968, 7, 1976.

18. Toma, H.E.; Stadler, E., Inorg. Chem. 1985, 24, 3085.
19. Johnson, C.R.; Shepherd, R.E., Inorg. Chem., 1983, 22, 1117.
20. Jackson, G.; Porter, G., Proc. Royal Soc. London, Ser. A., 1961, 260, 13.
21. Long, F.A.; Paul, M.A., Chem. Rev., 1957, 57, 1.
22. Toma, H.E.; Santos, P.S.; Lever, A.B.P., to be submitted to Inorg. Chem.
23. Lavalley, D.K.; Fleischer, E.B., J. Am. Chem. Soc., 1972, 94, 2583.
24. Foust, R.D.; Ford, P.C., J. Am. Chem. Soc., 1972, 94, 5686.
25. Lytle, F.E.; Petrosky, L.M.; Carlson, L.R., Anal. Chim. Acta, 1971, 57, 239.
26. Nicholson, R.S.; Shain, I., Anal. Chem., 1964, 36, 706.
27. Curtis, J.C.; Sullivan, B.P.; Meyer, T.J., Inorg. Chem., 1983, 22, 224.
28. Gaunter, R.; Taube, H., Inorg. Chem. 1970, 9, 2627.
29. Lever, A.B.P., "Inorganic Electronic Spectroscopy", 2nd Edn., Elsevier, Amsterdam, 1984.
30. von Kameke, A.; Tom, G.M.; Taube, H. Inorg. Chem., 1978, 17, 1790.
31. Creutz, C.; Taube, H., J. Am. Chem. Soc., 1969, 91, 3988.
32. Creutz, C.; Taube, H., J. Am. Chem. Soc., 1973, 95, 1086.

Figure 7

Spectroelectrochemistry of (A) $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ (1.3 mM) (B) $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ (1.4 mM) and (C) $[\text{Ru}(\text{bpz})_3[\text{Ru}(\text{NH}_3)_5]_6]^{14+}$ (0.4 mM, in aqueous solution, 0.10 M NaCl, 25°C: (a) 0 (b) 0.48, (c) 0.52, (d) 0.56, (e) 0.70, (f) 0, (g) 0.62, (h) 0.67, (i) 0.72, (j) 0.85, (k) 0, (l) 0.57, (m) 0.62, (n) 0.67, (o) 0.72, (p) 0.90 V vs NHE.

Figure 8

Cyclic voltammograms of the (A) 1:1, (B) 1:3 and (C) 1:6 tris(bipyrazine)-ruthenium(II)-pentaammineruthenium(II) polynuclear complexes, 0.8, 1.0, and 0.6 mM, respectively, in aqueous solution, 25°C, 0.10 M NaCl. Dotted lines refer to tentative deconvolution of the voltammograms in a series of reversible mono-electronic waves, black circles = theoretical values.

Figure 9

Cyclic voltammograms of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ complexes with (A) $n=1$, (B) $n=3$ and (C) $n=6$, in acetonitrile (saturated solution) in the presence of 0.10 M tetraethylammonium hexafluorophosphate, 25°C.

Figure Caption

Figure 1

Electronic spectra of (A) $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and (B) $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ (0.05 mM) in (a,d) water (pH 7), (b,e) 0.12 M HCl and (c,f) 3 M HCl.

Figure 2

Electronic spectra of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_n]^{2+2n}$ polynuclear complexes in aqueous solution with $n=1$ (A), 3 (B) and 6 (C), around 0.037, 0.033 and 0.017 mM, respectively.

Figure 3

(A) Electronic spectra of the $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ complex (0.033 mM) in (a) water (pH 7), (b) 5M HCl, (c) 8 M HCl, (d) 12 M HCl, and (B) in the presence of (e) 0, (f) 1, (g) 2 and (h) 3 equivalents of the aquapentacyano-ferrate(II) ion.

Figure 4

^1H NMR spectra of (A) bpz, (B) $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ and (C) $[(\text{NH}_3)_5\text{Ru}-\text{bpz}-\text{Ru}(\text{NH}_3)_5]^{4+}$, 0.03-0.05 M in D_2O , 30°C. (Chemical shifts versus TMS).

Figure 5

^1H NMR spectra of (A) $[\text{Ru}(\text{bpz})_3]^{2+}$, (B) $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_3]^{8+}$ and (C) $[\text{Ru}(\text{bpz})_3\{\text{Ru}(\text{NH}_3)_5\}_6]^{14+}$ 0.03 M in D_2O , 30°C. (Chemical shifts versus TMS).

Figure 6

Cyclic voltammograms of $[\text{Ru}(\text{NH}_3)_5(\text{bpz})]^{2+}$ (...) (3 mM) and $[\{\text{Ru}(\text{NH}_3)_5\}_2(\text{bpz})]^{4+}$ (—) (1.5 mM) at several potential scan rates, in aqueous solution, 0.10 M NaCl, 25°C.




Figure 2





Figure 3






Figure 3




Figure 5




Figure 6




Figure 5




Fig. 8




Figure 1

DL/1113/87/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Dauda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 92132	1

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Manfred Breiter
Institut für Technische Elektrochemie
Technischen Universität Wien
9 Getreidemarkt, 1160 Wien
AUSTRIA

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. R. Sutula
The Electrochemistry Branch
Naval Surface Weapons Center
Silver Spring, Maryland 20910

Dr. R. A. Marcus
Department of Chemistry
California Institute of Technology
Pasadena, California 91125

Dr. J. J. Auborn
AT&T Bell Laboratories
600 Mountain Avenue
Murray Hill, New Jersey 07974

Dr. M. S. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. Mel Miles
Code 3852
Naval Weapons Center
China Lake, California 93555

Dr. P. P. Schmidt
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Dr. Roger Belt
Litton Industries Inc.
Airtron Division
Morris Plains, NJ 07950

Dr. Ulrich Stimming
Department of Chemical Engineering
Columbia University
New York, NY 10027

Dr. Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Michael J. Weaver
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. R. David Rauh
EIC Laboratories, Inc.
Norwood, Massachusetts 02062

Dr. Edward M. Eyring
Department of Chemistry
University of Utah
Salt Lake City, UT 84112

Dr. M. M. Nicholson
Electronics Research Center
Rockwell International
3370 Miraloma Avenue
Anaheim, California

Dr. Nathan Lewis
Department of Chemistry
Stanford University
Stanford, California 94305

Dr. Hector D. Abruna
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. B. P. Lever
Chemistry Department
York University
~~Downsview~~, Ontario M3J 1P3

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5H UNITED KINGDOM

Dr. John Wilkes
Department of the Air Force
United States Air Force Academy
Colorado Springs, Colorado 80840-6528

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. Steven Greenbaum
Department of Physics and Astronomy
Hunter College
695 Park Avenue
New York, New York 10021

Dr. Donald Sandstrom
Boeing Aerospace Co.
P.O. Box 3999
Seattle, Washington 98124

Mr. James R. Moden
Naval Underwater Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. D. Rolison
Naval Research Laboratory
Code 6171
Washington, D.C. 20375-5000

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick
Department of Chemistry
The University of Southampton
Southampton, SO9 5NH UNITED KINGDOM

Dr. Edward Fletcher
Department of Mechanical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. Richard Pollard
Department of Chemical Engineering
University of Houston
Houston, Texas 77004

Dr. M. Philpott
IBM Research Division
Mail Stop K 33/801
San Jose, California 95130-6099

Dr. Martha Greenblatt
Department of Chemistry, P.O. Box 939
Rutgers University
Piscataway, New Jersey 08855-0939

Dr. Anthony Sammells
Eltron Research Inc.
4260 Westbrook Drive, Suite 111
Aurora, Illinois 60505

Dr. C. A. Angell
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymers Division
National Bureau of Standards
Gaithersburg, Maryland 20899

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Stanislaw Szpak
Naval Ocean Systems Center
Code 633, Bayside
San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. John Fontanella
Department of Physics
U.S. Naval Academy
Annapolis, Maryland 21402-5062

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Joel Harris
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Dr. J. O. Thomas
University of Uppsala
Institute of Chemistry
Box 531 Baltimore, Maryland 21218
S-751 21 Uppsala, Sweden

Dr. John Owen
Department of Chemistry and
Applied Chemistry
University of Salford
Salford M5 4WT UNITED KINGDOM

Dr. O. Stafsudd
Department of Electrical Engineering
University of California
Los Angeles, California 90024

Dr. Boone Owens
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Albert R. Landgrebe
U.S. Department of Energy
M.S. 68025 Forrestal Building
Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. Milos Novotny
Department of Chemistry
Indiana University
Bloomington, Indiana 47405

Dr. Mark A. McHugh
Department of Chemical Engineering
The Johns Hopkins University
Baltimore, Maryland 21218

Dr. D. E. Irish
Department of Chemistry
University of Waterloo
Waterloo, Ontario, Canada
N2L 3G1

Q/1113/87/2

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Henry S. White
Department of Chemical Engineering
and Materials Science
151 Amundson Hall
421 Washington Avenue, S.E.
Minneapolis, Minnesota 55455

Dr. Daniel A. Buttry
Department of Chemistry
University of Wyoming
Laramie, Wyoming 82071

Dr. W. R. Fawcett
Department of Chemistry
University of California
Davis, California 95616

Dr. Peter M. Blonsky
Eveready Battery Company, Inc.
25225 Detroit Road, P.O. Box 45035
Westlake, Ohio 44145

ABSTRACTS DISTRIBUTION LIST, 051A

Dr. M. A. El-Sayed
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. E. R. Bernstein
Department of Chemistry
Colorado State University
Fort Collins, Colorado 80521

Dr. J. R. MacDonald
Chemistry Division
Naval Research Laboratory
Code 6110
Washington, D.C. 20375-5000

Dr. G. B. Schuster
Chemistry Department
University of Illinois
Urbana, Illinois 61801

Dr. J. B. Halpern
Department of Chemistry
Howard University
Washington, D.C. 20059

Dr. M. S. Wrighton
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. W. E. Moerner
I.B.M. Corporation
Almaden Research Center
650 Harry Rd.
San Jose, California 95120-6099

Dr. A. B. P. Lever
Department of Chemistry
York University
Downsview, Ontario
CANADA M3J1P3

Dr. George E. Walrafen
Department of Chemistry
Howard University
Washington, D.C. 20059

Dr. Joe Brandelik
AFWAL/AADO-1
Wright Patterson AFB
Fairborn, Ohio 45433

Dr. Carmen Ortiz
Consejo Superior de
Investigaciones Cientificas
Serrano 121
Madrid 6, SPAIN

Dr. Kent R. Wilson
Chemistry Department
University of California
La Jolla, California 92093

Dr. G. A. Crosby
Chemistry Department
Washington State University
Pullman, Washington 99164

Dr. Theodore Pavlopoulos
NOSC
Code 521
San Diego, California 91232

Dr. John Cooper
Code 6173
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Joseph H. Boyer
Department of Chemistry
University of New Orleans
New Orleans, Louisiana 70148

Dr. Harry D. Gafney
Department of Chemistry
Queens College of CUNY
Flushing, New York 11367-0904

DATE
L MED
8