



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU - MANDARDS .964 A

8

Proceeding and the second

| Q | | PEROPE DOCU | MENTATION | PAGE | | | |
|---|--|--|---|--|--|--|--|
| 2 | | | 16. RESTRICTIVE | MARKINGS | JIII. | FILF | |
| ~ | CLASSIFICATION AUT OF | 3. DISTRIBUTION / AVAILABILITY OF REPORT | | | | | |
| 19 | -ICATION / DOWNGRADIN CHE | FED 1 8 1988 | | Approved for public release Distribution is unlimited | | | |
| A- | NG ORGANIZATION REPORT NUMBER(S) al Report No. 2 | | S. MONITORING ORGANIZATION REPORT NUMBER(S) | | | | |
| | PERFORMING ORGANIZATION | 6b. OFFICE SYMBOL | 7a. NAME OF MC | DNITORING ORG | ANIZATION | | |
| 4 | t. Chem. Eng | (ir applicable) | ONR | | | | |
| | (City, State, and ZIP Code) | <u>↓</u> | 7b. ADDRESS (Cit | y, State, and ZIF | P Code) | | |
| | ge, MA 02139 | | 800 N. Qui Arlington, | ncy St. VA 22217 | | | |
| 3a. NAME ORGAI | OF FUNDING / SPONSORING | 8b. OFFICE SYMBOL (If applicable) | 9. PROCUREMENT | -0517 | DENTIFICATION | NUMBER | |
| UTT | 1CE OI Naval Research | | | UNDING NUMBE | RS | | |
| 800 1-1 | N. Quincy Street | | PROGRAM ELEMENT NO. | PROJECT NO. | TASK NO. | | |
| 12. PERSO P. | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT | E. Cohen E COVERED | 14. DATE OF REPO | RT (Year, Month | n, Day) 15. PAG | GE COUN | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT I35. TIME hnical Report FROM_ EMENTARY NOTATION | E. Cohen E COVERED TO | 14. DATE OF REPO January 28 | RT (Year, Month , 1988 | n, Day) 15. PAC | GE COUN 19 | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT I35. TIME Anical Report FROM_ EMENTARY NOTATION pared for publication i | E. Cohen E COVERED TO In Macromolecules | 14. DATE OF REPO January 28 | RT (Year, Month , 1988 | n, Day) 15. PAC | GE COUM 19 | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELD | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP | E. Cohen E COVERED TO In Macromolecules 18. SUBJECT TERMS Block Copoly | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme | RT (Year, Month , 1988 e if necessary an r Chain Con | n, Day) 15. PAC nd identify by b nfiguration | GE COUN 19 block num | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC 19. ABSTR | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. | E. Cohen E COVERED TOTO In Macromolecules 18. SUBJECT TERMS Block Copol ary and identify by block | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) | RT (Year, Month , 1988 e if necessary an r Chain Cor | n, Day) 15. PAG nd identify by b nfiguration | GE COUN 19 block num | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene polyb ttering (SANS) experime | E. Cohen E COVERED TO | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer | RT (Year, Month , 1988 e if necessary and r Chain Cor e examined | n, Day) 15. PAG nd identify by b nf iguration in small a | GE COUN 19 block num 1 ingle 1 | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELD 19. ABSTR sca end | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP GROUP SUB-GROUP Two polystyrene polyb ttering (SANS) experime s inside the polybutadi | E. Cohen E COVERED TO IN Macromolecules 18. SUBJECT TERMS Block Copoly ary and identify by block outadiene block co ents designed to p ene microspheres | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or | n, Day) 15. PAC nd identify by b nfiguration in small a ne polybuta ne of the c | GE COUN 19 block num n ingle n idiene copolyt | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELO 19. A8STR sca end SBm and | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene polyb ttering (SANS) experime s inside the polybutadi , was phase-matched; th normal butadiene monom | E. Cohen E COVERED TO In Macromolecules II. SUBJECT TERMS Block Copoly any and identify by block butadiene block co ents designed to p ene microspheres the B moiety was sy ther. The other co | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer. SBm | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair | n, Day) 15. PAG nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short n | GE COUN 19 block num ingle m idiene copolytice of co perdent | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELO 19. ABSTR sca end SBm and poly | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene polyb ttering (SANS) experime s inside the polybutadi , was phase-matched; th normal butadiene monom vbutadiene sequences at | E. Cohen E COVERED TO In Macromolecules Block Copol: ary and identify by block butadiene block co ents designed to p ene microspheres the moiety was sy the other co the end of the p | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair block of e | n, Day) 15. PAG nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short p each of the | diene opolytice of d engle diene opolytice of d erdeu | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC 19. ABSTR 19. ABSTR 19. ABSTR and poly Cons by S | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report I3b. TIME FROM | E. Cohen E COVERED TO IN Macromolecules 18. SUBJECT TERMS Block Copol ary and identify by block outadiene block co ents designed to p ene microspheres the B molety was sy the coher co the end of the p intensities and the Bd chain ends a | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene i the form of are essential | RT (Year, Month , 1988 e if necessary au r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair block of e the scatte ly uniform | nd identify by b nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte ly distribu | GE COUN 19 block nur ingle i ingle i i ingle i i i i i i i i i i i i i i | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELD 19. ABSTR 19. ABSTR | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene-polyb ttering (SANS) experimes inside the polybutadi , was phase-matched; th normal butadiene monom ybutadiene sequences at sideration of scatterin SBmBd indicated that th the volume of the sphe | E. Cohen E COVERED TO In Macromolecules 18. SUBJECT TERMS Block Copoly any and identify by block butadiene block co ents designed to p ene microspheres the moiety was sy her. The other co the end of the p intensities and the Bd chain ends a tres. A model in | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene l the form of are essential which the Bd | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of the cimens. Or ing a suita Bd, contain block of e the scatte ly uniformi ends clust | n, Day) 15. PAG nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte by distribu | angle diene opolytie of o erdeu mole rn exi ted t | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC 19. ABSTR sca end SBm and poly Cons by S out sphe | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT Inical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene polyb ttering (SANS) experime s inside the polybutadi , was phase-matched; th normal butadiene monom vbutadiene sequences at sideration of scatterin SBmBd indicated that th the volume of the sphe eres failed to reproduc | E. Cohen E COVERED TO IN Macromolecules 18. SUBJECT TERMS Block Copol ary and identify by block butadiene block co ents designed to p ent designed to p ent designed to p ent of the p intensities and the end of the p intensities and the sent of the p intensities and the end of the p intensities and the sent of the p intensities and the end of the p intensities and the sent of the p intensities and t | 14. DATE OF REPC January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene i the form of are essential which the Bd features of t | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair block of e the scatte ly uniformal ends clust he observed | n, Day) 15. PAG nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte ly distribu ter at the i SANS patt | angle diene opolytie erdeu ern exited ti core of ern. | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELD 19. A8STR 19. A8STR 19. ASSTR 19. A | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT hnical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene-polyb ttering (SANS) experimes s inside the polybutadi , was phase-matched; th normal butadiene monom ybutadiene sequences at sideration of scatterin SBmBd indicated that th the volume of the sphe eres failed to reproduc | E. Cohen E COVERED TO In Macromolecules 18. SUBJECT TERMS Block Copoly any and identify by block butadiene block co ents designed to p ene microspheres the moiety was sy her. The other co the end of the p intensities and the end of the p intensities and the essential f | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene I the form of are essential which the Bd features of t | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of the cimens. Or ing a suita Bd, contain block of e the scatte ly uniform ends clust he observed | n, Day) 15. PAG nd identify by b nfiguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte by distributer at the sANS patt | angle : ingle | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 20. DISTR 20. DISTR | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT ISD. TIME FROM EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene-polyb ttering (SANS) experime s inside the polybutadi , was phase-matched; th normal butadiene monom vbutadiene sequences at sideration of scatterin SBmBd indicated that th the volume of the sphe eres failed to reproduc | E. Cohen E COVERED TO TO IN Macromolecules 18. SUBJECT TERMS Block Copol: ary and identify by block butadiene block co ents designed to p ene microspheres the end of the p intensities and the end of the p intensities and the end of the p intensities and the essential f | 14. DATE OF REPC January 28 (Continue on revers ymers, Polyme number) opolymers wer probe the loc of these spe ynthesized us opolymer, SBm polybutadiene i the form of are essential which the Bd features of t | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair block of e the scatte ly uniform ends clust he observed | n, Day) 15. PAG nd identify by b nf iguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte ly distribu ter at the 1 SANS patt | angle diene opolytice of o erdeu mole trn exited ti core o ern. | |
| 12. PERSO P. 13a. TYPE Tec 16. SUPPL Pre 17. FIELC 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 19. ABSTR 20. DISTR 20. DISTR 20. DISTR 22a. NAM | NAL AUTHOR(S) Cheng, C.V. Berney, R.E OF REPORT Inical Report EMENTARY NOTATION pared for publication i COSATI CODES GROUP SUB-GROUP ACT (Continue on reverse if necess. Two polystyrene polyb ttering (SANS) experime s inside the polybutadi , was phase-matched; th normal butadiene monom ybutadiene sequences at sideration of scatterin SBmBd indicated that th the volume of the sphe eres failed to reproduc IBUTION/AVAILABILITY OF ABSTRA CLASSIFIED/UNLIMITED SAME A SECOR RESPONSIBLE INDIVIDUAL | E. Cohen E COVERED TO IN Macromolecules IS. SUBJECT TERMS Block Copol any and identify by block butadiene block co ents designed to p ene microspheres the end of the p is function of the | 14. DATE OF REPO January 28 (Continue on revers ymers, Polyme probe the loc of these spe ynthesized us polymer, SBm polybutadiene i the form of are essential which the Bd features of t 21. ABSTRACT SE Unclassi 22b. TELEPHONE | RT (Year, Month , 1988 e if necessary and r Chain Con e examined ation of th cimens. Or ing a suita Bd, contair block of e the scatte ly uniform ends clust he observed | n, Day) 15. PAC nod identify by b infiguration in small a ne polybuta ne of the c able mixtur ned short p each of the ering patte ly distribu cer at the d SANS patt | dick nur angle : angle : adiene opolyn e of co ern exi core co ern exi core co ern. | |

OFFICE OF NAVAL RESEARCH Contract N00014-87-K-0517 R&T Code 4132001---02

TECHNICAL REPORT NO. 2

Distribution of Chain Ends Inside the Polybutadiene Microspheres of SB Diblock Copolymers

by

P. Cheng, C.V. Berney and R. E. Cohen
 Department of Chemical Engineering
 Massachusetts Institute of Technology
 Cambridge, Massachusetts 02139

Prepared for Publication in Macromolecules

January 28, 1988

Reproduction in whole or in part is permitted for any purpose of the U.S. government.

This document has been approved for public release and sale; its distribution is unlimited.

| Accesion For | | | | | |
|----------------|-------------|-----------|--------------------|--|--|
| NTIS | CRA&I | N. | | | |
| DTIC | TAB | | - 1 | | |
| Unanas | enced. | | | | |
| j Justific | 4100 | | | | |
| By East ib: | tien/ | | امہ ۔۔۔ ا ام | | |
| ,b | Vithal Ling | a nanga s | | | |
| 1 to 1 | A ta da | i of | | | |
| 11463 | - 11 C | *** | | | |
| A-1 | | | | | |
| | i i | | | | |

88 2 17 029

Distribution of Chain Ends Inside the Polybutadiene Microspheres of SB Diblock Copolymers

> P. Cheng¹, C.V. Berney and R.E. Cohen* Department of Chemical Engineering
> Massachusetts Institute of Technology
> Cambridge, Massachusetts 02139 USA

<u>Abstract</u>

Two polystyrene-polybutadiene block copolymers were examined in small angle neutron scattering (SANS) experiments designed to probe the location of the polybutadiene chain ends inside the polybutadiene microspheres of these specimens. One of the copolymers, SBm, was phase-matched; the B moiety was synthesized using a suitable mixture of deutero and normal butadiene monomer. The other copolymer, SBmBd, contained short perdeuterated polybutadiene sequences at the end of the polybutadiene block of each of the molecules. Consideration of scattering intensities and the form of the scattering pattern exhibited by SBmBd indicated that the Bd chain ends are essentially uniformly distributed throughout the volume of the spheres. A model in which the Bd ends cluster at the core of the spheres failed to reproduce the essential features of the observed SANS pattern.

¹ DuPont Company, Parkersburg, West Virginia 26181

Introduction

Diblock copolymers of polystyrene (S) and polybutadiene (B) which display a bulk morphology of B spheres in S matrix have been examined at various levels of structural detail in our laboratory. Among the important features characterized in our previous work using small angle neutron scattering (SANS) experiments were the BCC packing of the microspheres (1), the relationship of sphere size to B molecular weight (2,3), the nature of the interfacial zones between the B spheres and the S matrix (2), and the effective radius of gyration of the B chain segments of the diblock copolymer (4,5). Recently we reported on the distribution of S homopolymer added to the S matrix of the diblock (6).

In this paper we address the following question: Where are the ends of the B chain segments located inside the microspheres? Two possible answers to this question are (a) The ends are essentially uniformly distributed throughout the B spheres; (b) The polybutadiene chain ends are preferentially clustered toward the center of each sphere. Figure 1 demonstrates that it should be relatively easy to discriminate between these two first-order hypotheses if an appropriate pair of diblock copolymers were available for SANS experiments. The necessary pair of copolymers is a phase-matched (7) diblock copolymer, SBm, and a triblock copolymer, SBmBd, in which the first two blocks S and Bm are identical to the phase-matched diblock and the third, very short endblock, Bd, is a perdeuterated moiety. In the phase-matched diblock the polybutadiene block, Bm, is a random copolymer consisting of an isotopic mixture of proteo- and perdeutero-butadiene repeat units, the ratio of which is carefully selected (7, 8) to exactly match the neutron scattering length density of polystyrene. This ideal diblock produces no structural scattering in a SANS experiment as suggested by Figure 1a. However, the triblock containing the short perdeuterated endblocks will yield substantial amounts of structual scattering because the phase-matching is no longer preserved. The detailed nature of the SANS scattering pattern observed for SBmBd will depend dramatically upon the spatial distribution of the Bd chain ends inside the microspheres; thus SANS experiments can be used to determine whether either Figure 1b or Figure 1c is a reasonable representation of the chain end distribution in the SBmBd copolymer.

In what follows we describe the synthesis and molecular characterization of copolymers SBm and SBmBd and the examination of these copolymers using SANS experiments and transmission electron microscopy (TEM). Results of small angle

X-ray scattering (SAXS) experiments, which are described in detail elsewhere (8), are also used to support the main conclusions drawn from SANS and TEM results.

ACCURATE PARTICLES INTRODUCE RECEASE

Experimental

Polymer Synthesis (8)

High vacuum anoinic polymerization techniques, similar to those reviewed by Morton and Fetters (9), were employed in the synthesis of the block copolymers. A three-stage Kontes/Martin silicone oil diffusion pump facilitated vacuum conditions in the range of 10^{-5} Torr in the primary manifold.

Benzene was stored over fresh, concentrated sulfuric acid (3/1 volume ratio of benzene/acid) for several months during which time the bottom acid layer gradually turned dark brown, indicating that some polar impurities had been oxidized. Acid-treated benzene was washed twice with distilled water, neutralized by washing with dilute sodium bicarbonate solution followed by two washings with distilled water, and finally dried over anhydrous magnesium sulfate. The benzene was further dried by calcium hydride and the addition of a sodium dispersion. The appearance of silvery flecks of sodium indicated that the benzene was dry. It was then distilled on the vacuum line into a storage flask containing poly(styryl) lithium (PS-Li). The retention of the orange-red color of the polystyryl anion indicated that the solvent contained no impurities which would terminate living polymer chains.

Styrene was stirred on the vacuum line with a slurry of ground calcium hydride for several days with periodic degassing. The styrene was then distilled into a flask containing 5 ml of dibutylmagnesium (73 wt% in heptane) and stirred for several hours at room temperature: the solution turned bright yellow as the result of chain initiation by the dibutylmagnesium. The styrene was then degassed, vacuum distilled into calibrated ampules and diluted with benzene. The ampules, equipped with break-seals for attachment to polymerization reactors, were stored in a freezer. The density for styrene monomer at roomtemperature under its own vapor pressure was determined to be 0.897 g/ml using a calibrated ampule.

1,3-butadiene was stirred over lithium aluminum hydride (-20°C), degassed, and distilled onto sec BuLi. After stirring at -10°C for 30 min., the butadiene was distilled into a flask containing dibutyl magnesium and stored under vacuum at -20°C in the freezer. Before distilling into the delivery ampule, the monomer was further purified over sodium mirrors. The density of 1,3-butadiene at -78°C under its own vapor pressure was measured as 0.705 g/ml. 1,3 butadiene-d6 was obtained from two sources, Merck, Sharp & Dohme Ltd. and Cambridge Isotope Inc. The purification procedure for the perdeuterated monomer was identical to that of the normal 1,3-butadiene. Halogenated impurities are removed effectively as lithium salts when reacted with sec-BuLi. The density of 1,3-butadiene-d6 was measured as 0.780 g/ml at -78°C.

Diblocks SBm and SBmBd were synthesized in one reactor (8), the triblock being formed by extending a portion of the living SBm diblock with a third block of perdeuterated butadiene. The mixed ("m") block contained 83% normal polybutadiene (8). Secondary butyllithium initiator was titrated using techniques developed by Gilman and Cartledge (10). Polystyrene synthesis proceeded at room temperature in benzene for 8 hours; polybutadiene blocks were produced over 24 hour periods. The living polymers were terminated at appropriate stages of the synthesis with degassed methanol yielding approximately 1.9 grams of the polystyrene block, 4.6 grams of SBm and 17.1 grams of SBmBd. Total yield for the synthesis was greater than 99%.

Molecular Characterization (8)

Homopolymers and diblock copolymers were characterized using size exclusion chromatography, ultraviolet absorption, proton/carbon 13 NMR experiments, and considerations of the overall stoichiometry of the reaction. The waters GPCII apparatus was calibrated with 9 polystyrene standards and 2 polybutadiene standards obtained from American Polymer Standards Corp. Calibration of the Perkin Elmer 3B UV/VIS spectrophotometer at 262nm was accomplished using four carefully prepared solutions of polystyrene in 1,4 dioxane; at this wavelength there is no contribution to the UV absorption from polybutadiene, so the styrene content of SB block copolymers is readily obtained. NMR experiments (Bruker WM250) also provided information on the copolymer compositions and on the polybutadiene microstructures (8).

Table 1 Molecular Characterization of Polymers

| <u>Sample</u> | <u>M(b)(S)</u> | <u>M(Bm)</u> | <u>M(Bd)</u> | <u>Bm</u> | Microstructure | (^c) | <u>Bd</u> |
|------------------|----------------|---------------------|--------------|-----------|----------------|------------------|-----------|
| S ^(a) | 83.4 | | | | | - | - |
| SBm | 83.4 | 10.1 ^(d) | | //10 | | - | - |
| SBmBd | 83.4 | 10.1 | 1.2(*) | //10 | 5 | 1/44, | /5 |

(a) Polystyrene block of the copolymers

(b) Kg/mole; Mw/Mn <1.1

(c) cis 1,4/trans 1,4/vinyl

(d) Random copolymer; degree of polymerization 183 (152 B and 31 Bd)

(e) Degree of polymerization 20

Morphological Characterization

Transmission electron micrographs were obtained on a Phillips 300 electron microscope operated at 80 kV and calibrated against a diffraction grating with 54,864 lines per inch. Samples were stained in a 1 wt% osmium tetroxide aqueous solution, and cut into sections ~300 Å thick using a LKB ultramicrotome fitted with a freshly prepared glass knife.

Small-angle neutron scattering was performed at the National Bureau of Standards SANS spectrometer. The alignment, calibration, and transmission measurement was performed with the assistance of Dr. Charles C. Han; the NBS instrumentation has been described in detail in the literature(11,12). The important characteristics of the SANS instruments as used in this work are: wavelength λ -6Å, $\Delta\lambda/\lambda$ -0.25, sample to detector distance 3.6m. Silica and a PS/PSd 50/50 blend were used as standards in the calibration procedures (8).

Specimen Preparation

Polymer films were cast on a mylar film in an aluminum cup from a 5 wt% solution in toluene; solutions were passed through a Millipore Millex-SR 0.5 μ m filter prior to casting. During the first 24hrs of the film forming period at 60°, nitrogen was purged through the cup at a flow rate of 0.1 - 0.2 standard ft³/hr. A subsequent 10 hour drying period at 75°C employed the same nitrogen flow rate.

Films were cut into 1.5 cm x 1.5 cm square pieces with a thickness of . 0.3 mm. An additional drying process under vacuum employed a heating program starting at room temperature and progressing at a rate of 20° - 30° C per day up to 120°C. Then, the films were cooled, cleaned with methanol, stacked between two glass plates held together by a small binder clip, and transferred into the vacuum oven at 120°C. No appreciable compression of the film stack was observed during this 10 minute heating process. Final film thickness was about 2mm. Before neutron scattering, all the samples were simultaneously heated without the binder clip to 120°C in vacuum overnight and cooled down at a rate of 10° C/hr.

<u>Results</u>

Transmission electron micrographs of SBm and SBmBd are shown in Figures 2a and 2b respectively. Spheres of polybutadiene in a polystyrene matrix are revealed in both photographs. In both cases the sphere diameter is somewhat less than 200Å.

Results of SANS experiments on SBm and SBmBd appear in Figures 3a and 3b respectively. Unlike the micrographs of Figure 2 which are essentially indistinguishable, the SANS patterns for these two samples are strikingly different. For SBm the intensity of scattered neutrons is very close to zero over the entire range of Q $(-4\pi \sin\theta/\lambda)$ covered by the experiment. Intensities for SBmBd are orders of magnitude larger over the same Q range and significant features are seen in the scattering pattern; a large peak appears near Q = 0.02 and a pronounced shoulder appears in the Q range immediately above the peak. Results of SAXS experiments presented elsewhere(8), covering the same Q-range, show exactly the same features seen in SANS (a pronounced peak at the same location as the SANS main peak followed by a higher Q shoulder on this peak); in addition a broad rolling maximum is seen in the SAXS pattern near Q=0.06.

Discussion

We first note that the synthesis of SBm was successful in producing the desired phase-matched (7) result shown schematically in Figure 1a. The featureless scattering pattern with essentially zero intensity (Figure 3a) provides unequivocal evidence that the structural scattering has been eliminated by matching the neutron scattering length densities of the two moieties of the SBm diblock. When the short Bd endblocks are added, thereby producing SBmBd, the structural scattering reappears dramatically (Figure3b). It is now our task to determine whether or not this re-emergent scattering can be used to discriminate between the two extreme cases for Bd chain end distribution mentioned earlier and shown schematically in Figures 1b and 1c; i.e. we hope to determine from the information contained in Figure 3b whether the Bd chain ends distribute themselves uniformly throughout the polybutadiene spheres or cluster near the centers of the spheres.

A first consideration of Figure 3b involves the magnitude of the intensity exhibited by the large peak near Q=0.02. This peak intensity reflects the magnitude of the difference in scattering length density between the polystyrene matrix and the polybutadiene spheres. When this difference is zero the peak intensity is zero as seen in Figure 3a and in our earlier studies (4,5) of composition matched (7) mixtures of two diblocks. Koberstein (7) and Jahshan and Summerfield (13) have developed nearly equivalent expressions which allow for a calculation of the variation of the SANS contrast factor (and therefore scattering intensity) with mole fraction of deuterated repeat units in the polybutadiene regions of an SB diblock, assuming that these deuterated units are uniformly distributed within polybutadiene regions. The solid curve of Figure 4 shows this expected variation of intensity with mole fraction of deuterated B units in a SB block copolymer system. Intensity drops with added deutero-B units until contrast matching is achieved near 16 mole percent. The intensity is expected to rise with further addition of deuterated B units beyond the point of contrast matching because the scattering length density of the spheres now exceeds that of the polystyrene matrix. The three data points shown in Figure 4 correspond to the normalized scattering intensities (integrated between Q=0.012 and 0.032 \dot{A}^{-1}) for three samples: a normal SB diblock (8) with spherical B microdomains for which $I_N = 1.0$; phase matched diblock SB_m, $I_N = 0$; and SBmBd, $I_y=0.3$. Note that sample SBmBd contains about 25% deutero-polybutadiene repeat

units, 16% from the Bm moiety and another 9% from the Bd endblocks. The essentially perfect agreement between the observed normalized peak intensity for SBmBd and that expected on the basis of uniform distribution of deuterated segments in the polybutadiene spheres is strong support for the hypothesis (Figure 1b) of essentially uniform distribution of Bd chain ends in the polybutadiene sphere of SBmBd.

This tentative conclusion is supported by the results of modelling of the form of the scattering pattern exhibited by SBmBd. Previous papers from our laboratory (6) and elsewhere (14) have shown decisively that scattering from block copolymers of spherical morphology can be modelled by an appropriate combination of a Percus-Yevick (15) interparticle interference function and a Bessel function intraparticle form factor. Because the magnitude and distribution of center-to-center distances among spheres do not change when considering either Figure 1b or 1c, the parameters of the interparticle interference function cannot be expected to discriminate between these two cases. On the other hand, the key parameter of the intraparticle form factor, the effective polybutadiene sphere radius, changes dramatically from Figure 1b to 1c. For Figure 1b, representing uniformly distributed ends, the appropriate value of the radius of the scattering spheres is the overall radius of the polybutadiene spheres which from TEM (Figure 2b) is known to be about 100 Å and from SAXS data (8) a value of $R=92\dot{A}$ was determined. In Figure 5, curve b shows the degree to which the SANS data are described by Percus-Yevick interparticle scattering combined with an intraparticle form factor based on a value of 92 Å for the sphere radius. The essential characteristics of the SANS data in this region of Q are described by this model further supporting the earlier conclusion regarding uniform chain end distribution in the block copolymer spheres. We note further that SAXS and SANS spectra for the case of uniformly distributed ends should be similar, as observed (8), since both bulk density (SAXS) and neutron scattering length (SANS) vary in the same way throughout the sample. However if chain end clustering occurred, SAXS and SANS spectra would be quite different; the SANS pattern would be dictated by a contrast profile like that of Figure 1c whereas SAXS would still essentially see a profile like that of Figure 1b.

It remains to be shown that the SANS scattering pattern expected for the case of clustered Bd ends (Figure 1c) is significantly different from experimental observations. Based on the fact that the Bd segments have about 10% of the Bm chain length, we take the neutron scattering contrast of ~90% of the

sphere volume to be matched with polystyrene; the contrast then rises sharply in the inner 10% of each sphere. The radius of this hypothetical perdeuterated polybutadiene inner sphere is thus about 45 Å. The expected scattering pattern for this case, modelled using Percus-Yevick parameters as before and an intraparticle form factor with sphere radius of 45 Å, is shown as the solid curve a in Figure 5. Clearly this model does not reproduce the essential features of the observed SANS data. Thus the hypothesis of chain ends clustered at the sphere centers can be rejected.

Summary and Conclusions

A perfectly phase-matched diblock copolymer SBm was suitably fitted with short perdeuterated end sections (Bd ends of 10% the chain length of Bm) to form a triblock copolymer SBmBd. While TEM micrographs of the copolymers SBm and SBmBd were essentially identical, SANS patterns were remarkably different. The amount of SANS intensity displayed by the SBmBd was consistent with essentially uniform distribution of deuterated material throughout the polybutadiene spheres. The form of the SANS pattern of SBmBd was very similar to that observed in SAXS experiments; the SANS curve was well described by a Percus Yevick interparticle interference function and an intraparticle form factor which incorporated the full polybutadiene sphere radius (ca 92 Å). A similar analysis which incorporated only the inner core of the spheres failed to reproduce the essential features of the SANS data for SBmBd. From these observations it is concluded that the Bd ends of SBmBd are essentially uniformly distributed throughout the polybutadiene spheres and that models which envision the chain ends as being localized at the sphere cores are incorrect.

Acknowledgements

The work was supported in part by the Office of Naval Research and by the National Science Foundation. SANS data were obtained at NBS with a great deal of assistance from Dr. Charles Han, whose help is gratefully acknowledged. The SAXS data referred to here and which appear in (8) were obtained by Dr. John D. Barnes of NBS.

<u>References</u>

- (1) F.S. Bates, R.E. Cohen and C.V. Berney, <u>Macromolecules</u>, <u>15</u>, 598 (1982)
- (2) F.S. Bates, C.V. Berney and R.E. Cohen, Macromolecules, 16, 1101 (1983)
- (3) C.V. Berney, R.E. Cohen and F.S. Bates, Polymer, 23, 1222 (1982)
- (4) F.S. Bates, C.V. Berney, R.E. Cohen and G. Wignall, <u>Polymer</u>, <u>24</u>, 519 (1983)
- (5) C.V. Berney, P. Kofinas and R.E. Cohen, <u>Polymer Communications</u>, <u>27</u>, 330 (1986)
- (6) P. Cheng, C.V. Berney and R.E. Cohen, <u>Macromolecules</u>, <u>21</u>, 0000 (1988)
- (7) X. Quan and J.T. Koberstein, J. Polymer Sci., Physics, 25, 1381 (1987)
- (8) P. Cheng, Sc.D. Thesis, Massachusetts Institute of Technology 1988.
- (9) M. Morton and L. Fetters, <u>Rubber Chem. Tech.</u>, <u>48</u>, 359 (1975)
- (10) H. Gilman and F.K. Cartledge, J. Organomet_Chem., 2, 447 (1964)
- (11) C.J. Glinka, AIP Conference Proceedings, <u>89</u>, 395 (1981)
- (12) E.J. Amis, C.J. Glinka, C. Han, H. Hasagawa, T. Hashimoto,
 T.P. Lodge, Y. Matsushita, <u>Polymer Preprints</u>, <u>24</u>, 215 (1983)
- (13) S.N. Jahshan and G.C. Summerfield, <u>J. Polymer Sci.</u>, Physics, <u>18</u>, 1859 (1980)
- (14) D.J. Kinning and E.L. Thomas, <u>Macromolecules</u>, <u>17</u>, 1712 (1984)
- (15) J.K. Percus and G.J. Yevick, Phys. Rev., 110, 1 (1958)

<u>Figure</u>

- Scattering length density profiles for: (a) SBm; (b)SBmBd assuming uniform distribution of Bd ends in the polybutadiene sphere; (c) SBmBd assuming that the Bd chain ends cluster at the core of the sphere. Polystyrene matrix is crosshatched and polybutadiene sphere is unshaded.
- Transmission electron micrographs of osmium tetroxide stained sections of:
 (a) SBm; (b) SBmBd
- 3. SANS data for: (a) SBm and (b) SBmBd
- 4. SANS intensity as a function of mole fraction of perdeuterated repeat units within the polybutadiene spheres assuming uniform distribution of the perdeuterated material. Solid line: expected behavior based on expressions developed in references (7) and (13). Data points for samples indicated, normalyzed to SB intensity.
- 5. Fits of a combined interparticle interference function and an intraparticle form factor to the SBmBd SANS data. Curve a uses 45 Å for the radius of the intraparticle form factor; curve b, R = 92Å. Parameters of the interparticle interference function are identical for the two curves:





h

Figure 2



56353 La

(A - 2000)(333

Figure 3a







