UNCLASSIFIED	THE THO-DIMENS PERIODIC BACKG DEPT OF PHYSIC N00014-81-0776	IONAL ONE CO Round (U) S F Cornu e	PUERTO RICO T AL 24 JA	ISNA IN A DO UNIV RIO F IN 88 TR-30 F/G	TEDRAS 20/9 NL	
	1. P					



a Secretaria

MICROCOPY RESOLUTION TEST CHART NATIONA BUREAU CANDARDS (963.4

	ATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
REPORT NUMBER	2. GOVT ACCESSI	ON NO. 3. RECIPIENT'S CATALOG NUMBER
Technical Report #30		
The Two-Dimonsional One	-Component Pla	S. TYPE OF REPORT & PERIOD COVERED
in a Doubly Periodic Ba	ckground: Exac	t Interim Technical Report
Results	~	6. PERFORMING ORG. REPORT NUMBER
AUTHOR:		8. CONTRACT OR GRANT NUMBER(#)
Francoise Cornu, Bernar Lesser Blum	d Jancovici an	d N 00014-81-07/6
PERFORMING ORGANIZATION NAME AND A	10. PROGRAM ELEMENT PROJECT, TASK	
Physics Department, Uni Rico, Río Piedras, Puer	versity of Pue to Rico 00931	rto Task No. NR-051-0776
CONTROL ING OFFICE NAME AND ADORE	255	12. REPORT DATE
Code 572-Office of Nava	1-28-88	
Arlington, VA 22217	i noocaron	13. NUMBER OF PAGES
MONITORING AGENCY NAME & ADDRESS	it different from Controlling C	Difice) 15. SECURITY CLASS. (of this report)
		Unclassified
		154. DECLASSIFICATION DOWNGRADING
		FEB 2 2 1988
DISTRIBUTION STATEMENT (of the obstrac	t entered in Block 20, 11 dill	FEB 2 2 1988
DISTRIBUTION STATEMENT (of the observed)	t entered in Block 20, if diff	FEB 2 2 1988
DISTRIBUTION STATEMENT (of the observed SUPPLEMENTARY NOTES Prepared for publicatic	en in J. Stat.	FEB 2 2 1988
DISTRIBUTION STATEMENT (of the observed SUPPLEMENTARY NOTES Prepared for publicatio	en in J. Stat.	FEB 2 2 1988
DISTRIBUTION STATEMENT (of the observed SUPPLEMENTARY NOTES Prepared for publicatio <ev (continue="" *ords="" if="" nee<br="" on="" reverse="" side="">Plasmas structured inte</ev>	on in J. Stat.	Phys.
DISTRIBUTION STATEMENT (of the observe SUPPLEMENTARY NOTES Prepared for publicatio <ey #oros="" (continue="" elde="" if="" not<br="" on="" reverse="">Plasmas structured inte</ey>	on in J. Stat.	FEB 2 2 1988
DISTRIBUTION STATEMENT (of the observe SUPPLEMENTARY NOTES Prepared for publicatio <ev #ords="" (continue="" elde="" if="" nec<br="" on="" reverse="">Plasmas structured inte</ev>	on in J. Stat.	Phys.
DISTRIBUTION STATEMENT (of the observed SUPPLEMENTARY NOTES Prepared for publicatio <ev (continue="" if="" moros="" nee<br="" on="" reverse="" side="">Plasmas structured inte y of Continue on reverse side if nee</ev>	on in J. Stat. Trfaces	Phys.
DISTRIBUTION STATEMENT (of the ebetree SUPPLEMENTARY NOTES Prepared for publicatio <ey #0805="" (continue="" elde="" if="" nec<br="" on="" reverse="">Plasmas structured inte y #600 ABSTRACT Continue on reverse elde if nec Using a new method, we one-component plasma in model, in equilibrium of special value f = 2 of t of background shapes: t computed. In particula background; this is a c</ey>	on in J. Stat. en in J. Stat. erfaces find that the a inhomogeneo classical stati the coupling co the n-body dens ar, we can deal classical model	Phys. Two-dimensional bus background is a solvable stical machanics, for the onstant, for a larger class ities can be explicitly with a doubly periodic for a crystal mode of fixed
DISTRIBUTION STATEMENT (of the ebetree SUPPLEMENTARY NOTES Prepared for publicatio <ev (continue="" elde="" if="" nec<br="" on="" reverse="" roros="">Plasmas structured inte y for the Using a new method, we one-component plasma in model, in equilibrium of special value f = 2 of t of background shapes: t computed. In particula background; this is a c</ev>	on in J. Stat. en in J. Stat. erfaces find that the a inhomogeneo classical stati the coupling co he n-body dens ir, we can deal classical model source te	FEB 2 2 1988 Phys. Phys. Phys. number: two-dimensional bus background is a solvable stical machanics, for the enstant, for a larger class ities can be explicitly with a doubly periodic for a crystal mode of fixed Unclassified

.*

2. 6 1. 2 6 UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (The Date Entered) ions and mobile electrons. At Γ = 2, this system is conductor: the correlations have a fast decay, and the Stillinger-Lovett - ions and mobile elctrons. screening sum rule is obeyed. DTIC COPY SPECTED 6 Accession For NTIS GRAŁI DTIC TAB Unannounced Justification By__ Distribution/ Availability Codes Avail and/or Special Dist UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Date Batered) ,

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-81-0776

TASK No. NR-051-0776

TEHCNICAL REPORT #30

The Two-Dimensional One-Component Plasma in a Doubly Periodic Background: Exact Results

by

Francoise Cornu*, Bernard Jancovici*, and Lesser Blum**

*Laboratoire de Physique Theorique et Hautes Energies, Universite Paris-Sud, Bat 211, 91405 Orsay, France (This Laboratory is associated with the Centre National de la Recherche Scientifique).

**Department of Physics, University of Puerto Rico, Rio Piedras, Puerto Rico 00931

Prepared for Publication in Journal of Statistical Physics

Reproduction in whole or in part is permitted for any purpose of the United States Goverment

*This document should also appear in Item 10 of Document Control Data--DD Form 1473. Copies of form available from cognizant contract administrator. The Two-Dimensional One-Component Plasma in a Doubly Periodic Background : Exact Results

Françoise Cornu¹, Bernard Jancovici¹, and Lesser Blum²

Abstract

TATA AND DECOMPOSITE TATA AND A

Using a new method, we find that the two-dimensional one-component plasma in an inhomogeneous background is a solvable model, in equilibrium classical statistical mechanics, for the special value $\Gamma = 2$ of the coupling constant, for a larger class of background shapes : the n-body densities can be explicitly computed. In particular, we can deal with a doubly periodic background; this is a classical model for a crystal made of fixed ions and mobile electrons. At $\Gamma = 2$, this system is conductor : the correlations have a fast decay, and the Stillinger-Lovett screening sum rule is obeyed.

KEY WORDS : One-component inhomogeneous plasma; doubly periodic background; conducting phase.

LPTHE ORSAY 87/69

November 1987

¹ Laboratoire de Physique Théorique et Hautes Energies, Université Paris-Sud, Bât. 211, 91405 Orsay, France (This Laboratory is associated with the Centre National de la Recherche Scientifique).

² Department of Physics, POB AT, Faculty of Natural Sciences, University of Puerto Rico, Rio Piedras, Puerto Rico 00931.

1. INTRODUCTION

In statistical mechanics, it is obviously of interest to have exactly solvable models for Coulomb systems (plasmas, electrolytes, metals etc...). The simplest model of a Coulomb system is the one-component plasma (jellium) : identical charged particles move in a rigid charged background which ensures overall neutrality. In two dimensions, the Coulomb potential between two particles of charge e at a distance r from one another is $-e^2$ $ln(r/\lambda)$, where λ is an arbitrary length scale, and the dimensionless coupling constant is $\Gamma = \beta e^2$, where β is the inverse temperature; for the special value $\Gamma = 2$, it has been previously found that the equilibrium classical statistical mechanics of the two-dimensional one-component plasma can be worked out exactly for several kinds of background charge distributions : one is able to obtain the n-particle densities. Besides the simplest case of a uniform $background^{(1,2)}$, essentially one could deal with a background charge density depending on one space coordinate $^{(3)}$; this covers a variety of charged interfaces (electrical double layers) of interest to electrochemists.

This previous work left unsolved the important case of a doubly periodic background. In the present paper, we solve this case, using a more general new method; a preliminary account has been given by two of $us^{(4)}$. Thus we have an explicit solution for a model which can be understood as made of mobile (classical) "electrons" interacting between themselves and with a lattice of extended fixed "ions"; this one-component plasma in a periodic background can also be regarded as a two-component plasma in which the particles of one species have been fixed on a lattice. Like the symmetric two-component plasma, the present model is expected to undergo a Kosterlitz-Thouless phase transition between a low-temperature dielectric phase and a high-temperature conducting phase, and this transition is actually seen in computer simulations^(5,6). Here, we show

that, in our system, criteria which characterize a conductor are satisfied the correlations have a fast decay at large separations, and the Stillinger-Lovett sum rule⁽⁷⁾ is obeyed. According to these criteria, at $\Gamma = 2$, the system is in its conducting phase.

The paper is organized as follows. In Section 2, the general method is reviewed. In Section 3, it is shown that this method provides a simpler approach to the known case where the background density is inhomogeneous in one direction only. The doubly periodic background is discussed in Section 4 : we compute the n-particle densities and discuss sum rules.

2. METHOD

2.1. n-perticle densities

We start with N particles of charge -e in some background. The position of the ith particle is $r_i = (x_i, y_i)$; we shall also use the complex number $z_i = x_i + iy_i$. The Hamiltonian is

$$H = e^{2} \sum_{i=1}^{N} V(\mathbf{r}_{i}) - e^{2} \sum_{1 \leq i < j \leq N} ln(|z_{j} - z_{j}|/\lambda)$$
(2.1)

where $e^2 V$ is the background-particle interaction, and therefore, for an inverse temperature § such that $\Gamma = ge^2 = 2$, the Boltzmann factor is

$$exp(-\beta H) = C \left[det \left\{ exp\left[-V(r_j) \right] z_1^{j-1} \right\}_{1, j=1, ..., N} \right]^2$$
(2.2)

where C is a constant.

In the simple case of a background potential of circular symmetry, V(r) = V(r), the functions $exp[-V(r)] z^{j-1}$ are mutually orthogonal, and (2.2) has the same form as the squared wavefunction of a system of independent fermions; to deal with the Slater determinant which appears in (2.2) is a The determinant in (2.2) is just a slater determinant where is the standard, and carry to incompute the n-particle densities. All the previously solved cases could be obtained by starting with a circular geometry and taking a suitable limit.

In the present paper, we want to consider more general forms of the background potential, and the functions $\exp[-\nabla(\mathbf{r})] z^{j-1}$ are not necessarily \sim mutually orthogonal. However, we can choose an orthogonal basis $\psi_j(\mathbf{r})$ for the space of these functions, and rewrite (2.2) as

$$exp(-\beta H) = C \left| det \{ \Psi_{j}(r_{j}) \}_{j,j=1,...,N} \right|^{2}$$
 (2.3)

since the new determinant is proportional to the former one. It is then easy to show that the n-particle truncated densities can be expressed in terms of the projector

$$< r_1 |P| r_2 > = \sum_j \frac{\Psi_j(r_1) \Psi_j^*(r_2)}{\int d r |\Psi_j(r)|^2}$$
 (2.4)

83

$$\varrho(\mathbf{r}) = \langle \mathbf{r} | \mathbf{P} | \mathbf{r} \rangle
\varrho^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = -|\langle \mathbf{r}_{1} | \mathbf{P} | \mathbf{r}_{2} \rangle|^{2}$$

$$\varrho^{(n)}(\mathbf{r}_{1}, \mathbf{r}_{2}, ..., \mathbf{r}_{n}) = -\sum_{\substack{(i_{1} \ i_{2} ... i_{n})}} \langle \mathbf{r}_{1} | \mathbf{P} | \mathbf{r}_{1_{2}} \rangle ... \langle \mathbf{r}_{1_{n}} | \mathbf{P} | \mathbf{r}_{1_{1}} \rangle$$
(2.5)

where the summation runs over all cycles $(i_1 \ i_2 \dots i_n)$ built with $\{1, 2, \dots, n\}$. In the thermodynamic limit, the functions $\exp[-V(r)]z^{j-1}$ span the subspace of $\tau \vdash \mathcal{L}$ Hilbert space defined by the entire functions of z = x + iy times $\exp[-V(r)]$, and P becomes the projector on that subspace (of course, this is an intrinsic definition of P, independent of the choice of the orthogonal basis Ψ_i). Thus, the problem of obtaining the n-particle densities is reduced to computing the projector P.

In the simplest case of a uniform background density ρ_0 , the background potential V(r) can be chosen as $1/2 \pi \rho_0 r^2$ (plus some irrelevant constant), $\Psi_j = \exp(-\Psi) z^{j-1}$ and

$$<\mathbf{r}_{1}|\mathbf{P}|\mathbf{r}_{2}> \approx \varrho_{0} \exp[-\frac{\pi \varrho_{0}}{2} (|z_{1}|^{2} + |z_{2}|^{2} - 2z_{1}z_{2}^{*})]$$

$$\varrho(\mathbf{r}) \approx \varrho_{0}$$

$$\varrho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) \approx -\varrho_{0}^{2} \exp(-\pi \varrho_{0}|\mathbf{r}_{1} - \mathbf{r}_{2}|^{2})$$
(2.6)

In the general case, the background density $\rho_{\rm B}(\mathbf{r})$ can be considered as being the sum of a uniform contribution ρ_0 plus a non uniform modulation $\tilde{\rho}(\mathbf{r})$. Correspondingly, the background potential V(r) can be chosen of the form V₀(r) + $\mathscr{O}(\mathbf{r})$, where V₀(r) = (1/2) $\pi \rho_0 r^2$ and $\Delta \mathscr{O}(\mathbf{r}) = 2\pi \tilde{\rho}(\mathbf{r})$. As a first step towards the computation of the projector P on the space of the functions $\exp[-\mathscr{O}(\mathbf{r}) - V_0(\mathbf{r})]z^j$, $j \in \mathbb{N}$, it will turn out to be convenient to replace the z^j by another basis for the entire functions $\varphi_k(z) =$ $\exp\{-(1/2)\pi\rho_0[z-(k/\pi\rho_0)]^2\}$, $k \in \mathbb{R}$. The φ_k are indeed such a basis, since

$$z^{h} = \pi^{-1/2} 2^{-n} \int_{-\infty}^{\infty} dt H_{n}(t) \exp[-(z-t)^{2}]$$
 (2.7)

where the H_n(t) are Hermite polynomials, and (2.7) becomes a superposition of φ_k functions through a rescaling of z and t. The basis $exp[-V_0(r)]z^j$ is then replaced by

Actually, since (2.8) will be used for defining the projector P and thereafter computing the densities (2.5), we can omit in (2.8) the normalization factor $\exp(-k^2/4\pi\varrho_0)$ (this leaves the projector unchanged) and the phase factor $\exp(-i\pi\varrho_0 xy)$ (this leaves the densities unchanged because of their cyclic structure).

Therefore, an alternative definition of P is to take it as the projector on the space of the functions

$$\psi_{k}(\mathbf{r}) = \exp[-\mathscr{G}(\mathbf{r})] \exp[-\pi \varrho_{0} (\mathbf{x} - \frac{\mathbf{k}}{2\pi \varrho_{0}})^{2} + i\mathbf{k}\mathbf{y}]$$
(2.9)

For the potentials $\mathscr{P}(\mathbf{r})$ which will be considered here, strictly speaking the functions (2.9) do not belong to Hilbert's space, because $|\exp(iky)|$ does not decrease at infinity. However, these functions do form a basis in the sense of distributions, just like the plane waves in quantum mechanics.

2.2. Arbitrariness in the choice of V(r)

Let us remark that the background potential V(r) is not uniquely determined by the background density $\varrho_{\rm B}(r)$. In the thermodynamic limit, V(r) keeps a memory of the boundary conditions even after these boundaries have receded to infinity. The n-particle densities however should depend only on $\varrho_{\rm B}(r)$ for a system with screening properties which prevent the bulk from being affected by infinitely remote charged boundaries. It is satisfactory to check explicitly this independence upon the choice of V(r), to which it should be always possible to add an arbitrary harmonic function. Actually, since the confinement of the particles must be preserved, the total background potential must increase fast enough at infinity, and we shall only consider the addition to V(r) of a harmonic function of the form $f(\mathbf{r}) = a_2(x^2-y^2) + b_2xy + a_1x + b_1y + c$, with coefficients a_2 and b_2 of sufficiently small absolute value; furthermore, the term b_2xy can be removed by a rotation of the axes. The potential $f(\mathbf{r})$ can be interpreted as determined by suitable external electrodes.

It is then easy to see that $\varphi_{k}(z)$ can be chosen again of the form $\exp\{-\alpha[z - (2\alpha)^{-1}(k + \gamma + i\delta)]^{2}\}$, where α, γ, δ are real constants, and by a suitable choice of these constants, f(r) can be cancelled, except for irrelevant normalization and phase factors. Therefore, the projector P can be left unchanged.

incidentally, the basis (2.9) can be directly obtained by choosing the potential contribution from the uniform background as $V_0 = \pi \varrho_0 x^2$ and taking exp(kz) as the basis for the entire functions. Such a V_0 will be obtained if we reach the infinite system limit starting from a strip geometry rather than from a circular one.

2.3. Magnetic analogy

The arbitrariness in the choice of our background potential V(r) has a quantum-mechanical analog : a gauge transformation in a magnetic problem. Let us consider a particle of mass m and charge q moving in the xy plane and submitted to a uniform magnetic field B parallel to the z axis^(B). The ground state is infinitely degenerate. In the gauge where the vector potential is A = $(1/2)B \times r$, a basis for the ground-state wavefunctions is $exp[-(qB/4)r^2]z^j$, $j \in N$; in the gauge where the vector potential is A = j Bx (j is the unit vector along the y-axis), a basis for the ground-state wave functions is $exp[-(qB/2)[x - (k/qB)]^2 + iky]$, $k \in \mathbb{R}$. Obviously, our change from the basis $exp[-(1/2)\pi q_0 r^2]z^j$ to the basis $exp[-\pi q_0 [x - (k/2\pi q_0)]^2 + iky]$ is exactly of the same form. If a Slater determinant is built with the ground-state wave functions, the corresponding n-body densities must be gauge-independent, just as our n-body densities are independent of our choice of basis.

3. BACKGROUND INHOMOGENEOUS IN ONE DIRECTION.

We now revisit the case of a background density depending on one coordinate only⁽³⁾: $\rho_B(x)$. The potential \mathscr{P} can be chosen as $\mathscr{P}(x)$, and we have at hand the orthogonal basis : (2.9) is orthogonal because of the plane-wave factor exp(iky). Adapting (2.4) to the case of a continuous index k, with the usual prescription that the summation becomes $(L/2\pi) \int dk$ as the length L of the system in the y direction becomes infinite, we find

$$<\mathbf{r}_1 |\mathbf{P}| \mathbf{r}_2 > = \exp[-\mathscr{O}(\mathbf{x}_1) - \mathscr{O}(\mathbf{x}_2)]$$

$$x \int_{-\infty}^{\infty} \frac{dk}{2\pi} = \frac{\exp[ik(y_{1}-y_{2})]\exp\{-\pi\varrho_{0}[(x_{1}-\frac{k}{2\pi\varrho_{0}})^{2}+(x_{2}-\frac{k}{2\pi\varrho_{0}})^{2}]\}}{\int_{-\infty}^{\infty} dx \exp[-2\mathscr{D}(x)-2\pi\varrho_{0}(x-\frac{k}{2\pi\varrho_{0}})^{2}]}$$
(3.1)

Using (3.1) in (2.5), we retrieve at once the densities of Ref.3.

Some special care must be exercised for dealing with the case where the particles are confined to the half space x > 0 by an impenetrable wall at x = 0. Then, in (3.1), the range of x must be restricted to x > 0, and the range of k must be restricted to k > 0. This is shown as follows. Since we have already taken the limit of an infinite system, we shall start⁽³⁾ with a system in which an impenetrable barrier occupying the region - l < x < 0separates the plasma into two regions x < -l and x > 0. The impenetrable wall system will be obtained by taking the limit $l \rightarrow \infty$ in such a way that the remote regions x > 0 and x < -l no longer see each other. The reason for which the values k < 0 are suppressed in (3.1) will be seen to be that the norm in the denominator has a contribution from the remote region x < -lwhich becomes infinite for k < 0, in the limit $l \rightarrow \infty$. The independence of the regions x > 0 and x < -l can be achieved by requiring that each of them

 (\mathcal{O})

be globally neutral, and this will be the case if we choose a background potential symmetric \mathbf{n} with respect to the barrier, $V(x) = \pi \rho_0 x^2 + \mathscr{O}(x)$ for x > 0, $V(x) = +\infty$ for -L < x < 0, V(x) = V(-x - L) for x < -L; Then, remembering that $\mathscr{O}(x)$ is defined in <u>every</u> region as $V(x) - \pi \rho_0 x^2$, and changing -x - L into x, we can rearrange the denominator of (3.1) as

$$f(k) = \exp(-2k!) \int_{0}^{\infty} dx \exp[-2\mathscr{O}(x) - 2\pi\varrho_{0}(x + \frac{k}{2\pi\varrho_{0}})^{2}] + \int_{0}^{\infty} dx \exp[-2\mathscr{O}(x) - 2\pi\varrho_{0}(x - \frac{k}{2\pi\varrho_{0}})^{2}]$$
(3.2)

In the limit $l \rightarrow \infty$, the first term of f(k) (which is the contribution from the region x < -l) diverges if k < 0, vanishes if k > 0. As a consequence, (3.1) becomes

$$< \mathbf{r}_{1} | \mathbf{P} | \mathbf{r}_{2} > = \exp[-\mathscr{A}(\mathbf{x}_{1}) - \mathscr{A}(\mathbf{x}_{2})]$$

$$\times \int_{0}^{\infty} \frac{d\mathbf{k}}{2\pi} \frac{\exp[i\mathbf{k}(\mathbf{y}_{1} - \mathbf{y}_{2})] \exp\{-\pi\varrho_{0}\left[(\mathbf{x}_{1} - \frac{\mathbf{k}}{2\pi\varrho_{0}})^{2} + (\mathbf{x}_{2} - \frac{\mathbf{k}}{2\pi\varrho_{0}})^{2}\right]\}}{\int_{0}^{\infty} d\mathbf{x} \exp[-2\mathscr{A}(\mathbf{x}) - 2\pi\varrho_{0}(\mathbf{x} - \frac{\mathbf{k}}{2\pi\varrho_{0}})^{2}]} , \mathbf{x}_{1}, \mathbf{x}_{2} > 0$$
(3.3)

(3.3) is in agreement with the results⁽²⁾ about a hard wall carrying a surface charge density eo, in which case $\mathscr{O}(x) = 2\pi\sigma x, x > 0; \ \mathscr{O}(x) = 0, x < 0.$

4. DOUBLY PERIODIC BACKGROUND

In this Section, which is the core of the present paper, we study the case where the background density is doubly periodic. Thus, we consider a doubly periodic background potential modulation $\mathscr{P}(\mathbf{r})$:

$$\mathscr{O}(\mathbf{r} + n \, \mathbf{a} + m \, \mathbf{b}) = \mathscr{O}(\mathbf{r}), \quad n, m \in \mathbb{Z}$$

$$(4.1)$$

The unit cell is a parallelogram built with the vectors **a** and **b**, of area ab sin φ , where φ is the angle between **a** and **b**. The system is neutral, with a particle density ϱ_0 equal to the average of the total background density $\varrho_0 + (2\pi)^{-1}\Delta \mathscr{O}(\mathbf{r})$. In order to mimic a simple crystal of extended fixed ions and mobile electrons, of opposite charges, we take $\varrho_0 = (ab \sin \varphi)^{-1}$, which means there is one particle per unit cell.

4.1. n-particle densities

Although the functions (2.9) now are not orthogonal, they are a good starting point for computing the projector P. Choosing the y axis along the period vector **b**, and defining $\zeta \in [0, 1]$ and n integer by $k = 2\pi(\zeta + n)/b$, we can rewrite Ψ_{ν} as

$$\Psi_{\zeta,n}(\mathbf{r}) = \exp[-\mathscr{G}(\mathbf{r})] \exp[-\pi \varrho_0 (\mathbf{x} - \frac{\zeta + \mathbf{n}}{\varrho_0 \mathbf{b}})^2 + 2\pi i (\zeta + \mathbf{n}) \frac{\mathbf{y}}{\mathbf{b}}] \qquad (4.2)$$

As a consequence of the perioducity of \emptyset along the y axis,

$$\int_{-\infty}^{\infty} dy \, \Psi_{\zeta', n'}^{\bullet}(\mathbf{r}) \, \Psi_{\zeta, n}(\mathbf{r}) \propto \delta(\zeta - \zeta') \tag{4.3}$$

Furthermore, if the unit cell is a rectangle ($\varphi = \pi/2$), $(\varrho_0 b)^{-1} = a$, and the periodicity of \mathscr{P} along the x axis ensures that $\Psi_{\zeta,n}$ depends on x only through x - na; this suggests to introduce the Bloch functions

$$\Psi_{\zeta,\eta}(\mathbf{r}) = \sum_{n} \exp(-2\pi i \eta n) \Psi_{\zeta,n}(\mathbf{r}), \zeta, \eta \in [0, 1]$$
(4.4)

which do have the desired orthogonality property

$$\int_{-\infty}^{\infty} dx \, \Psi_{\zeta,\eta'}^{*}(\mathbf{r}) \, \Psi_{\zeta,\eta'}(\mathbf{r}) \propto \delta(\eta - \eta')$$
(4.5)

The argument can be easily extended to the more general case where the unit cell is a parallelogram, by introducing (dimensionless) oblique coordinates (X, Y) defined by $\mathbf{r} = X\mathbf{a} + Y\mathbf{b}$. Multiplying $\Psi_{\zeta,n}$ by an irrelevant phase factor exp{-i $\pi(a/b) \cos \phi[X^2 + (\zeta + n)^2]$ }, we obtain

$$\Psi_{\zeta,n}(\mathbf{r}) = \exp[-\mathscr{G}(\mathbf{r})] \exp\left[-\frac{\pi}{\tau} (X - \zeta - n)^2 + 2\pi i (\zeta + n)Y\right]$$
(4.6)

where $\tau = (b/a) \exp\{i[\phi - (\pi/2)]\}$; with these $\Psi_{\zeta,n}$ functions, which have the same form as (4.2), we can proceed as above, in oblique coordinates in the general case.

Thus, the $\Psi_{\zeta,\eta}$ are orthogonal :

$$\int d\mathbf{r} \, \Psi_{\zeta',\eta'}^{*}(\mathbf{r}) \, \Psi_{\zeta,\eta}(\mathbf{r}) = \delta(\zeta - \zeta') \, \delta(\eta - \eta') \, f(\zeta, \eta) \tag{4.7}$$

where

$$f(\zeta, \eta) = \frac{1}{\rho_0} \sum_{N} \exp(2\pi i \eta N) \int_{-\infty}^{\infty} dX \exp[-\frac{\pi}{\tau} (X - \zeta)^2 - \frac{\pi}{\tau^*} (X - \zeta - N)^2] \\ \times \int_{0}^{1} dY \exp[-2\mathscr{B}(X,Y)] \exp(-2\pi i NY)$$
(4.8)

These orthogonal $\Psi_{\zeta,\eta}$ can be used for building the projector (2.4), with the result

<
$$r_1$$
 iPi r_2 > = exp[- $\mathscr{G}(r_1) - \mathscr{G}(r_2)$] $\int_{0}^{1} d\zeta \int_{0}^{1} d\eta \frac{1}{f(\zeta,\eta)} \sum_{n=1}^{\infty} exp[2\pi i\eta(m-n)]$

$$x \exp\{-\frac{\pi}{\tau} (X_1 - \zeta - n)^2 - \frac{\pi}{\tau^*} (X_2 - \zeta - m)^2 + 2\pi i [(\zeta + n)Y_1 - (\zeta + m)Y_2]\}$$
(4.9)

A more compact form can be obtained by using the Poisson identity

$$\sum_{N} \exp[-\frac{\pi}{\tau^*} N^2 + 2\pi i z N] = \sqrt{\tau^*} \sum_{N} \exp[-\pi \tau^* (z - N)^2]$$
(4.10)

in (4.8), and also in (4.9) where we set m = n + N. The result is

$$f(\zeta, \eta) = \frac{1}{\varrho_0} \sqrt{\tau^*} \int_{-\infty}^{\infty} dX \int_{-\infty}^{\infty} dY \exp[-2\mathscr{A}(X, Y)]$$

× $\exp[-\frac{\pi}{\tau} (X-\zeta)^2 - \pi \tau^*(Y-\eta)^2 - 2\pi i (X-\zeta)(Y-\eta)]$ (4.11)

and

and a subscription

STATES S

$$<\mathbf{r}_{1} |\mathbf{P}|\mathbf{r}_{2} > = \exp[-\mathscr{G}(\mathbf{r}_{1}) - \mathscr{G}(\mathbf{r}_{2})]$$

$$\times \sqrt{\tau^{*}} \int_{-\infty}^{\infty} d\zeta \int_{-\infty}^{\infty} d\eta \frac{1}{f(\zeta,\eta)} \exp[-\frac{\pi}{\tau} (X_{1}-\zeta)^{2} - \pi\tau^{*}(Y_{2}-\eta)^{2} - 2\pi i (X_{2}-\zeta)(Y_{2}-\eta)$$

$$+ 2\pi i \zeta(Y_{1}-Y_{2})] \qquad (4.12)$$

Thus, we have obtained an integral representation of the projector P; when used in (2.5) it gives the n-body densities. Some symmetries of (4.12) are hidden; for instance $\langle \mathbf{r}_1 | \mathbf{P} | \mathbf{r}_2 \rangle = \langle \mathbf{r}_2 | \mathbf{P} | \mathbf{r}_1 \rangle^*$. Other ones are apparent; for instance, the function $f(\zeta, \eta)$ is doubly periodical with period 1, and this ensures that the densities have the same periodicity properties as the background.

As an illustration, we consider the case of a square unit cell (a = b, ϕ = $\pi/2$, τ = 1) with the simplest choice

$$\exp[-2\emptyset(\mathbf{r})] = 1 + \lambda(\cos 2\pi X + \cos 2\pi Y) , |\lambda| \le 1/2$$
(4.13)

The total background density is

$$\varrho_{\mathsf{R}}(\mathsf{r}) = \varrho_{\mathsf{0}} + (2\pi)^{-1} \Delta \mathscr{G} \tag{4.14}$$

and from (4.11) and (4.12), we find for the particle density

$$p(\mathbf{r}) = < \mathbf{r}|\mathbf{P}|\mathbf{r} >$$

$$= \rho_0 \sqrt{2} \exp[-2\mathscr{G}(\mathbf{r})] \int_{-\infty}^{\infty} d\zeta \int_{-\infty}^{\infty} d\eta \frac{\exp[-\pi (X-\zeta)^2 - \pi (Y-\eta)^2] \cos[2\pi (X-\zeta)(Y-\eta)]}{1 + \lambda e^{-\pi/2} (\cos 2\pi \zeta + \cos 2\pi \eta)}$$
(4.15)

The potential modulation $\mathscr{P}(X, Y)$, the background density $\varrho_{B}(X, Y)$ and the particle density $\varrho(X, Y)$ are displayed in Fig.1, for $\lambda = .49$. With this choice of a large amplitude λ , the background density is a rather tormented lanscape; the particle density tries to follow, but it does not quite succeed and it exhibits much smoother oscillations.

Another representation of $\rho(r)$ might be of interest. The normalization factor (4.8) can be written as

$$f(\zeta, \eta) = \frac{1}{\varrho_0} \int_0^1 dx \int_0^1 d\gamma |\Psi_{\zeta, \eta}(x, \gamma)|^2$$
(4.16)

and therefore

$$\varrho(\mathbf{X}, \mathbf{Y}) = \varrho_0 \int_0^1 d\zeta \int_0^1 d\eta \frac{|\Psi_{\zeta, \eta}(\mathbf{X}, \mathbf{Y})|^2}{\int_0^1 dX_1 \int_0^1 dY_1 |\Psi_{\zeta, \eta}(\mathbf{X}_1, \mathbf{Y}_1)|^2}$$
(4.17)

If we restrict ourselves to the case of a rectangular unit cell, τ is real, and

$$\Psi_{\zeta,\eta}(X, \mathbf{V}^{2} = \exp[-2\mathscr{O}(X, Y)] \sum_{m,n} \exp[-\frac{\pi}{\tau}(\mathbf{X} - \zeta - n)^{2} - \frac{\pi}{\tau}(\mathbf{X} - \zeta - m)^{2} + 2\pi i(n-m)(\mathbf{Y} - \eta)]$$
(4.18)

The sum on n and m can be replaced by a sum on y = m - n and v = m + n, with y and v of the same parity. The contributions from (y, v) even and (y, v)odd, respectively, can be expressed in terms of Jacobi theta functions. The result is

$$\begin{split} \left|\Psi_{\zeta,\eta}(X, y)^{2} &= \exp[-2\mathscr{O}(X, Y)](\frac{T}{2})^{1/2}[\theta_{3}(2\eta - 2y, \frac{2}{\tau})\theta_{3}(\chi - \zeta, \frac{T}{2}) + \theta_{2}(2\eta - 2y, \frac{2}{\tau}) \\ \theta_{4}(\chi - \zeta, \frac{T}{2})\right] \end{split} \tag{4.19}$$

where the 8 functions are defined by

$$\theta_{3}(x, t) = \sum_{n} \exp[-\pi tn^{2} + 2\pi inx] = t^{-1/2} \sum_{n} \exp[-(\pi/t)(x - n)^{2}]$$

$$\theta_{2}(x, t) = \sum_{n} \exp[-\pi t(n + \frac{1}{2})^{2} + 2\pi i(n + \frac{1}{2})x] = t^{-1/2} \sum_{n} (-1)^{n} \exp[-(\pi/t)(x - n)^{2}]$$

$$\theta_{4}(x, t) = \sum_{n} \exp[-\pi tn^{2} + 2\pi in(x + \frac{1}{2})] = t^{-1/2} \sum_{n} \exp[-(\pi/t)(x - n - \frac{1}{2})^{2}]$$
The particle density is obtained by using (4.19) in (4.17).

4.2. Decay of the correlations

The decay of the truncated densities (2.5) at large separations is faster than any inverse power law. This can be seen as follows. The decay of the densities is governed by the decay of P; the expression (4.12) of which we can study this mean (4.12) which is the form can be rewritten as (4.12) which is the form

$$< \mathbf{r}_1 | \mathbf{P} | \mathbf{r}_2 > = \exp[-\mathscr{G}(\mathbf{r}_1) - \mathscr{G}(\mathbf{r}_2)] \exp[2\pi i X_1(Y_1 - Y_2)]$$

$$\times \sqrt{\tau^{*}} \int_{-\infty}^{\infty} d\zeta \int_{-\infty}^{\infty} d\eta \frac{1}{f(\zeta + X_{1}, \eta + Y_{2})} \exp[-\frac{\pi}{\tau} \zeta^{2} - \pi \tau^{*} \eta^{2} - 2\pi i \zeta \eta + 2\pi i \zeta (Y_{1} - Y_{2}) - 2\pi i \eta (X_{1} - X_{2})]$$

$$(4.21)$$

Let us study the behavior of (4.21), for a fixed value of \mathbf{r}_1 , as \mathbf{r}_2 recedes to infinity. Since f is periodicy, in $\eta + Y_2$, we can define \overline{Y}_2 by $Y_2 = \overline{Y}_2 + \eta$, $\overline{Y}_2 \in$ [0, 1], n integer, and replace Y_2 by \overline{Y}_2 in f. If we content ourselves to keep a fixed value for \overline{Y}_2 , i.e. if \mathbf{r}_2 recedes to infinity by integer steps of Y_2 , the integral in (4.21) is the Fourier transform of a well-behaved function of ζ and η , and it decays faster than any inverse power law.

The resulting fast decay of the correlations is one of the criteria which is believed to characterize the conducting phase.

4.3. Sum rules

The one- and two-particle densities can be shown to obey several sum rules which characterize a conductor.

Neutrality

The averages, on a unit cell, of the particle density and of the background density are equal :

$$\int_{0}^{1} dX \int_{0}^{1} dY \varrho(X, Y) = \varrho_{0}$$
(4.22)

This sum rule is satisfied by (4.17).

Screening of a particle of the system

This screening rule means that a particle of the system induces a polarization cloud of exactly opposite charge :

$$\int d\mathbf{r}_{2} \varrho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = -\varrho(\mathbf{r}_{1})$$
(4.23)

The structure of (2.5) ensures that this rule is obeyed, because of the closure property

$$\int dr_{2} < r_{1} |P| r_{2} > < r_{2} |P| r_{1} > = < r_{1} |P| r_{1} >$$
(4.24)

The rule can also be checked explicitly on the representation (4.9), after some algebra.

<u>Screening of an infinitesimal test charge (Stillinger-Lovett rule)</u>

This screening rule means that an external infinitesimal test charge induces in the system a polarization cloud of exactly opposite charge. Through linear response theory, this statement becomes a sum rule for the truncated two-body density, the Carnie and Chan generalization⁽¹⁰⁾ of the Stillinger-Lovett rule⁽⁷⁾ to an inhomogeneous system, written here for two dimensions :

$$-\delta \int d\mathbf{r}_{1} \int d\mathbf{r}_{2} \ln \mathbf{r}_{2} S(\mathbf{r}_{1}, \mathbf{r}_{2}) = 1$$
 (4.25)

where S is the total charge structure factor

$$S(\mathbf{r}_{1}, \mathbf{r}_{2}) = \theta^{2} \left[\varrho^{(2)}_{T} (\mathbf{r}_{1}, \mathbf{r}_{2}) + \varrho(\mathbf{r}_{1}) \delta(\mathbf{r}_{2} - \mathbf{r}_{1}) \right]$$
(4.26)

in the present case of a periodic system, (4.25) can be written in other forms. Because of its periodicity properties, $S(r_1, r_2)$ is completely described by its double Fourier transform

$$\widetilde{S}_{6}(\mathbf{k}) = \frac{1}{A} \int_{U} d\mathbf{r}_{1} \int d\mathbf{r}_{2} \exp[1 \mathbf{6} \cdot \mathbf{r}_{1} + 1\mathbf{k} \cdot (\mathbf{r}_{2} - \mathbf{r}_{1})] S(\mathbf{r}_{1}, \mathbf{r}_{2})$$
(4.27)

where U means that the integration domain of \mathbf{r}_{1} is the unit cell, of area A = ab sinp; G is a vector of the reciprocal lattice. From (4.23), one derives $\beta = 0.5$ $S_{G}(G) = 0.$ Then, (4.25) can be reexpressed in terms of $S_{G}(\mathbf{k})$ and of the Fourier transform $2\pi/k^{2}$ of $-\ln r$ as

$$2\pi\beta \lim_{k\to 0} \frac{S_0(k)}{k^2} = 1$$
 (4.28)

In other words

$$\frac{1}{A} \int_{U} d\mathbf{r}_{1} \int d\mathbf{r}_{2} \exp[i\mathbf{k} \cdot (\mathbf{r}_{2} - \mathbf{r}_{1})] S(\mathbf{r}_{1}, \mathbf{r}_{2}) \underset{\mathbf{k} \to 0}{\sim} \frac{\mathbf{k}^{2}}{2\pi \beta}$$
(4.29)

Expanding the exponential in (4.29), we find the generalization to a periodic system of the Stillinger-Lovett second moment rule

$$\frac{1}{A} \int_{U} d\mathbf{r}_{1} \int d\mathbf{r}_{2} (\mathbf{r}_{2} - \mathbf{r}_{1})_{\alpha} (\mathbf{r}_{2} - \mathbf{r}_{1})_{\gamma} \varrho^{(2)} (\mathbf{r}_{2} - \mathbf{r}_{1}) = - \frac{\delta_{\alpha \gamma}}{\pi \varrho^{2}} \qquad (4.30)$$

where $\alpha, \gamma = 1, 2$ are cartesian components Eqs(4.29) and (4.30) are of the same form as in a homogeneous system, except for the average of r_1^{torust} the unit cell.

(4.30) becomes (using $\beta e^2 = 2$)

$$I_{X} = \int_{0}^{1} dX_{1} \int_{0}^{1} dY_{1} \int_{-\infty}^{\infty} dX_{2} \int_{-\infty}^{\infty} dY_{2} (X_{2} - X_{1})^{2} \varrho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = -\frac{\varrho_{0}^{2} b}{2\pi e \sin \varphi}$$
(4.31a)

$$I_{Y} = \int_{0}^{1} dX_{1} \int_{0}^{1} dY_{1} \int_{-\infty}^{\infty} dX_{2} \int_{-\infty}^{\infty} dY_{2} (Y_{2} - Y_{1})^{2} \varrho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = -\frac{\varrho_{0}^{2} \vartheta}{2\pi b \sin \vartheta} \quad (432b)$$

$$I_{XY} = \int_{0}^{1} dX_{1} \int_{0}^{1} dY_{1} \int_{-\infty}^{\infty} dX_{2} \int_{-\infty}^{\infty} dY_{2}(X_{2} - X_{1})(Y_{2} - Y_{1}) \rho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = \frac{\rho_{0}^{2} \cos \rho}{2\pi \sin \rho} (4.3 \text{ sc})$$

In Appendix A, we show that these sum rules (4.31) are indeed obeyed. For this purpose, it is convenient to express $p_T^{(2)}$ in terms of the projector P in its form (4.9).

The above mentioned proof applies to any periodic**M**, background, including the special case of a lattice of fixed charges of negligible size. For this limiting case, however, we can also give an alternative proof, which is described in Appendix B.

4.4. Irrational values of the number of particles per cell

of mobile particles per unit cell. If $p_0A = p/q$ (p, q integers), choosing a unit cell q times larger reduces the problem to $p_0A = p$, and it is easy to see that $\pi r_{1} = r_{2}$ the latter is solved by diagonalizing a q x q matrix.

What about irrational values of g_0A ? In terms of a model of a crystal, it would be a rather academic situation, since $-g_0A$ is the ratio between the charge of an ion and the charge of an electron. Nevertheless, this is a mathematically interesting situation. Furthermore, our problem is closely related to a magnetic analog⁽¹¹⁾ of importance for the theory of the quantum Hall effect⁽¹²⁾ and also to problems which arise in the theories of incommensurate structures⁽¹³⁾.

We have not been able to compute the densities for irrational values of q_0A . We only want to point out that, in the simplest case, the problem reduces to studying the solutions of an almost-Mathieu equation.

We start again with the functions (4.2), for a square unit cell of side a, and a potential modulation of the form (4.13):

$$\Psi_{\zeta,n}(\mathbf{r}) = \left[1 + \frac{(\cos 2\pi X + \cos 2\pi z)}{2}\right]^{1/2} \exp\left[-\pi y(X - \frac{\zeta + n}{2})^2 + 2\pi i(\zeta + n)Y\right]$$
(4.32)

where $\mu = \rho_0 A = \rho_0 a^2$. Computing the projector P on the space of these functions amounts to diagonalize the matrix formed by the scalar products

$$\int d\mathbf{r} \, \Psi_{\zeta,n}(\mathbf{r}) \, \Psi_{\zeta',m}(\mathbf{r}) = \delta(\zeta - \zeta') \, \mathbf{A}_{nm} \tag{4.33}$$

A simple calculation gives

$$A_{nm} = (2\nu)^{-1/2} \{ \left[1 + \lambda e^{-\pi/2\nu} \cos \frac{2\pi(\zeta+n)}{\nu} \right] \delta_{mn} + \frac{1}{2} \lambda e^{-\pi/2\nu} (\delta_{n m-1} + \delta_{n m+1}) \}$$
(4.34)

The problem of diagonalizing $\mathbf{A}_{\boldsymbol{n}\boldsymbol{m}}$ leads to

$$\frac{1}{2}(u_{n+1} + u_{n-1}) + [\cos \frac{2\pi(\zeta + n)}{\nu}]u_n = su_n$$
(4.35)

If y=1, the solution is exp($-2\pi i\eta n$), and we retrieve (4.4). If y is irrational, (4.35) is the almost-Mathieu equation in its full glory, and we leave the computation of the projector P as an open problem.

5. CONCLUSION.

At $\Gamma = 2$, we have obtained an exact solution for the equilibrium statistical mechanics of the model of fixed ions and mobile electrons introduced by Hansen et al^(5,6). We have shown that, at $\Gamma = 2$, the model exhibits the features of a conducting phase : the correlations at large separations decay faster than any inverse power law, and the system has good screening properties (the Stillinger-Lovett rule is obeyed).

On the basis of computer simulation results, it has been claimed by

Hansen and his collaborators that the coupling Γ_1 , at which the conductor-dielectric phase transition occurs, goes to 2 by upper values as the radius of the ions goes to zero. Our exact results at $\Gamma = 2$ are not in contradiction with this claim.

We check the sum rules (4.31), where

$$\rho_{T}^{(2)}(\mathbf{r}_{1}, \mathbf{r}_{2}) = -\langle \mathbf{r}_{1} | \mathbf{P} | \mathbf{r}_{2} \rangle \langle \mathbf{r}_{2} | \mathbf{P} | \mathbf{r}_{1} \rangle$$
(A.1)

using for $< \mathbf{r}_1 \mid \mathbf{P} \mid \mathbf{r}_2 >$ the representation (4.9).

We first consider

$$I_{Y} = -\int_{0}^{1} dX_{1} \int_{0}^{1} dY_{1} \exp[-2\mathscr{O}(\mathbf{r}_{1})] \sum_{\substack{n \ m' \ n' \ m'}} \int_{0}^{1} d\zeta \int_{0}^{1} d\eta \int_{0}^{1} d\zeta' \int_{0}^{1} d\eta' [f(\zeta, \eta) f(\zeta', \eta')]^{-1}$$

$$x \exp\{-2\pi i [\eta(n-m) - \eta'(n'-m')] - \frac{\pi}{\tau} (X_{1} - \zeta - n)^{2} - \frac{\pi}{\tau^{*}} (X_{1} - \zeta' - n')^{2} + 2\pi i (n-n')Y_{1}\}$$

$$x \int_{-\infty}^{\infty} dX_{2} \exp[-\frac{\pi}{\tau^{*}} (X_{2} - \zeta - m)^{2} - \frac{\pi}{\tau} (X_{2} - \zeta' - m')^{2}]$$

$$x \int_{-\infty}^{\infty} dY_{2} (Y_{2} - Y_{1})^{2} \exp[-2\mathscr{O}(\mathbf{r}_{2}) - 2\pi i (\zeta - \zeta')(Y_{2} - Y_{1}) - 2\pi i (m-m')Y_{2}] \qquad (A2)$$

Using the periodicity of \mathscr{D} , we find for the integral over Y_2 in (A2)

$$-\frac{1}{(2\pi)^2} - \frac{\partial^2}{\partial \zeta^2} \delta(\zeta - \zeta) \sum_{M} \delta_{m', M+M} \int_{0}^{1} dY_2 \exp[-2\mathscr{D}(\mathbf{r}_2) + 2\pi i MY_2]$$

In (A2), we can replace m' by m + M and n' by n + N. Using the periodicity of \mathscr{O} , we can replace $X_2 = m$ by X_2 and perform the sum over m which gives a $\delta(\eta - \eta')$; we can also replace $X_1 = n$ by X_1 and $\sum_n \int_0^1 dX_1 \dots$ by $\int_{-\infty}^{\infty} dX_1 \dots$ We obtain

$$I_{\gamma} = \frac{\rho_0^2}{(2\pi)^2} \int_0^1 d\zeta \int_0^1 d\eta \frac{1}{f(\zeta, \eta)}$$

$$\times \frac{1}{\varrho_{0}} \sum_{N} \exp(2\pi i \eta N) \int_{-\infty}^{\infty} dx_{1} \exp[-\frac{\pi}{\tau^{*}} (X_{1} - \zeta - N)^{2}] \int_{0}^{1} dY_{1} \exp[-2\emptyset(r_{1}) - 2\pi i NY_{1}]$$

$$\times \frac{1}{\varrho_{0}} \sum_{M} \exp(-2\pi i \eta M) \int_{-\infty}^{\infty} dx_{2} \exp[-\frac{\pi}{\tau} (X_{2} - \zeta - M)^{2}] \int_{0}^{1} dY_{2} \exp[-2\emptyset(r_{2}) + 2\pi i MY_{2}]$$

$$\times \frac{\partial^{2}}{\partial \zeta^{2}} \left\{ \frac{1}{f(\zeta, \eta)} \exp[-\frac{\pi}{\tau} (X_{1} - \zeta)^{2} - \frac{\pi}{\tau^{*}} (X_{2} - \zeta)^{2}] \right\}$$
(A3)

In (A.3), as a function of X₁ the derivative $(\partial^2/\partial\zeta^2)$... is a combination of terms of the form $\exp[-(\pi/\tau)(X_1-\zeta)^2]$ or $(X_1-\zeta)\exp[-(\pi/\tau)(X_1-\zeta)^2]$ or $(X_1-\zeta)^2$ exp $[-(\pi/\tau)(X_1-\zeta)^2]$. These terms enter the integral over X₁ and generate

$$\frac{1}{\rho_0} \sum_{\mathbf{N}} \exp(2\pi i\eta \mathbf{N}) \int_{-\infty}^{\infty} d\mathbf{X}_1 \exp\left[-\frac{\pi}{\tau} (\mathbf{X}_1 - \zeta)^2 - \frac{\pi}{\tau^*} (\mathbf{X}_1 - \zeta - \mathbf{N})^2\right] \int_{0}^{1} d\mathbf{Y}_1 \exp\left[-2\mathscr{G}(\mathbf{r}_1) - 2\pi i\mathbf{N}\mathbf{Y}_1\right]$$

multiplied on the right by 1 or $X_1 = \zeta$ or $(X_1 = \zeta)^2$, i.e. either $f(\zeta, \eta)$ as given by (4.8), or combinations of $\partial f/\partial \zeta$, $\partial f/\partial \eta$, and $\partial^2 f/\partial \zeta \partial \eta$. The same identifications can be made with $exp[-(\pi/\tau^*)(X_2-\zeta)^2]$ and its derivatives. The result of these identifications is

$$I_{Y} = -\frac{\varrho_{0}^{2}}{(2\pi)^{2}} \int_{0}^{1} d\zeta \int_{0}^{1} d\eta \left\{ \frac{4\pi}{\tau + \tau^{*}} + \frac{\partial}{\partial \eta} \left[\frac{1}{(\tau + \tau^{*})^{2}} \frac{1}{f} \frac{\partial f}{\partial \eta} + \frac{2i(\tau^{*} - \tau)}{(\tau + \tau^{*})^{2}} \frac{1}{f} \frac{\partial f}{\partial \zeta} \right] \right\} (A4)$$

Since f is periodic in ζ and η with periods 1, the contribution from the derivative ($\partial/\partial \eta$)... vanishes, and we obtain (4.31b).

The computation of $I_{\chi\gamma}$ follows the same lines. Instead of (A3) we obtain

$$I_{XY} = \frac{i\rho_0^2}{2\pi} \int_0^1 d\zeta \int_0^1 d\eta \frac{1}{f(\zeta,\eta)}$$

$$\frac{1}{20} \sum_{N} \exp(2\pi i \eta N) \int_{-\infty}^{\infty} dX_{1} \exp[-\frac{\pi}{\tau^{*}} (X_{1} - \zeta - N)^{2}] \int_{0}^{1} dY_{1} \exp[-2\vartheta(\mathbf{r}_{1}) - 2\pi i NY_{1}]$$

$$\frac{1}{20} \sum_{M} \exp(-2\pi i \eta M) \int_{-\infty}^{\infty} dX_{2} \exp[-\frac{\pi}{\tau} (X_{2} - \zeta - M)^{2}] \int_{0}^{1} dY_{2} \exp[-2\vartheta(\mathbf{r}_{2}) + 2\pi i MY_{2}]$$

$$\times (X_{2} - X_{1} - \frac{1}{2\pi i} \frac{\partial}{\partial \eta}) \frac{\partial}{\partial \zeta} \left\{ \frac{1}{f(\zeta, \eta)} \exp[-\frac{\pi}{\tau} (X_{1} - \zeta)^{2} - \frac{\pi}{\tau^{*}} (X_{2} - \zeta)^{2}] \right\}$$
(A5)

Again, we manage to recognize derivatives of $f(\zeta, \eta)$, with the result (4.31c).

Exchanging the X and Y axes in (4.31b) proves (4.31a).

APPENDIX B : LATTICE OF POINT PARTICLES

We take as the periodic background a lattice of fixed charged particles and consider the limiting case of point particles. We give a direct proof that the Stillinger-Lovett rule is obeyed.

Let $z_j = x_j + iy_j$ be the complex number which defines the position of the ith mobile particle; similarly let Z_j define the position of the j th fixed particle. At $\Gamma = 2$, the Boltzmann factor of a system of N mobile and N fixed point particles is

$$exp(-QH) = L^{2N} \left| \frac{\frac{\pi}{1 < k} \frac{z_i - z_k}{j < k}}{\frac{\pi}{1 < j}} \right|^2 = L^{2N} \left| \det \left\{ \frac{1}{z_i - z_j} \right\}_{i,j=1,...,N} \right|^2$$
(B.1)

where the second form is obtained by using an algebraic identity of Cauchy. In order to avoid short distance divergences, we introduce a cutoff at some small distance or, and replace (B.1) by

$$\exp(-\varrho H) = L^{2N} \det \left\{ \frac{1 - \exp[-|z_1 - z_j|^2/2\sigma^2]}{z_1 - z_j} \right\}_{i,j=1,...,N}$$
(B.2)

The limit $\sigma \rightarrow 0$ will be taken at the end of the calculation. (B.2) is again a squared determinant, and therefore the n-body truncated densities are again of the form (2.5), where now P is the projector on the space spanned by the functions $(z-Z_j)^{-1}\{1-\exp[-|z-Z_j|^2/2\sigma^2]\}$. An orthogonal basis for this space is obtained by building Bloch functions. Assuming for simplicity that the fixed particles are on the square lattice $Z = m + in, m, n \in \mathbb{Z}$, we define the Bloch functions

$$\Psi_{\zeta \eta}(x,y) = \sum_{mn} \exp[2\pi i (\zeta m + \eta n)] - \frac{1 - \exp\{-[(x-m)^2 + (y-n)^2]/2\sigma^2\}}{x + iy - (m + in)}$$
(B 3)

Using the Fourier transform

$$g(\zeta,\eta) = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \exp[2\pi i(\zeta x + \eta y)] \frac{1 - \exp[-(x^2 + y^2)/2\sigma^2]}{x + iy}$$

= $i \frac{\exp[-2\pi^2 \sigma^2(\zeta^2 + \eta^2)]}{\zeta + i\eta}$ (BA)

and the Poisson summation formula, we can rewrite Ψ as

$$\Psi_{\zeta,\eta}(x, y) = -\sum_{mn} \exp\{2\pi i [(\zeta + m)x + (\eta + n)y]\} g(\zeta + m, \eta + n)$$
(B5)

and we find for the projector

$$< \mathbf{r}_{1} |\mathbf{P}| \, \mathbf{r}_{2} > = \int_{-1/2}^{1/2} d\zeta \int_{-1/2}^{1/2} d\eta \, \frac{\Psi_{\zeta, \eta}(\mathbf{x}_{1}, \mathbf{y}_{1}) \, \Psi_{\zeta, \eta}(\mathbf{x}_{2}, \mathbf{y}_{2})}{\int_{-1/2}^{1/2} dx \int_{-1/2}^{1/2} dy |\Psi_{\zeta, \eta}(\mathbf{x}, \mathbf{y})|^{2}} = \int_{-1/2}^{1/2} d\zeta \int_{-1/2}^{1/2} d\eta \\ \times \frac{\sum_{n_{1} n_{1} m_{2} n_{2}}^{n_{2}} \exp\{2\pi i [(\zeta + m_{1})\mathbf{x}_{1} + (\eta + n_{1})\mathbf{y}_{1} - (\zeta + m_{2})\mathbf{x}_{2} - (\eta + n_{2})\mathbf{y}_{2}]\}g(\zeta + m_{1}, \eta + n_{1})\bar{g}(\zeta + m_{2}, \eta + n_{2})}{\sum_{m_{1} m_{1} m_{2} m_{2}}}$$
(B.6)

The structure of (B.6) allows us at once to check the average density sum rule

$$\int_{-1/2}^{1/2} dx \int_{-1/2}^{1/2} dy \, \varrho(\mathbf{r}) = \int_{-1/2}^{1/2} dx \int_{-1/2}^{1/2} dy < \mathbf{r} |P| \, \mathbf{r} > = 1$$
(B.7)

and the neutrality sum rule

$$\int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dy_2 \rho_T^{(2)}(\mathbf{r}_1, \mathbf{r}_2) = -\int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dy_2 < \mathbf{r}_1 |P| \mathbf{r}_2 > < \mathbf{r}_2 |P| \mathbf{r}_1 >$$
$$= - < \mathbf{r}_1 |P| \mathbf{r}_1 > = -\rho(\mathbf{r}_1)$$
(B.8)

We turn to the Stillinger-Lovett second moment

$$I = 2 \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dy_2 \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \frac{g^{(2)}}{r} (\mathbf{r}_1, \mathbf{r}_2) (x_1 - x_2)^2$$

= $-2 \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dy_2 \int_{-1/2}^{1/2} \frac{g^{(2)}}{r} (\mathbf{r}_1, \mathbf{r}_2) (x_1 - x_2)^2$
(B.9)

Using the representation (B.6) of the projector, we perform the space integrals with variables r_1 and $r_2 - r_1$; we find

$$I = \frac{1}{2\pi^2} \int_{-1/2}^{1/2} d\zeta \int_{-1/2}^{1/2} d\eta \int_{-1/2}^{1/2} d\zeta' \int_{-1/2}^{1/2} d\eta' \, \delta^{-}(\zeta - \zeta') \, \delta(\eta - \eta') \frac{f(\zeta, \eta; \zeta', \eta')f(\zeta', \eta'; \zeta, \eta)}{f(\zeta, \eta; \zeta, \eta)f(\zeta', \eta'; \zeta', \eta')}$$
(B.10)

where

ĺ

$$f(\zeta, \eta; \zeta', \eta') = \sum_{mn} g(\zeta + m, \eta + n) \bar{g}(\zeta' + m, \eta' + n)$$
 (B.11)

Replacing $\delta^{*}(\zeta-\zeta')$ by $\delta(\zeta-\zeta')(\delta^{2}/\delta\zeta^{2}),$ we obtain

$$I = -\frac{1}{\pi^{2}} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left[\frac{1}{f(\zeta,\eta;\zeta,\eta)} - \frac{\partial^{2}f(\zeta,\eta;\zeta,\eta)}{\partial \zeta\partial \zeta'} - \frac{1}{f^{2}(\zeta,\eta;\zeta,\eta)} - \frac{\partial f(\zeta,\eta;\zeta',\eta)}{\partial \zeta} - \frac{\partial f(\zeta,\eta;\zeta',\eta)}{\partial \zeta'} - \frac{\partial f(\zeta,\eta;\zeta',\eta)}{\partial \zeta'} \right]_{\zeta'=\zeta}$$
(B.12)

The function $f(\zeta, \eta; \zeta, \eta)$ defined by (8.4) and (8.11) is

$$f(\zeta, \eta; \zeta, \eta) = \sum_{mn} \frac{\exp\{-4\pi^2 \sigma^2 [\zeta + m)^2 + (\eta + n)^2]\}}{(\zeta + m)^2 + (\eta + n)^2}$$
(B.13)

It is convenient to display the singularity at $\zeta = \eta = 0$, coming from the term m = n = 0, and to represent f, in the integration domain $|\zeta|$, $|\eta| < 1/2$, by its Laurent expansion

$$f(\zeta, \eta; \zeta, \eta) = \frac{1}{\zeta^2 + \eta^2} + A + ...$$
 (B.14)

In the point particle limit, $\sigma \rightarrow 0$, the sum in (B.13) diverges for large (m, n) and A becomes infinite; however the terms of higher order in (ζ , η) represented by dots in (B.14) remain finite and can be neglected. If a similar analysis is performed for the derivatives of f appearing in (B.12), it is easily seen that only the term m = n = 0 plays a role as $\sigma \rightarrow 0$, because the sums on (m, n) which define these derivatives do <u>not</u> diverge for large (m, n); therefore

$$\frac{\partial^2 f(\zeta,\eta;\zeta',\eta)}{\partial \zeta \partial \zeta'} \Big|_{\zeta'=\zeta} \sim \Big| \frac{\partial g(\zeta,\eta)}{\partial \zeta} \Big|_{2}^{2} \sim \frac{1}{(\zeta^{2}+\eta^{2})^{2}}$$
(B.15)

and

$$\frac{\partial f(\zeta,\eta;\zeta',\eta)}{\partial \zeta} \frac{\partial f(\zeta,\eta;\zeta',\eta)}{\partial \zeta'}\Big|_{\zeta'=\zeta} \sim \left|g(\zeta\eta)\frac{\partial g(\zeta,\eta)}{\partial \zeta}\right|^2 \sim \frac{1}{(\zeta^2+\eta^2)^3} (B.16)$$

Using (B.14), (B.15), and (B.16) in (B.12), we find

$$I \sim -\frac{1}{\pi^2} \int_{-1/2}^{-1/2} \int_{-1/2}^{-1/2} \frac{A}{\left[1 + A(\zeta^2 + \eta^2)\right]^2}$$
(B.17)

As A goes to infinity, the integration domain in (B.17) can be extended to infinity and the integral evaluated in polar coordinates. Thus

$$\lim_{\sigma \to 0} I = -\frac{1}{\pi}$$
(B.18)

This proves that the Stillinger-Lovett rule holds in the point-particle limit $\sigma \rightarrow 0$.

We have not been able to compute I for a finite value of σ . It should be remarked that the Boltzmann factor (B.2) corresponds, when σ is non zero, to a complicated many-body interaction which would become the Coulomb law only if all particles were far apart from one another. There is no obvious reason for believing that such a system obeys or does not obey the Stillinger-Lovett rule. This is in contrast with the case of a bona fide system with extended fixed particles, which does obey the rule, as shown in Appendix A.

ACKNOWLEDGEMENTS

We are indebted to A. Alastuey for stimulating discussions. Part of this work was done while one of us (L.B.) was Invited Professor at the University of Paris-Sud. L. B also a chancon ledges partial support from The of Norval festance

REFERENCES

- 1. B. Jancovici, Phys. Rev. Lett. <u>46</u> : 386 (1981).
- 2. Ph. Choquard, P.J. Forrester and E.R. Smith, J. Stat. Phys. 33 13 (1983).
- A. Alastuey and J.L. Lebowitz, J. Phys. (France) <u>45</u> 1859 (1984), and references quoted therein.
- 4. F. Cornu and B. Jancovici, to be published.
- 5. J. Clérouin and J.P. Hansen, Phys. Rev. Lett. <u>54</u> : 2277 (1985).
- 6. J. Clérouin, J.P. Hansen and B. Piller, Phys. Rev. A <u>36</u> : 2793 (1987).
- 7. F.H. Stillinger and R. Lovett, J. Chem. Phys. <u>49</u> : 1991 (1968).
- 8. See e.g. C. Cohen-Tannoudji, B. Diu and F. Laloë, Mécanique Quantique (Hermann, Paris, 1977).
- 9. B. Jancovici, J. Stat. Phys. 28 : 43 (1982).
- 10. S.L. Carnie and D.Y.C. Chan, Chem. Phys. Lett. 77 : 437 (1981).
- 11. D.R. Hofstadter, Phys. Rev. B 14: 2239 (1976).
- D.J. Thouless, M. Kohmoto, M.P. Nightingale and M. denNijs, Phys. Rev. Lett. <u>49</u>: 405 (1982).
- 13. See e.g. 8. Souillard, Phys. Reports, <u>103</u> : 41 (1984).

1

8

FIGURE CAPTION

Fig.1. Potential modulation (a), background density (b), and particle density (c).



und. This surface must be turned upside down shifted upwards by one unit. No other change is needed ash. L. p. - 6 g(r) as A stands have ×



