

2

AD-A196 556

DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

2a. SECURITY CLASSIFICATION AND AUTHORITY Unclassified			1b. RESTRICTIVE MARKINGS		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE JUN 01 1988			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
4. PERFORMING ORGANIZATION REPORT NUMBER H			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Scripps Clinic and Research Foundation		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) 10666 North Torrey Pines Road LaJolla, CA 92037			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-87-C-7074		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62772A	PROJECT NO. 3M1- 62772A874	TASK NO. AC
			WORK UNIT ACCESSION NO. 121		
11. TITLE (Include Security Classification) (U) Metabolic Regulation of 2,3-DPG in Stored Red Blood Cells					
12. PERSONAL AUTHOR(S) Shobhana Vora					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 6/1/87 to 12/31/87		14. DATE OF REPORT (Year, Month, Day) 1988 March 1	
15. PAGE COUNT 17					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
06	01		erythrocyte, storage, glyoxalate, glycolate, oxalate,		
06	04		adenosine triphosphate, 2,3-diphosphoglycerate		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
The effect of 42 different compounds on red cell ATP and 2,3-DPG levels was studied. Of these oxalate, glyoxalate, ethyl oxaloacetate and L-phenylalanyl-L-alanine were found to improve 2,3-DPG levels. Glyoxalate and ethyl oxaloacetate exert their effect by conversion to oxalate. The latter compound has been shown to inhibit pyruvate kinase and in this way modifies red cell metabolism so as to cause an increase in red cell 2,3-DPG levels. L-phenylalanyl-L-alanine is rapidly hydrolyzed to phenylalanine and alanine, both of which also inhibit pyruvate kinase.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Mary Frances Bostian			22b. TELEPHONE (Include Area Code) 301-663-7325		22c. OFFICE SYMBOL SGRD-RMI-S

AD _____

METABOLIC REGULATION OF 2,3-DPG IN STORED RED BLOOD CELLS

FINAL REPORT

SHOBHANA VORA

MARCH 1, 1988

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21701-5012

Contract No. DAMD17-87-C-7074

Scripps Clinic and Research Foundation
10666 North Torrey Pines Road
LaJolla, CA 92037

Approved for public release; distribution unlimited

The findings in this report are not to be construed as an official
Department of the Army position unless so designated by other
authorized documents.

FOREWORD

For the protection of human subjects the investigator has adhered to policies of applicable Federal Law 45CFR46.



Accession For		
DTIC GRA&I	☒	
DTIC TAB	☐	
Unannounced	☐	
Justification		
By		
Distribution/		
Availability Codes		
Dist	Avail and/or	Special
A-1		

TABLE OF CONTENTS

FOREWORD	1
PROBLEM UNDER STUDY	
BACKGROUND AND REVIEW	
RATIONALE USED IN CURRENT STUDY	
EXPERIMENTAL METHODS	
RESULTS	
DISCUSSION AND CONCLUSIONS	
REFERENCES	
DISTRIBUTION LIST	

TABLE

1. The Effect of Various Additives on the ATP and 2,3 DPG Levels of Stored Blood

FIGURES

1. The effect of storage of packed red cells with the addition of ethyloxaloacetate (Et-oxaloacetate) L-phenylalanyl-L-alanine (phenylalanyl-ala) and phenylalamine butyl ester (phenylala butyl ester) at the final concentrations shown on ATP and 2,3-diphosphate glycerate levels
2. The effect of storage of red cells with the addition of α -amino isobutyric acid (amino-butyrate) and glyoxalate at the final concentrations shown on red cell ATP and 2,3-DPG levels
3. The structural relationship between glycolate, glyoxalate, and oxalate

1. Problem under study. The investigations carried out under this contract were aimed at attempting to find better mechanisms for maintenance of red cell 2,3-DPG levels during storage.

2. Background and Review. Since the studies of Benesch and Benesch (1) and Chanutin and Curnish (2) showing that the oxygen affinity of hemoglobin was strongly influenced by 2,3 phosphoglycerate (2,3-DPG) levels there has been a great deal of interest in attempting to improve the stability of 2,3-DPG levels during storage. This problem has been reviewed extensively previously by the principal investigator (3-6) and by others (7-9).

3. Rationale used in current study. The basic premise underlying the current studies was that inhibitors of glycolysis, particularly at the pyruvate kinase step, could exert an affect on the maintenance of 2,3-DPG levels.

4. Experimental Methods. In the initial screening studies blood from normal volunteer donors was drawn into CPD (citrate-phosphate-dextrose) or CPDA1 (citrate-phosphate-dextrose-adenine) solution (Fenwal Laboratories). The blood was divided into 4 nearly equal portions and was distributed into plastic blood bags. The volume of each aliquot was measured by weighing the bag and minor differences in volume were corrected by addition of 0.9% NaCl solution. Various substances were added in the saline to each of the aliquots. Samples without additions served as controls. ATP and 2,3-DPG estimations were carried out weekly or semiweekly for at least 6 weeks using standard methods (10).

In subsequent studies, 450 ml blood was drawn into 63 ml CPDA1, divided into 95 or 99 ml portions and distributed into plastic blood bags. The substance under study was added in 1 ml or 5 ml 0.9% NaCl to each 95 or

99 ml aliquot of blood. Saline without preservative was added to the control.

The blood samples were incubated at 37°C for 1 to 1-1/2 hr. prior to storage. The internal pH of red cells was measured by freezing and thawing packed red cells and measuring the pH of the lysate at 0°C with a Corning model 150 pH meter. Oxalate determinations (11) were carried out on plasma ultrafiltrates made from blood with added glyoxalate before storage and after 1 week.

5. Results. Table 1 lists the 42 substances screened and an evaluation of their affect on ATP and 2,3-DPG levels during storage.

Inspection of these data indicated that oxalate, glyoxalate, ethyl oxaloacetate, L-phenylalanyl-L-alanine, phenylalanine t-butyl ester, and aminoisobutyric acid might be effective in the improving maintenance of 2,3 DPG concentrations.

Accordingly, the effects of these compounds was reinvestigated, and the effect of oxalate, glyoxalate, ethyl oxaloacetate, and L-phenylalanyl-L-alanine on the maintenance of 2,3 DPG were found to be reproducible. Representative experiments are shown in figures 1 and 2.

To determine whether the effect of glyoxalate might be due to its conversion to oxalate, previously known to exert an effect on 2,3-DPG levels (12,13), the oxalate contents of blood samples stored with and without oxalate were determined. The concentration of oxalate in control blood was less than 5 moles. In contrast, the concentration of oxalate in blood stored with glyoxalate rose rapidly, reaching a 500 mole concentration within one hour of collection. Conversion of glyoxalate to oxalate required the presence of red cells. Glyoxalate added to plasma would not convert it to oxalate.


Amino acid analysis of blood stored with L-phenylalanyl-L-alanine showed complete hydrolysis of the dipeptide to phenylalanine and alanine within a week of storage.

6. Discussion and Conclusions. A large number of compounds have been tested for their effect on erythrocyte 2,3-DPG and ATP levels of blood stored in citrate phosphate adenine solutions. Inhibitors of ATPase, such as ouabain did not affect ATP levels. This finding is not surprising, since red cell ATPase is scarcely active at 4°C (14,15). Neither was an effect observed with most of the other compounds tested.


Four compounds were found to be effective in raising 2,3-DPG levels. These were oxalate, glyoxalate, ethyl oxaloacetate and L-phenylalanyl-L-alanine. The effect of oxalate was first discovered by Kandler et al (13) when he investigated the putative effect of ascorbate on 2,3-DPG levels. Pure ascorbate, formerly thought to be a potent agent in preserving 2,3-DPG (16) was without effect, but oxalate, contaminating the ascorbate was found to be the active substance. Recently we have shown that even .05 mM oxalate has a substantial effect on red cell 2,3-DPG levels and that the effect of this substance is largely due to inhibition of pyruvate kinase (12). Ethyl oxaloacetate is known to hydrolyze to oxalate, acetate, and ethanol (17).

The effect of glyoxalate was more of a surprise. Tested because of its similarity to glycolate and oxalate (Fig. 3), glyoxalate differs from oxalate only by a single oxidation state. Glyoxalate is a 2-carbon analogue of pyruvate while glycolate may be regarded as the corresponding 2-carbon analogue of lactate. Red cell membranes are freely permeable to the glyoxalate anion (18). In the presence of NAD, glyoxalate is oxidized to oxalate by heart (19) and muscle (20) lactate dehydrogenase. The same enzymes also have the capability of reducing glyoxalate to glycolate in the


presence of NAD. Oxalate, increases 2,3-DPG levels of stored red cells, and the quantity formed from glyoxalate is clearly sufficient to account for the effect of glyoxalate on 2,3-DPG levels. Oxalate appeared rapidly in blood to which glyoxalate is added. Presumably it is formed by oxidation of glyoxalate in the reaction:



For this reaction to create 0.5 mM oxalate, the concentration that was achieved within an hour of blood collection, the NADH formed must be re-oxidized to NAD^+ . There is insufficient pyruvate in blood to serve as an oxidizing agent for the amount of NAD^+ required. It may well be that a portion of the glyoxalate itself regenerates the need for NAD^+ in the reaction:



The overall reaction, then, is:



Thus, the effect of glyoxalate seems to be due simply to its coupled oxidate to oxalate. L-phenylalanyl-L-alanine had a pronounced effect on 2,3-DPG levels after 1 week's storage, but none subsequently. It is of interest that high concentrations of L-alanine (21,22) and L-phenylalanine (22) have previously been shown to aid in 2,3-DPG preservation in stored red cells. The red cell is rich in peptidase activity (23) and is nearly

impermeable to those dipeptides that have been tested.

REFERENCES

1. Benesch R, Benesch RE: The effect of organic phosphates from the human erythrocyte on the allosteric properties of hemoglobin. *Biochem Biophys Res Commun* 26:162-167,1967.
2. Chanutin A, Curnish RR: Effect of organic and inorganic phosphates on the oxygen equilibrium of human erythrocytes. *Arch Biochem Biophys* 121:96-102,1967.
3. Beutler E: Experimental blood preservatives for liquid storage. Greenwalt TJ, Jamieson GA eds. In: *The Human Red Cell In Vitro*, Grune & Stratton, Inc., New York, 1974, pp. 189-216.
4. Chaplin H Jr, Beutler E, Collins JA, Giblett ER, Polesky HF: Current status of red-cell preservation and availability in relation to the developing national blood policy. *N Engl J Med* 291:68-74,1974.
5. Beutler E: Viability, function, and rejuvenation of liquid-stored red cells. Chaplin H Jr, Jaffe ER, Lenfant C, Valeri CR eds. In: *Preservation of Red Blood Cells*, National Academy of Sciences, Washington, 1973, pp. 195-214.
6. Beutler E: The maintenance of red cell function during liquid storage. Schmidt PJ eds. In: *Progress in Transfusion and Transplantation*, Washington, 1972, pp. 285-297.
7. Valeri CR: *Blood Banking and the Use of Frozen Blood Products*, Chemical Rubber Co., Cleveland, OH, 1976, p 109.
8. Finch CA: Adenine-supplemented blood. *Vox Sang* 48:319-322,1985.
9. Hoegman CF: Anticoagulants in current use. Cash J eds. In: *Anticoagulants*, 1st Edition. Churchill Livingstone, Edinburgh, 1986, pp. 60-77.

10. Beutler E: Red Cell Metabolism: A Manual of Biochemical Methods, 3rd Edition. Grune & Stratton, Inc., New York, NY, 1984.
11. Kohlbecker G, Butz M: Direct spectrophotometric determination of serum and urinary oxalate with oxalate oxidase. J Clin Chem Clin Biochem 19:1103-1106,1981.
12. Beutler E, Forman L, West C: Effect of oxalate and malonate on red cell metabolism. Blood 70:1389-1393,1987.
13. Kandler R, Grode G, Symbol R, Hickey G: Oxalate is the active component that produces increased 2,3-DPG in ascorbate stored red cells. Transfusion 26:563 (abstract),1986.
14. Wood L, Beutler E: Temperature dependence of sodium-potassium activated erythrocyte adenosine triphosphatase. J Lab Clin Med 70:287-294,1967.
15. Beutler E, Kuhl W, West C: Volume control of erythrocytes during storage: The role of mannitol. Transfusion, In press, 1988.
16. Wood LA, Beutler E: The effect of ascorbate on the maintenance of 2,3-diphosphoglycerate (2,3-DPG) in stored red cells. Br J Haematol 25:611-618,1973.
17. Windholz M, Budavari S, Blumetti RF, Otterbein ES eds. The Merck Index, 10th Edition. Merck & Co., Inc., Rahway, New Jersey, 1983, p 554.
18. Hubbard AR, Sprandel U, Chalmers RA: Transport of oxalic acid, glycollic acid, glyoxylic acid and benzoic acid by resealed erythrocyte 'ghosts' prepared by a dialysis technique. Biochem Soc Trans 7:958-960,1979.
19. Warren WA: Catalysis of both oxidation and reduction of glyoxylate by pig heart lactate dehydrogenase isozyme 1. J Biol Chem 245:1675-1681,1970.
20. Duncan RJS, Tipton KF: The oxidation and reduction of glyoxylate by

- lactic dehydrogenase. Eur J Biochem 11:58-61,1969.
21. Dawson RB, Ottinger WE, Chiu WE, Salib AA, Fagan DS, Keath JR: Control of red cell 2,3-DPG levels in vitro and a proposal for in vivo control in response to hypoxia and metabolic demand. In: Cellular and Molecular Aspects of Aging: The Red Cell as a Model, Alan R. Liss, Inc., New York, 1984, pp. 349-365.
 22. Vora S: Metabolic manipulation of key glycolytic enzymes: a novel proposal for the maintenance of red cell 2,3-DPG and ATP levels during storage. Biomed Biochim Acta 46:285-289,1987.
 23. Melloni E, Salamino F, Sparatore B, Michetti M, Morelli A, Benatti U, De Flora A, Pontremoli S: Decay of proteinase and peptidase activities of human and rabbit erythrocytes during cellular aging. Biochim Biophys Acta 675:110-116,1981.

Personnel

Ernest Beutler, M.D.
Linda Forman, B.Sc.
Susan Sackmann, B.Sc.
Carol West, A.A.

DISTRIBUTION LIST

4 copies	Commander Letterman Army Institute of Research (LAIR), Bldg. 1110 ATTN: SGRD-ULZ-RC Presidio of San Francisco, CA 94129-6815
1 copy	Commander US Army Medical Research and Development Command ATTN: SGRD-RMI-S Fort Detrick, Frederick, Maryland 21701-5012
2 copies	Defense Technical Information Center (DTIC) ATTN: DTIC-DDAC Cameron Station Alexandria, VA 22304-6145
1 copy	Dean School of Medicine Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, MD 20814-4799
1 copy	Commandant Academy of Health Sciences, US Army ATTN: AHS-CDM Fort Sam Houston, TX 78234-6100

TABLE I. The Effect of Various Additives on the ATP and 2,3 DPG Levels of Stored Blood

Preservative	Additive	Concentration (mM)	ATP	2,3-DPG
CPD	glyoxylate	1	↓	↑
CPD	α-aminoisobutyric acid	1	↑	↑
CPD	cysteine*	5	-	-
CPD	3-I-butyl-1-methyl-xanthene	3	-	-
CPD	oxalate	2	↓	↑
CPDA1	palmitic acid	0.2	-	-
CPDA1	myristic acid	0.2	-	-
CPDA1	hydroxypyruvate	2	↑	-
CPDA1	ethyl oxaloacetate	2	↑	↑
CPDA1	keto glutarate	5	-	-
CPDA1	phenyl pyruvate	2	-	-
CPDA1	isocitrate	1	-	↓
CPDA1	t-aconitate	1	↑	↓
CPDA1	malonate	5	-	↓
CPDA1	oxaloacetate	2	-	↓
CPDA1	phenylalanine methyl ester	4.3	-	-
CPDA1	phenylalanyl methionine	2	-	-
CPDA1	oxobutyrate	1	↑	-
CPDA1	cysteine methyl ester	2	-	-
CPDA1	maleate	2.5	-	-
CPDA1	malate	2.5	-	-

TABLE I (Continued)

Preservative	Additive (mM)	Concentration	ATP	2,3-DPG
CPDA1	tartrate	2.5	-	-
CPDA1	cupric chloride	0.002	-	-
CPDA1	nickel sulfite	0.005	-	-
CPDA1	cobalt sulfate	0.005	-	-
CPDA1	valine	5	-	-
CPDA1	proline	5	-	-
CPDA1	tryptophan	5	-	-
CPDA1	D-alanine	5	†	-
CPDA1	isoleucine	5	-	-
CPDA1	methionine	5	†	-
CPDA1	L-phenylalanyl-L-alanine	5	†	†
CPDA1	oleate	0.05	-	-
CPDA1	linoleate	0.05	-	-
CPDA1	nicotinate	1	-	-
CPDA1	nicotinaldehyde	1	-	-
CPDA1	quinolinate	1	-	-
CPDA1	glycolaldehyde	1	-	-
CPDA1	nicotinamide	40	-	-
CPDA1	methyl glyoxal	1	-	-
CPDA1	Quabain	0.2	-	-
CPDA1	Quercitin	0.005	-	-

† >25% increase

↓ >25% decrease

* amino acids are in the L-conformation unless otherwise indicated




FIGURE 1 The effect of storage of packed red cells with the addition of ethyloxalacetate (Et-oxalacetate) L-phenylalanyl-L-alanine (phenylalanyl-ala) and phenylalamine butyl ester (phenylala butyl ester) at the final concentrations shown on ATP and 2,3-diphosphate glycerate levels.








FIGURE 2 The effect of storage of red cells with the addition of α -amino isobutyric acid (Amino-butyrate) and glyoxalate at the final concentrations shown on red cell ATP and 2,3-DPG levels.



GLYCOLATE



GLYOXALATE



OXALATE

FIGURE 3 The structural relationship between glycolate, glyoxalate, and oxalate. All three compounds are 2-carbon carboxylic acids, differing from each other only in their state of oxidation. They are interconvertible through the lactate dehydrogenase reaction (see text).