

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR-89-0168		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION AFOSR/NC		
6a. NAME OF PERFORMING ORGANIZATION California Institute of Technology		6b. OFFICE SYMBOL (If applicable)		7b. ADDRESS (City, State, and ZIP Code) Building 410, Bolling AFB DC 20332-6448	
6c. ADDRESS (City, State, and ZIP Code) Pasadena, CA 91125		8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-87-0039	
8b. OFFICE SYMBOL (If applicable) NC		8c. ADDRESS (City, State, and ZIP Code) Building 410, Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61102F		PROJECT NO. 2303	
				TASK NO. B3	
				WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification) (U) (2+1) REMPI of NO via the $D^2\Sigma^+$ state: Rotational branching ratios					
12. PERSONAL AUTHOR(S) H. Rudolph, S. N. Dixit, V. McKoy and Winifred M. Huo					
13a. TYPE OF REPORT Reprint		13b. TIME COVERED FROM 11/1/87 TO 11/1/88		14. DATE OF REPORT (Year, Month, Day) 88/12/14	
15. PAGE COUNT 3 (three)					
16. SUPPLEMENTARY NOTATION Chemical Physics Letters, Vol. 137, No. 6, 3 July 1987, 521-523					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Recent photoelectron spectroscopic studies in a (2+1) REMPI of ¹⁵ N ₂ via the Rydberg $D_{n0}^2\Sigma^+$ state have revealed anomalous ionic rotational branching ratios. We have performed ab initio calculations of these branching ratios and find that the molecular nature of the ionization continuum plays an essential role in the dynamics. Even though the bound orbital is very atomic-like (>98% p-like), the photoelectron continuum wavefunction is quite sensitive to the non-spherical nature of the molecular ionic potential and causes a strong persistence of the p-partial wave which, in turn, leads to a large $N=0$ peak. reprints: Delta DTIC ELECTE 23 FEB 1989					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL L P DAVIS, MAJOR, USAF			22b. TELEPHONE (Include Area Code) (202) 767-4963		22c. OFFICE SYMBOL AFOSR/NC

AFOSR-TR- 89 - 0168

(2+1) REMPI OF NO VIA THE $D^2\Sigma^+$ STATE: ROTATIONAL BRANCHING RATIOS**H. RUDOLPH, S.N. DIXIT¹, V. McKOY***Arthur Amos Noyes Laboratory of Chemical Physics², California Institute of Technology, Pasadena, CA 91125, USA*

and

Winifred M. HUO*NASA-Ames Research Center MS230-3, Moffett Field, CA 94035, USA*

Received 9 March 1987

Recent photoelectron spectroscopic studies in a (2+1) REMPI of NO via the Rydberg $D^2\Sigma^+$ state have revealed anomalous ionic rotational branching ratios. We have performed ab initio calculations of these branching ratios and find that the *molecular nature* of the ionization continuum plays an essential role in the dynamics. Even though the bound orbital is very atomic-like (>98% p-like), the photoelectron continuum wavefunction is quite sensitive to the non-spherical nature of the molecular ionic potential and causes a strong persistence of the p-partial wave which, in turn, leads to a large $\Delta N=0$ peak.

High resolution studies of photoelectron spectra (PES) produced by resonant-enhanced multiphoton ionization (REMPI) of individual ro-vibronic states are an important probe of the dynamics of such photoionization processes. Whereas ionic rotational states have been resolved in the photoelectron spectrum of H_2 with relative ease [1,2], the resolution of ionic states in heavier molecules has been very difficult because of the smaller rotational constants. Such studies have only been recently performed in NO [3-5]. The (1+1) REMPI PES of NO by Wilson et al. [4] revealed a number of peaks corresponding to various values of the difference (ΔN) in the angular momentum quantum numbers for the excited state (N_i) and the ion (N_+) ($\Delta N \equiv N_+ - N_i$). Ab initio calculations [6] of ionic rotational branching ratios helped to clarify some features of these spectra such as the dominance of the $\Delta N=0$ peak and the suppression of the $\Delta N = \pm 1$ peaks. The recent (2+1) REMPI PES via the $D^2\Sigma^+$ ($3p\sigma$) state of NO by Viswanathan et al. [5] shows a strong $\Delta N=0$ peak.

Due to the selection rule [6,7] $\Delta N + l = \text{odd}$, valid for ionization of a Σ state leaving the ion in a Σ state, a $\Delta N=0$ peak, of significant intensity implies that the photoelectron orbital must have substantial odd-wave (p, f, ...) character. This, combined with the "pure" Rydberg p nature of the 7σ orbital in the D state, led the authors of ref. [5] to suggest that there may be a strong *l*-mixing in the ionization continuum. In this communication, we present results of ab initio calculations of these ionic rotational spectra and compare them with the data of ref. [5]. Indeed, we find the p wave to be quite significant in the composition of the photoelectron orbital, hence accounting for the strong $\Delta N=0$ peak. More importantly, these results illustrate the need for an adequate description of electronic continuum states in quantitative studies of low-energy photoionization dynamics.

Our calculations were done using the theoretical framework described earlier [8]. The photoelectron wavefunction is calculated in the frozen-core Hartree-Fock approximation using the iterative Schwinger variational technique for an internuclear separation of $2.008 a_0$. This is the equilibrium internuclear distance for the NO^+ ion and most of the

¹ Present address: Lawrence Livermore National Laboratory, L-421, P.O. Box 808, Livermore, CA 94550, USA.

² Contribution No. 7538.

89 2 15 192

Approved for public release;
distribution unlimited.

Rydberg states of NO. We use a Gaussian basis set of (10s6p1d) on each atom contracted to [6s4p1d], supplemented with a (7s7p) uncontracted Gaussian basis at the center of mass (c.m.) of the molecule. This large and diffuse basis set on the origin assures the proper diffuseness of the Rydberg states. The electronic wavefunction of the 7σ orbital of the $D^2\Sigma^+$ state is calculated in the improved virtual orbital (IVO) scheme [9], using the NO^+ core ($E(NO^+) = -128.94691$ au). The total energy of the $D^2\Sigma^+$ state in this approximation is -129.03866 au. The use of the NO^+ core is based on the large relaxation of the NO core in the D state, compared to the ground state. Further details will be given elsewhere [10].

The 7σ IVO orbital calculated using the NO^+ core has 0.5% s, 99.2% p, and 0.1% d character, in a single center expansion about the c.m., in agreement with literature values [5,11]. For a photoelectron kinetic energy of ≈ 0.66 eV [5] the relative transition moments which are proportional to $|r_{if}^{l\mu}|^2$, as defined by eq. (13) of ref. [6], are 0.0498 ($l=0$), 0.0566 ($l=1$), 0.0424 ($l=2$), 0.0047 ($l=3$), and 0.0002 ($l=4$) in the $k\sigma$ channel, and 0.0141 ($l=1$), 0.2093 ($l=2$), 0.0134 ($l=3$), and 0.0019 ($l=4$) in the $k\pi$ channel. These results support the conclusions of Viswanathan et al. [5], that the $l=1$ (p wave) is indeed present and very strong in the continuum. Note that an "atomic" like analysis would have predicted ionization into s and d waves only. The unusually large strength of the p wave is a consequence of the l -mixing in the photoelectron continuum, caused by the non-spherical potential of the molecular ion. The core potential for higher Rydberg states approaches the potential of the ion, and this is expected to cause similar anomalous features in photoionization out of higher Rydberg states.

In fig. 1 we compare the calculated photoelectron spectrum with the experimentally measured ones for two-photon resonant excitation via the S_{21} (11.5) line. This transition was chosen as it is unmixed and the calculation of relative M_l populations in the excited state is simple. Our calculated rotational branching ratios are convoluted with a Lorentzian detection function having a fwhm of about 6 meV. The agreement of the theoretical and experimental PES is quite good for detection parallel and excellent for detection perpendicular to the laser polarization.




Fig. 1. Experimental [5] (left) and calculated (right) photoelectron spectra with the laser tuned to the S_{21} (11.5) line of the $X^2\Pi \rightarrow D^2\Sigma^+$ transition of NO, for (a) laser light polarized parallel ($\parallel$) to detection, (b) laser light polarized perpendicular ($\perp$) to detection.

The $\Delta N=0$ peak is quite significant in the parallel detection and is small, as observed experimentally as well, in the perpendicular detection. Combined with the selection rule $\Delta N + l = \text{odd}$, these results point to significant odd- l partial waves. More importantly, the good qualitative agreement between the theory and experiment illustrates the ability of ab initio calculations to unravel subtle features in the photoionization dynamics.

This material is based on research supported by the National Science Foundation under grant No. CHE-8521391 and AFOSR grant No. 87-0039. One of us (HR) gratefully acknowledges the support from the Danish Natural Science Research Council.

References

- [1] J.E. Pollard, D.J. Trevor, J.E. Reutt, Y.T. Lee and D.A. Shirley, *J. Chem. Phys.* 77 (1982) 34.
- [2] S.T. Pratt, P.M. Dehmer and J.L. Dehmer, *J. Chem. Phys.* 78 (1983) 4315.
- [3] K. Müller-Detlefs, M. Sander and E.W. Schlag, *Chem. Phys. Letters* 112 (1984) 291.

- [4] W.G. Wilson, K.S. Viswanathan, E. Sekreta and J.P. Reilly, J. Phys. Chem. 88 (1984) 672.
- [5] K.S. Viswanathan, E. Sekreta, E.R. Davidson and J.P. Reilly, J. Phys. Chem. 90 (1986) 5078.
- [6] S.N. Dixit, D.L. Lynch, V. McKoy and W.M. Huo, Phys. Rev. A32 (1984) 1267.
- [7] S.N. Dixit and V. McKoy, Chem. Phys. Letters 128 (1986) 49.
- [8] R.R. Lucchese, G. Raseev and V. McKoy, Phys. Rev. A25 (1982) 2572.
- [9] W.J. Hunt and W.A. Goddard III, Chem. Phys. Letters 24 (1974) 464.
- [10] H. Rudolph, S.N. Dixit and V. McKoy, J. Chem. Phys., to be published.
- [11] K. Kaufmann, C. Nager and M. Jungen, Chem. Phys. 95 (1985) 385.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1 20	

QUALITY INSPECTED