

AD-A209 351

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 101

Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field

by

H. R. Lee, H. G. Oh, Thomas F. George and C. I. Um

Prepared for Publication

in

Journal of Applied Physics

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

June 1989

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
JUN 20 1989
S E D

89 6 19 124

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-101			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field					
12. PERSONAL AUTHOR(S) H. R. Lee, H. G. Oh, Thomas F. George and C. I. Um					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) June 1989	
15. PAGE COUNT 17					
16. SUPPLEMENTARY NOTATION Prepared for publication in Journal of Applied Physics					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	ELECTRON ENERGY LEVELS		
			CYCLOTRON ORBITS		
			QUANTUM WELL		
			QUANTUM CALCULATION, (1e)		
			IN-PLANE MAGNETIC FIELD		
			BULK LANDAU LEVELS, (1e)		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
The exact eigenenergy spectrum of an electron is calculated in a quantum well within an in-plane magnetic field. The numerical solutions for the excited energy states as well as the ground state energy are found for various quantum-well widths and barrier heights. The cyclotron orbits are considerably affected by the quantum well. The energy levels higher than the potential height of the quantum well and the energy levels lower than the potential height of the quantum well show quite different behaviors. These are explained with the properties of the combined potential and the wave function inherent to the system.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field

H. R. Lee, H. G. Oh and Thomas F. George
Department of Physics and Astronomy
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

C. I. Um
Department of Physics
College of Science
Korea University
Seoul 136, KOREA



Abstract

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The exact eigenenergy spectrum of an electron is calculated in a quantum well within an in-plane magnetic field. The numerical solutions for the excited energy states as well as the ground state energy are found for various quantum-well widths and barrier heights. The cyclotron orbits are considerably affected by the quantum well. The energy levels higher than the potential height of the quantum well and the energy levels lower than the potential height of the quantum well show quite different behaviors. These are explained with the properties of the combined potential and the wave function inherent to the system.

PACS Nos.: 73.60.-n, 73.20.Dx, 71.50.+t

I. Introduction

The properties of electrons confined in semiconductor heterostructures have been studied for different situations. The carriers in two-dimensional systems subjected to a magnetic field perpendicular to the layers are completely quantized into Landau levels, which have been extensively studied with respect to the quantized Hall effect[1] and Shubnikov-de-Haas oscillations [2]. In this case, since the magnetic field is in the same direction as the confining electric field, the Hamiltonian can be separated into an electric part leading to subbands and a magnetic part leading to Landau levels. For any other orientation, this separation is not possible any more. Thus, in an external magnetic field parallel to the interface, the situation becomes more complicated. By studying the effects of magnetic field on the optical transitions, we may obtain the details of the subband structure[3,4]. Although an in-plane magnetic field usually has little effect on two-dimensional properties, it can strongly affect the spectrum of intersubband optical transitions[5]. If the magnetic length $a_H = (\hbar/\mu\omega)^{1/2}$, where μ and ω are the effective mass and the cyclotron frequency of an electron, respectively, is made comparable to the quantum well width by increasing the magnitude of the magnetic field, the problem is even more complicated because the confining electric field and the magnetic field contribute almost the same weight to the energy levels of electrons. There are several papers in theory[6-9] as well as in experiment[10-12] which study the effect of in-plane magnetic fields on two-dimensional systems. Recently Klama[8] has reported the quantization rules and analytic expressions of the electron energy spectrum in a thin film within a

longitudinal magnetic field. In his paper, the electronic motion is always inside the quantum well because the potential height is taken to be infinite. The eigenenergies of two-dimensional electrons subjected to a tilted magnetic field are solved analytically in a parabolic potential well[10], which is useful for the analysis of transport measurements in their experiments. Oliveira et al[9] have found the electron binding energy in a quantum well of n-type GaAs-Al_xGa_{1-x}As heterostructures by a self-consistent solution of the Poisson and Schrödinger equations. Most of their calculations were done for a couple of the lowest subbands in various systems.

Since the harmonic oscillator is one of the exactly solvable problems[13], the exact eigenvalues for a quantum well under an in-plane magnetic field can be obtained numerically. In this paper we present the numerical solutions for the energy spectrum of electrons in a quantum well with a finite potential height V_0 and well width W comparable to the magnetic length. The eigenenergy spectrum of an infinite potential well is also calculated for comparison. In the calculations, the normalized potential height $(V_0/\hbar\omega)$ and well width $(\sqrt{2}W/a_H)$ are used to see the results for an arbitrary potential height V_0 and well width W .

II. Theory

Let us consider a quantum well with well width W and the potential height V_0 in an external applied longitudinal DC magnetic field. The uniform external magnetic field $\vec{B} = (0, 0, B_0)$ oriented in the plane of the layer can be described by the vector potential $\vec{A} = (0, B_0x, 0)$. In the direction parallel to $\vec{B}$ the electrons have a free motion, whereas in the

plane perpendicular to $\vec{B}$ they are subjected to the combined potential of the magnetic field and the quantum well. Then, the Hamiltonian for an electron in a potential well with a magnetic field in the xz -plane can be written as

$$H = \frac{1}{2\mu} (p_x^2 + p_z^2) + \frac{1}{2\mu} \left[p_y + eB_0 x \right]^2 + V(x) \quad (1)$$

where μ is the effective mass and $V(x)$ is the potential energy of the electron in the well. Correspondingly, the Schrödinger equation is given by

$$H\psi(\vec{r}) = E\psi(\vec{r}) \quad (2)$$

Due to the translational symmetry in the y - and z -directions, the wave function can be written as

$$\psi(\vec{r}) = \chi(x) e^{i(q_y y + q_z z)} \quad (3)$$

Substituting this wave function into the Schrödinger equation (2), we obtain the effective one-dimensional Schrödinger equation

$$\frac{d^2 \chi(x)}{dx^2} + \frac{2\mu}{\hbar^2} \left[\epsilon - \frac{\mu \omega^2}{2} (x - x_0)^2 + V(x) \right] \chi(x) = 0 \quad (4)$$

where

$$\epsilon = E - \frac{p_z^2}{2\mu}, \quad (5)$$

and we have used $\omega = eB_0/\mu$ for the cyclotron frequency, $x_0 = a_H^2 q_y$ for the position of the orbit center and $a_H = (\hbar/\mu\omega)^{1/2}$ for the magnetic length.

Now the two boundaries caused by the quantum well can be determined depending on the location of the orbit center x_0 . For the general case, we assume the left and right boundaries of the well are located at $x_L = a$ and $x_R = b$ ($a < b$), respectively, i.e.,

$$V(x) = \begin{cases} 0 & \text{for } a < x < b \\ V_0 & \text{for } x < a, \quad b < x \end{cases} \quad (6)$$

Introducing the dimensionless length scale $\xi = \sqrt{2}x/a_H$ for the x -coordinate, the Schrödinger equation for the combined potential is divided into three regions:

$$\frac{d^2 x_I(\xi)}{d\xi^2} + \left[\frac{\epsilon}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_I(\xi) = 0 \quad \text{for } \frac{\sqrt{2}a}{a_H} < \xi < \frac{\sqrt{2}b}{a_H} \quad (7a)$$

$$\frac{d^2 x_{II}(\xi)}{d\xi^2} + \left[\frac{\epsilon - V_0}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_{II}(\xi) = 0 \quad \text{for } \xi < \frac{\sqrt{2}a}{a_H} \quad (7b)$$

$$\frac{d^2 x_{III}(\xi)}{d\xi^2} + \left[\frac{\epsilon - V_0}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_{III}(\xi) = 0 \quad \text{for } \frac{\sqrt{2}b}{a_H} < \xi \quad (7c)$$

Comparing the above Schrödinger equations (7a), (7b), and (7c) with the Weber equation [14],

$$\frac{d^2\psi(z)}{dz^2} + \left(m + \frac{1}{2} - \frac{1}{4}z^2\right)\psi(z) = 0, \quad (8)$$

we can identify the eigenvalues of the system:

$$\frac{\epsilon}{\hbar\omega} = m + \frac{1}{2} \quad \text{for} \quad \frac{\sqrt{2a}}{a_H} < \xi < \frac{\sqrt{2b}}{a_H} \quad (9)$$

and

$$\frac{\epsilon - V_0}{\hbar\omega} = m' + \frac{1}{2} \quad \text{for} \quad \xi < \frac{\sqrt{2a}}{a_H}, \text{ or } \frac{\sqrt{2b}}{a_H} < \xi. \quad (10)$$

The quantum numbers m and m' are related each other by,

$$(m - m')\hbar\omega = V_0. \quad (11)$$

The solution of equation (8) is the well-known Weber function[14]:

$$D_m(z) = 2^{m/2} e^{-z^2/4} \left[\frac{\sqrt{\pi}}{\Gamma(1/2 - m/2)} F\left(-\frac{1}{2}m \middle| \frac{1}{2} \middle| \frac{1}{2}z^2\right) - \frac{\sqrt{2\pi} z}{\Gamma(-m/2)} F\left(\frac{1}{2} - \frac{1}{2}m \middle| \frac{3}{2} \middle| \frac{1}{2}z^2\right) \right] \quad (12)$$

where $F(a|b|x)$ is the confluent hypergeometric function[15] and $\Gamma(z)$ is the gamma function.

From the asymptotic properties of the Weber function[14], the general solutions of the Schrödinger equations (7a), (7b) and (7c) may take the following forms:

$$x_I(\xi) = A D_m(\xi) + B D_m(-\xi) \quad (13a)$$

$$x_{II}(\xi) = C D_{m'}(-\xi) \quad (13b)$$

$$x_{III}(\xi) = F D_{m'}(\xi) \quad (13c)$$

where A, B, C and F are the normalization constants in each region. Then the continuity of the logarithmic derivatives of the wave functions at both boundaries $\xi_L = \sqrt{2}a/a_H$ and $\xi_R = \sqrt{2}b/a_H$ of the quantum well provides the following equation for the energy eigenvalues:

$$\begin{aligned} & D_m(\xi_R) D'_m(\xi_R) - D'_{m'}(\xi_R) D_m(\xi_R) \\ & - \frac{D'_m(\xi_L) D_{m'}(-\xi_L) + D'_{m'}(-\xi_L) D_m(\xi_L)}{D'_m(-\xi_L) D_{m'}(-\xi_L) + D'_{m'}(-\xi_L) D_m(-\xi_L)} \left[D'_{m'}(\xi_R) D_m(-\xi_R) + D_{m'}(\xi_R) D'_m(-\xi_R) \right] = 0 \end{aligned} \quad (14)$$

where

$$D'_m(\xi) = \frac{d}{d\xi} D_m(\xi) \quad (15)$$

III. Results and Discussion

Making use of the recursion relations of the Weber function[14], the numerical solutions for m can be obtained from Eq. (14) if the values of V_0 ,

ξ_L and ξ_R are given. The numerical solutions of m are illustrated in Figs. (1) and (2) for different values of normalized potential height $V_0/\hbar\omega$ and well width W/a_H . We use the potential height $V_0 = 5\hbar\omega$ and the well width $W = a_H/\sqrt{2}$ in Fig. 1 and $V_0 = 10\hbar\omega$ and $W = \sqrt{2}a_H$ were used in Fig. 2. In these figures, the various energy levels are plotted against the position ξ_0 of the orbit center. In each graph the dotted line shows the boundaries of the quantum well on a normalized scale, and the quantum number m (y-axis) is related to the eigenenergy by $\epsilon = (m + 1/2)\hbar\omega$. These results clearly show two different types of energy states: (1) the states confined in the quantum well (the lowest level in Fig. 1 and the lowest two levels in Fig. 2) and (2) the extended states (the higher levels in Figs. 1 and 2).

For the confined states (1), most of the wavefunction is pushed into the quantum well because the eigenenergy is lower than the potential height. As the orbit center is shifted from the center of the potential well, the electron penetrates and spends increasingly more time inside the potential barrier and less time in the well, resulting thus in an increase of energy. If the orbit center is moved deep into the barrier and far away from the quantum well, the low-level electrons see essentially a magnetic harmonic well elevated by V_0 . This means that the energy versus ξ_0 curve will flatten out at large values of ξ_0 . Hence the usual bulk Landau levels take over except for a shift of V_0 in energy. Before the bulk Landau levels take over, we can see the crossing of the confined state and the extended states in the intermediate shift of orbit center ξ_0 in both Figs. 1 and 2. These two states are degenerate at the moment they cross each other. If the orbit center is shifted into the barrier, the quantum well would be elevated as

well as changed in shape by the presence of the parabolic magnetic potential. This means that the overall enhancement of the quantum well due to the magnetic potential should be parabolic as the orbit center ξ_0 is moving away from the quantum well. Thus the eigenenergies of the confined states in the quantum well increase parabolically as a function of the orbit center ξ_0 . Clearly the crossings of the confined states and the extended states in Figs.1 and 2 exhibit parabolic behavior. The changes in shape of the quantum well neglecting the elevation have little effect on the crossings because the eigenenergies of the confined states are always limited by the barrier height V_0 , which is small compared to the elevation due to the magnetic potential.

In the second case (2), the energy spacing between neighboring modes in the energy spectrum is about $\hbar\omega$, but each mode makes oscillations as the orbit center moves away from the center of the quantum well. In this case, the electron energies are higher than the potential height V_0 and the wavefunctions are located partly within the well and partly inside the barrier, but still confined by the magnetic potential well. The oscillatory behavior of the eigenmodes seems to be related to the oscillating property of the harmonic wavefunction. The harmonic wavefunction $[D_m(z)]$ has many nodes and peaks depending on the quantum number m , and the wavefunction may be classified as symmetric (even parity) or antisymmetric (odd parity) for the centered orbit ($\xi_0=0$). The oscillations of eigenenergy in each mode depend on the number of nodes on the wave function. Thus the number of oscillations increases one by one as we go up to the higher levels. We also observe that the oscillations of the eigenenergy are within the range of the

wavepacket size ($\Delta x = \sqrt{m+1/2} a_H$). The alternating downward and upward concavity in energy spectrum at the middle of the quantum well seems due to the symmetric or antisymmetric properties of the wavefunction. If the wavefunction is symmetric, the middle peak of the wavefunction is located at the center of the potential well for the centered orbit ($\xi_0=0$). Therefore the eigenenergy will increase if the orbit center is shifted slightly toward barrier, resulting in the downward concavity. On the other hand, we expect upward concavity for the antisymmetric wavefunction because its node is now located at the lowest spot of the combined potential well. If these arguments are correct, the wavefunction of the lowest mode should be symmetric and have no node, the second mode should be antisymmetric and have one node, and so on from Figs. 1 and 2. In conclusion, we can say that the energy levels with oscillating structures are direct images of the magnetic wave function itself in the restricted geometry.

For the practical example of $\text{GaAs-Al}_x\text{GaAs}_{1-x}$ with $x = 0.3$, we get $V_0 = 0.25$ eV. To match this potential height in fig.1 where $V_0 = 5\hbar\omega$, we need $B = 29$ T, which corresponds to $a_H = 48$ Å. If $V_0 = 10\hbar\omega$ like in Fig. 2, a magnetic field of $B = 14$ T is required. To observe these results at $B = 10$ T which has a magnetic length $a_H = 81$ Å, the potential height of Figs. 1 and 2 correspond to $V_0 = 87$ and 174 meV, respectively. Even if these results were calculated in a relatively strong magnetic field due to the computer limit, the general features could be observed in a practical samples given above.

Finally, the eigenenergies for an infinite potential well ($V_0 \rightarrow \infty$) can be calculated similarly with the proper boundary conditions ($\chi_{II} = \chi_{III} = 0$,

$\chi_I(\xi - \xi_R) - \chi_I(\xi - \xi_L) = 0$) and the results are presented in Fig. 3 with a well width $W = \sqrt{2}a_H$. In this case, since all the electronic motions are within the quantum well, we cannot see such extended states as for a finite potential height. As explained earlier, the parabolic behavior is almost the same except for higher energies than before due to the infinite well.

Acknowledgments

This research was supported by the Office of Naval Research and the Office of Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes, notwithstanding any copyright notation hereon.

References

- [1] E. von Klitzing, G. Dorda, and M. Pepper, Phys. Rev. Lett. 45, 494 (1980), D. C. Tsui, H. L. Störmer, and A. C. Gossard, Phys. Rev. Lett. 48, 1559 (1982), and R. B. Laughlin, Phys. Rev. B, 23, 5632 (1981).
- [2] A. B. Fowler, F. F. Fang, W. E. Howard and P. J. Stiles, Phys. Rev. Lett. 16, 901(1966), and K. F. Komatsubara, K. Narita, Y. Katayama, N. Kotera, and M. Kobayashi, J. Phys. Chem. Solids 35, 723 (1974).
- [3] W. Beinvogl and J. F. Koch, Phys. Rev. Lett. 40, 1736 (1978).
- [4] F. Koch, in Physics in High Magnetic Fields, edited by S. Chikazumi and M. Miura (Springer, Berlin, 1981).
- [5] T. Ando, A. B. Fowler, and F. Stern, Rev. Mod. Phys. 54, 437 (1982).
- [6] T. Ando, J. Phys. Soc. Japan 39, 411 (1975).
- [7] S. K. Bhattacharya, Phys. Rev. B 25, 3756 (1982).
- [8] S. Klama, J. Phys. C: Solid State Phys. 20, 551 (1987).
- [9] Glaucia M. G. Oliveira, Vivili M. S. Gomes, A. S. Chaves, J. R. Leite and J. M. Worlock, Phys. Rev. B 35, 2896 (1987).
- [10] J. C. Maan, in Two-dimensional Systems. Heterostructures and Superlattices, Vol.53 in Solid State Sciences, edited by G. Bauer, F. Kuchar and H. Heinrich, (Springer-Verlag, Berlin, 1984).
- [11] J. H. Caseman and U. Merkt, Solid State Commun. 47, 917 (1983).
- [12] Z. Schlesinger, J. C. M. Hwang and S. J. Allen Jr., Phys. Rev. Lett. 50, 2098 (1983)
- [13] E. Schrödinger, Naturwirs 14, 664 (1926).
- [14] P. M. Morse and H. Feshbach, Methods of Theoretical Physics (McGraw-Hill, New York, 1953), p. 1403.
- [15] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, (Dover, New York, 1964), p. 503.

Figure Captions

1. Numerical solution for electron eigenenergy as a function of the orbit center ξ_0 with well width $W = a_H/\sqrt{2}$ and potential height $V_0 = 5\hbar\omega$.
2. Numerical solution for electron eigenenergy as a function of the orbit center ξ_0 with well width $W = \sqrt{2}/a_H$ and potential height $V_0 = 10\hbar\omega$.
3. Numerical solution for electron eigenenergy as a function of the orbit center ξ_0 with well width $W = \sqrt{2}/a_H$ and potential height $V_0 = \infty$.

Fig. 1




Fig. 2





Fig. 3



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Renssaele Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853