

DTIC FILE COPY

4

AD-A214 208

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 114

Optical Phonon Modes in a Double Heterostructure of Polar Crystals

by

R. Chen, D. L. Lin and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

October 1989

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
NOV 08 1989
S B D

89 11 08 03f

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-114			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Optical Phonon Modes in a Double Heterostructure of Polar Crystals					
12. PERSONAL AUTHOR(S) R. Chen, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) October 1989	
				15. PAGE COUNT 27	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	POLAR CRYSTALS CONTINUUM MODEL		
			DOUBLE HETEROSTRUCTURE POLARIZATION VECTOR		
			OPTICAL PHONONS EQUATION OF MOTION		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The equation of motion for the polarization vector for a double heterostructure of polar crystals is solved exactly within the framework of the continuum model. There exist only two types of phonon modes, the interface modes and the confined bulk modes, whose eigenvectors are obtained explicitly. Dispersion relations are derived analytically for the interface modes, while the confined bulk modes are dispersionless, a fact consistent with the model. It is also found that in Raman scattering experiments, the symmetric interface modes are predominantly longitudinal (LO) and the antisymmetric interface modes transversal (TO). In the central region of the Brillouin zone, however, they both split into two branches oscillating at LO and TO frequencies, respectively. Possible reinterpretation of various experiments are briefly discussed.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Optical phonon modes in a double heterostructure of polar crystals

R. Chen, D. L. Lin and Thomas F. George
Department of Physics and Astronomy
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

The equation of motion for the polarization vector for a double heterostructure of polar crystals is solved exactly within the framework of the continuum model. There exist only two types of phonon modes, the interface modes and the confined bulk modes, whose eigenvectors are obtained explicitly. Dispersion relations are derived analytically for the interface modes, while the confined bulk modes are dispersionless, a fact consistent with the model. It is also found that in Raman scattering experiments, the symmetric interface modes are predominantly longitudinal (LO) and the antisymmetric interface modes transversal (TO). In the central region of the Brillouin zone, however, they both split into two branches oscillating at LO and TO frequencies, respectively. Possible reinterpretation of various experiments are briefly discussed.

1988 PACS Nos.: 63.20.Dj, 68.35.Ja, 78.65.Fa, 78.65.Gb

I. Introduction

There has been great interest in recent years in the study of various vibrational modes supported by semiconductor heterostructures. The patterns of normal modes of vibrations that determine the electronic properties in such structures are evidently different from those in the bulk. The presence of interfaces necessarily alters the phonon modes, and possibly even their interaction with electrons may be modified because of the reduced dimensionality.

Experimental investigations of the spectra of possible acoustic modes in superlattices¹⁻⁴ have shown fairly-good agreement with theory. On the other hand, the longitudinal optical (LO) modes in polar crystals are much less well understood in these contexts. In all the calculations such as free-carrier absorption of light,⁵⁻⁷ scattering rates⁸ and polaronic effects,⁹ etc., the usual bulk Frölich Hamiltonian is assumed for the electron-phonon interaction in confined systems. The only requirements are that the material elastic properties match at the interfaces and that the dielectric properties are equal. More recently, the surface optical (SO) mode has been included in the treatment of polarons confined in a slab¹⁰ or near the interface in semi-infinite systems.^{11,12} For these surface and interface situations, the bulk Frölich Hamiltonian is still employed for treating the LO-phonon contribution.

On the other hand, evidences of confined modes peculiar to different types of layered structures have been noticed in various experiments. Measurements of magneto-absorption and cyclotron resonance in GaInAs/InP superlattices¹³ and GaAs/GaAlAs heterostructures¹⁴ indicate that the electron-LO-phonon interaction in these structures can be fundamentally different from that in the bulk case. In a numerical study of possible

modes of the optical phonon in layered polar crystals, it is found that phonon modes tend to be confined in each layer and that the penetration of vibrations into the adjacent layer is negligible.^{15,16} Moreover, the existence of confined phonon modes has been directly observed in a GaAs single quantum well of GaAs/GaAlAs heterostructures.¹⁷

In theoretical investigations of the vibrational modes in an ionic slab, Fuchs and Kliever¹⁸ have found the bulk LO mode with the nodes at the surfaces as well as the SO modes of different symmetries. Interface modes have been derived by Wendler¹⁹ by considering the polarization field in a double layer system, and by Lassnig²⁰ using the energy-loss method in a double heterostructure (DHS) of polar semiconductors. An alternative treatment²¹ of the DHS predicts some peculiar phonon modes that have not been borne out by observation.

In this article we present solutions for optical phonon modes in a semiconductor DHS using the continuum model of Born and Huang.²² The method of solution has been developed by various authors.^{18,19,23} Apart from the long-wavelength limit in the model, no further approximation is made throughout our calculation. Dispersion relations and eigenvectors for all the normal modes of lattice vibration are derived analytically. It is found that there exist two types of phonon modes, the interface modes and confined bulk modes.

While the existence of interface modes has been well recognized experimentally,^{24,25} their eigenvectors and dispersion relations in a DHS are solved explicitly for the first time in this paper. Our results show that either the symmetric or the antisymmetric interface modes has two branches. Their frequencies at the center of the Brillouin zone are exactly the same as those of the bulk LO and TO phonon in each material.

For	☒
1	☐
2	☐
3	☐
4	☐
5	☐
6	☐
7	☐
8	☐
9	☐
10	☐
11	☐
12	☐
13	☐
14	☐
15	☐
16	☐
17	☐
18	☐
19	☐
20	☐
21	☐
22	☐
23	☐
24	☐
25	☐
26	☐
27	☐
28	☐
29	☐
30	☐
31	☐
32	☐
33	☐
34	☐
35	☐
36	☐
37	☐
38	☐
39	☐
40	☐
41	☐
42	☐
43	☐
44	☐
45	☐
46	☐
47	☐
48	☐
49	☐
50	☐
51	☐
52	☐
53	☐
54	☐
55	☐
56	☐
57	☐
58	☐
59	☐
60	☐
61	☐
62	☐
63	☐
64	☐
65	☐
66	☐
67	☐
68	☐
69	☐
70	☐
71	☐
72	☐
73	☐
74	☐
75	☐
76	☐
77	☐
78	☐
79	☐
80	☐
81	☐
82	☐
83	☐
84	☐
85	☐
86	☐
87	☐
88	☐
89	☐
90	☐
91	☐
92	☐
93	☐
94	☐
95	☐
96	☐
97	☐
98	☐
99	☐
100	☐

Dist

A-1

Availability Codes

A-11 and/or
Special

Experimental evidence of confined bulk modes has recently been reported.^{17 25,26} We find that both the bulk LO and transverse optical (TO) modes are strictly confined. Further investigation on implications of such confinements is being carried out and will be reported elsewhere.

In Sec. II, we outline the procedure for deriving the equation of motion for the polarization vector. The coupled integral equations are solved for the interface modes in Sec. III and the confined bulk modes in Sec. IV. Consequences and implications of our results are discussed in Sec. V.

II. Equation of motion of the polarization field

Consider a DHS of two different polar crystals as shown in Fig. 1. A layer of material 1 with thickness a is sandwiched between two thick layers of material 2. We take the z -axis to be perpendicular to the interfaces which are located at $z = 0$ and $z = a$, respectively. Following Born and Huang,²² we start in the continuum approximation with the equation of motion for the relative displacement $u(\vec{r}, t)$ of the ion pair in material ν ($\nu = 1, 2$),

$$\mu_{\nu} \ddot{\vec{u}}_{\nu}(\vec{r}, t) = -\mu_{\nu} \omega_{0\nu}^2 \vec{u}_{\nu}(\vec{r}, t) + e^* \vec{E}(\vec{r}, t) \quad , \quad (1)$$

where μ is the reduced mass of the pair of ions, $\mu \omega_0^2$ is the short-range force constant not including Coulomb fields, $\vec{E}(\vec{r}, t)$ is the local electric field, and e^* is the effective charge of the ions. The subscript ν labels the material considered. The oscillating ions produce a polarization field $\vec{P}(\vec{r}, t)$ given by

$$\vec{P}(\vec{r}, t) = n_{\nu} e^* \vec{u}(\vec{r}, t) + n_{\nu} \alpha_{\nu} \vec{E}(\vec{r}, t) \quad , \quad (2)$$

where n is the number of ion pairs per unit cell and α is the polarizability. The first term in (2) represents the contribution of the ion pair when the lattice vibrates, and the second term is the electronic polarization of the ions due to the electric field associated with the optical modes. The part of the polarization produced by the electron itself as it moves through the crystal is, however, not included in our consideration, since the continuum model is not valid for such an effect.²³ The local field in (2) is related, in the long-wavelength limit, to the polarization by

$$\vec{E}(\vec{r}, t) = \vec{E}_l(\vec{r}, t) + 4\pi \int d\vec{r}' \Gamma(\vec{r} - \vec{r}') \cdot \vec{P}(\vec{r}') \quad (3a)$$

where

$$\vec{E}_l(\vec{r}, t) = \frac{4\pi}{3} \vec{P}(\vec{r}, t) \quad , \quad (3b)$$

and Γ denotes the Green tensor with components

$$\Gamma_{\alpha\beta} = \frac{1}{4\pi} \frac{\partial^2}{\partial x_{\alpha} \partial x_{\beta}} \frac{1}{|\vec{r} - \vec{r}'|} \quad . \quad (3c)$$

The equation of motion for the polarization then follows by plugging Eqs. (2) and (3) into (1):

$$\begin{aligned}
& \left[1 - \frac{4\pi}{3} \alpha_{\nu} n_{\nu} \right] \ddot{\vec{P}}(\vec{r}, t) + \left[\omega_{o\nu}^2 - 4\pi(\alpha_{\nu} n_{\nu} \omega_{o\nu}^2 + \frac{n_{\nu} e_{\nu}^{*2}}{3\mu_{\nu}}) \right] \vec{P}(\vec{r}, t) \\
& = 4\pi \alpha_{\nu} n_{\nu} \int d\vec{r}' \Gamma(\vec{r} - \vec{r}') \cdot \ddot{\vec{P}}(\vec{r}', t) \\
& + 4\pi(\alpha_{\nu} n_{\nu} \omega_{o\nu}^2 + \frac{n_{\nu} e_{\nu}^{*2}}{3\mu_{\nu}}) \int d\vec{r}' \Gamma(\vec{r} - \vec{r}') \cdot \vec{P}(\vec{r}', t) \quad (4)
\end{aligned}$$

The time-dependent part of the polarization can be separated by assuming $\vec{P}(\vec{r}, t) = \vec{P}(\vec{r}) e^{i\omega t}$ which, after substituting into (4), yields the equation for $\vec{P}(\vec{r})$,

$$\left[\frac{\lambda_{\nu} - \lambda_{o\nu}}{\alpha_{\nu} n_{\nu} (\lambda_{\nu} - \lambda_{o\nu}) - 1} - \frac{4\pi}{3} \right] \vec{P}(\vec{r}) = 4\pi \int d\vec{r}' \Gamma(\vec{r} - \vec{r}') \cdot \vec{P}(\vec{r}') \quad (5a)$$

where we have defined the parameters

$$\lambda_{\nu}^2 = 4\pi\omega^2 / \omega_{p\nu}^2 \quad (5b)$$

$$\lambda_{o\nu}^2 = 4\pi\omega_{o\nu}^2 / \omega_{p\nu}^2 \quad (5c)$$

with the ion plasma frequency $\omega_{p\nu}^2 = 4\pi n_{\nu} e_{\nu}^{*2} / \mu_{\nu}$.

Since the translational invariance in the z-direction is destroyed by the interfaces, we introduce the two-dimensional vectors $\vec{\kappa}$ and $\vec{\rho}$ so that $\vec{k} = (\vec{\kappa}, q)$ and $\vec{r} = (\vec{\rho}, z)$. The two-dimensional Fourier transforms can now be written as

$$\vec{P}(\vec{r}) = \left(\frac{1}{2\pi}\right)^2 \int_{-\infty}^{\infty} d\vec{\kappa} e^{i\vec{\kappa} \cdot \vec{\rho}} \vec{P}(\vec{\kappa}, z) \quad (6)$$

$$\frac{1}{r} = \frac{1}{2\pi} \int \frac{d\vec{\kappa}}{\kappa} \exp(i\vec{\kappa} \cdot \vec{\rho} - \kappa|z|) \quad (7)$$

Differentiating (7) twice with respect to the coordinates, we obtain

$$\Gamma = -\frac{1}{4\pi} \int d\vec{\kappa} e^{i\vec{\kappa} \cdot \vec{\rho} - \kappa|z|} \vec{K}\vec{K}/2\pi\kappa \quad (8)$$

where

$$\vec{K} = (\vec{\kappa}, i\theta(z)\kappa) \quad (9)$$

with the step function $\theta(z) = \pm 1$ for $z \gtrless 0$. Substituting (6) and (8) into (5a), and moving the term $-P_z$ from the right-hand side to the left, we can write after some algebra the resulting equation in a more symmetric form as

$$4\pi \begin{bmatrix} \chi_{\nu}^{-1}(\omega) & 0 & 0 \\ 0 & \chi_{\nu}^{-1}(\omega) & 0 \\ 0 & 0 & \chi_{\nu}^{-1}(\omega)\epsilon_{\nu}(\omega) \end{bmatrix} \cdot \vec{P}(\vec{\kappa}, z) = \frac{2\pi}{\kappa} \int_{-\infty}^{\infty} dz' e^{-\kappa(z-z')} \vec{K}\vec{K} \cdot \vec{P}(\vec{\kappa}, z') \quad (10)$$

where $\chi_{\nu}(\omega)$ is defined by

$$4\pi\chi_{\nu}^{-1}(\omega) = \frac{\lambda_{\nu} - \lambda_{0\nu}}{\alpha_{\nu} n_{\nu} (\lambda_{\nu} - \lambda_{0\nu}) - 1} - \frac{4\pi}{3} \quad (11)$$

It turns out that $\chi_\nu(\omega)$ is the isotropic dielectric susceptibility and is related to the dielectric function by $\chi_\nu(\omega) = \epsilon_\nu(\omega) - 1$ with

$$\epsilon_\nu(\omega) = \epsilon_{\infty\nu} \frac{\omega_{L\nu}^2 - \omega^2}{\omega_{T\nu}^2 - \omega^2} \quad (12a)$$

$$\epsilon_{\infty\nu} = 1 + 4\pi\alpha_\nu n_\nu / (1 - \frac{4\pi}{3} \alpha_\nu n_\nu) \quad (12b)$$

where we have defined the LO and TO phonon frequencies

$$\omega_{L\nu}^2 = \omega_{\text{ov}}^2 + \frac{2}{3} \omega_{\text{p}\nu}^2 / (1 + \frac{8\pi}{3} \alpha_\nu n_\nu) \quad (13a)$$

$$\omega_{T\nu}^2 = \omega_{\text{ov}}^2 - \frac{1}{3} \omega_{\text{p}\nu}^2 / (1 - \frac{4\pi}{3} \alpha_\nu n_\nu) \quad (13b)$$

Since the interface phonons propagate in the xy-plane, it is more convenient to express the polarization vector as $\vec{P} = (\vec{\pi}, P_s)$ where $\vec{\pi}$ is a two-dimensional vector defined by $\vec{\pi} = (P_x, P_z)$. Thus

$$\vec{P}(\vec{\kappa}, z) = P_\kappa(\vec{\kappa}, z) \hat{\kappa} + P_z(\vec{\kappa}, z) \hat{z} + P_s(\vec{\kappa}, z) \hat{s} \quad (14)$$

where the unit vector $\hat{s}$ is defined by $\hat{s} = \hat{z} \times \hat{\kappa}$. Substituting (14) in (10), we can separate the s-component and decouple (10) into two equations:

$$\begin{bmatrix} \chi_\nu^{-1}(\omega) & 0 \\ 0 & \chi_\nu^{-1}(\omega) \epsilon_\nu(\omega) \end{bmatrix} \cdot \vec{\pi}(\vec{\kappa}, z) = \frac{1}{4\pi} \int_{-\infty}^{\infty} dz' M(z-z') \cdot \vec{\pi}(\vec{\kappa}, z') \quad (15a)$$

for the s -called p-polarization, and

$$\chi_{\nu}^{-1}(\omega) P_s(\vec{\kappa}, z) = 0 \quad (15b)$$

for the s-polarization, where M is a Hermitian matrix given by

$$M(z-z') = M^{\dagger}(z'-z) = -2\pi\kappa e^{-\kappa|z-z'|} \begin{bmatrix} 1 & i\theta(z-z') \\ i\theta(z-z') & -1 \end{bmatrix} \quad (16)$$

As we shall show in the following sections, Eq. (15a) defines an eigenvalue problem whose solutions describe the interface modes and the confined modes. Equation (15b) describes the s-polarization, which is not of concern in the present paper. The eigenvectors $\vec{\pi}(\kappa, z)$ form a complete orthonormal set. Here we just give without proof the orthonormality relation

$$\int_{-\infty}^{\infty} dz \frac{\eta_{\nu}^4(\omega_i) \eta_{\nu}^4(\omega_j)}{\omega_{p\nu}^2} \vec{\pi}_j^*(\vec{\kappa}, z) \vec{\pi}_i(\vec{\kappa}, z) = \delta_{ij} \quad (17)$$

with $\eta_{\nu}^4(\omega_i) = 1/[1 + \alpha_{\nu} n_{\nu}(\lambda_{o\nu} - \lambda_{\nu})]$. The completeness relation is given by

$$\sum_i \vec{\pi}_i^*(\kappa, z) \vec{\pi}_i(\kappa, z') = \frac{\omega_{p\nu}^2}{\eta_{\nu}^2} I \delta(z-z') \quad (18)$$

for the so-called p-polarization, and

$$\chi_{\nu}^{-1}(\omega) P_s(\vec{\kappa}, z) = 0 \quad (15b)$$

for the s-polarization, where M is a Hermitian matrix given by

$$M(z-z') = M^{\dagger}(z'-z) = -2\pi\kappa e^{-\kappa|z-z'|} \begin{bmatrix} 1 & i\theta(z-z') \\ i\theta(z-z') & -1 \end{bmatrix} \quad (16)$$

As we shall show in the following sections, Eq. (15a) defines an eigenvalue problem whose solutions describe the interface modes and the confined modes. Equation (15b) describes the s-polarization, which is not of concern in the present paper. The eigenvectors $\vec{\pi}(\kappa, z)$ form a complete orthonormal set. Here we just give without proof the orthonormality relation

$$\int_{-\infty}^{\infty} dz \frac{\eta_{\nu}^{\frac{1}{2}}(\omega_i) \eta_{\nu}^{\frac{1}{2}}(\omega_j)}{\omega_{p\nu}^2} \vec{\pi}_j^*(\vec{\kappa}, z) \vec{\pi}_i(\vec{\kappa}, z) = \delta_{ij} \quad (17)$$

with $\eta_{\nu}^{\frac{1}{2}}(\omega_i) = 1/[1 + \alpha_{\nu} n_{\nu}(\lambda_{o\nu} - \lambda_{\nu})]$. The completeness relation is given by

$$\sum_i \vec{\pi}_i^*(\kappa, z) \vec{\pi}_i(\kappa, z') = \frac{\omega_{p\nu}^2}{\eta_{\nu}} I \delta(z-z') \quad (18)$$

where I stands for the unit matrix. Finally, we note that the polarization vector must be real, and consequently

$$P(\vec{\kappa}, z) = P^*(-\vec{\kappa}, z) \quad (19)$$

Similarly, we have

$$\vec{\pi}(\vec{\kappa}, z) = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \vec{\pi}^*(-\vec{\kappa}, z) \quad (19b)$$

III. Interface Modes

It is easier to solve the coupled integral equations (15a) by first transforming them into differential equations. This can be done by differentiating (15a) with respect to z twice and at the same time requiring

$$\det \begin{bmatrix} \chi_{\nu}^{-1}(\omega) & 0 \\ 0 & \chi_{\nu}^{-1}(\omega) \epsilon_{\nu}(\omega) \end{bmatrix} \neq 0 \quad (20)$$

The resulting equations are

$$\frac{d}{dz} P_{\kappa}(\vec{\kappa}, z) = i\kappa P_z(\vec{\kappa}, z) \quad (21a)$$

$$\frac{d^2}{dz^2} \vec{\pi}(\vec{\kappa}, z) = \kappa^2 \vec{\pi}(\vec{\kappa}, z) \quad (21b)$$

The solutions to (21) take the form

$$P_{\kappa}(\vec{\kappa}, z) = \begin{cases} iA_2 e^{\kappa z} & , \quad z < 0 \\ i(A_1 e^{\kappa z} - B_1 e^{-\kappa z}) & , \quad 0 \leq z \leq a \\ -iB_2 e^{-\kappa z} & , \quad z > a \end{cases} \quad (22a)$$

$$P_z(\vec{\kappa}, z) = \begin{cases} A_2 e^{\kappa z} & , \quad z < 0 \\ A_1 e^{\kappa z} + B_1 e^{-\kappa z} & , \quad 0 \leq z \leq a \\ B_2 e^{-\kappa z} & , \quad z > a \end{cases} \quad (22b)$$

Substituting (22) in the integral equation (15a), one obtains a set of homogeneous equations for the amplitudes A_{ν} and B_{ν} of the p-polarization. The condition for the existence of a nontrivial solution then leads to the dispersion relation

$$\frac{\epsilon_1(\omega) - \epsilon_2(\omega)}{\epsilon_1(\omega) + \epsilon_2(\omega)} = \pm e^{\kappa a} \quad (23)$$

The + and - signs on the right-hand side of (23) correspond to the symmetric and antisymmetric modes of the interface phonons, respectively. The polarization amplitudes are found to satisfy the following relations:

$$B_1^{\pm}/A_1^{\pm} = B_2^{\pm}/A_2^{\pm} = \mp e^{\kappa a} \quad (24a)$$

$$A_2^{\pm} = \frac{\chi_2(\omega)}{\chi_1(\omega)} (1 \pm e^{\kappa a}) A_1^{\pm} \quad (24b)$$

$$B_2^\pm = \frac{\chi_2(\omega)}{\chi_1(\omega)} (1 \pm e^{\kappa a}) B_1^\pm \quad (24c)$$

Equations (24) are equivalent to the boundary conditions that the wave functions have to satisfy at the interfaces. Thus the differential equations (21) yield naturally the correct boundary conditions after substituting their solutions (22) into the integral equation (15a). On the contrary, the hydrodynamic terms introduced in Ref. 23 are inconsistent with the boundary conditions. Combining Eqs. (22)-(24), we find the eigenvectors for the antisymmetric interface phonon modes to be

$$\vec{\pi}_a = \begin{cases} C_a \left[\frac{\epsilon_2(\omega)-1}{\epsilon_1(\omega)-1} \right] e^{\kappa z} (-i, -1) \sinh\left(\frac{\kappa a}{2}\right) & , \quad z < 0 \\ C_a [i \sinh(\kappa(z-\frac{a}{2})), \cosh(\kappa(z-a/2))] & , \quad 0 \leq z \leq a \\ C_a \left[\frac{\epsilon_2(\omega)-1}{\epsilon_1(\omega)-1} \right] e^{-\kappa(z-a)} (i, -1) \sinh\left(\frac{\kappa a}{2}\right) & , \quad z > a \end{cases} \quad (25a)$$

and for the symmetric modes to be

$$\vec{\pi}_s = \begin{cases} C_s \left[\frac{\epsilon_2(\omega)-1}{\epsilon_1(\omega)-1} \right] e^{\kappa z} (i, 1) \cosh\left(\frac{\kappa a}{2}\right) & , \quad z < 0 \\ C_s [i \cosh(\kappa(z-a/2)), \sinh(\kappa(z-a/2))] & , \quad 0 \leq z \leq a \\ C_s \left[\frac{\epsilon_2(\omega)-1}{\epsilon_1(\omega)-1} \right] e^{-\kappa(z-a)} (i, -1) \cosh\left(\frac{\kappa a}{2}\right) & , \quad z > a \end{cases} \quad (25b)$$

where, according to (17), the normalization constants are given by

$$C_{a,s} = \sqrt{\frac{\kappa}{\sinh(\kappa a)} \left[\frac{\eta_1}{\omega_{p1}^2} - \frac{\eta_2}{\omega_{p2}^2} \frac{\epsilon_1}{\epsilon_2} \left(\frac{\chi_2}{\chi_1} \right)^2 \right]} \quad (25c)$$

A heterostructure composed of two media with dielectric functions given by (12) always supports four distinct interface modes of vibrations, two from each medium. The dispersion relations for these modes can then be calculated explicitly from (12) and (23). The results are

$$\begin{aligned} \omega_a^{\pm} = & \left\{ \epsilon_{\infty 2} (\omega_{T1}^2 + \omega_{L2}^2) + \epsilon_{\infty 1} (\omega_{T2}^2 + \omega_{L1}^2) \coth\left(\frac{\kappa a}{2}\right) \right. \\ & \pm \left(\epsilon_{\infty 2}^2 (\omega_{T1}^2 - \omega_{L2}^2)^2 + \epsilon_{\infty 1}^2 (\omega_{T2}^2 - \omega_{L1}^2)^2 \coth^2\left(\frac{\kappa a}{2}\right) \right. \\ & + 2\epsilon_{\infty 1} \epsilon_{\infty 2} [(\omega_{T1}^2 + \omega_{L2}^2)(\omega_{T2}^2 + \omega_{L1}^2) \\ & \left. \left. - 2(\omega_{T2}^2 \omega_{L1}^2 + \omega_{L2}^2 \omega_{T1}^2)] \coth\left(\frac{\kappa a}{2}\right) \right]^{1/2} \right\}^{1/2} / (2[\epsilon_{\infty 2} + \epsilon_{\infty 1} \coth\left(\frac{\kappa a}{2}\right)])^{1/2} \quad (26a) \end{aligned}$$

$$\begin{aligned} \omega_s^{\pm} = & \left\{ \epsilon_{\infty 2} (\omega_{T1}^2 + \omega_{L2}^2) + \epsilon_{\infty 1} (\omega_{T2}^2 + \omega_{L1}^2) \tanh\left(\frac{\kappa a}{2}\right) \right. \\ & \pm \left(\epsilon_{\infty 2}^2 (\omega_{T1}^2 - \omega_{L2}^2)^2 + \epsilon_{\infty 1}^2 (\omega_{T2}^2 - \omega_{L1}^2)^2 \tanh^2\left(\frac{\kappa a}{2}\right) \right. \\ & + 2\epsilon_{\infty 1} \epsilon_{\infty 2} [(\omega_{T1}^2 + \omega_{L2}^2)(\omega_{T2}^2 + \omega_{L1}^2) \\ & \left. \left. - 2(\omega_{T2}^2 \omega_{L1}^2 + \omega_{L2}^2 \omega_{T1}^2)] \tanh\left(\frac{\kappa a}{2}\right) \right]^{1/2} \right\}^{1/2} / (2[\epsilon_{\infty 2} + \epsilon_{\infty 1} \tanh\left(\frac{\kappa a}{2}\right)])^{1/2} \quad (26b) \end{aligned}$$

It is seen from (26) that the interface phonon energies depend explicitly on the dimensionless quantity κa .

Let us now look at the limiting cases. When $a \rightarrow \infty$, $\tanh\left(\frac{\kappa a}{2}\right) = 1$ and $\coth\left(\frac{\kappa a}{2}\right) = 1$. Therefore, both (26a) and (26b) approach the same limit, given by

$$\begin{aligned}
\omega_{\infty}^{\pm} = & \left\{ \epsilon_{\infty 1} (\omega_{L1}^2 + \omega_{T2}^2) + \epsilon_{\infty 2} (\omega_{T1}^2 + \omega_{L2}^2) \right. \\
& \pm \left[\epsilon_{\infty 1}^2 (\omega_{L1}^2 - \omega_{T2}^2)^2 + \epsilon_{\infty 2}^2 (\omega_{L2}^2 - \omega_{T1}^2)^2 \right. \\
& \left. \left. + 2 \epsilon_{\infty 1} \epsilon_{\infty 2} [(\omega_{T1}^2 + \omega_{L2}^2)(\omega_{T2}^2 + \omega_{L1}^2) - 2(\omega_{T2}^2 \omega_{L1}^2 + \omega_{T1}^2 \omega_{L2}^2)] \right] \right\}^{\frac{1}{2}} \\
& \times [2(\epsilon_{\infty 1} + \epsilon_{\infty 2})]^{-\frac{1}{2}}, \quad (27)
\end{aligned}$$

which is identical to the result of a bilayer system with only one interface,¹⁹ as it should. In the limit $a \rightarrow 0$, the system reduces to a bulk material 2 with frequencies ω_{L2} and ω_{T2} . When $\kappa \rightarrow 0$, $\tanh(\frac{\kappa a}{2}) = 0$ and $\coth(\frac{\kappa a}{2}) \rightarrow \infty$. We then find from (26) that

$$\omega_a^{\pm} = \sqrt{[\omega_{T2}^2 + \omega_{L1}^2 \pm (\omega_{T2}^2 - \omega_{L1}^2)]/2} = \omega_{T2}, \omega_{L1} \quad (28a)$$

$$\omega_s^{\pm} = \sqrt{[\omega_{T1}^2 + \omega_{L2}^2 \pm (\omega_{T1}^2 - \omega_{L2}^2)]/2} = \omega_{T1}, \omega_{L2} \quad (28b)$$

that is, the limiting frequencies are given by the bulk LO and TO frequencies of the two materials. It may be worth mentioning yet another limit at this point. When the characteristic parameters of the two dielectrics approach each other, or when $\epsilon_1 \rightarrow \epsilon_2$, we find for a given width that the amplitudes of the interface modes of vibration diminish continuously and become zero at $\epsilon_1 = \epsilon_2$.

The interface phonon modes have no connection with bulk polarization charges because $\vec{\nabla} \cdot \vec{P} = 0$. They are accompanied by the surface charges σ at the interfaces. These charge densities can easily be determined by

calculating the difference of the polarization eigenvectors in the z -direction on both sides of the interface concerned. Thus we find from the z -components of $\vec{\pi}_{s,a}$ in (25)

$$\sigma_s = \begin{cases} |\sigma_s| & z = 0 \\ |\sigma_s| & z = a \end{cases} \quad \text{at} \quad (29a)$$

for the symmetric mode and

$$\sigma_a = \begin{cases} -|\sigma_a| & z = 0 \\ |\sigma_a| & z = a \end{cases} \quad \text{at} \quad (29b)$$

for the antisymmetric mode, where

$$|\sigma_s| = \left(\frac{\kappa}{\sinh(\kappa a)} \right)^{1/2} \frac{x_1 \sinh(\frac{\kappa a}{2}) + x_2 \cosh(\frac{\kappa a}{2})}{\left[\frac{x_1^2 \eta_1}{\omega_{p1}^2} + \frac{x_2^2 \eta_2}{\omega_{p1}^2} \coth(\frac{\kappa a}{2}) \right]^{1/2}} \quad (30a)$$

$$|\sigma_a| = \left(\frac{\kappa}{\sinh(\kappa a)} \right)^{1/2} \frac{x_1 \cosh(\frac{\kappa a}{2}) + x_2 \sinh(\frac{\kappa a}{2})}{\left[\frac{x_1^2 \eta_1}{\omega_{p1}^2} + \frac{x_2^2 \eta_2}{\omega_{p2}^2} \tanh(\frac{\kappa a}{2}) \right]^{1/2}} \quad (30b)$$

IV. Confined bulk LO and TO modes

For the LO modes, $\omega = \omega_{L\nu}$, and from (12a) we have $\epsilon_\nu(\omega_{L\nu}) = 0$ and $\epsilon_{\nu'}(\omega_{L\nu}) \neq 0$ for $\nu' \neq \nu$. Thus $\vec{D} = 0$ in layer ν , and $\vec{E} = -\vec{P}/\epsilon_0$. The differential equations satisfied by the polarization field associated with the longitudinal phonon modes follow from (15a). In layer ν we have

$$\frac{d}{dz} P_{\kappa}^L(\kappa, z) = i\kappa P_z^L(\kappa, z) \quad , \quad (31a)$$

and in layer $\nu' \neq \nu$

$$\frac{d^2}{dz^2} P_{\kappa}^L(\kappa, z) = \kappa^2 P_z^L(\kappa, z) \quad . \quad (31b)$$

The solution to (31b) has the same form as (22) with the coefficients determined by the boundary conditions for the field vectors. In layer ν , $\vec{D} = 0$, and $\vec{E} = -\vec{P}/\epsilon_0$. From the continuity of $\vec{D}$ across the interface, we have $\vec{D} = 0$ in layer ν' . Since $\epsilon_{\nu'} \neq 0$, we must have $\vec{E} = 0$ and $\vec{P} = 0$. Therefore (31b) has a solution that is identically zero everywhere in the layer ν' . The boundary conditions in layer ν can be found from (15a) for $\epsilon_{\nu} = 0$ and $\chi_n = -1$. After some algebra, we find from the coupled equations that

$$\vec{P}^L(a) = \vec{P}^L(0) = 0 \quad . \quad (32)$$

Hence, Eq. (31a) is satisfied in layer 1 by the eigenvectors

$$\vec{\pi}_m^L = \begin{cases} 0 & z < 0 \\ C_m^L [i \sin(\frac{m\pi}{a}z), \frac{m\pi}{a\kappa} \cos(\frac{m\pi}{a}z)] & 0 \leq z \leq a \\ 0 & z > a \end{cases} \quad , \quad (33a)$$

where m is an integer, and in layer 2 by

$$\vec{\pi}_q^L = \begin{cases} C_q^L [i \sin(qz), \frac{q}{\kappa} \cos(qz)] & , \quad z < 0 \\ 0 & 0 \leq z \leq a \\ C_q^L [i \sin q(z-a), \frac{q}{\kappa} \cos q(z-a)] & , \quad z > a \end{cases} \quad (33b)$$

where q is real, with the corresponding eigenfrequencies ω_{L1} and ω_{L2} , respectively. These modes are highly degenerate vibrations. We remark that the dispersionless nature of the confined bulk modes is consistent with the long-wavelength limit which is implied in the continuum model. The bulk optical photons have constant energy as $k \sim 0$.

Equations (33) clearly show that the longitudinal optical phonons are completely confined by the interfaces. In the central layer (labeled 1), the confinement leads to the quantization $q = m\pi/a$, where $m = 0, \pm 1, \pm 2, \dots$, while in the semi-infinite side layers (labeled 2), the wave number q remains continuous. The state vectors are normalized according to (17) with the normalization constants given by

$$C_m^L = \frac{\omega_{L1}}{\sqrt{\eta_1}} \sqrt{\frac{2}{a}} \frac{\kappa}{\sqrt{\kappa^2 + m^2 \pi^2 / a^2}} \quad (34a)$$

$$C_q^L = \frac{\omega_{L2}}{\sqrt{\eta_2}} \sqrt{\frac{1}{\pi}} \frac{\kappa}{\sqrt{\kappa^2 + q^2}} \quad (34b)$$

The confined LO phonons are related to both bulk polarization charges and interface polarization charges. The former can be found from $\rho = \vec{\nabla} \cdot \vec{P}$ and the latter follows from the boundary conditions at the interfaces. The results are

$$\rho_q^L = C_q (\kappa + q^2/\kappa) \sin(qz) \quad (35b)$$

$$\sigma_m^L = C_m \frac{m\pi}{a\kappa} \begin{cases} (-1) & \text{at } z = 0 \\ (-1)^m & \text{at } z = a \end{cases} \quad (36a)$$

$$\sigma_q^L = \begin{cases} C_q \frac{q}{\kappa} & z = 0 \\ -C_q \frac{q}{\kappa} & z = a \end{cases} \quad \text{at} \quad (36b)$$

We now turn our attention to the TO phonons for which $\omega = \omega_{T\nu}$. Equation (12a) then implies that $\chi_\nu^{-1}(\omega_{T\nu}) = 0$ and $\chi_{\nu'}^{-1}(\omega_{T\nu}) \neq 0$ for $\nu' \neq \nu$. Hence $\vec{E} = 0$ and $\vec{D} = \vec{P}$ in layer ν . The same consideration and procedures as described above for the LO phonons lead to eigenfrequencies ω_{T1} and ω_{T2} with corresponding eigenvectors

$$\vec{\pi}_m^T = \begin{cases} 0 & z < 0 \\ C_m^T [i \frac{m\pi}{a\kappa} \cos(\frac{m\pi}{a}z), \sin(\frac{m\pi}{a}z)] & 0 \leq z \leq a \\ 0 & z > a \end{cases} \quad (37a)$$

where m is an integer, and

$$\vec{\pi}_q^T = \begin{cases} C_q^T [i \frac{q}{\kappa} \cos(qz), \sin(qz)] & z < 0 \\ 0 & 0 \leq z \leq a \\ C_q^T [i \frac{q}{\kappa} \cos q(z-a), \sin q(z-a)] & z > a \end{cases} \quad (37b)$$

where q is a real number. We see from (37) that the TO phonons are also strictly confined by the presence of interfaces. Once more, these

eigenvectors are normalized according to (17), and the normalization constants are given by the same expressions as (34) except for the replacement of ω_{Lv} by ω_{Tv} . The confined TO modes are, however, not associated with any polarization charge, neither bulk nor surface charge.

The s-polarization modes are given by the solution of (15b) with $\chi_v^{-1}(\omega) = 0$. This implies that the s-polarization modes exist only when the eigenfrequencies are those of the transverse optical phonons in either medium. Since these modes are completely decoupled from the other vibrational modes, they are not involved in the interaction with electrons and hence will not be discussed further.

V. Discussion

We have shown that there exist two types of phonon modes in a double heterostructure consisting of two semiconducting materials, the interface phonons and the confined bulk phonons. The interface modes may be either symmetric or antisymmetric with respect to the center of the system. They are dispersive in nature, and their frequencies for given materials depend solely upon the dimensionless quantity κa . In the center region of the Brillouin zone, these modes have the same frequencies as those of the bulk LO and TO phonons in each material. For this reason, we shall refer to them as "LO-like" and "TO-like" interface phonons.

Since the bulk frequencies are determined by the positions of the zeroes and poles of the dielectric functions as can be seen from (12), different compositions of the double heterostructure can result in different frequency combinations. However, only three distinct combinations as shown in Fig. 2 are possible, where we have assumed $\omega_{T2} > \omega_{T1}$ without loss of generality. It is observed that as the width of the central layer

increases, the four interface modes become two degenerate modes. In the limit of large a , these modes have the same frequencies as those in a bilayer heterostructure. In case (a), the degenerate modes are material-like, while in the other two cases they are LO-like and TO-like. Experimentally, only case (a) has been observed thus far. It is therefore interesting to carry out experiments on samples with $\omega_{L1}, \omega_{L2} > \omega_{T1}, \omega_{T2}$, such as GaAs/Ga_{0.3}Al_{0.7}As (GaAs-type) and InP/AlSb.

It should also be of great interest to note that the peculiar mode observed in the 90° Raman scattering experiment²⁷ may be understood, as has been pointed out in a recent communication,²⁸ in terms of the interface modes derived in Sec. III. In other words, the novel slab modes reported in Ref. 27 are in fact the interface modes. A detailed analysis of this experiment will be published elsewhere, and here we give only a qualitative account. Since Raman scattering experiments involve only phonons of very small κ , the dominant component of the polarization vector $\vec{\pi}_a$ is P_z according to (25a). Therefore the antisymmetric interface modes are predominantly TO modes. In the central layer, this TO mode oscillates at the LO frequency of GaAs, in agreement with the experimental result in the right-angle scattering configuration. Similarly, (25b) shows that the polarization $\vec{\pi}_s$ has a dominant P_κ component, or the symmetric interface mode in the central layer is predominantly longitudinal and oscillates at the bulk TO frequency of GaAs.

In addition to the Raman scattering experiments, the interesting pinning phenomenon has been reported in recent measurements of cyclotron resonance. That the electron interacts with optical phonons at the bulk TO frequency has been observed in the measurements of the magnetopolaron frequency in semiconductor quantum wells,²⁹⁻³¹ and it has been attributed to

the classical dielectric effect.²⁹ This is essentially a polariton effect rather than a polaronic one. When the 1s-2p transition energy of a hydrogenic impurity atom in a GaAs quantum well is measured in strong magnetic fields,³¹ the pinning is found at a frequency about 40 cm^{-1} below ω_L ($\sim 20 \text{ cm}^{-1}$ below ω_T). To our knowledge, there is no theory up to the present time that can account for this result.³² The existence of traveling LO phonons and the zone-folding effect has been suggested as a possible source of this phenomenon.³¹ We have solved the interface phonon modes in a superlattice, and our preliminary results indicate that probably the interface modes are responsible for this strange pinning phenomenon. More careful study is necessary, however, before any definite conclusion can be made. Work along this direction is also underway and will be discussed in forthcoming publications.

Acknowledgments

One of us (DLL) would like to thank Bernard A. Weinstein and J. P. Cheng for useful discussions about the experimental situation and Bruce D. McCombe for helpful comments. This research was partially supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to copy and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

References

1. A. S. Baker, J. L. Merz and A. C. Gossard, Phys. Rev. B 17, 3181 (1978).
2. C. Colvard, R. Merlin, M. V. Klein and A. C. Gossard, Phys. Rev. Lett. 45, 298 (1980).
3. M. V. Klein, C. Colvard, R. Fischer and J. Morkoe, J. Physique Coll. 5, C45, 131 (1984).
4. J. Sapriel, J. C. Michel, J. C. Toledano, R. Vacher, J. Kervarec and A. Regreny, Phys. Rev. B 28, 2007 (1983).
5. C. Trallero Giner and M. Anton, Phys. Status Solidi B 133, 563 (1986).
6. F. Comas, C. Trallero Giner and H. Leon, Phys. Status Solidi B 138, 219 (1986).
7. H. Adamska and H. N. Spector, J. Appl. Phys. 56, 1123 (1984).
8. B. K. Ridley, J. Phys. C: Solid State Phys. 15, 5899 (1982).
9. S. Das Sarma, Phys. Rev. B 27, 2590 (1983).
10. X. X. Liang, S. W. Gu and D. L. Lin, Phys. Rev. B 34, 1251 (1986); B 35, 4118(E) (1987).
11. C. Y. Chen, T. Z. Ding and D. L. Lin, Phys. Rev. B 35, 4398 (1987).
12. C. Y. Chen, D. L. Lin and T. Z. Ding, Phys. Rev. B 36, 9816 (1987).
13. M. A. Brummel, R. J. Nicholas, J. C. Portal, M. Rozeghi and M. A. Poisson, Physica B & C 117 & 118 B, 753 (1983).
14. T. Englert, D. C. Tsui, J. C. Portal, J. Berens and A. Gossard, Solid State Commun. 44, 1301 (1982).
15. B. K. Ridley, in Feskorperprobleme XXV (Advances in Modern Physics), ed. by H. J. Quiesser (Vieweg, Braunschwig, 1985), p. 449 ff.
16. N. Sawaki and I. Alasaki, Physica B & C 134B, 494 (1985).
17. A. K. Arora, E. K. Suh, A. K. Ramdas, F. A. Chambers and A. L. Moretti, Phys. Rev. B 36, 6142 (1987).
18. R. Fuchs and K. L. Kliewer, Phys. Rev. 140, A2076 (1965).
19. L. Wendler, Phys. Status Solidi B 129, 513 (1985).
20. R. Lassnig, Phys. Rev. B 30, 7132 (1984).
21. M. Babiker, J. Phys. C: Solid State 12, 683 (1986).

22. M. Born and K. Huang, Dynamical Theory of Crystal Lattices (Clarendon, Oxford, 1968).
23. J. J. Licari and R. Evrard, Phys. Rev. B 15, 2254 (1977).
24. A. K. Sood, J. Menendez, M. Cardona and K. Ploog, Phys. Rev. Lett. 54, 2115 (1985).
25. D. Gammon, R. Merlin and H. Morkoc, Phys. Rev. B 35, 2552 (1987).
26. A. K. Sood, J. Menendez, M. Cardona and K. Ploog, Phys. Rev. Lett. 54, 2111 (1985).
27. J. E. Zucker, A. Pinczuk, D. S. Chemla, A. Gossard and W. Wiegmann, Phys. Rev. Lett 53, 1280 (1984).
28. D. L. Lin, R. Chen and T. F. George, Solid State Commun. (in press).
29. M. Ziesmann, D. Heitmann and L. L. Chang, Phys. Rev. B 35, 4541 (1987).
30. M. A. Hopkins, R. J. Nicholas, M. A. Brummel, J. J. Harris and C. T. Foxon, Phys. Rev. B 36, 4789 (1987).
31. Y. H. Chang, B. D. McCombe, J. M. Mercy, A. A. Reeder, J. Ralston and G. A. Wicks, Phys. Rev. Lett. 61, 1408 (1988).
32. J. P. Cheng and B. D. McCombe, Phys. Rev. Lett. 62, 1925 (1989).

Figure Captions

1. Geometry of the double heterostructure.
2. Dispersion relations of the interface modes in the double heterostructure for different compositions: (a) GaAs/AlAs for which $\omega_{L2} > \omega_{T2} > \omega_{L1} > \omega_{T1}$, (b) GaAs/Al_{0.3}Ga_{0.7}As (GaAs-type) for which $\omega_{L2} > \omega_{L1} > \omega_{T2} > \omega_{T1}$, (c) InP/AlSb for which $\omega_{L1} > \omega_{L2} > \omega_{T2} > \omega_{T1}$.

Fig. 1








Fig. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853