

AD-A222 398

DTIC FILE COPY

2



Defense Nuclear Agency
Alexandria, VA 22310-3398



DNA-TR-89-45

Chernobyl Accident Fatalities and Causes

A. Laupa
G. H. Anno
Pacific-Sierra Research Corporation
12340 Santa Monica Boulevard
Los Angeles, CA 90025-2587

June 1990

Technical Report

CONTRACT No. DNA 001-87-C-0104

Approved for public release;
distribution is unlimited.

1000
Cb

Destroy this report when it is no longer needed. Do not return to sender.

PLEASE NOTIFY THE DEFENSE NUCLEAR AGENCY,
ATTN: CSTI, 6801 TELEGRAPH ROAD, ALEXANDRIA, VA
22310-3398, IF YOUR ADDRESS IS INCORRECT, IF YOU
WISH IT DELETED FROM THE DISTRIBUTION LIST, OR
IF THE ADDRESSEE IS NO LONGER EMPLOYED BY YOUR
ORGANIZATION.



DISTRIBUTION LIST UPDATE

This mailer is provided to enable DNA to maintain current distribution lists for reports. We would appreciate your providing the requested information.

- ☐ Add the individual listed to your distribution list.
- ☐ Delete the cited organization/individual.
- ☐ Change of address.

NOTE:
Please return the mailing label from the document so that any additions, changes, corrections or deletions can be made more easily.

NAME: _____

ORGANIZATION: _____

OLD ADDRESS

CURRENT ADDRESS

TELEPHONE NUMBER: () _____

SUBJECT AREA(S) OF INTEREST:

DNA OR OTHER GOVERNMENT CONTRACT NUMBER: _____

CERTIFICATION OF NEED-TO-KNOW BY GOVERNMENT SPONSOR (if other than DNA):

SPONSORING ORGANIZATION: _____

CONTRACTING OFFICER OR REPRESENTATIVE: _____

SIGNATURE: _____

CUT HERE AND RETURN



•
•

Director
Defense Nuclear Agency
ATTN: TITL
Washington, DC 20305-1000

•
Director
Defense Nuclear Agency
ATTN: TITL
Washington, DC 20305 1000

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 900601		3. REPORT TYPE AND DATES COVERED Technical 870729 - 881231
4. TITLE AND SUBTITLE Chernobyl Accident Fatalities and Causes			5. FUNDING NUMBERS C - DNA 001-87-C-0104 PE - 62715H PR - RM TA - RH WU - DH026130	
6. AUTHOR(S) A. Laupa; G. H. Anno				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Pacific-Sierrra Research Corporation 12340 Santa Monica Boulevard Los Angeles, CA 90025-2587			8. PERFORMING ORGANIZATION REPORT NUMBER PSR Report 1865	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Defense Nuclear Agency 6801 Telegraph Road Alexandria, VA 22310-3398 RARP/Young			10. SPONSORING/MONITORING AGENCY REPORT NUMBER DNA-TR-89-45	
11. SUPPLEMENTARY NOTES This work was sponsored by the Defense Nuclear Agency under RDT&E RMC Code B4662D RM RH 00038 STRP 3500A 25904D.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Based on available sources of information, an assessment is made of the Chernobyl accident fatalities and causes of death that resulted from acute injury effects. Accident victims grouped according to whole body radiation dose and biologic response were examined and reconciled based on comparing various sources of information. Fatalities are identified with the occurrence of acute radiation sickness (ARS) syndromes. A maximum likelihood regression analysis was performed on the available fatality and dose data based on five different statistical models to estimate the LD ₁₀ , LD ₅₀ , and LD ₉₀ values. The estimated LD ₅₀ is about a factor or two higher than previously published values which is attributed to post accident emergency medical and clinical care.				
14. SUBJECT TERMS Chernobyl Accident Clinical Classification Radiation Pulmonitis			15. NUMBER OF PAGES 34	
Radiation Injury Lethality Causes Radiation Doses			16. PRICE CODE	
Bone Marrow Skin Damage Fatalities				
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED
			20. LIMITATION OF ABSTRACT SAR	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

CLASSIFIED BY:

N/A since Unclassified

DECLASSIFY ON:

N/A since Unclassified

14. SUBJECT TERMS (CONTINUED)

Intestinal Syndrome

Acute Effects

Accession For	
NTIS GRA&I	☒
DTIC TAB	☒
Unannounced	☐
Justification	
By _____	
Date _____	
For _____ Codes	
/or	
1	

A-1



CONVERSION TABLE

Conversion factors for U.S. customary
to metric (SI) units of measurement

To Convert From	To	Multiply By
angstrom	meters (m)	$1.000\ 000 \times 10^{-10}$
atmosphere	kilo pascal (kPa)	$1.013\ 25 \times 10^2$
bar	kilo pascal (kPa)	$1.000\ 000 \times 10^2$
barn	meter ² (m ²)	$1.000\ 000 \times 10^{-28}$
British Thermal unit (thermochemical)	joule (J)	$1.054\ 350 \times 10^3$
calorie (thermochemical)	joule (J)	4.184 000
cal (thermochemical) : cm ²	mega joule/m ² (MJ/m ²)	$4.184\ 000 \times 10^{-2}$
curie	giga becquerel (GBq)*	$3.700\ 000 \times 10^4$
degree (angle)	radian (rad)	$1.745\ 329 \times 10^{-2}$
degree Fahrenheit	degree kelvin (K)	$T_K = (t^{\circ}F + 459.67) / 1.8$
electron volt	joule (J)	$1.602\ 19 \times 10^{-19}$
erg	joule (J)	$1.000\ 000 \times 10^{-7}$
erg/second	watt (W)	$1.000\ 000 \times 10^{-7}$
foot	meter (m)	$3.048\ 000 \times 10^{-1}$
foot-pound-force	joule (J)	1.355 818
gallon (U.S. liquid)	meter ³ (m ³)	$3.785\ 412 \times 10^{-3}$
inch	meter (m)	$2.540\ 000 \times 10^{-2}$
jerk	joule (J)	$1.000\ 000 \times 10^9$
joule kilogram (J Kg) (radiation dose absorbed)	Gray (Gy)**	1.000 000
kilotons	terajoules	4.183
kip (1000 lbf)	newton (N)	$4.448\ 222 \times 10^3$
kip/inch ² (ksi)	kilo pascal (kPa)	$6.894\ 757 \times 10^3$
klap	newton-second/m ² (N-s-m ²)	$1.000\ 000 \times 10^2$
micron	meter (m)	$1.000\ 000 \times 10^{-6}$
mil	meter (m)	$2.540\ 000 \times 10^{-5}$
mile (international)	meter (m)	$1.609\ 344 \times 10^3$
ounce	kilogram (kg)	$2.834\ 952 \times 10^{-2}$
pound-force (lbf avoirdupois)	newton (N)	4.448 222
pound-force inch	newton-meter (N-m)	$1.129\ 848 \times 10^{-1}$
pound-force/inch	newton/meter (N/m)	$1.751\ 268 \times 10^2$
pound-force/foot ²	kilo pascal (kPa)	$4.788\ 026 \times 10^{-2}$
pound-force/inch ² (PSI)	kilo pascal (kPa)	6.894 757
pound-mass (lbm avoirdupois)	kilogram (kg)	$4.535\ 924 \times 10^{-1}$
pound-mass-foot ² (moment of inertia)	kilogram-meter ² (kg-m ²)	$4.214\ 011 \times 10^{-2}$
pound-mass/foot ³	kilogram/meter ³ (kg/m ³)	$1.601\ 846 \times 10^{-1}$
rad (radiation dose absorbed)	Gray (Gy)**	$1.000\ 000 \times 10^{-2}$
roentgen	coulomb/kilogram (C/kg)	$2.579\ 760 \times 10^{-4}$
shake	second (s)	$1.000\ 000 \times 10^{-8}$
slug	kilogram (kg)	$1.459\ 390 \times 10^1$
torr (mm Hg, 0°C)	kilo pascal (kPa)	$1.333\ 22 \times 10^{-1}$

*The becquerel (Bq) is the SI unit of radioactivity: Bq = 1 event/s.

**The Gray (Gy) is the SI unit of absorbed radiation.

TABLE OF CONTENTS

Section	Page
CONVERSION TABLE	iii
FIGURES	v
TABLES	vi
1 INTRODUCTION	1
2 DATA SOURCES ON CHERNOBYL VICTIMS	2
3 CHERNOBYL FATALITIES	6
4 UNCONFIRMED CHERNOBYL FATALITIES	8
5 CAUSES OF FATAL OUTCOMES	9
6 FATALITY INCIDENCE VERSUS RADIATION DOSE	14
7 LIST OF REFERENCES	24

FIGURES

Figures	Page
1 Chernobyl accident fatalities: normal distribution, 238 individuals, 90% confidence interval	18
2 Chernobyl accident fatalities: log-normal distribution, 238 individuals, 90% confidence interval	19
3 Chernobyl accident fatalities: Weibull distribution, 238 individuals, 90% confidence interval	20
4 Chernobyl accident fatalities: logistic distribution, 238 individuals, 90% confidence interval	21
5 Chernobyl accident fatalities: log-logistic distribution, 238 individuals, 90% confidence interval	22

TABLES

Table		Page
1	Chernobyl victim follow-up reported by State Committee [1986]	3
2	Chernobyl victim follow-up reported by Guskova [1987]	4
3	Chernobyl victim follow-up reported by Fry [1987]	5
4	Chernobyl fatalities	7
5	ARS fatalities at Hospital No. 6 in Moscow	10
6	ARS syndromes identified as causes of fatalities	10
7	Chernobyl accident fatalities and causes, Hospital No. 6 in Moscow	12
8	Radiation doses of individual patients	15
9	Radiation doses of individuals and groups	16
10	Regression models	17
11	Fatality percentile doses	18

SECTION 1

INTRODUCTION

This report reviews current available Soviet data sources on Chernobyl accident victims hospitalized for acute radiation sickness (ARS). Patients are grouped by the whole-body gamma radiation dose indicating the severity of bone marrow syndrome and biologic response criteria used to place patients in the different dose groups. We compare and reconcile the different data sources according to the fatalities reported as a result of the Chernobyl accident. Included is a brief review of information on additional unconfirmed casualties attributed to the nuclear accident.

We further review the Soviet data sources in terms of the reported causes of fatal outcomes and the occurrence of ARS syndromes identified as causes of fatalities. In many cases, fatal outcomes are ascribed to the effect of several severe ARS

syndromes, each of which could have been an independent cause of a fatal outcome. All patient groups identified in the data sources are correlated in terms of the severity of bone marrow syndrome and ascribed causes of fatal outcome. Based on the available data, all but two of the Chernobyl fatalities can be accounted for according to patient category, dose level, and lethal cause considerations.

A maximum likelihood analysis of the available data on the Chernobyl accident survivors and fatalities was also performed. Five different statistical models were applied to develop estimates of the incidence of lethality with total body dose. The marked effect in survival owing to medical attention afforded the accident victims is demonstrated.

SECTION 2

DATA SOURCES ON CHERNOBYL VICTIMS

There are three sources of original data on Chernobyl patients with acute radiation sickness:

- USSR State Committee on the Utilization of Atomic Energy, (hereinafter referred to as State Committee). *The Accident at the Chernobyl Nuclear Power Plant And Its Consequences*, August 1986
- Guskova. *Early Acute Effects Among the Victims of the Accident at the Chernobyl Nuclear Power Plant*, April 1987
- Fry. *Foreign Trip Report*, October 1987

Dr. S. A. Fry reports on presentations by Dr. A. Barabanova, a Soviet physician associated with Dr. A. Guskova at the Moscow Hospital No. 6, and by Dr. D. P. Orsanov, an internationally known biophysicist who went to Chernobyl two to three days after the accident to supervise environmental monitoring activities.

The main clinical response and medical outcome data on ARS patients, as reported in the three references, are summarized in Tables 1, 2, and 3.

Table 1. Chernobyl victim follow-up reported by State Committee [1986].

Dose Group	Dose Range (Gy)	No. of Patients		Onset Primary Reaction (h)	Latent Period (days)	Lymphocyte Count at 3-6 Days (per mm ³)	Time Platelets $\leq 4000/\text{mm}^3$ (days)	No. of Deaths		Fatal Radiation Damage To Skin
		Moscow	Kiev					Moscow	Kiev	
1st	0.8-2.1	(105) ^a		2	30+	600-1000	25-68	--	--	--
2d	2-4	53		1-2	15-25	300-500	17-24	--	--	--
3d	4.2-6.3	21	2	0.5-1.0	8-17	100-200	10-16	7 ^b	--	6
4th	6-16	20	2	0.5	6-8	<100	8-10	17 ^b	2 ^c	14
Total		203						26 ^d		20 ^e

^aRemainder not specifically reported.

^bIn days 10 to 50.

^cOn days 4 and 10.

^dOne additional victim died from severe burns at 6:00 a.m. on 26 April; another victim's body was not discovered. which puts the total at 28 deaths for the first 50 days following the accident.

^eGiven as 20 on p. 35 and as 19 on p. 34 of State Committee [1986].

Table 2. Chernobyl victim follow-up reported by Guskova [1987].

Dose Group	Dose Range (Gy)	No. of Patients		No. of Patients with Radiation Burns ^a		No. of Deaths		Day of Death after Exposure
		Moscow	Kiev	% of body surface damaged	Total	from ARS	Moscow Kiev	
1st	0.8-2.1	31	109	--	1	2	3	--
2d	2.0-4.0	43	12	1	9	2	12	1 -- 96
3d	4.2-6.3	21	--	3	15	3	21	7 -- 16, 18, 21, 23, 32, 34, 48,
4th	6-16	20	1	9	10	1	20	14, 14, 14, 15, 17, 17, 18, 18, 18, 20, 21, 23, 24, 24, 25, 30, 48, 77, 91
Total		115	122			56 ^d	27	1 ^c
Overall		237 ^e						28 ^f

^aBased on patients in Moscow.

^bFrom text, this percentage should appear as 50-90 (or 50-100).

^cDied on day 10 from combined heat and radiation injuries.

^dThe radiation burns were unquestionably fatal at least in 19 cases.

^eA change from 203 to 237, announced on November 1986, occurred primarily as a result of persons with 1st degree ARS.

^fWith the first two (one at his work station, the other from heat burns in Pripjat in the first 5 h), the total is 30 deaths.

Table 3. Chernobyl victim follow-up reported by Fry [1987].

Chernobyl Fatalities		
Description		No. of Victims
Initial (on-site and in Pripjat)		2
Died as result of ARS		
In Moscow		27
In Kiev		1
Died as result of CVA		1
Total deaths		31
Radiation-induced Skin Injuries, Moscow Hospital No. 6		
Description		No. of Victims
Number of ARS patients		115
Radiation-induced skin injuries		56 ^a
Skin injuries incompatible with life		19
Waves of Erythema		
Wave	Onset	Remarks
1st	36 h	No obvious skin injury for 36 h. First wave lasted up to 24 h.
2d	5-10 days later	Erythema more widespread. Involved skin areas covered by clothing. Coincided with latent period of ARS.
3d	2-3 months	Included 28-30 patients. 8-10 had no previous manifestation of skin injury.
^a Two of 56 patients also had thermal burns.		

SECTION 3

CHERNOBYL FATALITIES

The number of fatalities from the Chernobyl accident totals 31. Table 4 shows the reported numbers of death and reconciles the different data sources. There were two initial fatalities: one victim died at his workplace, his body was never recovered; another victim died from severe burns at 6:00 a.m. in the morning of the accident in the Pripyat hospital. State Committee [1986] reports 28 fatalities as occurring within the first 50 days (Table 1). Three people subsequently died from ARS on Days 77, 96, and 91 (Table 2), resulting in a total of 31 fatalities.

Dr. Barabanova reported that one transfer patient from Chernobyl, with signs and symptoms of acute radiation sickness, suffered a cerebrovascular accident (CVA) enroute and died from that cause shortly after admission to the hospital [Fry, 1987]. Hence, the Chernobyl fatalities include 2 initial fatalities, 28 ARS fatalities (27 died in Moscow, and 1 in Kiev), and 1 CVA fatality (Table 3). In Table 4, notice that

Guskova did not include the CVA fatality among the fatalities reported. Only one fatality is listed for Kiev in Table 2, whereas two fatalities are listed in Table 1.

Dr. Barabanova further reported that the CVA victim was among the fatalities at Hospital No. 6 in Moscow [Fry, 1987]. This piece of information appears to be in error. State Committee [1986] reports that two patients died in Kiev on days four and ten after the accident. Guskova [1987] shows only one death in Kiev, a patient who died on the 10th day from combined heat and radiation injuries. The transfer patient who died shortly after hospital admission from CVA must have been the Kiev fatality on day four. The early death in Kiev has been confirmed by Deputy USSR Minister of Health, Ye. Vorobyev, who told a press conference on 9 May 1986 that a third person, the first victim of radiation exposure, died three days after the accident in a Kiev hospital [Goure, 1987].

Table 4. Chernobyl fatalities.

Classification	State Committee (August 1986)	Guskova (April 1987)	Fry October 1987	Census
Initial Fatalities	2	2	2	2
Died as result of ARS				
In Moscow	24	27	27	27
In Kiev	2 ^a	1 ^b	1	1
Died as result of CVA	--	--	1	1 ^c
Reported total	28 ^d	30 ^c	31	31
Other	3 ^d	1 ^c	--	--
Total Fatalities	31	31	31	31

^aOn days four and ten.

^bOn day ten.

^cOne ARS patient suffered a CVA while being transferred from Chernobyl to Kiev and died shortly thereafter from that cause on day four.

^dReported fatalities refer to the first 50 days: three patients subsequently died from ARS [Guskova, 1987].

SECTION 4

UNCONFIRMED CHERNOBYL FATALITIES

According to Dr. Barabanova's presentation, Fry [October 1987] reports the following: "A journalist/photographer who spent some time at the accident site shortly after the accident is reported to have died about 9 months later; the cause of his death is controversial according to Dr. Barabanova who did not elaborate further."

The Moscow weekly *Nedelya* cited *Novoye Russkoye Slovo* of 31 May 1987 on the death of the motion picture director Vladimir Shevchenko two months earlier from radiation sickness [Goure, 1987a]. Shevchenko produced the documentary, "Chernobyl, Chronicle of Difficult Days," which was filmed in May 1986 at the Chernobyl power station. *Nedelya* also reported that two cameramen received large radiation doses and are [then] currently hospitalized.

The documentary film on the Chernobyl accident has been described in the *Moscow News* No. 11, 22-29 March 1987 [Goure, 1987a]. Readers are left with the impression that a camera and its crew are taken on a tour through the damaged reactor, often close to high levels of radiation by the plant's chief engineer, who says, "That pipe

there (ten or so meters away) is roughly 1000 roentgens."¹ Later, in referring to Y. Velikhov, Vice President of the USSR Academy of Sciences, seen walking in front of the camera, the reporter adds: "A little bit earlier in the film, we were told why the famous scientists had to risk their lives. What about the invisible cameraman? The other members of the filming crew?"

Three Soviet sources appear to agree on at least one additional Chernobyl fatality. The *Moscow News* highlights the radiation risk to the camera crew. *Nedelya* reports the death of the movie director and the hospitalization of two cameramen. Dr. Barabanova mentions the reported but controversial death of a journalist/photographer.

Non-Soviet sources report additional fatalities and radiation injuries among the cleanup crews according to Radio Free Europe "Rad Background Report," 14 October 1986 [Goure, 1987b]. At the same time, official Soviet spokesmen deny that more people have died from radiation overdoses at the Chernobyl atomic power plant since the accident there last year killed 31 people [Los Angeles Times, 1987].

1. Probably refers to a dose rate of 1000 R/h.

SECTION 5

CAUSES OF FATAL OUTCOMES

All three Soviet data sources discuss the importance of radiation damage to skin and basically repeat the same message:

- "In 19 cases, the deaths of patients with third- and fourth degree acute radiation disease occurred only as a result of severe damage, which was incompatible with survival, to 50-90% of the surface of their bodies." [State Committee, 1986]
- "In the cases of at least 19 of the 56 patients with burns, the burns were unquestionably fatal." [Guskova, 1987]
- "Of the 27 patients who died of radiation injuries at Hospital No. 6, 19 had radiation induced skin injuries over 90-100% [sic] of their body surface. It was the Soviets's opinion that in these cases, the skin injuries were so severe as to be considered incompatible with life, even in the absence of bone marrow damage." [Fry, 1987]

All these statements appear to be true but incomplete. Guskova [1987] discusses the interrelationships between individual causes of lethal outcomes in somewhat more detail, summarized in Table 5. Although all patients with third- and fourth-degree bone-marrow syndrome had severe radiation burns (Table 2), only five lethal outcomes could be ascribed exclusively to radiation damage to vast areas of skin. These five cases did not involve radiation enteritis or irreversible myelodepression, and their whole-body doses did not exceed 6 Gy. In 14 other cases, radiation damage to skin was combined with other severe syn-

dromes, each of which could have been an independent cause of a fatal outcome. These two groups total 19 fatalities, which is the number of cases with severe skin injury commonly cited as being incompatible with life. Apparently, the remaining eight fatalities were caused by combinations of various syndromes without radiation skin damage being an independent cause of the fatal outcomes.

Various ARS syndromes reported by Guskova [1987] as apparent independent causes of fatal outcomes (or being capable of producing a fatal outcome) are listed in Table 6. As mentioned before, radiation skin damage was the exclusive or contributing cause in 19 fatal outcomes. Lethal intestinal syndrome, indicated by the appearance of diarrhea from day four to day eight, was noted in 10 patients. These patients were exposed to 10 Gy or higher gamma radiation; they all died within the first three weeks after exposure.

Acute radiation pneumonitis was observed in seven patients with third- and fourth-degree ARS. It was typified by a rapidly intensifying difficulty in breathing, by ventilation failure, and by the onset of lethal outcomes from hypoxemic coma. "Autopsy revealed large blue lungs with marked interstitial edema." [Guskova, 1987]

Six cases with lethal outcomes were ascribed to radiation damage causing irreversible hematopoietic aplasia or to complications caused by bone marrow transplants. Hemophilia achieved thanatogenic significance only in one case.

Table 5. ARS fatalities at Hospital No. 6 in Moscow.

Causes of Fatality	Number of Fatalities
Radiation skin damage exclusively	5
Skin damage combined with other ARS syndromes	14
Combinations of ARS syndromes other than skin damage	<u>8</u>
Total fatalities	27

Table 6. ARS syndromes identified as causes of fatalities.

ARS Syndrome	Fatalities with Syndrome
Radiation skin damage	19
Intestinal syndrome (10 Gy or more)	10
Acute radiation pneumonitis	7
Irreversible hematopoietic aplasia or bone-marrow transplant complications	6
Hemophilia	1
Thermal burns/internal contamination	2
Radiation induced vascular damage	1

In Moscow, 56 of the 115 ARS patients suffered from radiation induced skin injuries. Two of the 56 patients formed a unique subgroup: they also had severe thermal skin burns produced by steam and incurred a significant internal radionuclide dose estimated as 1.5 to 4 sieverts on the basis of postmortem radiometry. The radionuclide doses of all other patients did not exceed 1 to 3 percent of their external dose. The gamma dose of these two patients was on the order of 4 to 5 Gy. The patients died on days 23 and 18, respectively, from their combined injuries [Fry, 1987].

One patient in Moscow suffered a fatal CVA², probably associated with generalized radiation induced vascular damage [Fry, 1987]. The patient received an average bone marrow dose of 3 Gy and suffered severe radiation induced skin injuries, manifesting as three waves of erythema, with the third wave developing almost three months after the accident. Accordingly, this patient (case 3 presented by Dr. Barabanova) must be the one fatality in the second-degree ARS group who died on day 96 after the accident (Table 2).

Based on currently available Soviet data sources reviewed, Table 7 provides a summary of our effort to correlate the Chernobyl accident fatalities and causes grouped in terms of:

- Severity of the bone marrow syndrome (the whole-body gamma radiation doses).
- ARS syndrome identified as exclusive or contributing cause of fatal outcomes.

- Patient injury or medical treatment category.

Patient categories include one cerebrovascular accident, a patient in the medium gamma dose group who died from lethal radiation damage to the vascular system. Five fatal outcomes are ascribed solely to radiation damage to large skin areas. These patients fall in the severe gamma dose group: their estimated whole-body doses did not exceed 6 Gy.

Further, patient groups include bone marrow transplants and fetal liver cell transplants. These patients were selected on the basis of a whole-body dose of 6 Gy or greater, estimated from the lymphocyte count in peripheral blood and from cytogenetic analysis of chromosome aberrations. This dose level was understood to represent irreversible or extremely protracted and deep myelodepression.

The bone marrow transplants include 13 patients grouped in two subcategories:

- a. Six patients had radiation damage to skin and intestines at a level that was deemed not incompatible with life. Four patients died between days 27 and 29 after the transplants. Two patients survived their bone marrow transplants; they had gamma exposures of 5.8 and 9.0 Gy. Gamma doses for the four patients who died ranged from 4.3 to 10.7 Gy, placing one fatality in the severe and three fatalities in the extremely severe bone marrow syndrome groups. They died from "mixed virus-bacterial infections," also called as "complications caused by bone marrow

2. Note that this CVA fatality occurred in Moscow; a different ARS patient suffered a CVA enroute to the hospital and died on day four in Kiev.

Table 7. Chernobyl accident fatalities and causes. Hospital No. 6 in Moscow.

Patient Category	Bone Marrow Syndrome			Lethal Radiation Damage to				Lethal Infections/		Internal Contamination	Hemophilia
	Fatalities			Skin	Intestines	Lungs	Vascular System	GVH ^a	Thermal Burns		
	Medium 2-4 Gy	Severe 4-6 Gy	Extremely Severe 6-16 Gy								
Cerebrovascular accident	1			--	--	--	Yes	--	--	--	--
Radiation skin damage alone		5		Yes	--	--	--	--	--	--	--
Bone marrow transplants											
Radiation damage "not incompatible with life"		1	3	--	--	--	--	Yes	--	--	?
Radiation damage "incompatible with life"			7	Yes	Yes	Yes	--	?	--	--	?
Fetal liver cell transplants			6	Yes	Yes	--	--	--	--	--	--
Thermal burns/internal contamination			2 ^b	Yes	--	--	--	--	Yes	Yes	--
Unspecified		1	1	?	?	?	--	--	--	--	--
Total	1	7	19	19	10	7	1	6	2	2	1

^aGVH = graft versus host.

^bExternal gamma dose 4 to 5 Gy, internal radionuclide dose 1.5 to 4 Gy.

transplants" [Guskova, 1987]. State Committee [1986] reports that all six bone marrow transplants in this group had similar disorders that may have been caused by graft-versus-host (GVH) disease. Two of the four fatalities in this group occurred prior to the publication of the State Committee [1986] report. It is stated that the GVH reactions may have been contributing causes of these two deaths.

- b. Seven patients had radiation damage to their skin, intestines, and lungs at a level that was deemed incompatible with life. They died from acute radiation damage to the skin, gut, and lungs on days 2 to 19 after the transplants (days 15 to 25 after exposure).

In the category of fetal liver cell transplants, all six patients died from radiation damage to the skin and intestines [Guskova, 1987]. Patients for the transplantation of human embryo liver cells were selected on the basis of extremely severe damage to the skin and intestines and extremely unfavorable prognosis [State Committee, 1986].

Two patients in the thermal burns and internal contamination category died from their combined injuries, including bone marrow damage and radiation damage to skin.

In Table 7 all but two of the 27 fatalities in Moscow can be traced through patient categories, thus leaving only two in the

"unspecified" category. The totals of individual lethal causes have been taken from Table 6. Adding the numbers of patients in relevant patient categories, the corresponding totals are not always identical with those in Table 6, but are still in general agreement. For example, we would have 13 rather than 10 cases of lethal radiation damage to intestines. In cases where the fatal outcomes associated with a patient category are ascribed to multiple contributing ARS syndromes ("yes" in Table 7), all the syndromes may not necessarily be present for each and every fatality.

In other cases, the relevant patient categories identify fewer fatalities than the totals of Table 6. For example, only four rather than six total cases of lethal infections and GVH diseases are identified directly. In these cases, the question marks in Table 7 suggest other possible patient categories associated with a particular cause of death.

The individual gamma doses of the accident victims are generally unavailable. Exceptions are the cerebrovascular accident victim with an estimated dose of 3 Gy, the four bone marrow transplant fatalities who died from mixed virus and bacterial infections with estimated doses of 4.3, 5.0-7.9, 5.8-6.0, and 10.7 Gy, respectively, and the two thermal burn and internal contamination patients with their combined doses shown in Table 7. Guskova [1987] also states that the patients with lethal intestinal syndromes received a short-term whole-body gamma radiation dose on the order of 10 Gy or higher.

SECTION 6

FATALITY INCIDENCE VERSUS RADIATION DOSE

A total of 238 Chernobyl accident victims received estimated whole-body gamma radiation doses in excess of 0.8 Gy. They were diagnosed as suffering from acute radiation sickness (ARS). Based on the severity of ARS, the patients were grouped in four dose range categories. The radiation data published to date include the dose ranges and the number of patients and deaths in each dose range group.

Although individual case histories have not been published, individual dose estimates have been reported for some cases, including 11 survivors and 7 fatalities. Table 8 lists individual doses and reference followed by a number or letter of patient identification. The numbers associated with Guskova [1987] refer to the original patient identification system; the letter "x" refers to patients mentioned but not otherwise identified in reports. Some of the individual doses reported by Guskova [1987] have been updated in a later report (UNSCEAR, 1987); Table 8 contains the updated values.

Table 9 summarizes the available data on radiation doses for both individuals and groups of individuals. In most cases, the patients in a dose group divide into two subgroups: a few patients with individual dose estimates, and the remainder distributed over the given dose range. The fatalities of the fourth dose group have additional patient subgroups. Ten patients died as a result of lethal intestinal syndrome; they were exposed to a dose of 10 Gy or higher (Table 6). Two patients died from a combination of severe thermal skin burns, internal radionuclide contamination, and bone marrow syndrome. Their average combined dose was about 7.25 Gy (Table

7). Four individuals in the fourth dose group have specific dose estimates, but no specific values were available for the remaining five individuals.

Based on the method of analyzing binary data given by Cox (1983), a maximum likelihood analysis of the data given in Table 9 was performed to estimate the incidence of Chernobyl fatalities with dose. Along with the explicit dose estimates given in Table 9 for both survivors and fatalities, it was assumed that the remaining individual doses were uniformly distributed over the corresponding dose range. For example, there are five fatalities in the dose range of 4.2 to 6.3 Gy listed under the "3rd" dose group in Table 9; thus, for this analysis, doses of 4.41, 4.83, 5.25, 5.67, and 6.09 Gy each were assumed for the five fatalities.

For purposes of comparison, five different model forms were applied to analyze the survivor/fatality binary data: normal, log-normal, Weibull, logistic, and log-logistic. Table 10 gives the model forms along with parameter estimates obtained. Based upon the χ^2 goodness-of-fit statistic, a ranking would fall between the Weibull and log-normal models, with Weibull the best fit. However, all models fit the data well ($p < 0.005$, $v = 236$ dgf).

Table 11 gives a summary of the 10-, 50-, and 90-percentiles obtained for each model. The 10-percentile dose estimates range from 391 to 423 cGy, the 50-percentile from 590 to 616 cGy, and the 90-percentile estimates from 789 to 895 cGy.

Plots of the incidence of fatalities are shown in Figs. 1 through 5 for each of the

Table 8. Radiation doses of individual patients.

Dose Group	Dose Range (Gy)	Number of		Survivors			Fatalities		
		Patients	Deaths	Dose	Patient Reference		Dose	Patient Reference	
1st	0.8-2.1	140	--	0.9	Guskova [1987]	- 97			
				1.4		- 48			
				1.9	Geiger [1986]	- 6			
2nd	2.0-4.0	55	1	3.3	Guskova [1987]	- 39	3	Fry [1987]	- 3
				3.9		- 21			
				3.3	State Committee [1986]	- D			
				2.25	Fry [1987]	- 2			
				3.0	Telyatnikov [1987]				
3rd	4.2-6.3	21	7	5.6	Guskova [1987]	- x	5.2	Guskova [1987]	- 6
				4.9	Geiger [1986]	- 2	4.4		- 5
4th	6-16	22	21	8.7	Guskova [1987]	- x	6.4	Guskova [1987]	- 28
							10.2		- 16
							9		- 8
							7	Fry [1987]	- x 7
Totals		238	29	11 patients			7 patients		

Table 9. Radiation doses of individuals and groups.

Dose Group	Dose Range (Gy)	Number of		Survivors		Fatalities	
		Patients	Deaths	Number	Dose or Dose Range (Gy)	Number	Dose or Dose Range (Gy)
1st	0.8-2.1	140	--	3 ^a 137 ^b	0.9, 1.4, 1.9 0.8-2.1		
2nd	2.0-4.0	55	1	5 ^a 49 ^b	2.25, 3.0, 3.3, 3.3, 3.9 2.0-4.0	1 ^a	3.0
3rd	4.2-6.3	21	7	2 ^a 12 ^b	4.9, 5.6 4.2-6.3	2 ^a 5 ^b	4.4, 5.2 4.2-6.3
4th	6-16	22	21	1 ^a	8.7	4 ^a 10 ^c 2 ^d 5 ^b	6.4, 7, 9, 10.2 10-16 7.25, 7.25 6-16
Total		238	29	209		29	

^aFrom Table 8.

^bRemainder of patients in dose group.

^cPatients with lethal intestinal syndrome (Table 6).

^dPatients with thermal burns and internal contamination (Table 7).

Table 10. Regression models.

Model		Parameters						
Type	Form	μ^* (cGy)	σ^* (cGy)	α	β	η	λ	χ^{2**}
Normal	$\Phi(\alpha+\beta D)$	616.4	161.1	1.737	0.00621	--	--	65.60
Log-normal	$\Phi(\alpha+\beta \ln D)$	623.3	206.5	1.9186	0.00484	--	--	76.28
Weibull	$1-\exp[-(D/\lambda)^\eta]$	--	--	--	--	4.149	689.75	57.58
Logistic	$\{1+\exp[-(\alpha+\beta D)]\}^{-1}$	--	--	0.9940	0.00661	--	--	67.95
Log-logistic	$\{1+\exp[-(\alpha+\beta \ln D)]\}^{-1}$	--	--	1.896	0.00501	--	--	72.05

* μ and σ are mean and standard deviation, respectively

** $p < 0.005$, $\nu = 236$ dgf.

Table 11. Fatality percentile doses.

Model	Percentile Dose (cGy)			slope* (percent/cGy)
	LD ₁₀	LD ₅₀	LD ₉₀	
Normal	409.9	616.4	822.9	0.194
Log-normal	391.4	592.0	895.3	0.159
Weibull	401.0	631.4	843.3	0.181
Logistic	422.8	606.0	789.3	0.218
Log-logistic	403.3	590.1	863.5	0.174

*slope $\equiv \frac{(90 - 10)}{LD_{90} - LD_{10}}$ (percent/cGy)




Figure 1. Chernobyl accident fatalities: normal distribution, 238 individuals. 90% confidence interval.




Figure 2. Chernobyl accident fatalities; log-normal distribution, 238 individuals, 90% confidence interval.




Figure 3. Chernobyl accident fatalities: Weibull distribution, 238 individuals, 90% confidence interval.




Figure 4. Chernobyl accident fatalities: logistic distribution, 238 individuals, 90% confidence interval.




Figure 5. Chernobyl accident fatalities: log-logistic distribution, 238 individuals, 90% confidence interval.

five different models used to analyze the data: the 90% confidence bounds are given by the two dashed lines. In order to graphically illustrate the data from Table 9 used in fitting the various models, the data were grouped according to the horizontal lines shown in the plots. The data used in the maximum likelihood procedure is treated in binary form, and the asterisks on the horizontal line simply indicate the dose range midpoints of the data groups. Accordingly, while the log-normal (Fig. 2) and log-logistic (Fig. 5) models appear by inspection to fit the data better than the other models, they actually have the lowest χ^2 values (see Table 10); although as indicated above, all five of the models fit the data well.

A range of incidence for radiation lethality based on some prior estimates is also indicated in Figs. 1 through 5 by the broad shaded band. The left-most boundary is based on an LD_{50} midline tissue dose of 256 cGy and a slope estimated from dose response curves for large animals (swine, sheep, goats, and dogs) which tend to be parallel (Bond and Robertson 1957, Cronkite and Bond 1958, and Cronkite 1982). The right-most boundary is based on an LD_{50} value of 325 cGy, suggested by Lushbaugh (1969) for healthy young adults, with the slope also inferred from large animals.

One means of assessing the efficacy of medical care including the various treatment modalities of ARS afforded the Chernobyl accident victims is by comparing

the LD_{50} based on prior estimates with that estimated here from the Chernobyl data. That is on average, prior estimates of the LD_{50} would be about 290 cGy whereas the value is about 607 cGy according to our analysis of the Chernobyl data; this amounts to about a factor of two reflected by medical care.

Also, as is apparent from the plots of fatality incidence versus dose, in general there is a significant difference in the slope magnitude between the Chernobyl dose response curves and that according to prior estimates of lethality. As indicated in Table 11, depending upon the model, slope estimates based on analysis of the Chernobyl data range from about 0.16 to 0.22 percent/cGy; the corresponding slope value from prior lethality estimates is about 0.44 percent/cGy. This difference in the steepness of the dose response curves could also be due in part to medical attention. Although more than likely, it is due to factors other than strictly ARS in combination with medical attention such as injuries to skin, which as mentioned above was a very significant factor. For example, in the absence of medical attention, a larger proportion of lethality would be expected at lower doses, i.e., in the second and third dose groups. This certainly would tend to steepen the dose response curves. However, because of the imprecision of data regarding burn injuries and other non-ARS effects, we have not attempted to perform any isolated causal analysis with regard to fatality incidence utilizing truncated subsets of data.

SECTION 7

LIST OF REFERENCES

- Bond, V. P. and J. S. Robertson, *Ann. Rev. Nuc. Sci.*, Vol. 7, pp. 135-162, 1957.
- Cox, D. R., *Analysis of Binary Data*, Chapman and Hall Ltd., New York, NY, 1983.
- Cronkite, E. P., "The Impact of Estimates of Human Radiation Tolerance upon Radiation Emergency Management," *Proceedings of a Symposium on the Control of Exposure of the Public to Ionizing Radiation in The Event of Accident or Attack*, National Council on Radiation Protection and Measurement, Bethesda, Maryland, May 1982, pp. 21-27.
- Cronkite, E. P. and V. P. Bond, "Acute radiation syndrome in man," *U.S. Armed Forces Med. J.*, Vol. 9, 1958, pp. 313-324.
- Fry, S. A., *Trip Report: International Atomic Energy Agency (IAEA) Workshop on the Medical Handling of Skin Lesions Following High Level Accidental Irradiation*, Institute Curie, Paris, France, Center for Epidemiologic Research, Medical and Health Services Division, Oak Ridge Associated Universities, September 28-October 2, 1987.
- Geiger, H. Jack, *The Accident at Chernobyl and the Medical Response*, Journal of American Medical Association, Vol. 256, No. 5, pp. 609- 612, August 1, 1986.
- Goure, Leon, "Radiation Situation at the AES and in the Danger Zone," *The Chernobyl Accident Data Base*, Sec. V, Science Applications International Corp. La Jolla, California, October 1987a.
- , "Medical," *The Chernobyl Accident Data Base*, Sec. VIII, Science Applications International Corp. McLean, Virginia, October 1987b.
- Guskova, A. K., *Early Acute Effects among the Victims of the Accident at the Chernobyl Nuclear Power Plant*, Ministry of Health of the USSR, LN-818-87, April 1987.
- Los Angeles Times*, "The World," Part 1, p. 2, December 10, 1987.
- Lushbaugh, C. C., "Reflections on Some Recent Progress in Human Radiobiology," *Advances in Radiation Biology*, Vol. 3, Academic Press, New York, 1969, pp. 277-315.
- Telyatnikov, L. P., Personal interview with the Fire Brigade Commander, November 1987.
- UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation), A/AC.82/R. 473, Annex H, *Early Effects in Man of High Doses of Radiation*; Appendix 1, "Acute Radiation Effects in Victims of the Chernobyl Nuclear Power Plant Accident," A. G. Guskova, ed., 10 November 1987.
- USSR State Committee on the Utilization of Atomic Energy, "The Accident at the Chernobyl Nuclear Power Plant And Its Consequences: Annex 7, Medical and Biological Problems," trans. by IAEA, IAEA Experts Meeting, Vienna, Austria, August 25-29, 1986.

DISTRIBUTION LIST

DNA-TR-89-45

DEPARTMENT OF DEFENSE

ARMED FORCES RADIOBIOLOGY RSCH INST
ATTN: AFFRI-DEPT OF RAD BIOCHEM
ATTN: BHS
ATTN: DIRECTOR
ATTN: EXH
ATTN: MRA
ATTN: PHY
ATTN: RSD
ATTN: SCIENTIFIC DIRECTOR
ATTN: TECHNICAL LIBRARY

ASSISTANT SEC OF DEF (C3I)
ATTN: DIR (S&TNFC3)

ASSISTANT SECRETARY OF DEFENSE
INTERNATIONAL SECURITY POLICY
ATTN: NUC FOR & ARMS CNTRL PLCY

ASSISTANT TO THE SECRETARY OF DEFENSE
ATTN: EXECUTIVE ASSISTANT
ATTN: MIL APPL
ATTN: MIL APPL, C FIELD

DEFENSE INTELLIGENCE AGENCY
ATTN: DB
5 CYS ATTN: DB-4, RSCH RESOURCES DIV
ATTN: DB-5C
ATTN: DB-6B
ATTN: DIA/VPA-2 (FED RES DIV)
ATTN: DN
ATTN: DT
ATTN: OFFICE OF SECURITY
ATTN: OS
ATTN: RTS-2B

DEFENSE INTELLIGENCE COLLEGE
ATTN: DIC/RTS-2
ATTN: DIC/2C

DEFENSE LOGISTICS AGENCY
ATTN: COMMAND SECURITY OFC

DEFENSE NUCLEAR AGENCY
ATTN: CID
ATTN: DFRA
ATTN: DFSP G ULLRICH
ATTN: NANF
ATTN: NASF
ATTN: OPNA
ATTN: OPNS
ATTN: RAEE
20 CYS ATTN: RARP
4 CYS ATTN: TITL

DEFENSE NUCLEAR AGENCY
ATTN: TDNM
2 CYS ATTN: TDTT W SUMMA

DEFENSE TECHNICAL INFORMATION CENTER
2 CYS ATTN: DTIC/FDAB

FIELD COMMAND DEFENSE NUCLEAR AGENCY
ATTN: FCPR

3 CYS ATTN: FCPR
ATTN: NUC SECURITY

INTELLIGENCE CENTER, PACIFIC
ATTN: COMIPAC

INTERSERVICE NUCLEAR WEAPONS SCHOOL
ATTN: TTV
2 CYS ATTN: TTV 3416TH TTSQ

JOINT DATA SYSTEM SUPPORT CTR
ATTN: C-332
ATTN: JNSV

NATIONAL DEFENSE UNIVERSITY
ATTN: ICAF, TECH LIB
ATTN: NWCLB-CR
ATTN: LIBRARY
ATTN: STRAT CONCEPTS DIV CTR

NATIONAL SECURITY AGENCY
ATTN: CHIEF A4213

NET ASSESSMENT
ATTN: DOCUMENT CONTROL

OFFICE OF THE SEC OF DEFENSE
ATTN: NAVAL FORCES
ATTN: STRATEGIC PROG & TNF

STRATEGIC AND THEATER NUCLEAR FORCES
ATTN: DR E SEVIN
ATTN: DR SCHNEITER

STRATEGIC TARGET PLANNING
ATTN: JKC (ATTN: DNA REP)
ATTN: JLT
ATTN: JP
ATTN: JPTP

THE JOINT STAFF
ATTN: ED30
ATTN: J-3 SPECIAL OPERATIONS
ATTN: J-3 NUC OPERATIONS BR
ATTN: J-5 NUC & CHEMICAL DIV
ATTN: JAD/SFD
ATTN: JAD/SSD
ATTN: JSOA
ATTN: SAGA

U S EUROPEAN COMMAND/ECJ-3
ATTN: ECJ-3

U S EUROPEAN COMMAND/ECJ-5NPG
ATTN: ECJ-5

U S EUROPEAN COMMAND/ECJ-6-OJ
ATTN: ECJ-6

U S EUROPEAN COMMAND/ECJ2-T
ATTN: ECJ2-T (TGTS DIV)

U S EUROPEAN COMMAND/ECJ4-LW
ATTN: ECJ4-LW

DNA-TR-89-45 (DL CONTINUED)

U S EUROPEAN COMMAND/ECJ5-N
ATTN: ECJ5-N (NUC BRANCH)

U S EUROPEAN COMMAND/ECJ7/LW
ATTN: ECJ-7 LW

UNDER SEC OF DEFENSE FOR POLICY
ATTN: DUSP/P
ATTN: USD/P

UNDER SECRETARY OF DEFENSE
ATTN: CHAIRMAN PSEAG
ATTN: CHAIRMAN, DEF SCIENCE BD
2 CYS ATTN: C3I

DEPARTMENT OF THE ARMY

ARMY RESEARCH INSTITUTE
ATTN: COMMANDER

COMBAT MATERIAL EVAL ELEMENT
ATTN: SECURITY ANALYST

DEP CH OF STAFF FOR OPS & PLANS
ATTN: DAMO-FDQ
ATTN: DAMO-NCN
ATTN: DAMO-NCN(NUC CHEM DIR)
ATTN: DAMO-ZXA

DEPARTMENT OF THE ARMY
ATTN: DAMO-NCZ
ATTN: DAMO-OD
ATTN: DAMO-ODSO
ATTN: DAPE-HRE
ATTN: SARD-ZCS (MAJ ADAMS)

HARRY DIAMOND LABORATORIES
ATTN: SLCIS-IM-TL (TECH LIB)

JOINT SPECIAL OPERATIONS COMMAND
ATTN: J-2
ATTN: J-5

U S ARMY AIR DEFENSE SCHOOL
ATTN: COMMANDANT

U S ARMY ARMAMENT RSCH DEV & ENGR CTR
ATTN: DRDAR-LCN-F

U S ARMY ARMOR SCHOOL
ATTN: ATSB-CTD
ATTN: TECH LIBRARY

U S ARMY BALLISTIC RESEARCH LAB
ATTN: AMXBR-VLD-R (DR RAINIS)
ATTN: AMXBR-VLD, DR KLOPCIC
ATTN: DRDAR-BLA-S, TECH LIB
ATTN: DRDAR-BLT
ATTN: SLCBR-D

U S ARMY CHEMICAL SCHOOL
ATTN: ATZN-CM-M

U S ARMY COMD & GENERAL STAFF COLLEGE
ATTN: ATZL-SWJ-CA
ATTN: ATZL-SWT-A

U S ARMY CONCEPTS ANALYSIS AGENCY
ATTN: TECHNICAL LIBRARY

U S ARMY FIELD ARTILLERY SCHOOL
ATTN: ATSF-CD

U S ARMY FORCES COMMAND
ATTN: AF-OPTS

U S ARMY HUMAN ENGINEERING LAB
ATTN: DIRECTOR
ATTN: DR D HODGE

U S ARMY INFANTRY CENTER
ATTN: ATSH-CD-CSO

U S ARMY ITAC
ATTN: IAX-Z

U S ARMY MATERIEL COMMAND
ATTN: DRCDE-D
ATTN: DRCSS

U S ARMY NUCLEAR & CHEMICAL AGENCY
ATTN: MONA-NU
ATTN: MONA-ZB

U S ARMY TRAINING AND DOCTRINE COMD
ATTN: ATCD-AO
ATTN: ATCD-FA
ATTN: ATCD-N
ATTN: ATOO-NCO

U S ARMY WAR COLLEGE
ATTN: LIBRARY
ATTN: STRATEGIC STUDIES

US ARMY MATERIEL SYS ANALYSIS ACTVY
ATTN: DRXSY-DS

USA MILITARY ACADEMY
ATTN: DEPT OF BEHAV SCI & LDRSHIP
ATTN: DIR NATL SECURITY STUDIES

USA SURVIVABILITY MANAGMENT OFFICE
ATTN: SLCSM-SE J BRAND

USACACDA
ATTN: ATZL-CAD-N

DEPARTMENT OF THE NAVY

FLEET INTELLIGENCE CENTER
ATTN: FICPAC, CODE 21

MARINE CORPS
ATTN: CODE PPO
ATTN: PSI G/RASP

MARINE CORPS DEV & EDUCATION COMMAND
ATTN: LIBRARY

NAVAL OCEAN SYSTEMS CENTER
ATTN: CODE 9642-B

NAVAL PERSONNEL RES & DEV CENTER
ATTN: CODE P302

NAVAL POSTGRADUATE SCHOOL
ATTN: CODE 1424 LIBRARY

NAVAL RESEARCH LABORATORY
ATTN: CODE 1240
ATTN: CODE 2627 (TECH LIB)

NAVAL SEA SYSTEMS COMMAND
ATTN: PMS-423
ATTN: SEA-09G53 (LIB)
ATTN: SEA-6631

NAVAL SURFACE WARFARE CENTER
ATTN: CODE F-31
ATTN: G GOO

NAVAL TECHNICAL INTELLIGENCE CTR
ATTN: NTIC-DA30

NAVAL WAR COLLEGE
ATTN: CODE E-11 TECH SVC
ATTN: CTR FOR NAV WARFARE STDS
ATTN: DOCUMENT CONTROL
ATTN: LIBRARY
ATTN: STRATEGY DEPT

NAVAL WEAPONS EVALUATION FACILITY
ATTN: CLASSIFIED LIBRARY

NUCLEAR WEAPONS TNG GROUP, ATLANTIC
ATTN: CODE 222
ATTN: DOCUMENT CONTROL

NUCLEAR WEAPONS TNG GROUP PACIFIC
ATTN: CODE 32
ATTN: DOCUMENT CONTROL

OFFICE OF CHIEF OF NAVAL OPERATIONS
ATTN: NIS-22
ATTN: NOP 06D
2 CYS ATTN: NOP 403
ATTN: NOP 50
ATTN: NOP 60
ATTN: NOP 60D
ATTN: NOP 603
ATTN: NOP 91
ATTN: OP 654

OPERATIONAL TEST & EVALUATION FORCE
ATTN: COMMANDER

PLANS, POLICY & OPERATIONS
ATTN: CODE-P
ATTN: CODE-POC-30

TACTICAL TRAINING GROUP, PACIFIC
ATTN: COMMANDER

DEPARTMENT OF THE AIR FORCE

AFIA/INIS
ATTN: AFIA/INKD

AFIS/INT
ATTN: INT

AIR UNIVERSITY
ATTN: STRATEGIC STUDIES

AIR UNIVERSITY LIBRARY
ATTN: AUJL-LSE
ATTN: LIBRARY

ASSISTANT CHIEF OF STAFF
2 CYS ATTN: AF/SAMI

ASSISTANT CHIEF OF THE AIR FORCE
ATTN: SAF/ALR

DEPUTY CHIEF OF STAFF/XOX
ATTN: AFXOOS
ATTN: AFXOXFM

FOREIGN TECHNOLOGY DIVISION, AFSC
ATTN: CCN
ATTN: SDA

SPACE DIVISION/YH
ATTN: YH

STRATEGIC AIR COMMAND/ADWN
ATTN: ADWN

STRATEGIC AIR COMMAND/SPD
ATTN: SPD

STRATEGIC AIR COMMAND/STIC
ATTN: 544 SIW/DI (STIC)

STRATEGIC AIR COMMAND/XOXO
ATTN: XOXO

STRATEGIC AIR COMMAND/XPX
ATTN: XPZ

STRATEGIC AIR COMMAND/XRFS
ATTN: XRFS

TACTICAL AIR COMMAND/XPSC
ATTN: TAC/DOA

U S AIR FORCE ACADEMY
ATTN: LIBRARY
ATTN: USAFA/SP

USAF SCHOOL OF AEROSPACE MEDICINE
ATTN: RADIATION SCIENCES DIV

WEAPONS LABORATORY
ATTN: WL/SUL

DEPARTMENT OF ENERGY

LAWRENCE LIVERMORE NATIONAL LAB
ATTN: PAUL T HERMAN
ATTN: Z DIVISION LIBRARY

LOS ALAMOS NATIONAL LABORATORY
ATTN: REPORT LIBRARY
ATTN: T DOWLER

SANDIA NATIONAL LABORATORIES
ATTN: DIV 9013, W LING
ATTN: TECH LIB

OTHER GOVERNMENT

CENTRAL INTELLIGENCE AGENCY

ATTN: COUNTER-TERRORIST GP
ATTN: DIRECTOR OF SECURITY
ATTN: MEDICAL SERVICES
ATTN: NIO-T
ATTN: N10 - STRATEGIC SYS
ATTN: R & D SUBCOMMITTEE
ATTN: TECH LIBRARY

FEDERAL BUREAU OF INVEST ACADEMY

ATTN: BEHAVIORAL RSCH UNIT
ATTN: LIBRARY

FEDERAL EMERGENCY MANAGEMENT AGENCY

ATTN: ASST ASSOC DIR FOR RSCH
ATTN: CIVIL SECURITY DIVISION
ATTN: G ORRELL NP-CP
ATTN: OFC OF CIVIL DEFENSE

U S DEPARTMENT OF STATE

ATTN: PM/STM

U S NUCLEAR REGULATORY COMMISSION

ATTN: DIR DIV OF SAFEGUARDS
ATTN: S YANIV

DEPARTMENT OF DEFENSE CONTRACTORS

ADVANCED RESEARCH & APPLICATIONS CORP

ATTN: SECURITY OFFICE

AEROSPACE CORP

ATTN: LIBRARY ACQUISITION

BDM INTERNATIONAL INC

ATTN: C SOMERS
ATTN: J BRADDOCK

DATA MEMORY SYSTEMS, INC

ATTN: T DUPUY

KAMAN SCIENCES CORP

ATTN: DASAC
ATTN: E CONRAD

KAMAN SCIENCES CORPORATION

ATTN: DASAC

PACIFIC-SIERRA RESEARCH CORP

2 CYS ATTN: A LAUPA
2 CYS ATTN: G ANNO
ATTN: H BRODE

PACIFIC-SIERRA RESEARCH CORP

ATTN: D GORMLEY
2 CYS ATTN: G MCCLELLAN

R & D ASSOCIATES

ATTN: C McDONALD
2 CYS ATTN: DOCUMENT CONTROL

R & D ASSOCIATES

ATTN: J THOMPSON

SCIENCE APPLICATIONS INTL CORP

ATTN: DOCUMENT CONTROL
ATTN: E SWICK, MS 33
ATTN: J MARTIN
ATTN: J WARNER
ATTN: L HUNT
ATTN: R J BEYSTER

SCIENCE APPLICATIONS INTL CORP

ATTN: B BENNETT
ATTN: DOCUMENT CONTROL
ATTN: J FOSTER
ATTN: J PETERS
ATTN: L GOURE
ATTN: M FINEBURG
ATTN: W LAYSON

SCIENCE APPLICATIONS INTL CORP

ATTN: R CRAVER

SCIENCE APPLICATIONS INTL CORP

ATTN: JOHN A SHANNON

SYSTEMS RESEARCH & APPLICATION CORP

ATTN: R STEELE
ATTN: S GREENSTEIN