

AD.

TECHNICAL REPORT NATICK/TR-91/022

> INACTIVATION OF *GIARDIA* CYSTS BY IODINE WITH SPECIAL REFERENCE TO GLOBALINE: A REVIEW

> > EDMUND M. POWERS

April 1991

Final Report April 1989 - April 1991

Approved for public release, distribution unlimited.

US ARMY NATICK RESEARCH, DEVELOPMENT AND ENGINEERING CENTER NATICK, MA 01760-5000

SOLDIER SCIENCE DIRECTORATE

DISCLAIMERS

The findings contained in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

DESTRUCTION NOTICE

For Classified Documents:

Follow the procedures in DoD 5200.22-M, Industrial Security Manual, Section II-19 or DoD 5200.1-R, Information Security Program Regulation, Chapter IX.

For Unclassified/Limited Distribution Documents:

Destroy by any method that prevents disclosure of contents or reconstruction of the document.

Table memory bases bases for the second and a second bases of the second bases of t	REPORT	DOCUMENTATION P	AGE	Form Approved OMB No. 0704-0188						
1. AGENCY USE ONLY (Leave block) 2. REPORT DATE 3. REPORT TOPE Final Report April 1990 1. THE AND SUBTICE 1. REPORT TO ATE FINAL REPORT APRIL 1990 FINAL REPORT APRIL 1990 1. THE AND SUBTICE CLARDIA CYSTS BY IODINE WITH SPECIAL F. HUNDER MUMBERS REFERENCE TO GLOBALINE: A REVIEW 5. HUNDER MUMBERS F. AUTHOR(S) AUTHOR(S) E. HIPPORTMEC ORGANIZATION NAME(S) AND ADDRESS(ES) EDMURD M. POWERS 0. SPONSORING CORGANIZATION NAME(S) AND ADDRESS(ES) E. PIPPORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) IN. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SUPPLEMENTARY NOTES 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SUPPLEMENTARY NOTES 11. SUPPLEMENTARY NOTES 11. SUPPLEMENTARY NOTES 12. DISTRIBUTION AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE 13. ABSTRACT (MAXIMUM 200 MORES) 11. SUPPLEMENTARY NOTES 14. SUPPLEMENTARY NOTES 11. SUPPLEMENTARY NOTES 15. ADSTRACT (MAXIMUM 200 MORES) 11. SUPPLEMENTARY NOTES 15. ADSTRACT (MAXIMUM 200 MORES) 11. SUPPLEMENTARY NOTES 15. ADSTRACT (MAXIMUM 20	Public reporting burden for this collection o gathering and maintaining the data needed collection 31 information, including suggest Davis High way, Suite 1204, Arlington, VA 22	f information is estimated to average 1 hour pe and completing and reviewing the collection of ons for reducing this burden. to Washington He 202-4302, and to the Office of Management an	r response, including the time for review information. Send comments regardin adquarters Services, Directorate for Info I Budget, Paperwork Reduction Project (ving instructions, searching existing data sources, g this burden estimate or any other aspect of this ormation Operations and Reports, 1215 Jefferson 0704-0188), Washington, DC 20503.						
1. THE AND SUBTRIE SUBMONS NUMBERS INACTIVATION OF GLARDIA CYSTS BY IODINE WITH SPECIAL REFERENCE TO GLOBALINE: A REVIEW S. FUNDME NUMBERS F. AUTHOR(S) E. 612786 EDMUND M. POWERS R. Code: CUS 1211 T. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) B. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US ARMY NATICK RDAE CENTER BIOLOGICAL SCIENCES DIVISION, SSD ATTM: STRNC-THM NATICK/TR-91/022 NATICK, MA. 01760 NATICK, MA. 01760 3. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) ID. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SUPPLEMENTARY NOTES ID. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 12. DISTRIBUTION AVAILABILITY STATEMENT I2b. DISTRIBUTION CODE 13. ABSTRACT (MARMUM-200 WORGS) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 14. ABSTRACT (MARMUM-200 WORGS) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 15. ALBORNE ALBORNE / MONITORING AGENCY NAME(S) AND ADDRESS(ES) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 14. SUPPLEMENTARY NOTES ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 13. ABSTRACT (MARMUM-200 WORGS) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) ID. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) </td <td>1. AGENCY USE ONLY (Leave b</td> <td>lank) 2. REPORT DATE 8 April 1991</td> <td>3. REPORT TYPE AND C Final Report: F</td> <td>pares covered pril 1989 to April 1990</td>	1. AGENCY USE ONLY (Leave b	lank) 2. REPORT DATE 8 April 1991	3. REPORT TYPE AND C Final Report: F	pares covered pril 1989 to April 1990						
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) B. PERFORMING ORGANIZATION REPORT NUMBER US ARMY NATICK RDAE CENTER BIOLOGICAL SCIENCES DIVISION, SSD ATTM: STRNC-YMM NATICK, MA. 01760 NATICK/TR-91/022 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SUPPLEMENTARY NOTES 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION CODE 13. ABSTRACT (Maximum 200 words) 13. ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying Giardia especially in cold (3 to 15°C) water. Crysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodinn tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 1007 destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Clobaline tablets to destroy Giardia is not presently known. Based on the information presented in this review. Wo changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS 15. SUMMER OF PAGES 17. SUBJECT TERMS 10. SUBJECT TERMS 18. SUBJECT TERMS 1	4. TITLE AND SUBTITLE INACTIVATION OF GIA REFERENCE TO GLOBAL 6. AUTHOR(S) EDMUND M. POWERS	RDIA CYSTS BY IODINE W INE: A REVIEW	TH SPECIAL 5.	FUNDING NUMBERS PE: 612786 TA: 14 WU: BOO AG Code: CUS 1211						
US ARMY NATICK RDALE CENTER BIOLOGICAL SCIENCES DIVISION, SSD ATTM: STRNC-YHM NATICK, MA. 01760 NATICK/TR-91/022 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SUPPLEMENTARY NOTES 11. SUPPLEMENTARY NOTES 12b. DISTRIBUTION AVAILABILITY STATEMENT 12a. DISTRIBUTION AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE 13. ABSTRACT (Maximum 200 words) 11. SUPPLEMENTARY NOTES 13. ABSTRACT (Maximum 200 words) 11. SUPPLEMENTARY NOTES 14. SUBJECT (Maximum 200 words) 11. SUPPLEMENTARY NOTES 15. G. SO minutes at 25°C and 25 minutes at 45°C for 1002 destruction of Giardia egets at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for 1002 destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for 1002 destruction of Giardia cysts at residual iodine are recommended: (1) reformulate Clobaline tablets to destroy Giardia is not presently known. Based on the information presented in this review. Two changes for Globaline are recommended: (1) reformulate Clobaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS CLARDIA WATER TREATMENT TABLETS TORAGE 15. NUMBER OF PAGES 26 15. WATER PUBLIFICATION THERMAL STABILITY 15. SEUNITY CLASSIFICATION OF TREAS DESTACT UNCLASSIFI	7. PERFORMING ORGANIZATION	NAME(S) AND ADDRESS(ES)		PERFORMING ORGANIZATION						
3. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY REPORT NUMBER 11. SUPPLEMENTARY NOTES 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION AVAILABILITY STATEMENT 12. DISTRIBUTION CODE 13. ABSTRACT (Maximum 200 words) 13. ABSTRACT (Maximum 200 words) 13. DISTRIBUTION 12. DISTRIBUTION CODE 13. ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying Giardia especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodine tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for IOX destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy Giardia is not presently known. Based on the information presented in this review, two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS 15. NUMBER OF PAGES 14. SUBJECT TERMS 26 14. SUBJECT TERMS 15. SECURITY CLASSIFICATION OF ABSTRACT OF AB	US ARMY NATICK RD&E BIOLOGICAL SCIENCES ATTN: STRNC-YMM NATICK, MA. 01760	CENTER DIVISION, SSD		NATICK/TR-91/022						
11. SUPPLEMENTARY NOTES 12a. DISTRIBUTION AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION 13. ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying Giardia especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodint tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 100% destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy Giardia is not presently known. Based on the information presented in this review. Two changes for Globaline are recommended (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS CIARDIA WATER TREATMENT TABLETS GLARDIA WATER TREATMENT TABLETS STORAGE 14. SUBJECT TERMS COLD WATER CIARDIA WATER TREATMENT TABLETS STORAGE 15. NUMBER OF PAGES 26 IODINE GLOBALINE CIARDIA WATER PURIFICATION THERMAL STABILITY 17. SECURITY CLASSIFICATION 18. SECURITY CLASSIFICATION OF ABSTRACT 17. SECURITY CLASSIFICATION 19. SECURITY CLASSIFICATION OF ABSTRACT 17. SECURITY CLASSIFIED U	9. SPONSORING / MONITORING A	GENCY NAME(S) AND ADDRESS(E	5) 10	SPONSORING/MONITORING AGENCY REPORT NUMBER						
12a. DISTRIBUTION AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE APPROVED FOR PUBLIC RELEASE: DISTRIBUTION 12b. DISTRIBUTION CODE 13. ABSTRACT (Maximum 200 words) 13. ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying Giardia especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodine tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 100% destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy Giardia is not presently known. Based on the information presented in this review. Two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS CONLD WATER CJARDIA WATER TREATMENT TABLETS STORAGE 14. SUBJECT TERMS COLD WATER CJARDIA WATER TREATMENT TABLETS STORAGE 15. NUMBER OF PAGES 26 IODINE GLOBALINE CONLD WATER CJARDIA WATER TREATMENT TABLETS STORAGE 16. PRICE CODE 26 IODINE GLOBALINE CONLD WATER CJARDIA UNCLASSIFIED 19. SECURITY CLASSIFICATION OF ABSTRACT	11. SUPPLEMENTARY NOTES	<u></u>	, , L							
APPROVED FOR PUBLIC RELEASE: DISTRIBUTION 13 ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying <u>Giardia</u> especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodine tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 100% destruction of <u>Giardia</u> cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy <u>Giardia</u> is not presently known. Based on the information presented in this review. two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14 SUBJECT TERMS 15. NUMBER OF PAGES CIARDIA WATER TREATMENT TABLETS STORAGE 17. SECURITY CLASSIFICATION THERMAL STABILITY 15. NUMBER OF PAGES 17. SECURITY CLASSIFICATION OF ABSTRACT OF ABSTRACT OF ARDOR 19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	12a. DISTRIBUTION AVAILABILIT	STATEMENT	[12	b. DISTRIBUTION CODE						
13. ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying Giardia especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodine tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 100% destruction of Giardia cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy Giardia is not presently known. Based on the information presented in this review, two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes. 14. SUBJECT TERMS COLD WATER 15. NUMBER OF PAGES 26 16. PRICE TERMS 15. NUMBER OF PAGES 17. SECURITY CLASSIFICATION OF THIS PAGE 19. SECURITY CLASSIFICATION OF ABSTRACT 17. SECURITY CLASSIFICATION OF THIS PAGE 19. SECURITY CLASSIFICATION OF ABSTRACT 18. TORCASIFIED UNCLASSIFIED 19. SECURITY CLASSIFICATION OF ABSTRACT 10. UNCLASSIFIED	APPROVED FOR PUBLIC	RELEASE: DISTRIBUTION								
14. SUBJECT TERMS 15. NUMBER OF PAGES GIARDIA WATER TREATMENT TABLETS STORAGE 26 IODINE GLOBALINE COLD WATER 16. PRICE CODE CYSTS WATER PURIFICATION THERMAL STABILITY 16. PRICE CODE 17. SECURITY CLASSIFICATION OF REPORT 18. SECURITY CLASSIFICATION OF THIS PAGE 19. SECURITY CLASSIFICATION OF ABSTRACT 20. LIMITATION OF ABSTRACT UNCLASSIFIED UNCLASSIFIED UNCLASSIFIED 20. LIMITATION OF ABSTRACT	13 ABSTRACT (Maximum 200 words) This review of the literature indicates that iodine is not completely effective in destroying <u>Giardia</u> especially in cold (3 to 15°C) water. Cysticidal efficacy of iodine at a given concentration and temperature depends on contact time. The iodine tablet, Globaline, used by U.S. Military Forces since 1952, required 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C for 100% destruction of <u>Giardia</u> cysts at residual iodine concentration of 10, 8, and 6 mg per liter, respectively. The time required at temperatures below 15°C for Globaline tablets to destroy <u>Giardia</u> is not presently known. Based on the information presented in this review, two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes.									
CYSTS WATER PURIFICATION THERMAL STABILITY 17. SECURITY CLASSIFICATION OF REPORT 18. SECURITY CLASSIFICATION OF THIS PAGE 19. SECURITY CLASSIFICATION OF ABSTRACT 20. LIMITATION OF ABSTRACT UNCLASSIFIED UNCLASSIFIED UNCLASSIFIED 20. LIMITATION OF ABSTRACT	14. SUBJECT TERMS GIARDIA WATER TREAT IODINE GLOBALINE	MENT TABLETS COLD WATER	STORAGE	15. NUMBER OF PAGES 26 16. PRICE CODE						
17. SECURITY CLASSIFICATION OF REPORT 18. SECURITY CLASSIFICATION OF THIS PAGE 19. SECURITY CLASSIFICATION OF ABSTRACT 20. LIMITATION OF ABSTRACT UNCLASSIFIED UNCLASSIFIED UNCLASSIFIED 20. LIMITATION OF ABSTRACT	CYSTS WATER PURIF	ICATION THERMAL STABIL								
UNCLASSIFIED UNCLASSIFIED UNCLASSIFIED	17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATI OF ABSTRACT	ON 20. LIMITATION OF ABSTRACT						
	UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED							

Prescribed by ANSI Std Z39-18 298-102

. Ĭ

TABLE OF CONTENTS

Pac	<u>je</u>
List of Tables iv	7
Preface	J
Introduction 1	L
Cysticidal efficacy of different forms of iodine 2	2
Influence of contact time and forms of iodine on destruction of <u>Giardia</u>	2
Globaline 2	2
Saturated idine 3	3
Elemental icdine 3	3
Tincture of iodine 3	3
Chemical and physical properties of Globaline	3
Effect of pH and titratable iodine	}
Thermal stability of Globaline 4	ļ
Effect of temperature on dissolution time 5	5
Resistance of Globaline to humidity	;
Discussion	5
References 8	}
Appendix 1	.0
Hard organic water (HOW) suspending medium 1	0

LIST OF TABLES

Table		Page
1.	Summary of comparative contact times and concentrations for different forms of iodine to destroy <u>Giardia</u> cysts	11
2.	Destruction of protozoa cysts by Globaline at various temperatures	12
3.	Destruction of <u>Giardia</u> cysts (1000/mL) by saturated iodine in clear and cloudy water at 3°C and 20°C	13
4.	Destruction of protozoa cysts by elemental iodine	14
5.	Destruction of <u>Giardia</u> cysts (1000/mL) by two percent tincture of iodine in clear and cloudy water at 3°C and 20°C	15
6.	Effect of pH and concentration on hydrolysis of iodin in aqueous solution	e 16
7.	Residual iodine released by Globaline in water at 25°C	17
8.	Thermal stability of Globaline tablets stored at 60°C (140°F)	. 18
9.	Loss of iodine from Globaline tablets in different	. 19

PREFACE

Protocols for treatment of canteen water with iodine tablets, called $Globaline^{(R)}$, are presently based on obtaining the destruction of a resistant waterborne protozoan known as <u>Giardia lamblia</u>. Current procedures for treatment of field water with Globaline, including contact time are presented in Military Specification MIL-W-283H (11) and field manuals such as FM 21-10 (5).

This review summarizes current knowledge about the efficacy of iodine for destruction of <u>Giardia lamblia</u> with special reference to Globaline iodine tablets in cold water. In addition, this report updates and reaffirms information on the use of iodine tablets for the purification of field water and suggests revisions to relevant specifications and manuals. The text can serve as a basis for instructing soldiers on the proper treatment of field water, if necessary. Research investigators may use this report as a ready reference and guide for ongoing research, such as the development of other water purification tablets, development of compatible beverage powders or for the identification of additional areas for research.

The literature reviews include government contract reports, special studies and journal articles dating from 1945 through 1990. Results from the literature pertaining to iodine, particularly Globaline, are summarized in the following tables and appropriate references are listed.

This research was undertaken during the period April 1989 to April 1991. The funding was Program Element 612786, Task 14, Work Unit B00.

The citation of trade names in this report does not constitute an official endorsement or approval of the use of a product.

9

INACTIVATION OF <u>GIARDIA</u> CYSTS BY IODINE WITH SPECIAL REFERENCE TO GLOBALINE : A REVIEW

INTRODUCTION

The Globaline iodine tablet (11) was developed for the U.S. Armed Forces during World War II (6) to replace the Halazone Chlorine tablet, which had been used since World War I for disinfecting canteen water (14). A research and development program was sponsored by the Army Quartermaster General and conducted by Harvard University investigators (6). To overcome the inherent limitations of Halazone which affected its efficacy, the Harvard investigators proposed the following criteria for the ideal water purification tablet for field use (6, 9, 14). The tablet should:

- 1. Be effective as a single tablet and be simple to use.
- 2. Dissolve and liberate its active ingredient rapidly.
- 3. Destroy all waterborne pathogens and be effective in all kinds of water.
- 4. Provide treated water that is acceptable and nontoxic.
- 5. Be stable during storage and actual use.

An intensive search was undertaken by the Harvard investigators (6) for a compound that would satisfy these criteria. After screening more than 57 iodine - and chlorine - releasing compounds, Globaline was selected as the tablet that best satisfied military requirements (14). It was first issued to American soldiers in 1952.

Globaline (M1L-W-283H) is composed of tetraglycine hydroperiodide[(18) active ingredient], sodium acid pyrophosphate and talc (9, 11, 14). One tablet dissolves in a liter of water to yield from 7.6-9.0 ppm titratable iodine(11).

To purify field water the soldier is instructed (5, 11) to add two Globaline tablets to a full canteen of water (1 quart or 1.1 liter), wait 5 minutes for the tablets to dissolve, shake the water for an indeterminate time period and wait an additional 30 minutes. No reference is made to water temperature. Contradictory instructions on labels of bottled Globaline tablets manufactured as late as January, 1989, instructed the soldier to wait for only 20 minutes after shaking.

It is generally known that the germicidal and cysticidal activity of disinfectants is slowed by low temperatures (3, 4, 6, 7, 8, 9, 13, 15, 16, 17, 19). Therefore, there is great concern that two Globaline tablets may not effectively purify cold water (5°C) by destroying <u>Giardia</u> cysts within the recommended 20 to 35 minutes. <u>Giardia</u> represents a known hazard in cold water streams, such as those found in the United States, from New Hampshire to Washington (7).

The use of chlorine and iodine in water treatment is universally accepted and the effectiveness of both compounds in destroying pathogenic bacteria in public water supplies is well recognized. Accordingly, this report does not address the voluminous literature concerning bacteria. However, a review of the literature revealed that there is a paucity of information regarding the efficacy of iodine against <u>Giardia</u> cysts and even less information on the efficacy of Globaline, at temperatures below 15°C. Guidelines do not exist for chemical treatment of public drinking water to destroy <u>Giardia</u>, because little is known of the effect of halogens on <u>Giardia</u> cyst viability. Furthermore, guidelines used for Globaline were originally based on killing <u>Entamoeba</u> <u>histolytica</u> cysts (3, 6), not <u>Giardia</u>. Until 1980, none of the chemical methods available for disinfecting drinking water had ever been tested for destruction of <u>Giardia</u> cysts (7). Because protozoan cysts are the most resistant of the waterborne pathogens to disinfectants, treatments effective against them will also be effective against bacteria and other microorganisms.

It was recognized in 1986 (personal communication, Morris Rogers, Natick, memorandum for the record, subject: <u>Giardia lamblia</u> in military water supplies, 25 March 1986) that Globaline (iodine) had serious limitations in cold water regarding the destruction of <u>Giardia</u>. This conclusion was based on work presented by Jarroll (7).

Therefore, this review addresses the specific questions concerning the ability of Globaline to destroy <u>Giardia</u> cysts in cold water within the recommended time period.

CYSTICIDAL EFFICACY OF DIFFERENT FORMS OF IODINE

Influence of contact time and form of iodine on destruction of Giardia

Table 1 summarizes comparative data for the various forms of iodine, which are presented in more detail in Tables 2 thru 5. Although the test waters differed, the minimal contact times required for comparable residuals of various forms of iodine to destroy Giardia at various water temperatures are presented. Only Globaline achieved 100% destruction of Giardia, but it required 60 minutes contact time at 15°C and at least 10 mg/L of residual iodine (15). At 25°C only 30 minutes was required even though the residual iodine was less. The lowest temperature studied with Globaline was 15°C. Less residual iodine was required as temperature increased. Only 99% kill times were determined in the studies with elemental iodine (16,17). Contact time and the residual elemental iodine required decreased as temperature increased. Saturated iodine (8) and tincture of iodine (7) were more effective at 20°C than at 3°C (see tables 3 and 5), but did not completely kill Giardia even after 30 to 40 minutes. Iodine residuals were lower than they were in studies with elemental iodine and Globaline. In the studies with all four forms of iodine, it is clear that longer contact times and higher iodine residuals are required in cold water (< 25°C) to completely destroy Giardia. Test waters included hard organic water (HOW: see appendix), deionized, distilled, tap, and river water.

Globaline

Table 2 presents the efficacy of Globaline for destruction of <u>Giardia</u> and <u>E. histolytica</u> cysts at temperatures ranging from 3 to 45°C (6, 7, 9, 10, 13, 15). Cyst densities were 1000 <u>Giardia</u> per mL (6) and 60 <u>E. histolytica</u> per mL (9,13). It is evident that 20 or 30 minutes was not sufficient time to destroy <u>Giardia</u> at 3 to 15°C even with two tablets in natural river water or HOW (7,15). A 100% mortality was achieved in HOW only after 60 minutes at 15°C, 30 minutes at 25°C and 25 minutes at 45°C (15). Residual iodine levels were 10 ppm, 8 ppm and 5.7 ppm, respectively. The HOW was a worst case water

2

that was made up to have a high iodine demand, high pH (pH 9), and high total dissolved solids (>1500 mg/L). Cysticidal activity was most dependent on contact time since less residual iodine was required as the temperature of the water was increased (7). Chang (4) showed that cysticidal action of halogens decreased as temperature decreased. The contact time required to destroy Giardia at temperatures between 3 and 15°C and between 15 and 25°C is not known. Activity of Globaline appeared to be independent of pH over a range of pH 4 to 8. <u>Giardia</u> was more resistant to Globaline than <u>E</u>. <u>histolytica</u>.

Saturated Iodine

Table 3 shows that saturated iodine did not completely destroy <u>Giardia</u> cysts (1000/mL) at 3°C even after 40 minutes contact time whether the water was clear or cloudy (8). As indicated by the data, efficacy could be improved by increasing the contact time of the residual iodine. At 20°C in both clear and cloudy water, 99.8% of the <u>Giardia</u> cysts were destroyed after only 15 minutes even though residual iodine levels were lower than at 3°C (7, 8). As with Globaline, pH appeared not to be a factor at the ranges shown.

Elemental Iodine

Studies with elemental (aqueous) iodine are presented in Table 4. Higher concentrations of elemental iodine were required to destroy <u>Giardia</u> cysts (10,000/mL) at 5°C than at 15 to 25°C (16, 17) in distilled water. The complete destruction of <u>E</u>. <u>histolytica</u> cysts (30 to 60/mL) in sewage diluted tenfold in Cambridge tap water (SCIW) required twice as much contact time (20 minutes) at 5°C than at 23°C, at the same iodine concentration (2). At low temperatures (0 to 5°C), a contact time of 10 minutes was not sufficient(3). In studies with <u>E</u>. <u>histolytica</u>, the cysticidal activity of iodine was independent of pH in the range between pH 3 to 7 (3). High germicidal activity was maintained over the pH range of 3 to 8 in the presence of a variety of water contaminants (3, 19). Having less dependence on pH was considered to be one of the advantages of using iodine to disinfect water. Lower residuals were required to destroy <u>Giardia</u> as the contact time increased (16, 17).

<u>Tincture of Iodine</u>

Table 5 shows that inactivation of <u>Giardia</u> cysts (1000/mL) by 2% tincture of iodine was more effective at 20°C than at 3°C in clear (deionized) water (7, 10). The failure of tincture of iodine at 3°C in clear water may have been due to insufficient contact time, since lower residuals inactivated <u>Giardia</u> cysts at 20°C (7, 10). Complete inactivation of <u>Giardia</u> was not achieved at the contact time and residual iodine levels studied.

CHEMICAL AND PHYSICAL PROPERTIES OF GLOBALINE

Effect of pH and titratable Iodine

When elemental iodine(I_2) is dissolved in water, the I_2 may remain unchanged or it may hydrolyze into five different species depending on pH and the initial concentration of titratable iodine (1,4,9,19). Of the five species produced during hydrolysis of I_2 , only I_2 and hypoiodous acid (HOI) are important for disinfection. Both are known to be powerful disinfectants (4, 9, 13, 19). The I₂ is more cysticidal (4,9) and sporicidal (19) than HOI, but the viricidal efficacy of HOI is 44 times greater than I₂ (4, 9). The sporicidal action (99% killing time of <u>Bacillus metiens</u> spores) was not affected at pH values between 6 and 7.5, where sufficient I₂ was present, but it decreased at pH 8 and above as the HOI fraction increased (19). Therefore to be effective as a broad spectrum disinfectant, both I₂ and HOI should be present in solution (4, 9, 13).

Table 6 shows that an insignificant amount of I_2 is hydrolyzed below pH 6 (1, 4, 9). All the titratable iodine exists as I_2 below pH 6 when the total titratable iodine is 0.5 ppm or greater. The same is true below pH 7.0 when the total titratable iodine is 5 ppm or greater. At pH 7.0, the amount of I₂ hydrolyzed to HOI ranged from 48% at 0.5 ppm to 10% at 5.0 ppm titratable I2. The percent I2 and HOI present were about equal at pH 7.0 in a dilute iodine solution (0.5 ppm) and also at pH 8.0, at 5.0 ppm titratable I2. The decomposition of HOI to form iodate is slow and depends on the buffering system and the time of standing (4). Chang (3) demonstrated that the concentration of I_2 required for effective disinfection of \underline{E} . histolytica cysts was about 5 ppm at 23°C for 10 minutes in clear water. Therefore, to allow for the iodine demands of natural waters, it was concluded that a suitable dose of iodine for emergency disinfection was about 8 ppm (3, 4). Water with I2 demands greater than 4 ppm required twice the dose of I₂ and waters near 0°C required an increased treatment time of 20 minutes (3, 4). Consequently, Globaline tablets were formulated to provide 8 ppm of titratable I_2 per liter of water (6) which gave a sufficient I_2 residual to destroy 30 cysts of <u>E</u>. <u>histolytica</u> within 10 minutes in most natural waters at 23°C (3, 4).

The effectiveness of Globaline tablets for disinfecting cysts of \underline{E} . <u>histolytica</u> was presented by O'Connor (13) and Kapoor (9). As shown above in Table 2, one Globaline tablet at 10°C lowered the initial pH of tap water to pH 6.6, which left a residual of 6.9 ppm of I₂. Since Chang (4) showed that this resulted in only about 5% HOI (Table 6), the final HOI concentration would be about 0.35 ppm (6.9 x 0.05). This small amount of HOI may not be sufficient to be viricidal. Therefore, the high concentration of titratable I₂ and the use of acidic buffer result in a high cysticidal but low viricidal efficiency (9, 13).

Thermal Stability of Globaline

Military criteria for water purification tablets require that they be chemically stable during storage for five years as packaged. The tablets must also withstand extremes of temperature encounted in different parts of the world and in storage warehouses.

The stabilty of Globaline tablets at ambient temperatures over a six year period stored in the original, wax - sealed brown glass bottle was determined by measuring residual iodine levels for tablets produced in 1983 and in 1989, as shown in Table 7 (Powers, E. M., 1989, unpublished). Each of ten tablets was dissolved in one liter of halogen - free deionized water. The average tablet produced between 7.1 to 8.23 mg/L of residual iodine. Therefore, two tablets added to a liter of water will provide sufficient iodine residuals to destroy <u>Giardia</u> at the contact times indicated in Table 1. Globaline tablets exhibited excellent storage stability and product reliability since residual iodine levels measured for tablets in 1983 were approximately the same as tablets produced in 1989.

Table 8 shows the thermal stability of Globaline tablets during accelerated storage studies at $60^{\circ}C$ ($140^{\circ}F$) and ambient humidity. Tablets produced in 1989 were exposed in the original 20 x 50 mm screw - capped, sealed bottles, in which they are packaged 50 tablets to a bottle, as well as exposed to the air in open petri dishes (Powers, E.M., unpublished). Two tablets from each container were removed periodically over a 30 day period. Each tablet was dissolved in one liter of deionized water and after a 30 minute waiting period the total residual iodine was titrated amperometrically (Wallace and Tiernan Titrator). Results from the Harvard Study (6) in 1945, in which tablets were exposed to the air on watch glasses, are presented for comparison. Most of the iodine was retained over the 30 - day period by tablets stored in tightly capped bottles. However, when exposed to the air, tablets lost iodine rapidly after 7 days and 66% had dissipated after 25 days. Results in 1945 were similar (6).

In similar studies with powdered Globaline exposed to 60°C on open watch glasses, the Harvard Group (6) reported a 50% loss of iodine after 49 days and a 92% loss after 103 days. Therefore, bottles of Globaline tablets must be tightly sealed after removal of each tablet. Under these conditions, deterioration of Globaline will be slow even at elevated temperatures.

Effect of Temperature on Dissolution Time

Rapid solubility is an important criterion for acceptability of Globaline tablets. Field trials with soldiers indicated that they became impatient with agents that required a waiting period of more than 10 minutes (6, 9, 13). Therefore, since dissolving time must be added to the time required for germicidal activity to obtain a total treatment time (12), the germicide must be dispersed quickly and uniformly to be most effective and to achieve adequate treatment of water. Globaline has a solubility of 380 grams per liter of distilled water (5, 9, 13, 18).

In simulated field studies, a Globaline tablet was placed in a liter of tap water at 23°C (6, 9, 13). The flask containing the water was inverted end-over-end continuously until the tablet was dissolved. Dissolution times, observed visually, were as follows:

Temperature °C	Dissolution Time (seconds)
10	114
20	72
23	46
30	48
40	30

ņ

To be considered satisfactory for field use tablets must dissolve within one minute at 23°C (6, 12). As shown above, Globaline dissolved within 46 seconds at 23°C. Dissolution time becomes shorter as temperature is increased. It was also shown that storage of Globaline at 60°C and room humidity did not affect the dissolving properties (6).

Resistance of Globaline to Humidity

To determine the stability of Globaline in humid atmospheres, the Harvard Group conducted experiments at humidities encountered in jungle warfare (6). The Globaline tablets were placed on flat dishes in desiccator jars controlled at humidities of 100%, 79%, and 55%. Table 9 presents the results of these studies. The tablets were hyproscopic and unstable at 100% humidity, losing 63% of their iodine in only two days. They were completely dissolved after 4 days. At 79% humidity, tablets lost more than half of their iodine after only six days. Globaline tablets were much more stable at 55% humidity retaining 54.2% of the iodine after 111 days. In another study (6) of simulated field conditions, bottles of Globaline were placed in a control room held at 80 to 90% humidity and 26.7°C(80°F). During the day, bottles were opened for a minute and a tablet was withdrawn. None of the tablets showed an appreciable loss of active iodine after three weeks, and there was little variation between the bottles with or without cotton plugs. When compared to four other halogen compounds, Globaline was shown to be the most stable at normal (32%) relative humidity.

DISCUSSION

This review of the literature indicates that most procedures with Globaline and other iodine solutions did not achieve adequate or complete destruction of <u>Giardia</u> cysts, in cold water (7, 8, 10, 16, 17). The failure of these methods to inactivate <u>Giardia</u> appeared to be due to an insufficient contact time. It was shown that cysticidal and sporicidal action of halogens decreases as the temperature decreases (4, 19). Wyss and Strandskov (19), for example, showed that lowering the temperature 10°C increased the killing time of <u>Bacillus metiens</u> spores fourfold. Therefore, temperature must be considered when iodine is used for disinfection of drinking water.

Because disinfection follows a first - order chemical reaction (2, 3, 4, 16), there is an inverse relationship between concentration of iodine and contact time with microorganisms. This means that low concentrations of iodine will require longer periods of time to kill <u>Giardia</u> and other microorganisms. The disinfection reaction is also slowed by cold temperatures.

The development of the Globaline iodine tablet during World War II (6) was a major breakthrough in water purification and provided the soldier with an effective method of producing potable water under emergency conditions. It has satisfied most of the requirements established for a water purification tablet and was a much more efficient disinfectant than the Halazone tablet which it replaced. Although the development of Globaline by Harvard scientists (6) was a tremendous effort, they operated under some basic assumptions that are no longer upheld: The first assumption was that cysts of E. <u>histolytica</u> were the most disinfectant resistant waterborne pathogen and should be used as the test organism. It is now known that <u>Giardia</u> cysts are much more resistant to disinfectants, including iodine, as shown in Tables 2 and 4. Consequently, guidelines presently used for Globaline (two tablets per liter) are based on the destruction of <u>Giardia</u>. The second assumption was that molecular iodine alone was the germicidal species and not products produced by hydrolysis of iodine. We know now that while I_2 is a very good cysticidal agent, it is not as viricidal as HOI (9, 13). Therefore, both I_2 and HOI must be present in solution to be cysticidal as well as viricidal. The concentration of both species is dependent on pH and total residual iodine.

The Harvard scientists (6) assumed that the destruction of viruses by disinfectants was of the same order of magnitude as that of most pathogenic nonsporulating bacteria. Thus, as suggested by O'Connor (13) and Kapoor (9), it would be useful to evaluate the viricidal power of Globaline tablets. Because sodium acid pyrophosphate buffer present in Globaline (11) lowers the pH, the amount of HOI liberated may not be sufficient to be viricidal (9, 13). Kapoor (9) suggested that Globaline could be improved by substituting an alkaline buffer (pH 8), which would yield almost equal percentages of I_2 and HOI (Table 6).

The cysticidal efficacy of Globaline at temperatures below 15° C has not been sufficiently studied and it is uncertain at the present time. Therefore, there is great concern that Globaline tablets may not effectively disinfect some waters encountered by American soldiers within the recommended 35 minute treatment time. Studies are now underway under contract from the U.S. Army Natick R D and E Center, that will provide efficacy data at 5°C.

Based on the information presented in this review, two changes for Globaline are recommended: (1) reformulate Globaline tablets with an alkaline buffer and (2) increase the treatment time for one liter of water from 35 minutes to 60 minutes.

References

1. Black, A.P., R.N. Kinman, W.C. Thomas Jr., G. Freund and E.D. Bird. 1965. Use of iodine for disinfection. J. AM. Water Works Assoc. 57: 1401-1421.

Ą

2. Block, S.S. 1985. Disinfection, sterilization, and preservation. Lea and Febiger (Publishers), Philadelphia, Pa.

3. Chang, S.L. and J. C. Morris. 1953. Elemental iodine as a disinfectant for drinking water. Industrial and Engineering Chemistry. 45: 1009-1015.

4. Chang, S.L. 1958. The use of active iodine as a water disinfectant. J. Amer. Pharmaceutical Assoc. 47: 417-423.

5. Department of the Army. 1989. Unit field sanitation team. Field Manual FM 21-10-1. Headquarters, Department of the Army, Washington, D.C.

6. Harvard University. 1945. Disinfection of water and related substances. Final report to the Committee of Medical Research of the Office of Scientific Research and Development on the Investigations pursued from October 1, 1942 to December 31, 1945 under contract No. OEMcmr-251. Cambridge, MA.

7. Jarroll, E.L., Jr., A.K. Bingham and E.A. Meyer. 1980. <u>Giardia</u> cyst destruction: effectiveness of six small-quantity water disinfection methods. Am. J. Trop. Med. Hyg. 29:8-11.

8. Jarroll, E.L., Jr., A.K. Bingham, and E.A. Meyer. 1980. Inability of an iodination method to destroy completely <u>Giardia</u> cysts in cold water. Western Journal of Medicine. 32: 567-569.

9. Kapoor, S.K. 1969. A critique on disinfection of drinking water. Department of Civil Engineering, series no. 50. University of Illinois, Urbana, Illinois.

10. Meyer, E.A. 1981. Effect of halogens on <u>Giardia</u> cyst viability. Municipal Environmental Research Laboratory, Office of Research and Development, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268. EPA-600/52-81-174.

11. Military Specification, MIL-W-283H. 1987. Water purification tablet, iodine. Standardization Document Order Desk, 700 Robin Avenue, 4 D, Philadelphia, PA 19111 - 5094.

12. Morris, J.C., S.L. Chang, G.M. Fair, and G.H. Conant, Jr. 1953. Disinfection of drinking water under field conditions. Industrial and Engineering Chemistry. 45: 1013-1015.

13. O'Connor J.T. and S.K. Kapoor. 1970. Small quantity field disinfection. J. Am. Water Works Assoc. 62:80-84.

14. Rogers, M.R., J.J. Vitaliano, A.M. Kaplan, and E. Pillion. 1977. Military individual and small group water disinfecting systems: An assessment. Military Medicine 141:268-277. 15. Rogers, M. 1988. Cysticidal efficiency of iodine water purification tablets. Unpublished report to Fort Drum 12 May 88. Natick RD&E Center, STRNC - YMA, Natick, MA 01760.

16. Rubin, A.J. 1989. C.T. products for the inactivation of Giardia cysts by chlorine, chloramine, iodine, ozone and chlorine dioxide. Personal communication, Water Resources Center, Ohio State University, Columbus, Ohio.

17. Rubin, A.J. and Y.F.R. Chen. 1989. Inactivation of <u>Giardia muris</u> cysts by elemental iodine. Personal communication, Water Resources Center, Ohio State University, Columbus, Ohio.

18. Windholz, M. (ed.). 1983. The Merck Index, tenth edition: 9032. Merck and Co., Inc., Rahway, N.J.

19. Wyss, O. and F.B. Strandskov. 1945. The germicidal action of iodine. Arch. Biochem, 6: 261-267.

> This document reports research undertaken at the US Army Natick Research, Development and Engineering Center and has been assigned No. NATICK/TR-91/02.2 in the series of reports approved for publication.

APPENDIX

Hard Organic Water (HOW) Suspending Medium.

The final composition of the hard organic water was as follows:

NaHCO3	0.0042	М
KHCO3	0.0024	М
CaCl ₂	0.0019	М
MgCl ₂	0.0011	М

Tea infusion 0.15%

The sodium bicarbonate and potassium carbonate were made up as 100 x solutions (0.42 and 0.24 M, respectively). The calcium and magnesium chloride solution were made up as $1,000 \times 1.1$ m, respectively). One commercial tea bag was boiled in one liter of water for 30 minutes. All solutions were autoclaved and diluted before use.

Iodine	Temp. °C	Water	рН	Residual icdine (mg/L)	Contact time (minutes)	Percent mortality	Reference
Globaline	15	HOWb	9	10	60	100	15, Table 2
(2 tablets)	25	HOW	9	8	30	100	15, Table 2
	45	HOW	9	5.7	25	100	15, Table 2
	5	Distilled	7	22.0	24	>99	16,17, Table 4
Elemental	15	Distilled	7	11.4	21	>99	16,17, Table 4
	25	Distilled	7	11.2	6.3	>99	16,17, Table 4
Saturated	20	River	7	3.01	15-40	>99.8	7,8,10,Table 3
	20	Deionized	7.1	3.3	15-40	>99.8	7,8,10,Table 3
Tincture(2%)	3	River	7.4	4.1	30	>99.8	7,10,Table 5
	20	River	7.1	3	30	>99.8	7,10,Table 5

Table 1. Summary^a of comparative contact times and concentrations for different forms of iodine to destroy <u>Giardia</u> cysts.

^aData taken from following Tables 2 through 6. ^bHard organic water plus tea infusion (see Appendix for composition)

t i

Temp °C	Water	No. of tablets per liter	Contact time (minutes)	Residual I ₂ (mg/L)	Final pH	Percent mortality	Reference					
	<u>Giardia</u> (1000/mL)											
3 15 15 20 25 25 45 45 45 45	Clear ^b Cloudy ^C HOW HOW Clear Cloudy HOW HOW HOW HOW HOW HOW HOW	1 2 2 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	20 20 30 45 60 20 20 20 20 30 5 10 15 20 25	6.8 9.88 10.4 8.9 10.0 5.44 9.47 8.73 8.03 7.24 4.18 5.2 5.6 5.7	4.6 5.8 8.1 8.1 5.1 5.7 8.1 8.1 8.1 8.1 8.1 8.1 8.3 8.4	97.5 >99.8 99.34 99.98 100 >99.8 >99.8 99.94 100 99.6 99.7 99.8 99.98 99.98 99.98 100	7,10 7,10 15 15 15 7,10 7,10 15 15 15 15 15 15 15 15 15 15					
			Entamoeba	<u>histolytica</u> (60/mL)							
3 10 23 23 28	Tap Tap Tap Tea Infusion Tap	1 1 1 2 1	25 15 10 5 5	7.5 6.9 7.5 8.7 7.5	6.5 6.6 6.6 6.4 6.6	100 100 100 100 100	6,9,13 6,9,13 6,9,13 6,9,13 6,9,13 6,9,13					

Table 2. Destruction of protozoa cysts by Globaline^a at various temperatures.

^aTetraglycine hydroperiodide ^bClear - deionized, halogen free ^CCloudy - Willamette River water, Portland, Oregon ^dHard organic water plus tea infusion

Temp. C	Water condition ^a	Stock iodine solution (mL)	Contact time (minutes)	Residual I ₂ (mg/L)	Final pH	Percent mortality	Reference
3	clear	26	15	6.87	7.1	56.5	8
			20	7.09	5.7	96.5	7,10
			30	6.87	7.1	93.3	8
	,		40	6.87	7.1	95.9	8
1	cloudy	26	15	4.58	7.3	37.2	8
	-		20	3.22	7.1	77.3	7,10
			30	4.58	7.3	57.5	.8
			40	4.58	7.3	81.5	8
20	clear	13	15	3.15	6.0	>99.8	7,10
			15	3.29	7.1	>99.85	8
		<i>i</i>	30	3.29	7.1	>99.85	8
			40	3.29	7.1	>99.85	8
	cloudv	13	15	3.01	7.3	>99.85	8
	7		20	3.01	7.0	>99.84	7.10
		ta in a	30	3.01	7.3	99.85	8
			40	3.01	7.3	>99.85	8

Ł

Table 3. Destruction of <u>Giardia</u> cysts (1000/mL) by saturated iodine in clear and cloudy water at $3^{\circ}C$ and $20^{\circ}C$.

¢

ŝ

^aClear water was halogen free deionized water; cloudy water was from the Willamette River, Portland, Oregon.

Temp °C	Water	рH	Contact time (minutes)	Residual I ₂ (mg/L)	Protozoa	Percent mortality	Reference
5 16,17	DWb	7	4.7	60.0	GC	99	
	DŴ	7	10.4	36.4	G	99	16,17
	DW .	7	23.6	22.2	G	99	16,17
	SCIW	3–7	20	8.0	Ee	100	3
6	SCIW	8	5	14.1	E	100	4
15	DW	7	13.0	15.2	G	99	16,17
	DW	7	21.2	11.4	G	99	16,17
23	SCIW	3–7	10	8.0	E	100	3
25	DW	8	5	7.1	Е	100	4
	DW	5	3.8	9.2	G	99	16,17
	DW	5	10.0	5.9	G	99	16,17
	DW	5 💡	38.0	3.6	G	9 9	16,17
	DW	7	3.2	15.4	G	99	16,17
	DW	7	6.3	11.2	G	99	16,17
	DW	7	17.8	6.2	G	99	16 , 17
	DW	9	2.0	37.0	G	99	16 , 17
	DW	9	10.0	23.0	G	99	16,17
	DW	9	11.0	19.0	G	99	16,17
	DW	9	44.6	10.0	G	99	16,17

Table 4.	Destruction	of	protozoa	cvsts	bv	elemental	iodine ^a
70070 1 0		~-		~y~~~~	- L Y		

^aIodine dissolved in distilled water ^bDW - Distilled water ^CG - <u>Giardia</u> cysts, 10,000/mL ^dSCIW - 10% sewage in Cambridge tap water ^eE - <u>Entamoeba histolytica</u> cysts, 30 to 60/mL

Table 5. Destruction of <u>Giardia</u> cysts (1000/mL) by two percent tincture of iodine in clear and cloudy water at 3 and $20^{\circ}C$

Temp. °C	Water condition	2% Tincture of iodine (mL)	Contact time (minutes)	Residual I ₂ (mg/L)	Final pH	Percent mortality	Ref.
3	Clear ^a	0.25	30	3.37	5.8	74.6	7,10
	Cloudy ^D	0.5	30	4.08	7.4	99.8	7,10
20	Clear	0.25	30	3.22	5.8	99.8	7,10
	Cloudy	0.5	30	2.79	7.1	99.8	7,10

1

^aClear water was deionized, halogen free water. ^bCloudy water was obtained from the Willamette River at Portland, Oregon.

		Titratable iodine									
	0.5	5 ppm ^a	m ^a 1 ppm ^b		3 ppm ^b		5 ppm ^C				
рН	I ₂	HOI	1 ₂	HOI	1 ₂	HOI	1 ₂	HOI			
5	99	1	99	1	100	0	100	0			
6	90	10	95	5	98	2	98	2			
7	52	48	67	33	82	18	90	10			
8	12	88	20	80	40	60	50	50			

Table 6. Effect of pH and concentration on hydrolysis of iodine in aqueous solution

^aAdapted from reference 1 (Black, 1965) ^bAdapted from reference 4 (Chang, 1958) ^CAdapted from reference 9 (Kapoor, 1969)

			Residual I2(mg/L) ²		
	lear of manufacture	NO. tablets	Range	Average ³	
	1983	10	7.16 - 9.31	8.23	<u>.</u>
	1989	10	5.01 - 7.88	7.1	
<u></u>	1989	10	5.01 - 7.88	7,1	

Table 7. Residual iodine released by Globaline in water at $25^0 \mbox{C}^1$

¹Powers, E.M. 1989. U.S. Army Natick RD&E Center, Natick, MA 01760-5020.

²One tablet dissolved in 1 liter of iodine demand-free deionized water for 30 minutes.

á

³Average of 10 tablets.

	Average percent loss of residual iodine			
Days at 60°C	Screw capped bottles ^a	Open ^a Petri dish	Open watch glass ^b	
1	15	0.5	5	
2	17	8.5	2	
3	15	19	9	
7	22	29	37	
9	22		42	
25	27	66		
30	27	66		
	•			

Thermal stability of Globaline tablets stored at $60^{\circ}C$ (140°F) Table 8.

a Powers E.M, 1989 b Adapted from the Harvard study, 1945 (6)

		Percentage of iodine lost				
	Time in days	100% humidity	79% humidity	55% humidity		
	0	0	0	0		
	1	32.3	12.3	11.1		
	2	63	ND	13.9		
	3	84.6	9.6	ND		
	4	b	31.5	22.2		
	5	ND^{C}	38.4	19.4		
	6	ND	53.4	19.4		
	7	ND	ND	20.8		
:	100	ND	ND	54.1		
	111	ND	ND	54.2		

Table 9. Loss of iodine from Globaline tablets in different humidities at room temperature.^a

^aAdapted from reference 6 ^bTablet dissolved after 4 days ^CNot done