

AD-A246 665



2

TECHNICAL REPORT BRL-TR-3315

BRL

THE TENSOR EQUATION $AX + XA = G$

MIKE SCHEIDLER

FEBRUARY 1992

DTIC
ELECTE
MAR 02 1992
S B D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.

U.S. ARMY LABORATORY COMMAND

BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

92-04995



92 2 26 008

NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE February 1992	3. REPORT TYPE AND DATES COVERED Final March-July 1991		
4. TITLE AND SUBTITLE The Tensor Equation $AX + XA = G$		5. FUNDING NUMBERS PR: 1L161102AH43		
6. AUTHOR(S) Mike Scheidler				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Ballistic Research Laboratory ATTN: SLCBR-DD-T Aberdeen Proving Ground, MD 21005-5066		10. SPONSORING / MONITORING AGENCY REPORT NUMBER BRL-TR-3315		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) We study the second-order tensor equation $AX + XA = G$ for symmetric, positive-definite A and arbitrary G . Motivated by applications in the continuum mechanics literature, we also examine some special cases where G depends on A and another tensor H . For arbitrary dimensions, we establish relations between the solutions X for various forms of G . These results, together with Rivlin's identities for tensor polynomials in two variables, are applied in two and three dimensions to obtain explicit formulas for X in direct (component-free) notation. The results include new formulas as well as new derivations of previously known formulas. An application to the kinematics of rigid motions is considered.				
14. SUBJECT TERMS continuum mechanics, kinematics, strain rate, tensors, linear algebraic equations		15. NUMBER OF PAGES 31		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT SAR	

INTENTIONALLY LEFT BLANK.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENT	v
1. INTRODUCTION	1
2. ALGEBRAIC PRELIMINARIES	4
3. THE TENSORS L_A AND M_A	6
4. SOME USEFUL IDENTITIES	13
5. FORMULAS FOR THE THREE-DIMENSIONAL CASE	16
6. FORMULAS FOR THE TWO-DIMENSIONAL CASE	22
7. AN APPLICATION TO THE KINEMATICS OF RIGID MOTIONS	25
8. REFERENCES	27
DISTRIBUTION LIST	29

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Date	
Availability Codes	
Dist. Statement	
A-1	

INTENTIONALLY LEFT BLANK.

ACKNOWLEDGMENT

I would like to thank Joseph Santiago for his criticism of an earlier draft.

INTENTIONALLY LEFT BLANK.

1. INTRODUCTION

Tensor equations of the form

$$AX + XA = G \quad (1.1)$$

arise in a variety of problems in continuum mechanics; cf. Sidoroff (1978), Guo (1984), Hoger and Carlson (1984), Leonov (1976), Stickforth (1982–1983), Mehrabadi and Nemat-Nasser (1987), Dienes (1979), Stickforth and Wegener (1988), Hoger (1986), Nemat-Nasser (1990), Truesdell (in press), Wheeler (1990), and Scheidler (to be published) for applications to kinematics and constitutive modeling. Here the second-order tensors A and G are regarded as known, and we wish to solve (1.1) for the second-order tensor X . For the applications considered in the aforementioned references, A is symmetric and positive-definite. This restriction, which will be imposed here also, guarantees that a solution X of (1.1) exists and is unique. Indeed, relative to any principal basis for A , this solution is given by the simple component formula

$$X_{ij} = \frac{G_{ij}}{a_i + a_j}, \quad (1.2)$$

where a_i are the (necessarily positive) eigenvalues of A . Of course, to actually compute the components of X by means of (1.2), we must first solve for the eigenvalues and eigenvectors of A . This has motivated several authors to seek explicit formulas for the tensor X directly in terms of the tensors A and G . An example is the formula of Sidoroff (1978) and Guo (1984) for the case where G is skew and the underlying vector space is three-dimensional:

$$X = \frac{1}{I_A II_A - III_A} \left[(I_A^2 - II_A) G - (A^2 G + G A^2) \right]. \quad (1.3)$$

Here I_A , II_A , and III_A denote the principal invariants of A . Since equations of the form (1.3) are often said to be displayed in direct notation, we will refer to formulas of this type as *direct formulas* for X or *direct solutions* of (1.1).

Although the derivation of the component formula (1.2) is a trivial exercise, the derivation of direct formulas for X is a nontrivial problem. Furthermore, whereas the component formula (1.2) is valid for any finite-dimensional vector space, the direct formulas become more complex as the

dimension of the underlying vector space increases. For example, in the two-dimensional case the solution of (1.1) for skew G is

$$X = \frac{1}{I_A} G, \quad (1.4)$$

which is substantially simpler than its three-dimensional counterpart (1.3). The formula (1.4) is a special case of the second of two direct formulas obtained by Hoger and Carlson (1984) for arbitrary G in the two-dimensional case. For the three-dimensional case, Sidoroff (1978), Hoger and Carlson (1984), Leonov (1976) and Stickforth (1982–1983) have obtained direct solutions of (1.1) for arbitrary G . For the two-dimensional and three-dimensional cases, Mehrabadi and Nemat-Nasser (1987) obtained direct formulas for X when $G = AH - HA$ with H arbitrary. For the three-dimensional case with G (and thus X) skew, Sidoroff (1978), Guo (1984) and Dienes (1979) obtained formulas for the axial vector* of X in terms of A and the axial vector of G .

The direct solutions of (1.1) have been used to obtain direct formulas for various tensors of interest in continuum mechanics. Examples include formulas for:

- (1) the material time derivatives of the stretch and rotation tensors in terms of either the velocity gradient (Guo 1984; Stickforth and Wegener 1988) or the stretching and spin tensors (Mehrabadi and Nemat-Nasser 1987; Dienes 1979; Stickforth and Wegener 1988; Hoger 1986) or the material time derivatives of the Cauchy-Green tensors (Hoger and Carlson 1984);
- (2) the stretching tensor in terms of the Jaumann rate of the left Cauchy-Green tensor (Sidoroff 1978; Leonov 1976);
- (3) the derivative of the square root of a tensor (Hoger and Carlson 1984);
- (4) the elastic and plastic spin tensors in terms of the corresponding stretching tensors (Nemat-Nasser 1990);

*The *axial vector* of a skew tensor G is the unique vector g such that $Gv = g \times v$ for every vector v , where $g \times v$ denotes the vector cross product of g and v .

- (5) the spin tensor in terms of the rotational momentum and the Euler tensor, for a body undergoing a rigid motion (Truesdell, in press).

The derivatives of the stretch and rotation tensors with respect to the deformation gradient (Wheeler 1990; Scheidler, to be published) also satisfy tensor equations of the form (1.1).

Now the tensor G on the right-hand side of (1.1) may depend on A and some other tensor, say H , which is regarded as known. Expressions for G of the form

$$\begin{aligned} AH - HA, \quad AHA, \quad HA - AH^T, \quad AH - H^T A, \\ AH + H^T A, \quad HA + AH^T, \quad A^2 H + HA^2 \end{aligned} \quad (1.5)$$

occur in the applications listed above; here H^T denotes the transpose of H . Of course, to obtain a direct solution of (1.1) in these cases we can simply substitute the expression for G in terms of A and H into one of the direct solutions which are valid for any G . However, in many of these cases there exist direct formulas for X which are simpler than (but necessarily equivalent to) the formulas obtained by this approach. One of the goals of this paper is to develop methods which yield these simpler formulas with a minimal amount of computational effort.

The paper is organized as follows. Section 2 contains some preliminary material. In Sections 3 and 4 we study the general properties of the solutions of the equation (1.1) and the equation obtained from (1.1) by setting $G = AH - HA$, i.e., the tensor equation

$$AX + XA = AH - HA. \quad (1.6)$$

By "general" properties we mean properties which are independent of the dimension of the underlying vector space. We show that when G is given in terms of A and some tensor H by one of the expressions in (1.5), the solution of (1.1) has a simple expression in terms of the solution of (1.6). Furthermore, we show that the solution of (1.1) for arbitrary G can be expressed in terms of the solution of (1.6) with $H = G$.

In Sections 5 and 6 we restrict attention to the three-dimensional and two-dimensional cases, respectively, since these are the cases of most interest in continuum mechanics. We utilize

Rivlin's (1955) identities for tensor polynomials in two variables to derive direct solutions of (1.1) for the special case where G is orthogonal to every polynomial $\perp A$. These formulas are then used to derive direct solutions of (1.6) for arbitrary H , and those formulas are in turn used to derive direct solutions of (1.1) for arbitrary G . When G is given by one of the expressions in (1.5), simpler direct solutions of (1.1) follow from the identities in Section 4 and our direct solutions of (1.6). Several of the direct formulas derived in this paper have been obtained previously by some of the authors mentioned above. However, our derivations differ from theirs, and we derive some new formulas as well. Our emphasis on the special case (1.6) of the tensor equation (1.1) was motivated by some results in the paper of Mehrabadi and Nemat-Nasser (1987); see the comments following equation (4.5) in this paper.

In Section 7 we use one of our formulas to derive a new direct formula for the spin tensor in terms of the rotational momentum and the tensor of inertia for a body undergoing a rigid motion. In a follow-up paper (Scheidler, to be published) we use the general properties and the direct formulas obtained here to derive direct formulas for the derivatives of the stretch and rotation tensors with respect to the deformation gradient. We apply those results to the derivation of direct formulas for the material time derivatives of the stretch and rotation tensors and also to the derivation of direct formulas for the stress in a hyperelastic material in terms of the derivative of the strain energy with respect to the right stretch tensor. The latter problem was solved recently by Wheeler (1990) for the isotropic case.

2. ALGEBRAIC PRELIMINARIES

The underlying vector space is a finite-dimensional inner product space $\mathcal{V}$ over the reals. By a second-order tensor we mean a linear transformation from $\mathcal{V}$ into $\mathcal{V}$. The vector space of all second-order tensors is denoted by Lin . Sym and Skw are the subspaces of Lin consisting of symmetric and skew tensors. Throughout this paper A denotes an arbitrary symmetric positive-definite tensor with eigenvalues $a_i > 0$, $i = 1, 2, \dots, N$, where N is the dimension of $\mathcal{V}$. The identity tensor is denoted by I ; B , G , H , and X denote second-order tensors which are arbitrary unless specified otherwise. The inner product of B and H is

$$B \cdot H := \text{tr}(B^T H). \quad (2.1)$$

$$\text{sym } H := \frac{1}{2} (H + H^T), \quad \text{skw } H := \frac{1}{2} (H - H^T). \quad (2.2)$$

We record the following identities for later use:

$$\begin{aligned} 2 \text{ sym } (AH) &= AH + H^T A = AH + HA - 2 (\text{skw } H) A \\ &= A \text{ sym } H + (\text{sym } H) A + A \text{ skw } H - (\text{skw } H) A, \end{aligned} \quad (2.3)$$

$$\begin{aligned} 2 \text{ sym } (HA) &= HA + AH^T = AH + HA - 2A \text{ skw } H \\ &= A \text{ sym } H + (\text{sym } H) A + (\text{skw } H) A - A \text{ skw } H, \end{aligned} \quad (2.4)$$

$$\begin{aligned} 2 \text{ skw } (AH) &= AH - H^T A = AH - HA + 2 (\text{skw } H) A \\ &= A \text{ sym } H - (\text{sym } H) A + A \text{ skw } H + (\text{skw } H) A, \end{aligned} \quad (2.5)$$

$$\begin{aligned} 2 \text{ skw } (HA) &= HA - AH^T = HA - AH + 2 A \text{ skw } H \\ &= (\text{sym } H) A - A \text{ sym } H + A \text{ skw } H + (\text{skw } H) A. \end{aligned} \quad (2.6)$$

A tensor H is a polynomial in A if $H = c_m A^m + c_{m-1} A^{m-1} + \dots + c_1 A + c_0 I$ for some nonnegative integer m and scalars $c_0, c_1, \dots, c_m$. The subspace of Lin consisting of all polynomials in A is denoted by $\mathcal{P}(A)$. The Cayley-Hamilton theorem implies that

$$\begin{aligned} \mathcal{P}(A) &= \text{span } \{I, A, \dots, A^{N-1}\} \\ &= \text{span } \{A, A^2, \dots, A^N\} \\ &= \text{span } \{\dots, A^{-2}, A^{-1}, I, A, A^2, \dots\}. \end{aligned} \quad (2.7)$$

Let $\mathcal{P}(A)^\perp$ denote the orthogonal complement of $\mathcal{P}(A)$, i.e., $H \in \mathcal{P}(A)^\perp$ iff H is orthogonal to every polynomial in A . Then from (2.7) and (2.1) it follows that

$$\begin{aligned}
H \in \mathcal{P}(A)^\perp &\Leftrightarrow \text{tr}(A^k H) = 0, \quad \forall k \in \{0, 1, \dots, N-1\} \\
&\Leftrightarrow \text{tr}(A^k H) = 0, \quad \forall k \in \{1, 2, \dots, N\} \\
&\Leftrightarrow \text{tr}(A^k H) = 0, \quad \forall \text{ integer } k.
\end{aligned} \tag{2.8}$$

In particular, if $H \in \mathcal{P}(A)^\perp$ then H is deviatoric, i.e., $\text{tr } H = 0$. Also note that every skew tensor belongs to $\mathcal{P}(A)^\perp$.

By a fourth-order tensor we mean a linear transformation from Lin into Lin . K denotes an arbitrary fourth-order tensor; the image of $H \in \text{Lin}$ under K is denoted by $K[H]$. It is easily verified that *the following two conditions are equivalent*:

$$\begin{aligned}
K[H^T] &= (K[H])^T; \\
K[\text{sym } H] &= \text{sym } K[H] \quad \text{and} \quad K[\text{skw } H] = \text{skw } K[H].
\end{aligned} \tag{2.9}$$

If these conditions hold for each $H \in \text{Lin}$ we say that K is *even*. Similarly, it is easily verified that *the following two conditions are equivalent*:

$$\begin{aligned}
K[H^T] &= - (K[H])^T; \\
K[\text{sym } H] &= \text{skw } K[H] \quad \text{and} \quad K[\text{skw } H] = \text{sym } K[H].
\end{aligned} \tag{2.10}$$

If these conditions hold for every $H \in \text{Lin}$, we say that K is *odd*. The transpose of K is the unique fourth-order tensor K^T with the property

$$B \cdot K^T[H] = K[B] \cdot H, \quad \forall B, H \in \text{Lin}. \tag{2.11}$$

We say that K is symmetric if $K^T = K$.

3. THE TENSORS L_A AND M_A

In this section we study the properties of five fourth-order tensors associated with any symmetric

positive-definite second-order tensor A . The two most important of these are denoted by L_A and M_A and characterized by the conditions

$$X = L_A [G] \Leftrightarrow AX + XA = G , \quad (3.1)$$

$$X = M_A [H] \Leftrightarrow AX + XA = AH - HA . \quad (3.2)$$

The other fourth-order tensors are denoted by N_A , B_A and C_A . N_A is introduced primarily to simplify the statement of some of the identities relating L_A and M_A . B_A and C_A are introduced to facilitate the definition of the other tensors and the derivation of their properties.

The fourth-order tensor B_A is defined by

$$B_A [X] := AX + XA . \quad (3.3)$$

Note that B_A is invertible. Indeed, the condition $B_A [X] = G$ is just the tensor equation (1.1) which, as pointed out in the introduction, has a unique solution X for any given G . We denote the inverse of B_A by L_A :

$$L_A := (B_A)^{-1} . \quad (3.4)$$

Then $B_A [X] = G$ iff $X = L_A [G]$, which is equivalent to the statement (3.1). If we let I denote the fourth-order identity tensor on Lin , then*

$$L_A B_A = B_A L_A = I , \quad (3.5)$$

which is equivalent to the identities

$$L_A [AH + HA] = A L_A [H] + L_A [H] A = H . \quad (3.6)$$

*The product of two fourth-order tensors is defined to be their composition: $(L_A B_A) [X] := L_A [B_A [X]]$ for each $X \in \text{Lin}$.

The fourth-order tensor $\mathbf{C}_A$ is defined by

$$\mathbf{C}_A [H] := AH - HA . \quad (3.7)$$

It is easily verified that $\mathbf{C}_A$ and $\mathbf{B}_A$ commute; indeed,

$$\mathbf{C}_A \mathbf{B}_A = \mathbf{B}_A \mathbf{C}_A = \mathbf{C}_A^2 . \quad (3.8)$$

We define the fourth-order tensor $\mathbf{M}_A$ by

$$\mathbf{L}_A \mathbf{C}_A = \mathbf{C}_A \mathbf{L}_A , \quad (3.9)$$

where (3.9)₂ follows from pre-multiplying and post-multiplying (3.8)₁ by $\mathbf{L}_A$ and using (3.5). From (3.9) and (3.7) we see that

$$\mathbf{M}_A [H] = \mathbf{L}_A [AH - HA] = A \mathbf{L}_A [H] - \mathbf{L}_A [H] A . \quad (3.10)$$

Then (3.2) follows from (3.1) (with $G = AH - HA$) and (3.10)₁. From (3.9), (3.8), and (3.5) we obtain

$$\mathbf{M}_A \mathbf{B}_A = \mathbf{B}_A \mathbf{M}_A = \mathbf{C}_A = \mathbf{L}_A \mathbf{C}_A^2 = \mathbf{C}_A^2 \mathbf{L}_A , \quad (3.11)$$

which is equivalent to the identities

$$\begin{aligned} \mathbf{M}_A [AH + HA] &= A \mathbf{M}_A [H] + \mathbf{M}_A [H] A \\ &= AH - HA \\ &= \mathbf{L}_A [A^2 H - HA^2] = A^2 \mathbf{L}_A [H] - \mathbf{L}_A [H] A^2 . \end{aligned} \quad (3.12)$$

The fourth-order tensor $\mathbf{N}_A$ is defined by

$$\mathbf{N}_A := \mathbf{M}_A \mathbf{C}_A = \mathbf{C}_A \mathbf{M}_A = \mathbf{L}_A (\mathbf{C}_A)^2 , \quad (3.13)$$

where (3.13)_{2,3} follow from (3.9). From (3.13) and (3.7) we see that

$$\begin{aligned} N_A [H] &= M_A [AH - HA] = A M_A [H] - M_A [H] A \\ &= L_A [A^2 H + H A^2 - 2 AHA] . \end{aligned} \quad (3.14)$$

By setting $G = A^2 H + H A^2 - 2 AHA$ in (3.1) and using (3.14)₃, it follows that

$$X = N_A [H] \Leftrightarrow AX + XA = A^2 H + H A^2 - 2 AHA . \quad (3.15)$$

The motivation for introducing the tensor N_A is based not so much on the result (3.15) as on the frequent occurrence of the expressions in (3.14)_{1,2}.

It follows from the results above that the set of fourth-order tensors

$$\mathcal{S}_A := \{B_A, C_A, L_A, M_A, N_A\} \quad (3.16)$$

is commutative. Each tensor $K_A \in \mathcal{S}_A$ also has the following properties:

$$(K_A)^T = K_A ; \quad (3.17)$$

$$K_A [BH] = B K_A [H] , \text{ if } AB = BA ; \quad (3.18)$$

$$K_A [HB] = K_A [H] B , \text{ if } AB = BA ; \quad (3.19)$$

$$K_{QAQ^T} [QH Q^T] = Q K_A [H] Q^T , \forall Q \in \text{Orth} , \quad (3.20)$$

where Orth denotes the set of orthogonal tensors in Lin. In other words, $K_A [H]$ is an isotropic function of A and H which is linear in H . Furthermore, B_A , L_A and N_A are even, whereas C_A and M_A are odd. The easiest way to establish these properties is to first prove them for the tensors B_A and C_A , and then use the definitions (3.4), (3.9), and (3.13) to prove the properties for L_A , M_A , and N_A . For example, it is easily established that (3.18) holds for $K_A = B_A$ or C_A . To prove (3.18) for $K_A = L_A$, use the result $B_A [BH] = B B_A [H]$, apply L_A to obtain $BH = L_A [B B_A [H]]$, and then set $H = L_A [X]$ to obtain $B L_A [X] = L_A [BX]$. To prove (3.18) for $K_A = M_A$, use the results for $K_A = C_A$ and L_A :

$$\begin{aligned} M_A [BH] &= L_A [C_A [BH]] = L_A [B C_A [H]] \\ &= B L_A [C_A [H]] = B M_A [H] . \end{aligned}$$

The proof of (3.18) for $K_A = N_A$ is similar.

In addition to the expressions (3.10) for M_A in terms of L_A , we also have

$$M_A [H] = H - 2 L_A [H] A = 2 A L_A [H] - H . \quad (3.21)$$

To prove (3.21)₁, for example, use (3.10), (3.6), and (3.19) (with $K_A = L_A$ and $B = A$):

$$\begin{aligned} M_A [H] &= L_A [AH - HA] \\ &= L_A [AH + HA] - 2 L_A [HA] \\ &= H - 2 L_A [H] A . \end{aligned}$$

In addition to the expressions (3.14)_{1,2} for N_A in terms of M_A , we also have

$$\begin{aligned} N_A [H] &= AH - HA - 2 M_A [H] A \\ &= HA - AH + 2 A M_A [H] . \end{aligned} \quad (3.22)$$

To prove (3.22)₁, for example, use (3.14)₁, (3.12), and (3.19):

$$\begin{aligned} N_A [H] &= M_A [AH - HA] \\ &= M_A [AH + HA] - 2 M_A [HA] \\ &= AH - HA - 2 M_A [H] A . \end{aligned}$$

From (3.21) we obtain the following expressions for L_A in terms of M_A :

$$\begin{aligned} L_A [H] &= \frac{1}{2} (H - M_A [H]) A^{-1} \\ &= \frac{1}{2} A^{-1} (H + M_A [H]) \\ &= \frac{1}{4} (A^{-1}H + HA^{-1} + A^{-1} M_A [H] - M_A [H] A^{-1}) . \end{aligned} \quad (3.23)$$

Now from (3.14) it follows that

$$A^{-1} N_A [H] A^{-1} = M_A [H] A^{-1} - A^{-1} M_A [H] . \quad (3.24)$$

Then by (3.23)₃ and (3.24) we obtain the following expressions for L_A in terms of N_A :

$$\begin{aligned} L_A [H] &= \frac{1}{4} (A^{-1}H + HA^{-1} - A^{-1} N_A [H] A^{-1}) \\ &= \frac{1}{4} A^{-1} (AH + HA - N_A [H]) A^{-1} . \end{aligned} \quad (3.25)$$

To establish our next result we need the fact that

$$A^2H + HA^2 = 2 AHA \Leftrightarrow AH = HA . \quad (3.26)$$

This can be verified by taking components relative to a principal basis for A :

$$\begin{aligned} A^2H + HA^2 = 2 AHA &\Leftrightarrow (a_i^2 + a_j^2 - 2a_i a_j) H_{ij} = 0 \\ &\Leftrightarrow (a_i - a_j)^2 H_{ij} = 0 \\ &\Leftrightarrow (a_i - a_j) H_{ij} = 0 \\ &\Leftrightarrow AH - HA = 0 . \end{aligned}$$

Since L_A is nonsingular, from (3.7), (3.10), (3.12)₃, (3.14), (3.23), (3.25), and (3.26) we conclude that the following conditions are equivalent:

$$\begin{aligned} AH &= HA , \\ A L_A [H] &= L_A [H] A , \\ A M_A [H] &= \pm M_A [H] A , \\ C_A [H] &= 0 , \\ M_A [H] &= 0 , \\ N_A [H] &= 0 , \end{aligned}$$

$$\begin{aligned}
L_A [H] &= \frac{1}{2} A^{-1} H, \\
L_A [H] &= \frac{1}{2} H A^{-1}, \\
L_A [H] &= \frac{1}{4} (A^{-1} H + H A^{-1}). \tag{3.27}
\end{aligned}$$

Now suppose that $H \in \mathcal{P}(A)$. Then $AH = HA$, and thus $L_A [H] = A^{-1} H$. And since $A^{-1} \in \mathcal{P}(A)$, it follows that $L_A [H] \in \mathcal{P}(A)$. Conversely, if $L_A [H] \in \mathcal{P}(A)$ then by (3.6)₂ it follows that $H \in \mathcal{P}(A)$. Thus we have shown that

$$L_A [H] \in \mathcal{P}(A) \Leftrightarrow H \in \mathcal{P}(A). \tag{3.28}$$

In other words, if $AX + XA = H$, then X is a polynomial in A iff H is a polynomial in A . From (3.6)₂ we also see that

$$2 \operatorname{tr} (A^{k+1} L_A [H]) = \operatorname{tr} (A^k H), \quad \forall \text{ integer } k. \tag{3.29}$$

Then from (3.29) and (2.8) we conclude that

$$L_A [H] \in \mathcal{P}(A)^\perp \Leftrightarrow H \in \mathcal{P}(A)^\perp. \tag{3.30}$$

In other words, if $AX + XA = H$, then X is orthogonal to every polynomial in A iff H is orthogonal to every polynomial in A . Since $\operatorname{tr} (A^k (AH - HA)) = 0$ for any integer k , from (2.8) with $H \rightarrow AH - HA$ we conclude that

$$C_A [H] = AH - HA \in \mathcal{P}(A)^\perp. \tag{3.31}$$

Then from (3.10)₁, (3.14)₁, (3.30), and (3.31) we see that

$$M_A [H], N_A [H] \in \mathcal{P}(A)^\perp. \tag{3.22}$$

In particular, for any tensor H the solution X of (1.6) is orthogonal to every polynomial in A . Of course, since A and H commute for any $H \in \mathcal{P}(A)$, from (3.27) we also have

$$C_A[H] = M_A[H] = N_A[H] = 0, \quad \forall H \in \mathcal{P}(A). \quad (3.33)$$

We conclude this section with a description of our method for generating direct formulas for $L_A[H]$ and $M_A[H]$. Suppose that by some means we have obtained a direct formula for $L_A[G]$ which is valid for any $G \in \mathcal{P}(A)^\perp$ but not necessarily for other G . As we will see in Sections 5 and 6, such formulas are easily derived in the two-dimensional and three-dimensional cases. Since $AH - HA \in \mathcal{P}(A)^\perp$ for any tensor H , and since $M_A[H] = L_A[AH - HA]$, by setting $G = AH - HA$ in our formula for $L_A[G]$ we obtain a formula for $M_A[H]$ which is valid for any tensor H . Then from (3.23) we obtain formulas for $L_A[H]$ which are valid for any H . Alternatively, we can use the relation $N_A[H] = M_A[AH - HA]$ to obtain a formula for $N_A[H]$, and then use (3.25) to obtain formulas for $L_A[H]$ which are valid for any H . In these formulas for $L_A[H]$ we can, of course, replace the A^{-1} terms by a polynomial in A via the Cayley-Hamilton theorem.

4. SOME USEFUL IDENTITIES

Consider the tensor equation

$$AX + XA = G(A, H), \quad (4.1)$$

where $G(A, H)$ denotes one of the expressions listed in (1.5). In terms of the notation introduced in the previous section, the solution of (4.1) is $X = L_A[G(A, H)]$; and for the special case $G(A, H) = AH - HA$ this solution can also be written as $X = M_A[H]$. As we will see in the next two sections, the direct formulas for $M_A[H]$ (for arbitrary H) are much simpler than the direct formulas for $L_A[G]$ (for arbitrary G). This should not be too surprising in view of the method described in the previous section for generating these formulas. For the cases where $G(A, H)$ is given by one of the other expressions in (1.5), the existence of relatively simple direct formulas for X is due to the fact that there are simple expressions for $L_A[G(A, H)]$ in terms of M_A (or N_A), A and H . The purpose of this section is to derive these identities.

We begin with the case $G(A, H) = AHA$, that is, with the tensor equation

$$AX + XA = AHA, \quad (4.2)$$

which is equivalent to the tensor equation

$$A^{-1}X + XA^{-1} = H. \quad (4.3)$$

These equations arise in the problem of finding formulas for the material time derivative of the right stretch tensor in terms of the stretching tensor (Mehrabadi and Nemat-Nasser 1987; Hoger 1986).

Since the solution of (4.2) is $X = L_A[AHA]$, whereas the solution of the equivalent equation (4.3) is $X = L_{A^{-1}}[H]$, we see that

$$L_{A^{-1}}[H] = L_A[AHA] = A L_A[H] A, \quad (4.4)$$

where (4.4)₂ follows from (3.18) and (3.19) (with $B = A$). By substituting the expressions for $L_A[H]$ in (3.23) and (3.25) into (4.4), we obtain the identities

$$\begin{aligned} L_{A^{-1}}[H] &= \frac{1}{2} A (H - M_A[H]) \\ &= \frac{1}{2} (H + M_A[H]) A \\ &= \frac{1}{4} (AH + HA - N_A[H]). \end{aligned} \quad (4.5)$$

In other words, the solution X of (4.2) and (4.3) is also given by any of the three expressions on the right-hand side of (4.5). The second and third of these identities are equivalent to ones obtained by Mehrabadi and Nemat-Nasser (1987) under the assumption that H is symmetric. For the three-dimensional case, they used these identities and their direct formula for $M_A[H]$ to obtain direct solutions of (4.3).*

*Their equations (8.8), (8.12)–(8.13) and (8.16) correspond to equations (4.3), (4.5)₂ and (4.5)₃ in this paper. In their derivation of (8.13), and apparently of (8.16) as well, they utilized the symmetry of H ($= 2\dot{D}$ in their notation). However, these results, as well as their direct formulas for $L_{A^{-1}}[H]$, are valid for arbitrary H ; see also the comments following equation (5.26) in this paper.

Compared with (4.4), the formula for $M_{A^{-1}}$ in terms of M_A is much simpler:

$$M_{A^{-1}} = -M_A. \quad (4.6)$$

Indeed, from (3.2) we see that $X = M_A [H]$ iff $A^{-1} X + X A^{-1} = A^{-1} (-H) - (-H) A^{-1}$ iff $X = M_{A^{-1}} [-H]$, which yields (4.6). Note that the identities (4.5) can also be obtained by replacing A with A^{-1} in (3.23) and then using (4.6) and (3.14).

Now consider the case $G(A, H) = A^2 H + H A^2$, that is, the tensor equation

$$AX + XA = A^2 H + H A^2. \quad (4.7)$$

This equation arises in the problem of finding formulas for the Jaumann rate of the left stretch tensor in terms of the stretching tensor (Scheidler, to be published). Alternate expressions for the solution $X = L_A [A^2 H + H A^2]$ of (4.7) follow from the identities

$$\begin{aligned} L_A [A^2 H + H A^2] &= A^2 L_A [H] + L_A [H] A^2 \\ &= 2 L_A [H] A^2 + (AH - HA) \\ &= 2 A^2 L_A [H] - (AH - HA) \\ &= \frac{1}{2} (AH + HA + N_A [H]). \end{aligned} \quad (4.8)$$

(4.8)₁ follows from (3.18) and (3.19); (4.8)_{2,3} follow from (4.8)₁ and (3.12); (4.8)₄ follows from (3.14)₃, (4.4)₁ and (4.5)₃.

Next, we derive alternate expressions for $L_A [G(A, H)]$ when $G(A, H) = HA - AH^T$, $AH - H^T A$, $AH + H^T A$ or $HA + AH^T$; for applications of the tensor equation (4.1) in these cases see Guo (1984) and Scheidler (to be published). We claim that

$$\begin{aligned} L_A [HA - AH^T] &= 2 L_A [\text{skw} (HA)] \\ &= \text{skw} H - M_A [\text{sym} H] \\ &= \text{skw} (H - M_A [H]), \end{aligned} \quad (4.9)$$

$$\begin{aligned}
L_A [AH - H^T A] &= 2 L_A [\text{skw} (AH)] \\
&= \text{skw} H + M_A [\text{sym} H] \\
&= \text{skw} (H + M_A [H]) ,
\end{aligned} \tag{4.10}$$

$$\begin{aligned}
L_A [AH + H^T A] &= 2 L_A [\text{sym} (AH)] \\
&= H - 2 L_A [\text{skw} H] A \\
&= \text{sym} H + M_A [\text{skw} H] \\
&= \text{sym} (H + M_A [H]) ,
\end{aligned} \tag{4.11}$$

$$\begin{aligned}
L_A [HA + AH^T] &= 2 L_A [\text{sym} (HA)] \\
&= H - 2 A L_A [\text{skw} H] \\
&= \text{sym} H - M_A [\text{skw} H] \\
&= \text{sym} (H - M_A [H]) .
\end{aligned} \tag{4.12}$$

These identities follow easily from (2.3)–(2.6), (3.6), (3.10), (3.18), (3.19) and the fact that M_A is odd (see (2.10)).

We conclude this section with two identities which are used in Scheidler (to be published):

$$\begin{aligned}
M_A [AH - H^T A] &= 2 M_A [\text{skw} (AH)] \\
&= A \text{skw} H - (\text{skw} H) A + N_A [\text{sym} H] ,
\end{aligned} \tag{4.13}$$

$$\begin{aligned}
M_A [HA - AH^T] &= 2 M_A [\text{skw} (HA)] \\
&= A \text{skw} H - (\text{skw} H) A - N_A [\text{sym} H] .
\end{aligned} \tag{4.14}$$

These identities follow easily from (2.5), (2.6), (3.12), and (3.14).

5. FORMULAS FOR THE THREE-DIMENSIONAL CASE

In this section we derive direct formulas for the fourth-order tensors L_A , M_A , and N_A under the assumption that the underlying vector space $\mathcal{V}$ is three-dimensional. We begin by listing some identities for second-order tensors which will be used in the derivation of these formulas. The principal invariants of a second-order tensor B are denoted by I_B , II_B and III_B :

$$I_B := \text{tr } B, \quad II_B := \frac{1}{2} [(\text{tr } B)^2 - \text{tr } (B^2)], \quad III_B := \det B. \quad (5.1)$$

From the Cayley-Hamilton theorem we obtain

$$A^3 = I_A A^2 - II_A A + III_A I, \quad (5.2)$$

and

$$III_A A^{-1} = A^2 - I_A A + II_A I. \quad (5.3)$$

Since the expression $I_A I - A$ occurs frequently below, it is convenient to introduce a special symbol for this expression:

$$\tilde{A} := I_A I - A. \quad (5.4)$$

The eigenvalues of $\tilde{A}$ are $a_2 + a_3$, $a_3 + a_1$, $a_1 + a_2$; in particular, $\tilde{A}$ is symmetric positive-definite. Also note that (Sidoroff 1978; Guo 1984; Stückforth and Wegener 1988)

$$III_{\tilde{A}} = I_A II_A - III_A = (a_1 + a_2)(a_2 + a_3)(a_3 + a_1), \quad (5.5)$$

and that (Guo 1984)

$$III_{\tilde{A}} \tilde{A}^{-1} = A^2 + II_A I. \quad (5.6)$$

The following identities are due to Rivlin (1955):

$$\begin{aligned} A^2 H A^2 &= II_A A H A - III_A (A H + H A) \\ &\quad + I_{A^2 H} A^2 + \alpha_1(A, H) A + III_A I_{A H} I, \end{aligned} \quad (5.7)$$

$$\begin{aligned} A^2 H A + A H A^2 &= I_A A H A - III_A H \\ &\quad + I_{A H} A^2 + \alpha_2(A, H) A + III_A I_H I, \end{aligned} \quad (5.8)$$

$$\begin{aligned} A^2 H + H A^2 &= -A H A + I_A (A H + H A) - II_A H \\ &\quad + I_H A^2 + \alpha_3(A, H) A + \alpha_4(A, H) I, \end{aligned} \quad (5.9)$$

where

$$\begin{aligned}\alpha_1(A, H) &= \text{tr} [(\text{III}_A I - \text{II}_A A) H] \\ &= \text{III}_A I_H - \text{II}_A I_{AH} = -I_{\tilde{A}A^2H}\end{aligned}\quad (5.10)$$

$$\begin{aligned}\alpha_2(A, H) &= \text{tr} [(A^2 - I_A A) H] \\ &= I_{A^2H} - I_A I_{AH} = -I_{\tilde{A}AH},\end{aligned}\quad (5.11)$$

$$\begin{aligned}\alpha_3(A, H) &= -\text{tr} [(I_A I - A) H] \\ &= I_{AH} - I_A I_H = -I_{\tilde{A}H},\end{aligned}\quad (5.12)$$

$$\begin{aligned}\alpha_4(A, H) &= \text{tr} [(A^2 - I_A A + \text{II}_A I) H] \\ &= I_{A^2H} - I_A I_{AH} + \text{II}_A I_H = \text{III}_A I_{A^{-1}H}.\end{aligned}\quad (5.13)$$

Rivlin's identities, as well as (5.2)–(5.5)₁, are valid for any second-order tensors A and H . However, we will continue to assume that A is symmetric positive-definite.

Observe that if $H \in \mathcal{P}(A)^\perp$ (see (2.8)) then the second line on the right-hand side of Rivlin's identities (5.7)–(5.9) vanishes. In this case we also have the identity

$$\tilde{A} (AH + HA) \tilde{A} = \text{III}_{\tilde{A}} H, \quad \forall H \in \mathcal{P}(A)^\perp. \quad (5.14)$$

This follows by substituting (5.4) into the left-hand side of (5.14), expanding, and then using (5.5)₁, (5.8), and (5.9). Similarly, from (5.5)₁, (5.7), and (5.9) we obtain the identity

$$(A^2 + \text{II}_A I) H (A^2 + \text{II}_A I) = \text{III}_{\tilde{A}} (AH + HA), \quad \forall H \in \mathcal{P}(A)^\perp. \quad (5.15)$$

In view of (5.6) we see that the identities (5.14) and (5.15) are equivalent.

Now suppose that $G \in \mathcal{P}(A)^\perp$, in which case $L_A[G] \in \mathcal{P}(A)^\perp$ also (see (3.30)). If we set $H = L_A[G]$ in (5.14) we obtain

$$\tilde{A} (A L_A[G] + L_A[G] A) \tilde{A} = \text{III}_{\tilde{A}} L_A[G].$$

But since $A L_A [G] + L_A [G] A = G$ for any tensor G (see (3.6)), we have the following simple formula for $L_A [G]$:

$$\text{III}_{\tilde{A}} L_A [G] = \tilde{A} G \tilde{A}, \quad \forall G \in \mathcal{P}(A)^\perp. \quad (5.16)$$

By substituting the definition (5.4) of $\tilde{A}$ into (5.16) and expanding, we obtain the alternate formula

$$\text{III}_{\tilde{A}} L_A [G] = I_A^2 G - I_A (AG + GA) + AGA, \quad \forall G \in \mathcal{P}(A)^\perp. \quad (5.17)$$

By using Rivlin's identity (5.9) with $H = G$, we see that (5.17) is equivalent to the formula

$$\text{III}_{\tilde{A}} L_A [G] = (I_A^2 - \Pi_A) G - (A^2 G + G A^2), \quad \forall G \in \mathcal{P}(A)^\perp. \quad (5.18)$$

In view of (3.1), the formulas (5.16)–(5.18) yield simple direct solutions of the tensor equation (1.1) when $G \in \mathcal{P}(A)^\perp$. Since any skew tensor belongs to $\mathcal{P}(A)^\perp$, the formulas (5.16)–(5.18) hold in particular for any $G \in \text{Skw}$. For this special case the formula (5.18) was obtained by Sidoroff (1978) and Guo (1984). Their derivations employ the axial vector associated with a skew tensor and thus do not carry over to the more general case where G need only belong to $\mathcal{P}(A)^\perp$.

Since $M_A [H] = L_A [AH - HA]$, and since $AH - HA \in \mathcal{P}(A)^\perp$ for any tensor H (see (3.31)), by setting $G = AH - HA$ in (5.16) and (5.17) we obtain the following simple formulas for $M_A [H]$:

$$\begin{aligned} \text{III}_{\tilde{A}} M_A [H] &= \tilde{A} (AH - HA) \tilde{A} \\ &= I_A^2 (AH - HA) - I_A (A^2 H - HA^2) + A^2 HA - AHA^2. \end{aligned} \quad (5.19)$$

Furthermore, since

$$AH - HA = H\tilde{A} - \tilde{A}H, \quad (5.20)$$

by (5.19), we also have

$$\begin{aligned} \text{III}_{\tilde{A}} M_A [H] &= \tilde{A} (H\tilde{A} - \tilde{A}H) \tilde{A} \\ &= \tilde{A} H \tilde{A}^2 - \tilde{A}^2 H \tilde{A}. \end{aligned} \quad (5.21)$$

The formula (5.19)₂ was obtained by Mehrabadi and Nemat-Nasser (1987) by repeated applications of the Cayley-Hamilton theorem. We emphasize that (5.19) and (5.21) are valid for any tensor H . In view of (3.2), these formulas yield simple direct solutions of the tensor equation (1.6). Furthermore, in view of (3.1) and the identities (4.5)_{1,2}, (4.9), (4.10), (4.11)_{2,4}, and (4.12)_{2,4}, the formulas (5.16)–(5.18), (5.19) and (5.21) yield simple direct solutions of (4.1) when $G(A, H) = AHA$, $HA - AH^T$, $AH - H^T A$, $AH + H^T A$ or $HA + AH^T$.

Since $N_A[H] = M_A[AH - HA]$, by replacing H with $AH - HA$ in (5.19)₁ we obtain the formula

$$\text{III}_{\tilde{A}} N_A[H] = \tilde{A} (A^2 H + HA^2 - 2 AHA) \tilde{A}. \quad (5.22)$$

By substituting (5.4) into (5.22), expanding, and then using (5.2) to reduce the A^3 terms, we obtain

$$\begin{aligned} \text{III}_{\tilde{A}} N_A[H] = & -2A^2 HA^2 + 2I_A (A^2 HA + AHA^2) - 2(I_A^2 + \text{II}_A) AHA \\ & + (I_A \text{II}_A + \text{III}_A) (AH + HA) - 2I_A \text{III}_A H. \end{aligned} \quad (5.23)$$

Substitution of Rivlin's identities (5.7) and (5.8) into (5.23) yields

$$\begin{aligned} \text{III}_{\tilde{A}} N_A[H] = & -4\text{II}_A AHA + (I_A \text{II}_A + 3\text{III}_A) (AH + HA) - 4I_A \text{III}_A H \\ & - 2\alpha_2(A, H) A^2 + 2\alpha_5(A, H) A - 2\text{III}_A \alpha_3(A, H) I, \end{aligned} \quad (5.24)$$

where α_2 and α_3 are defined by (5.11) and (5.12), and

$$\begin{aligned} \alpha_5(A, H) = & \text{tr} [(I_A A^2 + (\text{II}_A - I_A^2) A - \text{III}_A I) H] \\ = & I_A I_{A^2 H} + (\text{II}_A - I_A^2) I_{AH} - \text{III}_A I_H. \end{aligned} \quad (5.25)$$

By using $N_A[H] = M_A[AH - HA]$, (5.20) and (5.21) with $H \rightarrow H\tilde{A} - \tilde{A}H$, we also obtain

$$\begin{aligned} \text{III}_{\tilde{A}} N_A[H] = & \tilde{A} (\tilde{A}^2 H + H\tilde{A}^2 - 2\tilde{A}H\tilde{A}) \tilde{A} \\ = & \tilde{A}^3 H\tilde{A} + \tilde{A}H\tilde{A}^3 - 2\tilde{A}^2 H\tilde{A}^2 \\ = & -2\tilde{A}^2 H\tilde{A}^2 + I_{\tilde{A}} (\tilde{A}^2 H\tilde{A} + \tilde{A}H\tilde{A}^2) - 2\text{II}_{\tilde{A}} \tilde{A}H\tilde{A} + \text{III}_{\tilde{A}} (\tilde{A}H + H\tilde{A}), \end{aligned} \quad (5.26)$$

where the last formula follows from the Cayley-Hamilton theorem for $\bar{A}$. In view of (4.5)₃ and (4.8)₄, the formulas (5.22)–(5.26) yield direct solutions of the tensor equations (4.2), (4.3), and (4.7). The direct solution of (4.2) or (4.3) obtained by substituting (5.23) into (4.5)₃ is equivalent to the first of two formulas obtained by Mehrabadi and Nemat-Nasser (1987); cf. equation (8.8) and the formula preceding (8.17) in their paper.* Their second formula is similar to the one obtained by substituting (5.24) into (4.5)₃; Rivlin's identity (5.9) may be used to obtain their result from ours and vice versa.

Now we derive some formulas for $L_A [H]$ which are valid for any tensor H . Substitution of (5.23) into the expression (3.25)₁ for L_A in terms of N_A yields

$$\begin{aligned} 2 \text{III}_{\bar{A}} L_A [H] = & AHA - I_A (AH + HA) + (I_A^2 + \text{II}_A) H \\ & + I_A \text{III}_A A^{-1} HA^{-1} - \text{III}_A (A^{-1}H + HA^{-1}). \end{aligned} \quad (5.27)$$

This formula was stated without proof by Leonov (1976) and Stickforth and Wegener (1988). Leonov attributes (5.27) to L. M. Zubov; Stickforth and Wegener refer the reader to Stickforth (1982–1983) for a proof. Substitution of the formula (5.3) for A^{-1} into (5.27) yields

$$\begin{aligned} 2 \text{III}_A \text{III}_{\bar{A}} L_A [H] = & I_A A^2 HA^2 - I_A^2 (A^2 HA + AHA^2) \\ & + \text{III}_{\bar{A}} (A^2 H + HA^2) + (I_A^3 + \text{III}_A) AHA \\ & - I_A^2 \text{II}_A (AH + HA) + (I_A^2 \text{III}_A + \text{II}_A \text{III}_{\bar{A}}) H. \end{aligned} \quad (5.28)$$

This formula is due to Hoger and Carlson (1984); they derived it under the assumption that H is symmetric and then observed that the formula is valid for any H . In view of (3.1), the formulas (5.27) and (5.28) yield direct formulas for the solution X of the tensor equation $AX + XA = H$ for arbitrary H .

Observe that when $G \in \mathcal{P}(A)^\perp$, the simple formulas (5.16)–(5.18) for $L_A [G]$ do not follow immediately from (5.27) or (5.28). However, Hoger and Carlson (1984) noted that substitution of Rivlin's identities (5.7)–(5.9) into (5.28) yields

*The last term on the right-hand side of their first formula contains a misprint; the term should read $I_V \text{III}_V \hat{D}$.

$$\begin{aligned} \text{III}_{\tilde{A}} \mathbf{L}_A [H] = & - (A^2 H + H A^2) + (\text{I}_A^2 - \text{II}_A) H \\ & + \beta_2(A, H) A^2 + \beta_1(A, H) A + \beta_0(A, H) I ; \end{aligned} \quad (5.29)$$

cf. their equation (2.7) for expressions for the isotropic scalar functions β_i . They observed that if $H \in \text{Skw}$ then the β_i are zero, and consequently, (5.29) reduces to (5.18) for skew G . Indeed, it is clear from their expressions for the $\beta_i(A, H)$ that they reduce to zero for any $H \in \mathcal{P}(A)^\perp$, and thus that (5.18) is a special case of (5.29). By using a representation for isotropic tensor functions of two symmetric tensors, together with his formula for $\mathbf{L}_A [G]$ when $G \in \text{Skw}$ (see (5.18)), Sidoroff (1978) also obtained (5.29). His expressions for the β_i are more compact than those of Hoger and Carlson; however, Sidoroff's expressions involve A^{-1} and A^{-2} and thus may be less useful for some applications. Other direct formulas for $\mathbf{L}_A$ in terms of A and A^{-1} follow from (3.23)_{1,2} and (5.19)₁:

$$\begin{aligned} 2 \mathbf{L}_A [H] &= [H - \frac{1}{\text{III}_A} \tilde{A} (AH - HA) \tilde{A}] A^{-1} \\ &= HA^{-1} + \frac{1}{\text{III}_A} \tilde{A} (H - AHA^{-1}) \tilde{A} \\ &= A^{-1} [H + \frac{1}{\text{III}_A} \tilde{A} (AH - HA) \tilde{A}] \\ &= A^{-1} H + \frac{1}{\text{III}_A} \tilde{A} (H - A^{-1} HA) \tilde{A} \end{aligned} \quad (5.30)$$

6. FORMULAS FOR THE TWO-DIMENSIONAL CASE

In this section we derive direct formulas for the fourth-order tensors $\mathbf{L}_A$, $\mathbf{M}_A$, and $\mathbf{N}_A$ under the assumption that the underlying vector space $\mathcal{V}$ is two-dimensional. We begin by listing some identities for second-order tensors which will be used in the derivation of these formulas. The principal invariants of a second-order tensor B are denoted by I_B and II_B :

$$\text{I}_B := \text{tr } B, \quad \text{II}_B := \det B. \quad (6.1)$$

From the Cayley-Hamilton theorem we obtain

$$A^2 = I_A A - \Pi_A I, \quad \Pi_A A^{-1} = I_A I - A. \quad (6.2)$$

The following identities are due to Rivlin (1955):

$$AH + HA = I_A H + I_H A + \beta_1(A, H) I, \quad (6.3)$$

$$AHA = \Pi_A H + I_{AH} A - \Pi_A I_H I, \quad (6.4)$$

where

$$\begin{aligned} \beta_1(A, H) &= \text{tr} [(A - I_A I) H] \\ &= I_{AH} - I_A I_H = -\Pi_A I_A^{-1} H. \end{aligned} \quad (6.5)$$

Observe that if $H \in \mathcal{P}(A)^\perp$, then all but the first terms on the right-hand side of Rivlin's identities (6.3) and (6.4) vanish:

$$AH + HA = I_A H, \quad AHA = \Pi_A H, \quad \forall H \in \mathcal{P}(A)^\perp. \quad (6.6)$$

Now suppose that $G \in \mathcal{P}(A)^\perp$, in which case $L_A[G] \in \mathcal{P}(A)^\perp$ also. If we set $H = L_A[G]$ in (6.6)₁ and use the fact that $AL_A[G] + L_A[G]A = G$ for any tensor G , we obtain the following simple formula for $L_A[G]$:

$$I_A L_A[G] = G, \quad \forall G \in \mathcal{P}(A)^\perp. \quad (6.7)$$

In particular, the formula (6.7) holds for any $G \in \text{Skw}$. Since $M_A[H] = L_A[AH - HA]$, and since $AH - HA \in \mathcal{P}(A)^\perp$ for any tensor H , by setting $G = AH - HA$ in (6.7) we obtain the following simple formula for $M_A[H]$:

$$I_A M_A[H] = AH - HA. \quad (6.8)$$

This formula was obtained by Mehrabadi and Nemat-Nasser (1987) using a different method.

Since $N_A[H] = M_A[AH - HA]$, by replacing H with $AH - HA$ in (6.8) we obtain the formula

$$I_A N_A[H] = A^2 H + HA^2 - 2AHA. \quad (6.9)$$

Substitution of (6.2)₁ into (6.9) yields

$$I_A N_A [H] = -2 AHA + I_A (AH + HA) - 2 \Pi_A H . \quad (6.10)$$

Recall that $X = L_A^{-1} [H]$ is the solution of the tensor equations (4.2) and (4.3). As an application of the formula (6.10) for $N_A [H]$, observe that substitution of (6.10) into the identity (4.5)₃ yields the formulas

$$\begin{aligned} 2 I_A L_A^{-1} [H] &= AHA + \Pi_A H \\ &= 2 \Pi_A H + I_{AH} A - \Pi_A I_H I . \end{aligned} \quad (6.11)$$

The second formula follows from the first by Rivlin's identity (6.4). A formula equivalent to (6.11)₁ was obtained by Hoger (1986) using a different method; cf. equation (4.5) and the equation preceding (4.3) in her paper.

Finally, we derive some formulas for $L_A [H]$ which are valid for any tensor H . Substitution of (6.10) into (3.25)₂ yields

$$2 I_A L_A [H] = H + \Pi_A A^{-1} H A^{-1} . \quad (6.12)$$

This also follows from (6.11)₁ with $A \rightarrow A^{-1}$ and the identities $I_{A^{-1}} = I_A / \Pi_A$ and $\Pi_{A^{-1}} = 1 / \Pi_A$. Recall that A^{-1} can be expressed in terms of A by the simple formula (6.2)₂. By substituting (6.2)₂ into (6.12) and expanding, we obtain

$$2 I_A \Pi_A L_A [H] = AHA - I_A (AH + HA) + (I_A^2 + \Pi_A) H . \quad (6.13)$$

Substitution of Rivlin's identities (6.3) and (6.4) into (6.13) yields

$$2 I_A \Pi_A L_A [H] = 2 \Pi_A H + \beta_1(A, H) A + \beta_0(A, H) I , \quad (6.14)$$

where β_1 is given by (6.5), and

$$\begin{aligned} \beta_0(A, H) &= \text{tr} [(\Pi_A^2 - \Pi_A) I - I_A A] H \\ &= (\Pi_A^2 - \Pi_A) I_H - I_A I_{AH} . \end{aligned} \quad (6.15)$$

The formulas (6.13) and (6.14) were obtained by Hoger and Carlson (1984).

7. AN APPLICATION TO THE KINEMATICS OF RIGID MOTIONS

The material in this section, with the exception of the last equation, is taken from § I.10 of the textbook by Truesdell (in press). Let W denote the spin tensor. Let M , E and J denote the rotational momentum, the Euler tensor and the tensor of inertia of a body relative to the (possibly time - dependent) point x_0 . J and E are symmetric positive-definite tensors which are related as follows:

$$J = (\text{tr } E) I - E . \quad (7.1)$$

If the body is undergoing a rigid motion, and if x_0 is either the location of the center of mass of the body or a fixed point of the body, then the rotational momentum relative to x_0 is related to the spin tensor and the Euler tensor relative to x_0 by the formula

$$M = - (EW + WE) . \quad (7.2)$$

Truesdell (in press) utilized Guo's formula (see (5.18)) to solve (7.2) for the spin tensor in terms of the rotational momentum and the Euler tensor:

$$W = \frac{-1}{I_E II_E - III_E} \left[(I_E^2 - II_E) M - (E^2 M + M E^2) \right] . \quad (7.3)$$

The purpose of this section is to point out that W has a simpler expression in terms of the tensor of inertia. Indeed, since the solution of (7.2) is $W = L_E [-M]$, by setting $A = E$ and $G = -M$ in (5.4), (5.5) and (5.16) and using (7.1), we find that

$$W = - \frac{1}{III_J} J M J . \quad (7.4)$$

INTENTIONALLY LEFT BLANK.

8. REFERENCES

- Dienes, J. K. "On the Analysis of Rotation and Stress Rate in Deforming Bodies." Acta Mechanica, vol. 32, pp. 217-232, 1979.
- Guo, Z. - H. "Rates of Stretch Tensors." Journal of Elasticity, vol. 14, pp. 263-267, 1984.
- Hoger, A. "The Material Time Derivative of Logarithmic Strain." International Journal of Solids Structures, vol. 22, pp. 1019-1032, 1986.
- Hoger, A., and D. E. Carlson. "On the Derivative of the Square Root of a Tensor and Guo's Rate Theorems." Journal of Elasticity, vol. 14, pp. 329-336, 1984.
- Leonov, A. I. "Nonequilibrium Thermodynamics and Rheology of Viscoelastic Polymer Media." Rheologica Acta, vol. 15, pp. 85-98, 1976.
- Mehrabadi, M. M., and S. Nemat-Nasser. "Some Basic Kinematical Relations for Finite Deformations of Continua." Mechanics of Materials, vol. 6, pp. 127-138, 1987.
- Nemat-Nasser, S. "Certain Basic Issues in Finite-Deformation Continuum Plasticity." Meccanica, vol. 25, pp. 223-229, 1990.
- Rivlin, R. S. "Further Remarks on the Stress-Deformation Relations for Isotropic Materials." Journal of Rational Mechanics and Analysis, vol. 4, pp. 681-701, 1955.
- Scheidler, M. "The Derivatives of the Stretch and Rotation Tensors." U.S. Army Ballistic Research Laboratory, Aberdeen Proving Ground, MD, to be published.
- Sidoroff, F. "Sur L'équation Tensorielle $AX + XA = H$." C. R. Acad. Sc. Paris A286, pp. 71-73, 1978.
- Stickforth J., and K. Wegener. "A Note on Diene's and Aifantis' Co-Rotational Derivatives." Acta Mechanica, vol. 74, pp. 227-234, 1988.
- Stickforth, J. "Vector- und Tensorrechnung für Ingenieure." Übungen und Anwendungen (Lecture Notes). Techn. Univ. Braunschweig, 1982-1983.
- Truesdell, C. A First Course in Rational Continuum Mechanics, vol. 1, 2d ed., Academic Press (in press).
- Wheeler, L. "On the Derivatives of the Stretch and Rotation With Respect to the Deformation Gradient." Journal of Elasticity, vol. 24, pp. 129-133, 1990.

INTENTIONALLY LEFT BLANK.

No. of
Copies Organization

- 2 Administrator
Defense Technical Info Center
ATTN: DTIC-DDA
Cameron Station
Alexandria, VA 22304-6145
- 1 Commander
U.S. Army Materiel Command
ATTN: AMCAM
5001 Eisenhower Avenue
Alexandria, VA 22333-0001
- 1 Commander
U.S. Army Laboratory Command
ATTN: AMSLC-DL
2800 Powder Mill Road
Adelphi, MD 20783-1145
- 2 Commander
U.S. Army Armament Research,
Development, and Engineering Center
ATTN: SMCAR-IMI-I
Picatinny Arsenal, NJ 07806-5000
- 2 Commander
U.S. Army Armament Research,
Development, and Engineering Center
ATTN: SMCAR-TDC
Picatinny Arsenal, NJ 07806-5000
- 1 Director
Benet Weapons Laboratory
U.S. Army Armament Research,
Development, and Engineering Center
ATTN: SMCAR-CCB-TL
Watervliet, NY 12189-4050
- (Unclass. only) 1 Commander
U.S. Army Armament, Munitions
and Chemical Command
ATTN: AMSMC-IMF-L
Rock Island, IL 61299-5000
- 1 Director
U.S. Army Aviation Research
and Technology Activity
ATTN: SAVRT-R (Library)
M/S 219-3
Ames Research Center
Moffett Field, CA 94035-1000

No. of
Copies Organization

- 1 Commander
U.S. Army Missile Command
ATTN: AMSMI-RD-CS-R (DOC)
Redstone Arsenal, AL 35898-5010
- 1 Commander
U.S. Army Tank-Automotive Command
ATTN: ASQNC-TAC-DIT (Technical
Information Center)
Warren, MI 48397-5000
- 1 Director
U.S. Army TRADOC Analysis Command
ATTN: ATRC-WSR
White Sands Missile Range, NM 88002-5502
- 1 Commandant
U.S. Army Field Artillery School
ATTN: ATSF-CSI
Ft. Sill, OK 73503-5000
- (Class. only) 1 Commandant
U.S. Army Infantry School
ATTN: ATSH-CD (Security Mgr.)
Fort Benning, GA 31905-5660
- (Unclass. only) 1 Commandant
U.S. Army Infantry School
ATTN: ATSH-CD-CSO-OR
Fort Benning, GA 31905-5660
- 1 Air Force Armament Laboratory
ATTN: WL/MNOI
Eglin AFB, FL 32542-5000
- Aberdeen Proving Ground
- 2 Dir, USAMSAA
ATTN: AMXSY-D
AMXSY-MP, H. Cohen
- 1 Cdr, USATECOM
ATTN: AMSTE-TC
- 3 Cdr, CRDEC, AMCCOM
ATTN: SMCCR-RSP-A
SMCCR-MU
SMCCR-MSI
- 1 Dir, VLAMO
ATTN: AMSLC-VL-D
- 10 Dir, BRL
ATTN: SLCBR-DD-T

No. of
Copies Organization

- 1 U.S. Army Research Office
ATTN: SLCRO-MA, Julian Wu
P.O. Box 12211
Research Triangle Park, NC 27709-2211
- 1 Los Alamos National Laboratory
ATTN: John K. Dienes, MS B214
Los Alamos, NM 87545

USER EVALUATION SHEET/CHANGE OF ADDRESS

This laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers below will aid us in our efforts.

1. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

2. How, specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

3. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. _____

4. General Comments. What do you think should be changed to improve future reports? (indicate changes to organization, technical content, format, etc.) _____

BRL Report Number BRL-TR-3315 Division Symbol _____

Check here if desire to be removed from distribution list. _____

Check here for address change. _____

Current address: Organization _____
Address _____

DEPARTMENT OF THE ARMY
Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-5066

OFFICIAL BUSINESS

BUSINESS REPLY MAIL

FIRST CLASS PERMIT No 0001, APG, MD

Postage will be paid by addressee.

Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-5066



NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

