

AD-A258 403



2

COLLOQUE INTERNATIONAL

Convergences en analyse multivoque et unilatérale

22-26 juin 1992
MARSEILLE - LUMINY




DTIC
ELECTE
DEC 02 1992
S E D

RÉSUMÉS DES CONFÉRENCES

ÉDITÉS PAR

HEDY ATTOUCH

MICHEL THÉRA



92-30616



2028

CONVERGENCES EN ANALYSE MULTIVOQUE ET UNILATERALE

22 - 26 JUIN 1992

CIRM, Marseille Luminy

Accession For	
NTIS	CRA&I ☒
DTIC	TAB ☐
Unannounced	☐
Justification	<i>per form 50</i>
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
<i>A-1</i>	

Résumés des Conférences

Convergences en Analyse Multivoque et Unilatérale

Organisateurs: H. Attouch (Montpellier)
M. Théra (Limoges)

Congrès satellite du Congrès Européen

Avec le patronage de:

Commission pour le Développement et les Echanges
Conseil Général des Bouches du Rhône
Conseil National de la Recherche Scientifique
Département des Sciences Mathématiques de l'Université Montpellier II
Direction des Etudes et Recherches Doctorales
European Research Office
Laboratoire "Analyse non-linéaire et Optimisation" Limoges
Ministère de la Recherche et de la Technologie
Ministère des Affaires Etrangères
Pôle Montpellier-Méditerranée d'Analyse non-linéaire Appliquée et Optimisation
Société de Mathématiques et Applications Industrielles (groupe MODE)
Société Mathématique de France
Université de Limoges
Ville de Marseille

Thèmes abordés

Convergences d'ensembles (Painlevé-Kuratowski, Mosco, Hausdorff et Hausdorff bornée, slice,...). G-convergence des fonctions (épiconvergence, epi-hypo convergence,...). Convergence en graphes des opérateurs. Topologies sur les hyperespaces.

Applications à l'approximation, la perturbation, l'analyse sensitive en :

- optimisation et analyse non-lisse (épi-dérivées des fonctions, proto-différentiation des opérateurs);
- optimisation stochastique (schémas numériques basés sur des méthodes d'approximation) et statistique (loi des grands nombres pour les variables aléatoires semi-continues);
- calcul des variations et mécanique des milieux continus, homogénéisation des matériaux composites, relaxation des problèmes de contrôle,...)
- analyse asymptotique des fonctions et des ensembles (fonctions de récession,...).
- Géométrie des espaces de Banach et problèmes d'optimisation bien posés.

Conférences plénières

Z. Artstein (Rehovot)
E. Balder (Utrecht)
G. Beer (Los Angeles)
A. Cellina (Trieste)
E. De Giorgi (Pisa)
U. Mosco (Rome)
S. M. Robinson (Madison)
R. T. Rockafellar (Seattle)
R. J.-B. Wets (Davis)

Conférenciers et Auteurs

Arstein, Zvi (Rehovot)
Attouch, Hedy (Montpellier)
Auslender, Alfred (Clermont-Ferrand)
Azé, Dominique (Perpignan)
Bahraoui, Mohamed (Montpellier)
Baillon, Jean-Bernard (Lyon 1)
Balder, Erik (Utrecht)
Beer, Gerald (Los Angeles)
Benoist, Joël (Limoges)
Blum, E (Lima)
Bouchitté, Guy (Toulon)
Bougeard, Mireille (Paris 10)
Cellina, Arrigo (Trieste)
Chabi, El Hassan (Montpellier)
Cominetti, Roberto (Santiago)
Corradini, Pietro Poggi (Milan)
Correa, Raphael (Santiago)
Costantini, Camillo (Milan)
Crouzeix, Jean-Pierre (Clermont-Ferrand)
De Giorgi, Ennio (Pise)
Deville, Robert (Besançon)
Dinh-The Luc (Hanoi)
Flam, Sjur (Bergen)
Gautier, Serge (Pau)
Hess, Christian (Paris-Dauphine)
Hiriart-Urruty, Jean-Baptiste (Toulouse 3)
Jalby, Vincent (Montpellier)
Jofré, Alejandro (Santiago)
Jongen, Bert (Aachen)
Kebbour (Clermont-Ferrand)
Klatte, Diethard (Zürich)
Lemaire, Bernard (Montpellier)
Levi, Sandro (Milan)

Loridan, Pierre (Dijon)
Lucchetti, Roberto (Milan)
Malivert, Christian (Limoges)
Marino, Antonio (Pise)
Martinez-Legaz, Juan-Enrique (Barcelone)
Michaille, Gérard (Montpellier)
Michelot, Christian (Paris 1)
Morgan, Jacqueline (Naples)
Mosco, Umberto (Rome)
Moussaoui, Mohamed (Avignon)
Ndoutoune, James (Libreville)
Noll, Dominikus (Stuttgart)
Oettli, Werner (Mannheim)
Papini, Pier-Luigi (Bologne)
Penot, Jean-Paul (Pau)
Phelps, Robert (Seattle)
Poliquin, René (Edmonton)
Qi, Li Qun (Sydney)
Revalski, Julian (Sofia)
Riahi, Hassan (Marrakech)
Ricceri, Biagio (Catane)
Robinson, Stephen (Madison)
Rockafellar, Tyrrel (Seattle)
San Martin, Jaime (Santiago)
Seeger, Alberto (Avignon)
Simons, Stephen (Santa Barbara)
Sonntag, Yves (Marseille)
Théra, Michel (Limoges)
Thibault, Lionel (Pau)
Torre, Anna (Pavie)
Valadier, Michel (Montpellier)
Vitolo, Paolo (Potenza)
Volle, Michel (Avignon)
Wets, Roger J.-B (Davis)
Zalinescu, Constantin (Cluj)
Zolezzi, Tullio (Gênes)

Présentation scientifique

H. Attouch et M. Théra

Introduction. Les convergences de suites d'ensembles et les topologies sur les fermés d'un espace topologique se sont révélées au cours des dernières années comme étant la clé de la compréhension de nombreux problèmes de convergence, approximation, perturbation en analyse non-linéaire, optimisation, statistique... Le congrès du CIRM du 22 au 26 Juin 1992, est consacré à ces questions. Nous présentons dans les pages qui suivent ces nouveaux concepts et quelques unes de leurs applications.

Les notions fondamentales de l'analyse mathématique, telles que continuité, dérivation, intégration, approximation, développements (en série...) reposent sur le choix d'un concept de limite. Dans l'approche classique, c'est la convergence simple des fonctions et des opérateurs qui sous-tend la plupart de ces notions, avec bien sûr des adaptations pour chaque domaine d'utilisation (convergence presque sûre, convergence au sens des distributions, convergence uniforme sur les bornés, sur les compacts...).

Les développements récents en analyse non-linéaire et optimisation ont amené une rupture par rapport à cette vision classique: déjà, l'analyse convexe nous a familiarisé avec l'idée que la convexité d'une fonction se formule naturellement en termes de convexité de son **épigraphe**, et sa semicontinuité inférieure en termes de fermeture de ce dernier. En analyse linéaire, les propriétés des opérateurs se formulent souvent simplement à l'aide de leurs **graphes** (théorème du graphe fermé...). L'analyse des opérateurs à partir de leurs graphes s'est imposée tout naturellement avec la théorie des opérateurs maximaux monotones. Ces exemples nous font sentir le rôle géométrique clé que jouent l'épigraphe pour les fonctions et le graphe pour les opérateurs dans l'analyse des problèmes unilatéraux (minimisation...). Il est naturel de les retrouver au coeur des notions de convergences adaptées à l'analyse variationnelle et l'optimisation.

L'epi-convergence d'une suite de fonctions se définit géométriquement par la convergence de la suite des épigraphes correspondants. La graph-convergence d'une suite d'opérateurs se définit en termes de convergence des graphes correspondants. A la base de ces concepts se trouve donc la notion de **convergence pour une suite d'ensembles**.

L'epi-convergence, la graph-convergence possèdent bien sûr, des versions analytiques qui, contrairement à la convergence simple où la variable est gelée, prennent en compte le comportement des fonctions ou des opérateurs en des points voisins. Ils sont particulièrement utiles lorsque l'on considère des fonctions ou des opérateurs de domaines variables ou lorsque l'on rencontre de fortes variations ou oscillations dans les données d'un problème. Ils ont ainsi permis de traiter de nombreux problèmes de convergences, approximation, perturbation, sensibilité échappant à une analyse utilisant les concepts classiques de convergence basés sur la convergence simple. Ces développements récents se rattachent à un courant mathématique dont nous retraçons brièvement l'histoire.

Un bref historique. La théorie des **applications multivoques** (semicontinues, mesurables), et les questions de différentiation (cônes tangents, dérivées généralisées) ou d'intégration (multivoque) qui s'y rattachent, ainsi que que l'étude des **ensembles aléatoires** et la théorie de la décision statistique ont longtemps été les moteurs principaux de la théorie des convergences d'ensembles et des hypertopologies (topologies sur les fermés d'un espace topologique). Ces concepts se sont progressivement dégagés grâce aux travaux de P. Painlevé (1909), L. Vietoris (1923), G. Choquet (1947), K. Kuratowski (1948), G. Bouligand (1950), E. Michael (1951), J. M. G. Fell (1962), G. Matheron (1975), et, plus directement inspirés par l'étude des ensembles aléatoires et la théorie de la décision statistique, R. A. Wijsman (1964), E. Effros (1965), D. Walkup & R. Wets (1967), B. Van Cutsem (1971). Son champ d'applications a été en s'élargissant avec la théorie des jeux C. Berge (1959), l'économie

mathématique R. Aumann (1964), W. Hildenbrand (1979), J.-P. Aubin (1979), la théorie des points fixes de type Ky-Fan, la viabilité et le contrôle des systèmes dynamiques J.-P. Aubin (1981), B. Cornet (1981), G. Haddad (1981), J.-P. Aubin & A. Cellina (1984), H. Frankowska (1984), l'étude des fonctionnelles intégrales et le contrôle optimal avec R. T. Rockafellar (1975), Ch. Castaing & M. Valadier (1977), Z. Artstein (1972), C. Olech (1976), E. Balder (1984).

Une étape décisive dans le développement de ces concepts a été leur mariage avec l'**analyse variationnelle**. La compréhension des méthodes d'approximation variationnelle (Galerkin, éléments finis) ou de perturbation (en liaison avec la capacité et la théorie du potentiel) a amené U. Mosco en Italie (1969) et J. L. Joly en France (1970) à faire le lien avec les inéquations variationnelles et l'analyse convexe en dimension infinie, et à introduire la **Mosco-convergence**. En liaison avec l'étude de problèmes liés aux hypersurfaces minima, E. De Giorgi dégageait vers 1975 les concepts topologiques généraux de convergence pour des suites de fonctions (non nécessairement convexes), qui permettent de passer à la limite sur les problèmes de minimisation correspondants, à savoir la théorie de la **Γ -convergence**.

Dans cette orientation variationnelle, on trouve un grand nombre de travaux d'**analyse convexe** portant sur les convergences de suites de convexes fermés, de fonctions convexes s.c.i. et de leurs sous-différentiels dans les espaces de Banach. La **bicontinuité de la transformation de Fenchel** relativement à la Mosco epi-convergence dans les espaces de Banach réflexifs est démontrée par Mosco en 1971, cette propriété fondamentale étant en grande partie à l'origine de l'introduction de ce concept. En 1976, la continuité de l'opération de sous-différentiation $f \rightarrow \partial f$ lorsque l'on munit les fonctions convexes s.c.i. de la Mosco epi-convergence et les opérateurs sous-différentiels de la graph-convergence est mise en évidence par H. Attouch. Ce dernier résultat montre clairement le lien entre epi-convergence des fonctions et graph-convergence des opérateurs. Plus récemment, G. Beer montre l'existence d'une topologie induisant la Mosco convergence (1988), qu'il étend ensuite aux Banach généraux sous le nom de **slice convergence**. Outre la continuité de la polarité, une étude systématique des propriétés de continuité des opérations sur les convexes (intersection, addition vectorielle...) et sur les fonctions convexes (addition, inf-convolution...) est entreprise dans les travaux R. C. Bergstrom & L. Mc Linden (1981), H. Attouch & D. Azé & R. Wets, R. T. Rockafellar & R. Wets, D. Azé & J.-P. Penot, G. Beer & R. Lucchetti, M. Volle, J.-P. Aubin & H. Frankowska, S. Dolecki, H. Rihai...

Des progrès décisifs ont été également accomplis dans le cadre non convexe avec l'introduction de versions "localisées" des métriques de type Hausdorff par H. Attouch & R. Wets avec comme corollaires les notions d'**epi-distance** et de **graph-distance** (1988). Une analyse quantitative des questions de convergence et d'approximation pour les problèmes de minimisation, point-selles... est ainsi rendue possible; citons dans cette orientation les travaux de G. Beer, R. Lucchetti, D. Azé & J. P. Penot, D. Pai & P. Shunmugaraj, H. Attouch & J.-L. Ndoutoume & M. Théra, M. Soueycatt, J. Lahrache.

Les travaux récents de G. Beer, R. Lucchetti, Y. Sonntag & C. Zălinescu, A. Lechicki & S. Levi ont permis de classer et de comprendre de façon unifiée les nombreuses convergences d'ensembles et hypertopologies apparues ces dernières années.

Ces travaux théoriques ont été accompagnés de nombreuses **applications**:

a) *Mécanique des milieux continus, calcul des variations et E.D.P.* (problèmes à petits paramètres, **homogénéisation** de milieux composites, couches minces, transition de phase, contrôle et relaxation) avec les travaux en Italie de l'école de Pise animée par E. De Giorgi (G. Buttazzo, G. Dal Maso, P. Marcellini, A. Marino, L. Modica, C. Sbordone, S. Spagnolo), en France de H. Attouch, D. Azé, G. Bouchitte, A. Brillard, A. Damlamian, G. Michaille, F. Murat, C. Picard, P. Suquet, L. Tartar, en Russie de A. V. Marchenko & E. Y. Hruslov et aux USA de B. Kohn.

b) *Optimisation stochastique et théorie de la décision statistique* (approximation numérique, lois des grands nombres pour des ensembles ou des fonctions s.c.i. aléatoires, problème variationnel moyen) avec les travaux de R. Wets, G. Salinetti & R.

Wets, R. T. Rockafellar & R. Wets, Z. Artstein, H. Attouch & R. Wets, Ch. Hess, F. Hiai, P. Kall, Ch. Castaing, A. Choukairi, A. Truffert.

c) *Analyse non régulière, optimisation et contrôle* (cône tangent, dérivées généralisées, epi-dérivées directionnelles du premier et du deuxième ordre, théorème d'inversion locale pour une application multivoque) avec les travaux de J. P. Aubin, A. Auslender, D. Azé, R. Cominetti, L. Contesse & J.-P. Penot, R. Correa, S. Dolecki, H. Frankowska, D. Klatte, P. Michel & J.-P. Penot, J. L. Ndoutoume, D. Noll, J.M. Borwein & D. Noll, J. P. Penot, R. A. Poliquin, S.M. Robinson, R. T. Rockafellar, A. Seeger, M. Soueycatt, L. Thibault.

d) *Analyse numérique et optimisation* (approximation, stabilité et conditionnement numérique, combinaison d'algorithmes avec des méthodes d'approximation, sensibilité, étude de problèmes non coercifs et fonctions de récession) avec les travaux de A. Auslender, A. Auslender & J.-P. Crouzeix, H. Attouch & R. Wets, S. Flam & R. Wets, J.-B. Hiriart-Urruty, B. Lemaire, A. Moudafi, P.-L. Papini, Y. Sonntag, S. Flam & R. Wets, A. Seeger, P. Tossing, T. Zolezzi, méthodes d'homotopie et de continuation en optimisation paramétrique avec les travaux de J. Guddat, Th. Jongen, H. Attouch & J.-P. Penot & H. Rihai.

e) *Principes variationnels* (Ekeland, Borwein-Preiss), géométrie des Banach, problèmes bien posés (au sens de Hadamard, de Tychonov), propriétés génériques et différentiabilité, avec les travaux de R.R. Phelps, G. Beer, R. Deville & G. Godefroy & V. Zizler, R. Lucchetti, J.M. Borwein, J.M. Borwein & S. Fitzpatrick, P. Kenderov, H. Attouch & H. Riahi, F. De Blasi, P.-L. Papini, J. Mijak, J. P. Revalski, S. Simons.

f) *Théorie de l'approximation* où le rôle fondamental de l'approximation épigraphique (inf-convolution) et ses propriétés de régularisation ont été mis progressivement en évidence dans un cadre de plus en plus général: approximation de **Baire-Wijsman** puis **approximation Moreau-Yosida** introduite par J.-J. Moreau et H. Brezis dans le cas de fonctions convexes et un cadre Hilbertien (1973) suivis par les travaux de R. Wets, H. Attouch, J.-P. Penot, M. Bougeard, J.-B. Hiriart-Urruty. Très récemment, l'approximation-régularisation de **Lasry-Lions** a permis de s'affranchir de toute hypothèses de convexité ou de croissance sur les fonctions et ce, en dimension infinie (H. Attouch & D. Azé, M. Volle, J. Benoist).

g) *Convergences de suites d'opérateurs maximaux monotones* ou d'opérateurs accréatifs et des semi-groupes associés. Initialement introduites par H. Brezis, A. Pazy, M. Crandall, Ph. Benilan, H. Attouch, L. Mc Linden ces convergences ont été utilisées dans l'étude de problèmes d'évolution avec opérateurs dépendant du temps par J.-J. Moreau (problème de rafle en élastoplasticité), H. Attouch & A. Damlamian, et en les combinant avec des perturbations multivoques semicontinues par A. Cellina, B. Cornet, Ch. Castaing, N.S. Papageorgiou. Combinée avec l'approximation Yosida, la graph-convergence de suites d'opérateurs maximaux monotones a permis de définir une notion de somme étendue (variationnelle) H. Attouch & J.-B. Baillon & M. Thera.

h) *Théorie des jeux, problèmes de min-max et économie mathématique* où l'étude des questions de stabilité et approximation pour les problèmes de min-max, de points selles ont amené l'introduction de nouvelles notions de convergences, H. Attouch & R. Wets (epi-hypo convergence pour les problèmes de point selle), G. Greco (Γ -convergence), E. Cavazzuti (équilibre de Nash), J. Morgan et P. Loridan (équilibre de Stackelberg), B. Lemaire (optima de Pareto), B. Cornet (cônes tangents et viabilité), S. Simons, S. Flam.

Outre les ouvrages de référence classiques C. Berge (1959), C. Kuratowski (1958), G. Matheron (1975), Ch. Castaing & M. Valadier (1977), plusieurs ouvrages récents traitent des convergences d'ensembles: Y. Sonntag (1982), H. Attouch (Pitman; 1984), E. Klein & A. C. Thomson (John-Wiley; 1984) J.-P. Aubin & H. Frankowska (Birkhäuser; 1990), R. T. Rockafellar & R. Wets, G. Beer, Y. Sonntag & C. Zalinescu, (à paraître).

1. Convergences de suites d'ensembles.

1.1 Convergence au sens de Painlevé-Kuratowski.

Soit (X, τ) un espace topologique, que, pour la simplicité de l'exposé, on supposera métrisable. Etant donnée une suite de parties $A_1, A_2, A_3, \dots$ de l'espace topologique X , les limites inférieures et supérieures (topologiques) de la suite $\{A_n; n \in \mathbb{N}\}$ sont définies par les formules

$$\text{Li } A_n = \{x \in X : \exists (a_n) \rightarrow x \text{ avec pour tout } n \in \mathbb{N}, a_n \in A_n\}$$

$$\text{Ls } A_n = \{x \in X : \exists n(1) < n(2) < n(3) < \dots \text{ et } \forall k \in \mathbb{N} \quad a_k \in A_{n(k)} \text{ avec } (a_k) \rightarrow x\}.$$

En d'autres termes, $\text{Li } A_n$ est l'ensemble formé par toutes les limites possibles de suites $\{a_n; n \in \mathbb{N}\}$ avec $a_n \in A_n$ pour tout n , alors que $\text{Ls } A_n$ est formé par toutes les valeurs d'adhérence de telles suites. Etant donnée une suite $\{A_n; n \in \mathbb{N}\}$, on peut toujours définir ces ensembles $\text{Li } A_n$ et $\text{Ls } A_n$, qui peuvent être éventuellement vides, et l'on a l'inclusion

$$\text{Li } A_n \subset \text{Ls } A_n.$$

Lorsque l'égalité a lieu, on dit que la suite $\{A_n; n \in \mathbb{N}\}$ converge ou plus précisément converge au sens de Painlevé-Kuratowski, et l'on note

$$\text{Lim } A_n = \text{Li } A_n = \text{Ls } A_n.$$

La référence à la topologie τ est implicite dans ces formules; lorsque l'on veut la mettre en évidence on écrit $A = \tau\text{-Lim } A_n$. Dans les formules suivantes "cl" désigne l'opération de fermeture et $\mathcal{N}_\infty^\#$ désigne la grille du filtre de Fréchet $\mathcal{N}_\infty$ sur $\mathbb{N}$:

$$\text{Li } A_n = \bigcap_{N \in \mathcal{N}_\infty^\#} \text{cl} \bigcup_{k \in N} A_k$$

$$\text{Ls } A_n = \bigcap_{N \in \mathcal{N}_\infty^\#} \text{cl} \bigcup_{k \in N} A_k.$$

Ces formules sont intéressantes à double titre: elles mettent clairement en évidence que les ensembles $\tau\text{-Li } A_n$, $\tau\text{-Ls } A_n$ et donc $\tau\text{-Lim } A_n$ (lorsque la limite existe) sont des ensembles fermés pour la topologie τ . De plus, ces limites restent inchangées lorsque l'on remplace A_n par sa fermeture. C'est la raison pour laquelle, sans diminuer la généralité des résultats, on peut raisonner sur des ensembles fermés. Ces formules permettent d'étendre très simplement les concepts précédents aux cas d'ensembles indexés par un paramètre continu $\{A_\varepsilon; \varepsilon \rightarrow 0\}$.

Donnons quelques exemples élémentaires en dimension finie:

a) Une suite de boules $\mathbb{B}(x_n, \rho_n)$ converge vers la boule $\mathbb{B}(x, \rho)$ si et seulement si $\{x_n; n \in \mathbb{N}\}$ converge vers x et $\{\rho_n; n \in \mathbb{N}\}$ converge vers ρ . Si ρ_n tend vers l'infini, alors ces boules convergent vers l'espace tout entier, et leurs complémentaires vers l'ensemble vide.

b) Considérons une suite qui prend alternativement deux valeurs, $A_n = A_0$ pour n pair et $A_n = A_1$ pour n impair, où A_0 et A_1 désignent deux fermés distincts. Alors cette suite ne converge pas, on a $\text{Li } A_n = A_0 \cap A_1$, $\text{Ls } A_n = A_0 \cup A_1$.

c) Dans l'exemple suivant, les ensembles A_n sont des épigraphes: $A_n = \text{epi } f_n$ avec

$$f_n(x) = \begin{cases} 0 & \text{si } x \leq 0 \\ -nx & \text{si } 0 \leq x \leq \frac{1}{n} \\ nx-2 & \text{si } \frac{1}{n} \leq x \leq \frac{2}{n} \\ 0 & \text{si } x \geq \frac{2}{n} \end{cases}$$

La suite des ensembles A_n converge, sa limite est encore un épigraphe à savoir ,

$$\text{Lim } A_n = A = \text{epi } f$$

où f est la fonction qui vaut zéro partout, sauf à l'origine où elle vaut -1 . Cet exemple élémentaire d'épi-convergence illustre bien la différence avec la convergence simple, puisque sur cet exemple la suite $\{f_n; n \in \mathbb{N}\}$ converge simplement vers la fonction identiquement nulle.

d) Considérons à présent une suite d'opérateurs monotones $\mathcal{A}_n$ de $\mathbb{R}$ dans $\mathbb{R}$ et prenons pour ensembles A_n dans $\mathbb{R}^2$ leurs graphes: $A_n = \text{graph } \mathcal{A}_n$ (dans la pratique, on identifie l'opérateur et son graphe qui sont alors désignés par le même symbole).

$$\mathcal{A}_n(x) = \begin{cases} -1 & \text{si } x \leq -\frac{1}{n} \\ nx & \text{si } -\frac{1}{n} \leq x \leq \frac{1}{n} \\ 1 & \text{si } x \geq \frac{1}{n} \end{cases}$$

La suite des ensembles A_n converge et sa limite est l'ensemble $A = \text{graph } \mathcal{A}$ où

$$\mathcal{A}(x) = \begin{cases} -1 & \text{si } x < 0 \\ [-1, +1] & \text{si } x = 0 \\ +1 & \text{si } x > 0 \end{cases}$$

Cet exemple illustre l'approximation Yosida de l'opérateur maximal monotone multivoque $\mathcal{A}$: les opérateurs $\mathcal{A}_n$ sont maximaux monotones lipschitziens, et la convergence a bien lieu en graphe, la convergence simple des opérateurs $\mathcal{A}_n$ ne permettant de récupérer que $\mathcal{A}^0$ la section de norme minimale de l'opérateur $\mathcal{A}$.

Lorsque la topologie τ est métrisable, notant d une distance induisant cette topologie, et posant:

$$d(x, A) = \inf_{y \in A} d(x, y),$$

on peut reformuler les limites topologiques de suites d'ensembles:

$$\text{Li } A_n = \{ x \in X : \limsup_{n \rightarrow \infty} d(x, A_n) = 0 \} = \{ x \in X : \lim_{n \rightarrow \infty} d(x, A_n) = 0 \},$$

$$\text{Ls } A_n = \{ x \in X : \liminf_{n \rightarrow \infty} d(x, A_n) = 0 \}.$$

1.2 Convergence au sens de Wijsman et Mosco convergence.

Les formules ci-dessus suggèrent naturellement la notion de convergence suivante: la suite $\{A_n; n \in \mathbb{N}\}$ converge au sens de Wijsman vers A , on note $A = W\text{-}\lim A_n$, si pour tout $x \in X$,

$$d(x, A) = \lim_{n \rightarrow \infty} d(x, A_n).$$

La convergence au sens de Wijsman entraîne la convergence au sens de Kuratowski-Painlevé. Elle est, en général, strictement plus forte que cette dernière. La convergence au sens de Wijsman permet de topologiser de façon simple la convergence topologique d'ensembles: identifiant un ensemble A à la fonction distance $d(\cdot, A)$, on considère la topologie de la convergence simple des fonctions distances. Si X est un espace métrisable séparable, on obtient ainsi une topologie métrisable séparable sur l'ensemble $\mathcal{F}(X)$ des fermés de X , dont la tribu borélienne est précisément la **tribu d'Effrös** $\mathcal{E}$ (Ch. Hess; 1986). Cette tribu joue un rôle clé dans l'étude des application multivoques mesurables, elle est engendrée par les ensembles

$$U^- = \{ F \in \mathcal{F}(X); F \cap U \neq \emptyset \}$$

où U décrit l'ensemble des ouverts de X . Une application multivoque mesurable n'est autre qu'une application mesurable à valeurs dans $\mathcal{F}(X)$ muni de la tribu $\mathcal{E}$ (R.T. Rockafellar; 1977).

La topologie de Wijsman se révèle d'un usage délicat en dimension infinie car elle dépend du choix de la métrique (ou de la norme) induisant la topologie τ . Lorsque l'on se restreint aux convexes fermés d'un espace de Hilbert et que l'on prend pour d la distance associée à la norme, on obtient la Mosco-convergence:

Soit $\{A_n; n \in \mathbb{N}\}$ une suite de fermés d'un espace normé X . La suite $\{A_n; n \in \mathbb{N}\}$ **Mosco-converge** vers A , et l'on notera $A = M\text{-}\lim A_n$, si on a les deux propriétés suivantes:

(i) pour tout x dans A il existe une suite x_n convergeant **fortement** vers x telle que x_n appartienne à A_n pour tout $n \in \mathbb{N}$.

(ii) pour toute sous-suite $n(1) < n(2) < n(3) < \dots$ et $\forall k \in \mathbb{N} \quad y_k \in A_{n(k)}$ avec y_k convergeant **faiblement** vers y , on a $y \in A$.

Théorème (H. Attouch-Y. Sonntag-M. Tsukada). Soit X un espace de Banach réflexif muni d'une norme strictement convexe et Kadec ainsi que la norme duale (la convergence faible et la convergence des normes entraînent la convergence forte), ce qui est toujours possible après renormage. Soit $\{A_n; n \in \mathbb{N}\}$, A une suite de convexes fermés non vides de X .

Les propositions suivantes sont équivalentes:

(i) la suite $\{A_n; n \in \mathbb{N}\}$ Mosco-converge vers A ;

(ii) $\forall x \in X, \quad d(x, A) = \lim_{n \rightarrow \infty} d(x, A_n);$

(iii) $\forall x \in X, \quad \text{proj}_A x = \lim_{n \rightarrow \infty} \text{proj}_{A_n} x.$

où $\text{proj}_A x$ désigne la projection de x sur A .

La convergence au sens de Mosco joue un rôle clé en analyse variationnelle convexe dans les espaces réflexifs. Nous verrons plus loin ses propriétés remarquables relativement à la dualité. Lorsque l'espace X n'est pas réflexif, elle perd une bonne partie de ses propriétés.

1.3 Convergences de type Hausdorff.

$(X, \|\cdot\|)$ désigne un espace normé de boule unité fermée $\mathbb{B}$. Etant donnés $C, D \subset X$, l'excès de l'ensemble C sur D est défini par

$$e(C, D) := \sup_{x \in C} d(x, D)$$

avec la convention $e = 0$ si $C = \emptyset$. La distance de Hausdorff entre C et D est définie par

$$\text{haus}(C, D) := \max \{ e(C, D), e(D, C) \}.$$

On a la formule suivante:

$$\text{haus}(C, D) = \sup_{x \in X} |d(x, C) - d(x, D)|.$$

Cette métrique qui joue un rôle fondamental en analyse est souvent trop forte lorsque l'on manie des ensembles non bornés tels que épigraphes, cônes, graphes... A cet effet, une version "localisée" a été introduite par H. Attouch & R. Wets (1987):

Pour tout $\rho \geq 0$, la ρ -Hausdorff distance entre C et D est donnée par :

$$\text{haus}_\rho(C, D) := \max \{ e(C_\rho, D), e(D_\rho, C) \}$$

où C_ρ (resp. D_ρ) est défini par $C_\rho := C \cap \rho \mathbb{B}$ (resp. $D_\rho := D \cap \rho \mathbb{B}$).

Une suite $\{C_n \subset X; n \in \mathbb{N}\}$ de parties de X converge au sens des ρ -Hausdorff distances vers un ensemble C , si pour tout $\rho > 0$,

$$\lim_{n \rightarrow \infty} \text{haus}_\rho(C_n, C) = 0.$$

Cette convergence est associée à une topologie métrisable; lorsque l'on travaille avec les fermés non vides d'un espace normé X , on peut prendre comme métrique D

$$D(A, B) := \sum_{n=1}^{+\infty} 2^{-n} \frac{d_n(A, B)}{1 + d_n(A, B)}$$

où

$$d_n(A, B) := \sup_{\|x\| \leq n} |d(x, A) - d(x, B)|.$$

Cette topologie, associée à la convergence uniforme sur les bornés des fonctions distances est donc intermédiaire entre la topologie de Wijsman (associée à la convergence simple des fonctions distances) et la topologie de Hausdorff (associée à la convergence uniforme des fonctions distances). En dimension finie, la topologie de Wijsman coïncide avec la topologie des ρ -Hausdorff distances et ces topologies induisent la convergence au sens de Painlevé-Kuratowski.

1.4 Hypertopologies

L'étude des topologies sur les fermés d'un espace topologique (X, τ) (hypertopologies) induisant les convergences topologiques d'ensembles est un thème central dans toutes les questions évoquées précédemment. Historiquement, ce sont les hypertopologies du type "hit and miss" qui sont apparues les premières. Etant donné $E \subset X$, on introduit

$$E^- = \{ A \in \mathcal{F}(X); A \cap E \neq \emptyset \}$$

$$E^+ = \{ A \in \mathcal{F}(X); A \subset E \}.$$

(1) La topologie de **Vietoris** sur $\mathcal{F}(X)$ a pour sous-base les parties de la forme V^- et les parties de la forme W^+ où V et W sont des ouverts de (X, τ) .

(2) La topologie de **Fell** sur $\mathcal{F}(X)$ a pour sous-base les parties de la forme V^- où V est un ouvert de (X, τ) et les parties de la forme W^+ où W est de complémentaire compact.

(3) Si X est un espace vectoriel normé, la **Mosco** topologie (introduite par G. Beer) sur les fermés faibles de X a pour sous-base les parties de la forme V^- où V est ouvert de X muni de la topologie de la norme et les parties de la forme W^+ où W est de complémentaire faiblement compact.

Cette approche géométrique se marie bien avec la théorie des multiapplications mesurables et l'analyse stochastique. L'approche "moderne" pour définir des hypertopologies est plus fonctionnelle, elle consiste à associer à un ensemble une famille de fonctions et de considérer la topologie initiale qui leurs est associée:

On a vu que la **Wijsman** topologie est la topologie la moins fine (topologie initiale) rendant continue les applications $A \rightarrow d(x, A)$ pour tout x dans X .

La **Slice topologie de Beer** sur les convexes fermés d'un espace normé peut être définie comme topologie initiale associée aux fonctions de saut

$$A \rightarrow D(A, B) := \inf \{ \|x - y\| : x \in A, y \in B \}$$

où B parcourt l'ensemble des convexes fermés bornés de X . Elle est associée à la famille de semi-métriques

$$p_F(A, B) = |D(A, F) - D(B, F)|$$

où F appartient à une famille de parties de X (ici les convexes fermés bornés de X). Y. Sonntag et C. Zalinescu obtiennent une classification des hypertopologies en distinguant celles associées à une famille de semi-métriques du type précédent (le type p) et celles associées à une famille de semi-métriques du type suivant (le type q)

$$q_F(A, B) = \sup_{x \in F} |d(x, A) - d(x, B)|.$$

Dans ce dernier type on reconnaît la métrique de **Hausdorff** en prenant pour F l'espace tout entier, la topologie de **Attouch-Wets** ("bounded Hausdorff topology") lorsque F décrit l'ensemble des parties bornées de X .

2. Epi-convergence de suites de fonctions.

De nombreuses questions liées à la minimisation de fonctions $f: X \rightarrow \mathbb{R} \cup \{+\infty\}$ peuvent être abordées de façon géométrique en identifiant la fonction et son épigraphe

$$\text{epi } f \equiv \{(x, \alpha) : x \in X, \alpha \in \mathbb{R}, \alpha \geq f(x)\}.$$

Les convergences épigraphiques (epi-convergences) de suites de fonctions se définissent géométriquement en termes de convergences topologiques de la suite de leurs épigraphes. On a donc les correspondances suivantes:

$$f = \text{epi-lim } f_n \Leftrightarrow \text{epi } f = \text{Lim epi } f_n$$

$$f = \text{W-epi-lim } f_n \Leftrightarrow \text{epi } f = \text{W-Lim epi } f_n$$

$$f = \text{M-epi-lim } f_n \Leftrightarrow \text{epi } f = \text{M-Lim epi } f_n$$

$$f = \text{epi-dist-lim } f_n \Leftrightarrow \text{epi } f = \text{haus}_\rho\text{-Lim epi } f_n \quad \forall \rho > 0$$

On dira alors que la suite de fonctions $\{f_n; n \in \mathbb{N}\}$ respectivement **epi-converge**, Wijsman epi-converge, Mosco epi-converge, epi-distance converge vers f . Une limite topologique d'épigraphes est encore un épigraphe, plus précisément:

$$\text{Ls epi } f_n = \text{epi} (\text{epi-liminf } f_n)$$

$$\text{Li epi } f_n = \text{epi} (\text{epi-limsup } f_n),$$

avec

$$(\text{epi-liminf } f_n)(x) = \min \{ \liminf f_n(x_n); x_n \rightarrow x \},$$

$$(\text{epi-limsup } f_n)(x) = \min \{ \limsup f_n(x_n); x_n \rightarrow x \}.$$

L'epi-convergence de la suite de fonctions $\{f_n; n \in \mathbb{N}\}$ se traduit par l'égalité entre les deux fonctions $\text{epi-liminf } f_n$ et $\text{epi-limsup } f_n$. On a donc que $\{f_n; n \in \mathbb{N}\}$ epi-converge vers f au point x , si les deux propriétés suivantes sont satisfaites:

- (i) pour toute suite x_n convergeant vers x , $\liminf f_n(x_n) \geq f(x)$
- (ii) il existe une suite $\hat{x}_n$ convergeant vers x telle que $f(x) \geq \limsup f_n(\hat{x}_n)$.

Lorsque l'on veut mettre en évidence la topologie τ prise sur X on note $f = \tau\text{-epi-lim } f_n$.

On peut noter que ces conditions entraînent que f est semicontinue inférieurement. L'epiconvergence n'est pas comparable en général avec la convergence simple et fait intervenir les valeurs des fonctions au voisinage du point considéré. Dans le cas de suites monotones, les deux notions coïncident modulo des opérations de fermeture:

Si la suite $\{f_n; n \in \mathbb{N}\}$ est croissante, alors $\text{epi-lim } f_n = \sup_n (\text{cl } f_n)$

Si la suite $\{f_n; n \in \mathbb{N}\}$ est décroissante, alors $\text{epi-lim } f_n = \text{cl inf}_n (f_n)$.

L'importance de l'epi-convergence tient au fait que cette notion est minimale parmi les notions permettant le passage à la limite sur des suites de problèmes de minimisation. Son utilisation combinée avec les méthodes de compacité est mise en évidence dans l'énoncé suivant où l'on désigne par $\varepsilon\text{-argmin } f = \{x \in X; f(x) \leq \inf f + \varepsilon\}$:

Théorème (E. De Giorgi; 1977). Soit $\{f, f_n: X \rightarrow \mathbb{R} \cup \{+\infty\}; n=1,2,\dots\}$ une suite de fonctions à valeurs réelles étendues. On suppose qu'il existe une suite $(\hat{x}_n; n \in \mathbb{N})$ τ -relativement compacte pour une topologie τ sur X et une suite de réels positifs $\{\varepsilon_n; n \in \mathbb{N}\}$ tendant vers zéro tels que $\hat{x}_n \in \varepsilon_n\text{-argmin } f_n$ pour tout $n \in \mathbb{N}$. Alors, l'epi-convergence pour la topologie τ de la suite $\{f_n; n \in \mathbb{N}\}$ vers f entraîne :

$$(i) \min f = \lim_{n \rightarrow \infty} \inf f_n$$

$$(ii) L_s(\varepsilon_n\text{-argmin } f_n) \subset \text{argmin } f.$$

Dans les applications, cette propriété variationnelle est complétée par le résultat de stabilité:

Si $f = \tau\text{-epi-lim } f_n$ et $g: X \rightarrow \mathbb{R}$ est τ -continue, alors $f + g = \tau\text{-epi-lim } (f_n + g)$.

La topologie τ intervient donc naturellement en liaison avec les propriétés de coercivité, d'inf-compactité de la suite f_n . Le fait de travailler avec des fonctions pouvant éventuellement prendre des valeurs infinies permet de formuler un résultat abstrait de compacité pour l'epi-convergence:

Théorème Soit (X, τ) un espace topologique métrisable séparable. Alors, de toute suite de fonctions $\{f_n: X \rightarrow \overline{\mathbb{R}}; n=1,2,\dots\}$ on peut extraire une sous suite τ -epi-convergente.

L'approximation epigraphique ou inf-convolution joue un rôle central en théorie de l'epi-convergence. Etant donné une fonction $f: X \rightarrow \mathbb{R} \cup \{+\infty\}$ où $(X, \|\cdot\|)$ est un espace normé, pour toute valeur strictement positive du paramètre λ , on définit

$$f_\lambda(x) := \inf_{u \in X} \left\{ f(u) + \frac{1}{2\lambda} \|x - u\|^2 \right\}$$

l'approximation Moreau-Yosida d'indice λ de f , et

$$f_{[\lambda]}(x) := \inf_{u \in X} \left\{ f(u) + \frac{1}{\lambda} \|x - u\| \right\}$$

l'approximation Baire-Wijsman d'indice λ de f . Ces fonctions sont respectivement localement lipschitziennes et lipschitziennes partout définies sur l'espace tout entier, et jouent le rôle des fonctions distances lorsque les ensembles considérés sont des épigraphes. De nombreuses propriétés d'epi-convergence se reformulent donc en termes de convergence simple des approximations épigraphiques. Par exemple, la Mosco epi-convergence d'une suite de fonctions convexes s.c.i. sur un espace de Hilbert est équivalente à la convergence simple de toutes les approximations Moreau-Yosida:

$$f = M\text{-epi-lim } f^n \Leftrightarrow \forall \lambda > 0 \quad \forall x \in X \quad f_\lambda(x) = \lim_{n \rightarrow \infty} (f^n)_\lambda(x).$$

Un résultat du même type pour les approximations Baire-Wijsman a été obtenu par D. Aze.

La suite $\{f_\lambda; \lambda \rightarrow 0\}$ converge simplement et de façon monotone croissante vers f lorsque l'on suppose f s.c.i.. En fait, cette convergence a lieu au sens de l'epi-distance, ce qui donne un contrôle métrique de cette approximation.

3. Epi-convergence de fonctions convexes et dualité.

Soit $(X, \|\cdot\|)$ un espace normé, $\Gamma(X)$ désigne l'ensemble des fonctions convexes semicontinues inférieurement propres sur X , et dualement $\Gamma^*(X^*)$ désigne l'ensemble des fonctions convexes semicontinues inférieurement pour la topologie $\sigma(X^*, X)$, propres définies sur X^* .

Etant donné $f \in \Gamma(X)$, sa *conjuguée* $f^* \in \Gamma^*(X^*)$ est définie par la formule

$$f^*(y) := \sup \{ \langle x, y \rangle - f(x) : x \in X \}.$$

où $\langle x, y \rangle$ désigne le couplage entre $x \in X$ et $y \in X^*$.

Etant donné $h \in \Gamma^*(X^*)$, on adopte la convention habituelle qui consiste à définir h^* seulement sur X plutôt que sur tout X^{**} , de sorte que $h^* \in \Gamma(X)$. La transformation de Fenchel $f \rightarrow f^*$ est alors une involution bijective entre $\Gamma(X)$ et $\Gamma^*(X^*)$. On dit que $y_0 \in X^*$ appartient au sous-différentiel de $f \in \Gamma(X)$ au point $x_0 \in X$, et l'on écrit $y_0 \in \partial f(x_0)$, si pour tout $x \in X$, on a $f(x) \geq f(x_0) + \langle x - x_0, y_0 \rangle$. En termes de fonctions convexes conjuguées, on a que $y_0 \in \partial f(x_0)$ si et seulement si l'égalité de Fenchel est satisfaite : $f(x_0) + f^*(y_0) = \langle x_0, y_0 \rangle$. Le *sous-différentiel* de f est l'opérateur dont le graphe dans $X \times X^*$ est donné par la formule:

$$\begin{aligned} \partial f &:= \{(x, y) \in X \times X^* : y \in \partial f(x)\} \\ &= \{(x, y) \in X \times X^* : f(x) + f^*(y) = \langle x, y \rangle\}. \end{aligned}$$

Rappelons la formulation analytique de la Mosco epi-convergence: Une suite $\{f_n; n=1,2,\dots\}$ de fonctions de $\Gamma(X)$ Mosco epi-converge vers f si en tout point $x \in X$ les deux propriétés suivantes sont satisfaites:

- (i) pour toute suite x_n convergeant **faiblement** vers x , $\liminf f_n(x_n) \geq f(x)$
- (ii) il existe une suite $\hat{x}_n$ convergeant **fortement** vers x telle que $f(x) \geq \limsup f_n(\hat{x}_n)$

Théorème (U. Mosco; 1971) Soit X un espace de Banach réflexif. Alors la transformation de Fenchel $f \rightarrow f^*$ est bicontinue pour la Mosco epi-convergence. En d'autres termes

$$f = M\text{-epi-lim } f_n \Leftrightarrow f^* = M\text{-epi-lim } (f_n)^*.$$

Ce résultat, qui explique le rôle important joué par la Mosco-epi-convergence en analyse convexe, tombe en défaut lorsque l'espace n'est plus réflexif. Pour pallier les défauts de la Mosco convergence dans un cadre non réflexif, G. Beer a introduit récemment la *slice convergence*:

Soit X un espace normé général et $\{f_n; n \in \mathbb{N}\}$ une suite de fonctions de $\Gamma(X)$. On dira que f_n **slice epi-converge** vers f et l'on notera $f = \tau_s\text{-lim } f_n$ si les deux propriétés suivantes sont satisfaites:

$$(ii) \left\{ \begin{array}{l} \forall x_0 \in X \exists \langle x_n \rangle \rightarrow x_0 \text{ tel que } f(x_0) = \lim_{n \rightarrow \infty} f_n(x_n) \\ \forall y_0 \in X^* \exists \langle y_n \rangle \rightarrow y_0 \text{ tel que } f^*(y_0) = \lim_{n \rightarrow \infty} f_n^*(y_n) \end{array} \right.$$

La slice epi-convergence coïncide avec la Mosco epi-convergence lorsque l'espace X est réflexif et la transformation de Fenchel $f \rightarrow f^*$ est bicontinue pour la slice epi-convergence dans les espaces normés généraux (Beer, 1991).

Le résultat suivant fait le lien entre la Mosco epi-convergence d'une suite de fonctions convexes s.c.i. et la graph-convergence de la suite de leurs sous-différentiels.

Théorème (H. Attouch; 1977). Pour toute suite $\{f, f_n: X \rightarrow \mathbb{R} \cup \{+\infty\}; n \in \mathbb{N}\}$ de fonctions convexes s.c.i. propres, où X désigne un espace de Banach réflexif, les propriétés suivantes sont équivalentes:

$$(i) f = M\text{-}epi\text{-}\lim_{n \rightarrow \infty} f_n$$

$$(ii) \text{gph } \partial f = s \times s \lim_{n \rightarrow \infty} \text{gph } \partial f_n \text{ plus la "condition de normalisation (NC)":}$$

La condition de normalisation provient du fait que f est déterminée par ∂f à une constante additive près et s'énonce ainsi:

$$(NC) \quad \begin{cases} \exists (x_0, x_0^*) \in \text{gph } \partial f, \exists (x_{0n}, x_{0n}^*) \in \text{gph } \partial f_n \text{ pour tout } n \in \mathbb{N} \text{ tel que} \\ x_{0n} \xrightarrow{s} x_0, x_{0n}^* \xrightarrow{s} x_0^* \text{ et } f_n(x_{0n}) \rightarrow f(x_0). \end{cases}$$

Ce résultat a été étendu au cas d'espaces de Banach généraux en remplaçant la Mosco epi-convergence par la slice epi-convergence (H. Attouch & G. Beer; 1991).

Les opérations de polarité et de sous-différentiation jouissent également de propriétés de continuité remarquable vis à vis des topologies associées aux ρ -Hausdorff distances:

Théorème (G. Beer; 1990). Soit X un espace normé arbitraire. La transformation de Fenchel $f \rightarrow f^*$ est bicontinue pour la topologie de l'epidistance. En d'autres termes

$$f = \text{epi-dist-lim } f_n \Leftrightarrow f^* = \text{epi-dist-lim } (f_n)^*.$$

Théorème (Attouch-Ndoutoume-Théra; 1991). Soit $\{f, f_n: X \rightarrow \mathbb{R} \cup \{+\infty\}; n \in \mathbb{N}\}$ une suite de fonctions convexes s.c.i. propres, où X désigne un espace de Banach général. Alors, l'implication (i) $\Rightarrow$ (ii) a lieu:

$$(i) f = \text{epi-dist } \lim f_n;$$

$$(ii) \partial f = \text{gph-dist } \lim \partial f_n \text{ plus la condition de normalisation (NC).}$$

$$(NC) \quad \begin{cases} \exists (x_0, x_0^*) \in \text{gph } \partial f, \exists (x_{0n}, x_{0n}^*) \in \text{gph } \partial f_n \text{ pour tout } n \in \mathbb{N} \text{ tel que} \\ x_{0n} \xrightarrow{s} x_0, x_{0n}^* \xrightarrow{s} x_0^* \text{ et } f_n(x_{0n}) \rightarrow f(x_0). \end{cases}$$

Si de plus X est super-réflexif, alors (i) et (ii) sont équivalents.

Bibliographie

- [Ar] Z. Artstein, Distribution of random sets and random selections, *Israel J. Math.* **46**, 313-324, 1983.
- [Ar] ———, A variational convergence that yields chattering systems, *Analyse non linéaire*, eds. H. Attouch, J.-P. Aubin, F. Clarke, I. Ekeland, 49-71, Gauthier-Villars, Paris, 1989.
- [Ar-H] Z. Artstein and S. Hart, Law of large numbers for random sets and allocation processes, *Mathematics of Operations Research* **6**, 482-492, 1981.
- [Ar-V] Z. Artstein and R.A. Vitale, A strong law of large numbers for random compact sets. *The Annals of Probability* **3**, 879-882, 1975.
- [Ar-W] Z. Artstein and R. J.-B. Wets, Approximating the integral of a multifunction, *J. Multivariate Analysis* **24**, 285-308, 1988.
- [At₁] H. Attouch, Convergence de fonctions convexes, des sous-différentiels et semi-groupes associés, *C. R. Acad. Sci. Paris* **284**, 539-542, 1977.
- [At₂] ———, Familles d'opérateurs maximaux monotones et mesurabilité, *Ann. Mat. Pura Appl.*, (4), **120**, 35-111, 1979.
- [At₃] ———, Convergence de fonctionnelles convexes, *Journées d'analyse non linéaire de Besançon*, *Lecture Notes in Mathematics* #665, P. Bénéilan et J. Robert eds., 1977.
- [At₄] ———, *Variational convergence for functions and operators*, *Applicable Maths. Series*, Pitman, London, 1984.
- [At-Az] H. Attouch and D. Azé, Approximation and regularization of arbitrary functions in Hilbert spaces by the Lasry-Lions method, *Séminaire d'analyse convexe de Montpellier*, **20**, exposé n°12, 1990.
- [A-A-W] H. Attouch, D. Azé and R. Wets, Convergence of convex-concave saddle functions: *Annales de l'Institut H. Poincaré, Analyse non linéaire*, **5**, 532-572, 1988.
- [A-A-B] H. Attouch, D. Azé and G. Beer, On some inverse stability problems for the epigraphical sum, *Nonlinear Analysis, Theory and Applications*, **16**, n°3, 241-254, 1991.
- [A-B-T] H. Attouch, J.-B. Baillon and M. Théra, Sum of maximal monotone operators revisited. The variational sum. Working paper, Université de Montpellier, 1992.
- [A-B₁] H. Attouch and G. Beer, On the convergence of subdifferentials of convex functions. *Séminaire d'analyse convexe de Montpellier*, **21**, exposé n°8, 1991.
- [A-B₂] ———, Stability of the geometric Ekeland variational principle: the convex case. *Séminaire d'analyse convexe de Montpellier* **21**, exposé n°11, 1991.
- [A-B-M] H. Attouch, G. Bouchitte and M. Mabrouk, Formulations variationnelles pour des équations elliptiques semi-linéaires avec second membre mesure, *C. R. Acad. Sci. Paris* **306**, 161-164, 1988.
- [A-D] H. Attouch and A. Damlamian, Strong solutions for parabolic variational inequalities. *Journal of Nonlinear Analysis*, **2**, n°3, 329-353, 1978.
- [A-L-W] H. Attouch, R. Lucchetti and R. Wets, The topology of the ρ -Hausdorff distance. *Annali Mat. Pura. Appl.*, (IV), Vol. CLX, 303-320, 1991.
- [A-M-R] H. Attouch, A. Moudafi and H. Riahi, Quantitative stability analysis for maximal monotone operators and semi-groups of contractions, *Séminaire d'analyse convexe de Montpellier*, **21**, exposé n°9, 1991.
- [A-N-T] H. Attouch, J. L. Ndoutoume and M. Théra, Epigraphical convergence of functions and convergence of their derivatives in Banach spaces, *Séminaire d'analyse convexe de Montpellier*, **20**, exposé n°9, 1990.
- [A-P-R] H. Attouch, J.-P. Penot and H. Riahi, The continuation method and variational convergence, in "Fixed point theory and applications", M.A. Théra and J. B. Baillon eds., *Pitman Research Notes in Mathematics* #252, Longman, 1991.
- [A-R] H. Attouch and H. Riahi, The epi-continuous method for minimization problems. Relation with the degree theory of F. Browder for maximal monotone operators, in

- "Partial differential equations and the calculus of variations" Vol I, F. Colombini and al. eds., Birkhäuser, Boston, 29-58, 1989.
- [A-W₁] H. Attouch and R. Wets, Isometries for the Legendre-Fenchel transform, Trans. Amer. Math. Soc., **296**, 33-60, 1986.
- [A-W₂] ———, Quantitative stability of variational systems: I. The epigraphical distance, Trans. Amer. Math. Soc., **328**, n^o2, 695-729, 1991.
- [A-W₃] ———, Quantitative stability of variational systems: II. A framework for nonlinear conditioning, IIASA Working paper 88-9, Laxenburg, Austria, 1988.
- [A-W₄] ———, Epigraphical analysis, *Analyse non linéaire*, eds. H. Attouch, J.-P. Aubin, F. Clarke, I. Ekeland, 73-100, Gauthier-Villars, Paris, 1989.
- [A-W₅] ———, Epigraphical processes: law of large numbers for random lsc. functions, Séminaire d'analyse convexe de Montpellier, **20**, exposé n^o13, 1990. To appear in Annales de l'Institut H. Poincaré, *Analyse non linéaire*.
- [Au] J.-P. Aubin, Lipschitz behaviour of solutions to convex minimizations problems, Math. Operations Research, **9**, 87-111, 1984.
- [Au-C] J.-P. Aubin and A. Cellina, *Differential inclusions*, Springer-Verlag, Grundlehren der Math. Wiss. #264, 1984.
- [Au-E] J.-P. Aubin and I. Ekeland, *Applied Nonlinear Analysis*, J. Wiley-Intersciences, 1984.
- [Au-F] J.-P. Aubin and H. Frankowska, *Set-valued Analysis*, Birkhäuser, 1990.
- [Au-W] J.-P. Aubin and R. Wets, Stable approximations of set-valued maps, Ann. Inst. Henri Poincaré, **5**, n^o 6, 519-535, 1988.
- [Aum] R.J. Aumann, Integrals of set-valued functions, J. Math. Anal. Appl. **12**, 1-12, 1965.
- [Aus] A. Auslender and J.-P. Crouzeix, Well behaved asymptotical convex functions, *Analyse non linéaire*, eds. H. Attouch, J.-P. Aubin, F. Clarke, I. Ekeland, 101-121, Gauthier-Villars, Paris, 1989.
- [Aus-Co] A. Auslender and R. Cominetti, A comparative study of multifunction differentiability with applications in mathematical programming, Math. Operations Research, **16**, n^o 2, 240-258, 1991.
- [Az] D. Azé, Convergences variationnelles et dualité, thèse, Université de Perpignan, 1986.
- [Az-P₁] D. Azé and J.-P. Penot, Recent quantitative results about the convergence of sets and functions, *Functional analysis and approximation*, P.-L. Papini ed., Pitagora Editrice, Bologna, 90-110, 1989.
- [Az-P₂] D. Azé and J.-P. Penot, Operations on convergent families of sets and functions, Optimization, **21**, 4, 521-534, 1990.
- [Ba₁] E. Balder, A unified approach to several results involving integrals of multifunctions, preprint University of Utrecht, n^o 657, April 1991.
- [Ba₂] ———, A general approach to lower semi continuity and lower closure in optimal control theory, SIAM J. Control Optim. **22**, 570-598, 1984.
- [B-G-K-T] B. Bank, J. Guddat, D. Klatte, B. Kummer and K. Tammer, *Nonlinear Parametric Optimization*, Akademie-Verlag, Berlin, 1982.
- [Be₁] G. Beer, On Mosco convergence of convex sets, Bull. Australian Math. Soc. **38**, 239-253, 1988.
- [Be₂] ———, On the Young-Fenchel transform for convex functions, Proc. Amer. Math. Soc. **104**, 1115-1123, 1988.
- [Be₃] ———, Conjugate convex functions and the epidistance topology, Proc. Amer. Math. Soc., **108**, 117-126, 1990.
- [Be₄] ———, Topologies on closed and closed convex sets and the Effros measurability of set valued functions, Séminaire d'analyse convexe Montpellier, **21**, exposé n^o3, 1991.
- [Be₅] ———, The slice topology: A viable alternative to Mosco convergence in nonreflexive spaces, Séminaire d'analyse convexe Montpellier, **21**, exposé n^o2, 1991.

- [BB] G. Beer and J. M. Borwein, Mosco convergence and reflexivity, *Proc. Amer. Math. Soc.* 109 (1990), 427-436.
- [BLLN] G. Beer, A. Lechicki, S. Levi, and S. Naimpally, Distance functionals and the suprema of hyperspace topologies, to appear, *Annali Mat. Pura Appl.*
- [Be-L₁] G. Beer and R. Lucchetti, The epi-distance topology: continuity and stability results with applications to convex optimization problems, to appear in *Math. Oper. Research*.
- [Be-L₂] ———, Convex optimization and the epi-distance topology, *Trans. Amer. Math. Soc.*, 327, n°2, 795-813, 1991.
- [Be-L₃] ———, Weak topologies for the closed subsets of a metrizable space, to appear in *Trans. Amer. Math. Soc.*
- [BP] G. Beer and D. Pai, On convergence of convex sets and relative Chebyshev centers, *J. Approximation Theory*, 62, 147-169, 1990.
- [Ben] Ph. Benilan, Equations d'évolution dans un espace de Banach quelconque et applications, thèse, Orsay, 1972.
- [BF] J. M. Borwein and S. Fitzpatrick, Mosco convergence and the Kadec property, *Proc. Amer. Math. Soc.* 106 (1989), 843-849.
- [Bo-Pr] J. M. Borwein and D. Preiss, A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions, *Trans. Amer. Math. Soc.*, 303, 517-527, 1991.
- [Bou] G. Bouchitte, Convergence et relaxation de fonctionnelles du calcul des variations à croissance linéaire. Application à l'homogénéisation en plasticité, *Annales Faculté des Sciences de Toulouse*, 8, n°1, 7-36, 1986-87.
- [Br] H. Brezis, *Opérateurs maximaux monotones et semi groupes de contractions dans les espaces de Hilbert*, North Holland, 1973.
- [B-C-P] H. Brezis, M.G. Crandall and A. Pazy, Perturbations of nonlinear maximal monotone sets in Banach spaces, *Comm. Pure and Appl. Math.*, 23, 123-144, 1970.
- [Bu₁] G. Buttazzo, Su una definizione generale dei Γ -limiti, *Boll. Un. Mat. Ital.*, 14-B, 722-744, 1977.
- [Bu₂] ———, *Semicontinuity, relaxation ad integral representation in the calculus of variations*, Pitman Research Notes in Math., 207, 1989.
- [CV] C. Castaing and M. Valadier, *Convex analysis and measurable multifonctions*, Springer-Verlag, Lecture Notes in mathematics, Berlin, 580, 1977.
- [C-E] C. Castaing and F. Ezzaki, Variational inequalities for integrands martingales and additive random sequences. Preprint Université de Montpellier.
- [Chb] Z. Chbani, Caractérisation de la convergence d'intégrales définies à partir d'opérateurs elliptiques par la convergence des coefficients, Séminaire d'analyse convexe de Montpellier, 21, exposé n°12, 1991.
- [C-D-D] V. Chiado Piat, G. Dal Maso and Defranceschi, G-convergence of monotone operators *Annales de l'Institut H. Poincaré, Analyse non linéaire* 7, n°3, 123-160, 1990.
- [Cho] G. Choquet, Convergences, *Annales de l'Institut Fourier* 23, 55-112, 1947/48.
- [Chouk] A. Choukairi-Dini, Contribution à l'étude des martingales et au lemme de Fatou multivoque, thèse Université de Montpellier, 1988.
- [Com] R. Cominetti, Metric regularity, tangent sets and second order optimality conditions, *Appl. Math. Opt.* 21, 265-288, 1990.
- [Cor] B. Cornet, Topologies sur les fermés d'un espace métrique, *Cahiers de Mathématiques de la Décision* #7309, Université de Paris-Dauphine, 1973.
- [C-J-T] R. Correa, A. Jofré and L. Thibault: *Characterization of lower semicontinuous convex functions*, *Proc. Amer. Math. Soc.* To appear.
- [Cl] F. Clarke, *Optimization and Nonsmooth Analysis*, Wiley-Interscience, 1983.
- [Da] G. Dal Maso and L. Modica, Nonlinear stochastic homogenization, *Annali di Matematica Pura ed Applicata* 144, 347-389, 1986.
- [De G₁] E. De Giorgi, Sulla convergenza di alcune successioni di integrali del tipo dell'area. *Rendiconti di Matematica* 4, 8, 277-294, 1975.

- [De G₂] _____, Convergence problems for functionals and operators. Proc. Int. Meeting on "*Recent Methods in nonlinear Analysis*", Rome 1978.ed. E. De Giorgi, E. Magenes, U. Mosco. Pitagora, Bologna, 131-188, 1979.
- [D-L] I. Del Prete et M.B. Lignola, On convergence of closed-valued multifunctions, Boll. Un. Mat. Ital., B-6 (1983).
- [DGZ] R. Deville, G. Godefroy and S. Zisler, Un principe variationnel utilisant des fonctions bosses, C.R. Acad. Sci. Paris **312**, série I, 281-286, 1991.
- [D] C. Do, Generalized second-order derivatives of convex functions in reflexive Banach spaces, Preprint 1991.
- [Do₁] S. Dolecki, Tangency and differentiation: some applications of convergence theory, Ann. Mat. Pura e Appl. **130**, 223-235, 1982.
- [Do₂] S. Dolecki et Ch. Maliver, Stability of efficient sets : Continuity of mobile polarities Nonlinear Analysis TMA, **12**, 1464-1486 (1988).
- [DSW] S. Dolecki, G. Salinetti and R. J.-B. Wets, Convergence of functions: equi-semicontinuity, Trans. Amer. Math. Soc., **276**, 409-429, 1983.
- [Fe] J.M.G. Fell, A Hausdorff topology for the closed subsets of a locally compact non-hausdorff space, Proc. Amer. Math. Soc. **13**, 472-476, 1962.
- [Fl] S.D. Fläm, On variational stability in competitive economies, preprint Bergen University, 1992.
- [Fl-Gr] S.D. Fläm and G.H. Greco, Minimax and Intersection Theorems, (J.B. Baillon & M. Théra éd.), Pitman Research Notes in Mathematics n° **252**, 123-140, 1991.
- [Fl-W] S.D. Fläm and R. J.-B. Wets, Existence results and finite horizon approximates for infinite horizon optimization problems, Econometrica, **55**, n°5, 1187-1209, 1987.
- [Fla] J. Flachsmeyer, Verschiedene Topologisierungen im Raum der abgeschlossenen Menglu Math. Nachr. **26**, 321-337, 1964.
- [FLL] S. Francaviglia, A. Lechicki, and S. Levi, Quasi-uniformization of hyperspaces and convergence of nets of semicontinuous multifunctions, J. Math. Anal. Appl. **112**, 347-370, 1985.
- [Gu] J. Guilleme, Convergences et Relations Compactantes. Application aux Equilibres de Stackelberg, à paraître dans Rendiconti di Matematica, Università di Padova.
- [G] G. H. Greco, Théorèmes des minimax locaux et fonctions topologiquement fermées. *Partial differential equations and the Calculus of variations*, Vol. 2, Essays in honor of E. De Giorgi, Colombini, Marino, Modica, Spagnolo eds., Birkhäuser, 1989.
- [He] C. Hess, Contributions à l'étude de la mesurabilité, de la loi de probabilité, et de la convergence des multifonctions, thèse, Montpellier, 1986.
- [Hia] F. Hiai, Strong law of large numbers for multivalued random variables, in "*Multifunctions and integrands*", G.Salinetti ed., Springer Verlag, Berlin, Lecture Notes in Math. **1091**, 160-172, 1984.
- [Hil] W. Hildenbrand, *Core and Equilibria of a large Economy*, North Holland-American elsevier, Amsterdam, 1979.
- [H-U₁] J.-B. Hiriart-Urruty, Lipschitz r -continuity of the approximate subdifferential of a convex function, Math. Scand. **47**, 123-134, 1980.
- [H-U₂] J.-B. Hiriart-Urruty, " ϵ -subdifferential calculus" in Research Notes in Math., Pitman **57**, 1982.
- [H-U₃] J.-B. Hiriart-Urruty and A. Seeger, The second order subdifferential and the Dupin indicatrices of a nondifferentiable convex function, Proc. London Math. Soc **58**, 351-365, 1989.
- [J] J.L. Joly, Une famille de topologies et de convergences sur l'ensemble des fonctionnelles convexes, thèse faculté des sciences de Grenoble, 1970.
- [Jo] H. Th. Jongen and G.-W. Weber, On parametric nonlinear programming, Annals of Operations research, **27**, 253-284, 1990.
- [Ka] T. Kato, *Perturbation theory for linear operators*, Springer, Berlin, 1966.
- [Ki] A.J. King, R.J.-B. Wets, Epi-consistency of convex stochastic programs, Stochastics and Stochastics Reports, **34**, 83-92, 1990.
- [KL] E. Klein and A.C. Thompson, *Theory of Correspondences, Including applications to Mathematical Economics*, John-Wiley & Sons, New York, 1984.
- [Ku] K. Kuratowski, *Topology, vol 1.*, Academic Press, New York, 1966.

- [La] J. Lahrache, Stabilité et convergence dans les espaces non réflexifs, Séminaire d'analyse convexe Montpellier, **21**, exposé n°10, 1991.
- [LL] A. Lechicki and S. Levi, Wijsman convergence in the hyperspace of a metric space, *Bull. Un. Mat. Ital.* **5-B**, 435-452, 1987.
- [Lem₁] B. Lemaire, Coupling optimization methods and variational convergence, *trends in mathematical optimization*. Series of Num. Math., K.H. Hoffmann, J.-B. Hiriart-Urruty, C. Lemarechal, J. Zowe eds., Birkhäuser, **84**, 163-179, 1987.
- [Lem₂] ———, Approximation in multiobjective optimization, *Journal of Global Optimization* **2**, 117-132, 1992.
- [Lo] P. Loridan and J. Morgan, New results on approximate solutions in two level optimization, *Optimization*, **20**, 819-836, 1989.
- [Ma] A. Marino, The calculus of variations and some semilinear variational inequalities of elliptic and parabolic type, *Partial differential equations and the Calculus of variations*, Vol. 2, Essays in honor of E. De Giorgi, Colombini, Marino, Modica, Spagnolo eds., Birkhäuser, 787-822, 1989.
- [Mat] M. Matzeu, Su un tipo di continuità dell' operatore subdifferenziale, *Boll. U.M.I.* **11**, 107-124, 1975.
- [Mc] L. McLinden, Successive approximation and linear stability involving convergent sequences of optimization problems, *J. Approximation Theory*, **35**, 311-354, 1982.
- [Me-Mi] K. Messaoudi and G. Michaille, Stochastic homogenization of non convex integral functionals. Duality in the convex case. Séminaire d'analyse convexe de Montpellier, **21**, exposé n°14, 1991.
- [Mi] E. Michael, *Topologies on spaces of subsets*, *Trans. Amer. Math. Soc.* **71**, 152-182, 1951.
- [Mor] J.-J. Moreau, On unilateral constraints, friction and plasticity, *C.I.M.E. New variational technics in Math. Physics*, Bressanone. Ed. Cremonese, Rome, 1974.
- [Mo₁] U. Mosco, Convergence of convex sets and of solutions of variational inequalities, *Advances in Math.* **3**, 510-585, 1969.
- [Mo₂] ———, On the continuity of the Young-Fenchel transform, *J. Math. Anal. Appl.* **35**, 518- 535, 1971.
- [Mol] K. Mouallif, Convergence variationnelle et méthodes perturbées pour les problèmes d'optimisation et de point selle, thèse d'état, Université de Liège, 1989.
- [Mod] A. Moudafi, Perturbation de la méthode de l'inverse partiel. Publication Avamac 87-05, Université de Perpignan, Vol. II, 1987.
- [Mou] M. Moussaoui, Approximations lipschitziennes de multifonctions et applications à la relaxation de certains problèmes de contrôle optimal et de calcul des variations, thèse Université de Montpellier, 1990.
- [N] J. L. Ndoutoume, Epi-convergence en calcul différentiel généralisé. Résultats de convergence et approximation, thèse, Université de Perpignan, 1987.
- [N-T] J. L. Ndoutoume and M. Théra, Generalized second order derivatives of convex functions in locally convex vector spaces, preprint 1992.
- [No] D. Noll, Generalized second fundamental form for Lipschitzian Hypersurfaces by way of second epi derivatives, to appear in *Bull. Can. Math.*
- [Pe₁] J.-P. Penot, Topologies and convergences on the space of convex functions. Preprint Univ. Pau., to appear, *Nonlinear Analysis*, T.M.A.
- [Pe₂] ———, The cosmic Hausdorff topology, the bounded-Hausdorff topology and continuity of polarity, *Proc. Amer. Math. Soc.*, **113**, 275-285, 1991.
- [Ph] R. Phelps, Convex functions, monotone operators, and differentiability, *Lecture Notes in Mathematics* #1364, Springer-Verlag, Berlin, 1989.
- [Po₁] R. A. Poliquin, An extension of Attouch's theorem and its application to second-order epi-differentiation of cconvex composite functions, to appear *Trans. Amer. Math. Soc.*
- [Po₂] ———, Subgradient monotonicity and convex functions, *Nonlinear Analysis TMA.*, **14** n° 4, 305-317, 1990.
- [Ria] H. Riahi, Quelques résultats de stabilité en analyse épigraphique: approche topologique et quantitative, thèse, Université de Montpellier, 1989.
- [Ric] B. Ricceri, Fixed points of lower semicontinuous multifunctions and applications: alternative and minimax theorems, *Memorie di Matematica*, **9**, 331-338, 1985.

- [Rob₁] S. Robinson, Local epi-continuity and local optimization, MRC University of Wisconsin Madison, 1985.
- [Rob₂] ———, Regularity and stability for convex multivalued functions, *Math. Oper. Res.* **1**, 130-143, 1976.
- [Rob₃] ———, An implicit-function theorem for nonsmooth functions, *Math. Oper. Res.* **16**, n° 2, 292-309, 1991.
- [Ro₁] R.T. Rockafellar, "Integral functionals, normal integrands and measurable selections" in *Nonlinear operators and the Calculus of Variations*, Springer-Verlag Lecture Notes in Math., **543**, 157-207, 1976.
- [Ro₂] ———, Maximal monotone relations and the second derivatives of nonsmooth functions, *Annales de l'Institut H. Poincaré, Analyse non linéaire*, **2**, 167-184, 1985.
- [Ro₃] ———, First and second-order epi-differentiability in nonlinear programming, *Trans. Amer. Math. Soc.* **307**, 75-108, 1988.
- [Ro₄] ———, Generalized second derivatives of convex functions and saddle functions, *Trans. Amer. Math. Soc.* **322**, 51-77, 1990.
- [Ro₅] ———, Second order optimality conditions in non-linear programming obtained by way of epi-derivatives, *Math. Oper. Res.* **14**, 462-484, 1989.
- [SW] G. Salinetti and R. Wets, On the relation between two types of convergence for convex functions, *J. Mathematical Analysis and Applications* **60**, 211-226, 1977.
- [Sc] R. Schultz, Estimates for Kuhn-Tucker points of perturbed convex programs, *Optimization* **19** (1), 29-43, 1988.
- [Se₁] A. Seeger, Analyse du second ordre de problèmes non différentiables, thèse Université P. Sabatier (Toulouse) 1986.
- [Se₂] ———, Limiting behaviour of the approximate second-order subdifferential of a convex function, *J. Optim. Th. Appl.* **74**, 527-544, 1992.
- [S-P] P. Shunmugaraj and D.V. Pai, On stability of approximate solutions of minimization problems, W.P. Indian Institute of Technology, Bombay, to appear in *Numerical Functional Analysis and Optimization*.
- [Si] S. Simons, The occasional distributivity of ? over ? and the change of variable formula for conjugate functions, *Nonlinear Analysis, Theory, Methods & Applications*, **14**, n° 12, 1111-1120, 1990.
- [So] Y. Sonntag, Convergence au sens de Mosco; théorie et applications à l'approximation des solutions d'inéquations, thèse de l' Université de Provence, Marseille, 1982.
- [SZ] Y. Sonntag and C. Zalinescu, Set convergences : An attempt of classification, in *Proceedings of Intl. Conf. on Diff. Equations and Control Theory*, Iasi, Romania, August, 1990.
- [Sou] M. Soueycatt, Analyse epi/hypo-graphique des problèmes de points selles, thèse Université de Montpellier, 1991.
- [Sp] S. Spagnolo, Sulla convergenza di soluzioni di equazioni paraboliche ed ellittiche, *Ann. Scu. Norm. Pisa*, **22**, 1968.
- [Th] L. Thibault, Subdifferentials of nonconvex vector-valued functions, *J. Math. Anal. Appl.*, **86**, 319-344, 1982.
- [To] P. Tossings, Convergence variationnelle et opérateurs maximaux monotones, W.P. No. **90014**, Université de Liège, 1990.
- [Tr] A. Truffert, Conditional expectations of integrands and random sets, *Annals of Operational Research* **30**, 117-156, 1991.
- [Ts] M. Tsukada, Convergence of best approximations in a smooth Banach space, *J. Approximation Theory* **40** (1984), 301-309.
- [Val₁] M. Valadier, Young measures, in "Methods of convex analysis" (ed. A. Cellina), *Lecture Notes in Math.* **1446**, Springer Verlag, Berlin, 152-188, 1990.
- [Val₂] ———, Sur l'espérance conditionnelle multivoque non convexe, *Ann. Inst. H. Poincaré, Sec. B* **16**, 1980.
- [Van] B. Van Cutsem, Eléments aléatoires à valeurs convexes compactes, thèse Grenoble 1971.

- [Vi] L. Vietoris, Kontinua Zweiter Ordnung, Monatshefte für Mathematik und Physik, Vol. 33, 49-62, 1923.
- [Vo1] M. Volle, Convergences en niveau et en épigraphes, C. R. Acad. Sci. Paris 299, 295-298, 1984.
- [Vo2] _____, Conjugaison par tranches, Ann. di Mat. pura ed applicata IV, 139, 279-312, 1985.
- [Wa-We] D. Walkup and R. Wets, Continuity of some convex-cone valued mappings, Proc. Amer. Math. Soc. 18, 229-235, 1967.
- [We1] R. Wets, Convergence of convex functions, variational inequalities and convex optimization problems, in "Variational inequalities and complementary problems", eds. P. Cottle, F. Gianessi, J.-L. Lions, Wiley, Chichester (UK), 375-403, 1980.
- [We2] _____, A formula for the level sets of epi-limits and some applications, in "Mathematical theory of optimization", eds. J.P. Cecconi and T. Zolezzi, Springer Verlag, Lecture Notes in Math 979, 256-268, 1983.
- [Wi] R. Wijsman, Convergence of sequences of convex sets, cones, and functions, II, Trans. Amer. Math. Soc. 123, 32-45, 1966.
- [Zo] T. Zolezzi, On stability analysis in mathematical programming, North-Holland, Mathematical Programming Study, 21, 227-242, 1984.

Programme des Conférences

Lundi 22 juin

Président de séance: R. J.-B. Wets (Davis)

9h-10h G.A. Beer (Los Angeles)
10h-10h30 S. Levi (Milan)

Pause Café

Président de séance: D. Klatte (Zürich)

10h45-11h15 J.-P. Penot (Pau)
11h15-11h45 R. Lucchetti (Milan)
11h45-12h15 Y. Sonntag (Marseille)-**C. Zălinescu** (Iasi)

Déjeuner

Président de séance: S. Simons (Santa Barbara)

14h30-15h30 R. J.-B. Wets (Davis)
15h30-16h Ch. Hess (Paris 9)

Pause Café

Président de séance: M. Valadier (Montpellier)

16h15-16h45 S. Gautier (Pau)-**L. Thibault** (Pau)
16h45-17h15 E. Blum (Lima)-**W. Oettli** (Mannheim)

Pause

Président de séance: R. Correa (Santiago)

17h30-18h P. Loridan (Dijon)-**J. Morgan** (Naples)
18h-18h30 A. Jofré (Santiago)

Mardi 23 juin

Président de séance: J.-P. Penot (Pau)

9h-10h R. T. Rockafellar (Seattle)
10h-10h30 **R. Cominetti**-J. San Martin (Santiago)

Pause Café

Président de séance: A. Marino (Pise)

10h45-11h15 **J.-L. Ndoutoume** (Libreville)-M. Théra (Limoges)
11h15-11h45 D. Noll (Stuttgart)
11h45-12h15 R. Poliquin (Edmonton)

Déjeuner

Président de séance: A. Damlamian (Paris)

14h30-15h30 A. Cellina (Trieste)
15h30-16h G. Bouchitté (Toulon)

Pause Café

Président de séance: J.-B. Baillon (Lyon)

16h15-16h45 H. Attouch (Montpellier)-**D. Azé** (Perpignan)
16h45-17h15 J. Benoist (Limoges)

Pause

Président de séance: G. Godefroy (Paris)

17h30-18h Th. Jongen (Aachen)
18h-18h30 H. Riahi (Marrakech)
18h30-19h L. Qi (Sydney)

Mercredi 24 juin

Président de séance: R. T. Rockafellar (Seattle)

9h-10h E. De Giorgi (Pise)
10h-10h30 **R. Deville** (Besançon)

Pause Café

Président de séance: S. M. Robinson (Madison)

10h45-11h15 A. Marino (Pise)
11h15-11h45 H. Attouch (Montpellier)-**J.-B. Baillon** (Lyon 1)-M. Théra
(Limoges)

Pause

Président de séance: L. Thibault (Pau)

12h-12h30 S. Simons (Santa Barbara)
12h30-13h B. Ricceri (Catane)

Déjeuner

14h Excursion dans les calanques de Cassis

Soirée bouillabaisse

Jeudi 25 juin

Président de séance: J.-B. Hiriart-Urruty (Toulouse)

9h-10h S. M. Robinson (Madison)
10h-10h30 A. Auslender (Clermont-Ferrand)

Pause Café

Président de séance: Th. Jongen (Aachen)

10h45-11h15 M. A. Bahraoui-**B. Lemaire** (Montpellier)
11h15-11h45 A. Seeger (Avignon)
11h45-12h15 D. Klatte (Zurich)

Déjeuner

Président de séance: G. Bouchitté (Toulon)

14h30-15h30 U. Mosco (Rome)
15h30-16h **J.-B. Hiriart-Urruty** (Toulouse)-R. R. Phelps (Seattle)

Pause Café

Président de séance: T. Zolezzi (Gênes)

16h15-16h45 P.-L. Papini (Bologne)
16h45-17h15 M. Volle (Avignon)

Pause

Président de séance: W. Oettli (Mannheim)

17h30-18h S. J. Flam (Bergen)
18h-18h30 J.-E. Martinez-Legaz (Barcelone)
18h30-19h **J.-P. Crouzeix**-R. Kebbour (Clermont-Ferrand)

Vendredi 26 juin

Président de séance: A. Cellina (Trieste)

9h-10h E. Balder (Utrecht)
10h-10h30 M. Valadier (Montpellier)

Pause Café

Président de séance: A. Auslender (Clermont-Ferrand)

10h45-11h15 **R. Correa**-A. Jofré (Santiago)-L. Thibault (Pau)
11h15-11h45 T. Zolezzi (Gênes)
11h45-12h15 J. P. Revalski (Sofia)

Déjeuner

Président de séance: P.-L. Papini (Bologne)

14h30-15h30 Z. Artstein (Rehovot)
15h30-16h **M. Bougeard** (Paris 10)-Ch. Michelot (Paris 1)

Fin du colloque

Table des matières /Contents

Arstein, Zvi Differentiating set-valued maps.	1
Attouch, Hedy - Azé, Dominique Approximation and Regularization of Arbitrary Functions in Hilbert Spaces by the Lasry-Lions Method.	2
Attouch, Hedy - Baillon, Jean-Bernard - Théra, Michel Sum of Maximal Monotone Operators revisited: The variational sum.	5
Auslender, Alfred Convergence of stationary sequences for variational inequalities with maximal monotone operators.	7
Balder, Erik A survey of Young measure theory and some applications.	8
Beer, Gerald Wijsman convergence of convex sets: a retrospective.	10
Benoist, Joël Approximation and regularization of arbitrary sets in finite dimension.	12
Blum, E. - Oettli, Werner Asymptotic Cones, Closed Images and a Theorem of Dieudonné.	14
Bouchitté, Guy Equicoercivité de problèmes variationnels. Le rôle des fonctions de récession.	16
Bougeard, Mireille - Michelot, Christian About the proximal determination of Huber's estimators .	17
Cellina, Arrigo A Eulogy on Nonconvexity.	20
Cominetti, Roberto -San Martin Jaime Convergence issues in penalty methods. Linear programming for instance.	21
Corradini, Pietro Poggi Suprema of Wijsman topologies in normed spaces.	23
Costantini, Camillo A splitting property of the upper bounded-Hausdorff convergence	25
Correa, Raphael Subdifferential monotonicity as a characterization of convex functions.	26

Crouzeix, Jean-Pierre-Kebbour, R Index multiplicatifs de convexité/concavité et applications.	28
De Giorgi, Ennio Titre à préciser.	29
Deville, Robert Stability of subdifferentials of non convex functionals.	30
Dinh-The Luc - Luchetti, Roberto - Malivert, Christian On the convergence of efficient Sets.	31
Flam, Sjur On variational stability in competitive economies.	33
Gautier, Segre - Thibault, Lionel Viability for constrained stochastic differential equations .	36
Hess, Christian Multivalued strong law of large numbers in the Wijsman topology and the slice topology.	37
Hiriart-Urruty, Jean-Baptiste -Phelps, Robert Subdifferential calculus without "Qualification" conditions.	39
Jalby, Vincent Lipschitz Approximation of Lattice Valued Functions	41
Jofré, Alejandro On b-subgradients and applications.	43
Jongen, Bert Semi-infinite optimization: structure and stability of the feasible set.	45
Klatte, Diethard Metric Regularity and Stability in semi-infinite optimization.	46
Lemaire, Bernard - Bahraoui, M. A. Convergence of diagonally stationary sequences in Convex Optimization.	48
Levi, Sandro Some new results on Wijsman topologies.	50
Loridan, Pierre - Morgan, Jacqueline Weak two-level optimization problems and Tykhonov regularization.	52
Lucchetti, Roberto Hypertopologies: old and new approaches.	53
Marino, Antonio Bifurcation for variational inequalities with constraints on the derivatives.	54
Martinez-Legaz, Juan-Enrique Dual representation of cooperative games.	57
Michaille, Gérard - Chabi E. Ergodic theory and the calculus of variation.	58
Mosco, Umberto Variational topological properties of the space of Dirichlet forms.	60

Ndoutoume, James - Théra, Michel Generalized second -order derivatives of convex functions in locally convex topological vector spaces.	61
Noll, Dominik Graphical convergence and generalized second derivative for nonsmooth functions.	63
Papini, Pier-Luigi Approximation by convex functions.	65
Penot, Jean-Paul Convergent and divergent concepts from convergence theory and their uses in optimization theory.	66
Poliquin, René Calculus of epi-derivatives and proto-derivatives.	68
Qi, Lijun Solving Nonsmooth equations via Generalized Jacobians and Iteration Functions.	70
Revalski, Julian Necessary and sufficient conditions for the existence of densely defined selections of multivalued mappings.	72
Riahi, Hassan Topological degree for maximal monotone operators.	74
Ricceri, Biagio Some topological Min-Max theorems via an alternative principle for multifunctions.	77
Robinson, Stephen Graph-convergence in mathematical programming.	79
Rockafellar, Tyrrel Proto-differentiability of solution mappings in optimization.	81
Seeger, Alberto Approximate Euler-Lagrange inclusion, approximate transversality condition, and sensitivity analysis of convex parametric problems of calculus of variations.	82
Simons, Stephen Swimming below icebergs, and the maximal monotonicity of subdifferentials.	83
Sonntag, Yves - Zălinescu, Constantin Set convergences: a survey and a classification.	85
Torre, Anna A new Topology for the Solid Sets of a Topological Space.	87
Valadier, Michel Strong convergence implied by weak in L^1 .	89
Vitolo, Paolo Remarks about the Infimum of Hausdorff Metric Topologies.	91
Volle, Michel The use of monotone norms in convex analysis.	92

Wets, Roger J.-B
Convergence of Integral Functionals.

93

Zolezzi, Tullio
On the convergence of the generalised gradients.

95

Differentiating set-valued maps

Zvi Artstein

The Weizmann Institute Rehovot 76100, Israel

Abstract

Derivatives of compact set-valued maps will be defined, with the goal of developing a calculus suitable to express evolutions of sets, and eventually differential equations for set evolution. The motivation arises in controlled differential inclusions. The talk will compare the suggested derivative to the available notions of derivatives for multifunctions

Approximation and Regularization of Arbitrary Functions in Hilbert Spaces by the Lasry-Lions Method

Hedy Attouch

U. S. T. L., 34095 Montpellier Cedex 5, France

and

Dominique Azé

Université de Perpignan 66860 Perpignan Cedex, France

Abstract

The Lasry-Lions's regularization method is extended to arbitrary functions. In particular, to any proper lower semicontinuous function $f : X \rightarrow \mathbb{M} \cup \{+\infty\}$ defined on a Hilbert space X and which is quadratically minorized (i.e. $f(x) \geq -c(1 + \|x\|^2)$ for some $c \geq 0$), is associated a sequence of differentiable functions with Lipschitz continuous derivatives which approximate f from below. Some variants of the method are considered including the case of non quadratic kernels.

Approximation methods play an important role in nonlinear analysis. A number of problems in variational analysis and in optimization theory give rise to nonsmooth functions with possibly infinite values defined on finite or infinite dimensional spaces. By using a regularization procedure based on the infimal convolution or epigraphical sum (see [1]),

these problems can be attacked with the help of classical analysis tools. Let us mention in this direction the pioneering works of K. Yosida [6], H. Brézis [2], J.-J. Moreau [5]. These authors deal with convex lower semicontinuous functions in Hilbert spaces and with the corresponding subdifferential operators which are maximal monotone. The regularized function is proved to be $C^{1,1}$ (continuously differentiable with Lipschitz continuous gradient). Some direct extensions have been recently obtained in [1] for more general kernels than the square of the norm. A difficult problem is to *extend these results to the non convex case*. A decisive step in this direction has been done recently by J.-M. Lasry and P.-L. Lions in [4]. They were motivated by the study of the Hamilton-Jacobi equations and worked with bounded uniformly continuous functions. In [3] Theorem 2.6, boundedness and uniform continuity assumptions are removed: the approximation/regularization result is obtained assuming that the absolute value of the function is quadratically majorized. Our main results state that, given any quadratically minorized function f defined on a Hilbert space X with values in $\mathbb{R} \cup \{+\infty\}$, the function $(f_\lambda)^\mu$ defined by the formula

$$(f_\lambda)^\mu(x) := \sup_{y \in X} \inf_{u \in X} \left\{ f(u) + \frac{\|u - y\|^2}{2\lambda} - \frac{\|y - x\|^2}{2\mu} \right\}$$

is $C^{1,1}$ whenever $0 < \mu < \lambda$ and approaches f from below as the parameters λ and μ go to 0. Observe that our growth assumption on f allows to treat the case of an indicator function. Clearly, by exchanging the order of the inf-sup operations, one obtains a corresponding approximation from above. The paper is organized with respect to increasing generality. We consider successively the convex, then the convex up to a square case, and finally the general case. A natural question that arises concerns the flexibility of the method. The case of non quadratic kernels is also considered. These results open new perspectives in nonsmooth analysis and ask for further developments: one may think defining generalized derivatives by relying on these approximation techniques. Regularization of sets can be considered too by using their indicator functions.

References

- [1] H. Attouch, R.J.B. Wets, Epigraphical analysis, in *Analyse non Linéaire*, eds. H. Attouch, J.-P. Aubin, F.H. Clarke, I. Ekeland, 73-100, Gauthier-Villars Paris et C.R.M. Montreal (1989).
- [2] M. Bougeard, J.-P. Penot and A. Pompelet, Towards minimal assumptions for the minimal convolution regularization, *J. of Approx. Theory*, 64, (1991), 245-270
- [3] H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland, Amsterdam, (1973).
- [4] J.-M. Lasry, P.-L. Lions, A remark on regularization in Hilbert spaces, *Israel Journal of Mathematics*, 55, (1986), 257-266.

- [5] J.-J. Moreau, Fonctionnelles convexes, Lecture notes, Collège de France, Paris, (1967).
- [6] K. Yosida, Functional analysis, third edition, Springer, Berlin, Heidelberg, New York, (1971).

Sum of Maximal Monotone Operators revisited: The variational sum

Hedy Attouch

U. S. T. L., 34095 Montpellier Cedex 5, France

Jean-Bernard Baillon

Université Lyon I, 69622 Villeurbanne Cedex, France and

Michel Théra

Université de Limoges, 87060 Limoges Cedex, France

Abstract

The sum of (nonlinear) maximal monotone operators is reconsidered from the Yosida approximation and graph-convergence point of view. This leads to a new concept, called *variational sum*, which coincides with the classical (pointwise) sum when the classical sum is maximal monotone. In the case of subdifferentials of convex functions, the variational sum is equal to the subdifferential of the epigraphical sum (inf-convolution) of the functions. A general feature of the variational sum is to involve not only the values of the two operators at the given point but also their values at nearby points.

References

- [1] H. Attouch, On the maximality of the sum of two maximal operators. *Nonlinear Analysis, Theory, Methods and Applications*, Vol.5, N°2, pp. 143-147, 1981.
- [2] H. Attouch, *Variational Convergence for Functions and Operators*, Applicable Maths. Series, Pitman, London, 1984.
- [3] H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland, Amsterdam, (1973).
- [3] H. Brézis, M. G. Crandall, A. Pazy, Convergence and Approximation of non-linear semi-group in Banach spaces, *J. Funct. Anal.* pp. 63-74, 1971.
- [4] Ph. Clément, P. Egberts, On the sum of two maximal monotone operators. *Differential and Integral Equations*, Vol3, N°6, pp. 1127-1138, 1990.
- [5] M. Lapidus, *Formule de Trotter et calcul opérationnel de Feynman*, Thèse de Doctorat d'Etat, Université Paris 6, 1986.

CONVERGENCE OF STATIONARY SEQUENCES FOR VARIATIONAL INEQUALITIES WITH MAXIMAL MONOTONE OPERATORS

A. AUSLENDER

Department of Applied Mathematics
University Blaise Pascal
63170 Aubière Cedex, France

Abstract. Let T be a maximal monotone operator defined on $\mathbb{R}^N$. In this paper we consider the associated variational inequality : $0 \in T(x^*)$ and stationary sequences $\{x_k^*\}$ for this operator, i.e., satisfying $T(x_k^*) \rightarrow 0$. The aim of this paper is to give sufficient conditions ensuring that these sequences converge to the solution set $T^{-1}(0)$ especially when they are unbounded. For this we generalize and improve the directionally local boundedness Theorem of Rockafellar to maximal monotone operators T defined on $\mathbb{R}^N$.

A survey of Young measure theory and some applications

Erik J. Balder

Mathematisch Instituut, Rijksuniversiteit Utrecht, Netherlands

Abstract

A survey will be given of Young measure theory and some of its applications. As for the theory, it has been discovered in the past decade that Young measure theory forms an extension of the classical weak convergence theory for probability measures on a topological space [Ba1,Ba2,Va], with a useful reinforcement in the form of K -convergence [Ba3,Ba4,Ba5]. As for the applications, I shall limit myself mainly to discussing Fatou-type lemmas [Ba1,Ba6,BaHe] and results à la Visintin [Ba7,Ca,Ba5,Ba8].

References

- [Ba1] E.J. Balder, A general approach to lower semicontinuity and lower closure in optimal control theory, SIAM J. Control Optim. **22** (1984), 570-598.
- [Ba2] —, On Prohorov's theorem for transition probabilities, Travaux Sémi. Anal. Convexe **19** (1989), 9.1-9.11.
- [Ba3] —, Unusual applications of a.e. convergence: *Almost Everywhere Convergence* (G. Edgar and L. Sucheston, eds.), Academic Press, New York, 1989, pp. 31-53.
- [Ba4] —, New sequential compactness results for spaces of scalarly integrable functions, J. Math. Anal. Appl. **151** (1990), 1-16.
- [Ba5] —, On equivalence of strong and weak convergence in L_1 -spaces under extreme point conditions, Isr. J. Math. **75** (1991), 21-47.
- [Ba6] —, Fatou's lemma in infinite dimensions, J. Math. Anal. Appl. **136** (1988), 450-465.

[Ba7] —, On weak convergence implying strong convergence in L_1 -spaces, Bull. Austral. Math. Soc. **33** (1986), 363-368.

[Ba8] —, A unified approach to several results involving integrals of multifunctions, forthcoming.

[BaHe] — and C. Hess, Fatou's lemma for multifunctions with unbounded values, preprint, May 1992.

[Ca] C. Castaing, Convergence faible et sections extrémales, Travaux Sém. Anal. Convexe **18** (1988), 2.1-2.18.

[Va] M. Valadier, Young measures, in: Methods of Nonconvex Analysis (A. Cellina, ed.), Lecture Notes in Math. **1446**, Springer-Verlag, Berlin, 1990, pp. 152-188.

Wijsman Convergence of Closed and Convex Sets: an Overview

G. Beer

California State University, Los Angeles Ca 90032 USA

Abstract

A natural way to define convergence for a sequence (or net) of closed sets in a metric space is to insist that the associated sequence (or net) of distance functionals converges pointwise to the distance functional of the limit set. This point of view stems from the seminal paper of R. Wijsman on convergence and convex duality, and has since been developed by numerous authors. In this talk, we give an overview of the major results involving this mode of convergence, indicating connections with convex analysis, Banach space geometry, and measurable multifunctions.

References

- [1] H. Attouch, *Variational convergence for functions and operators*.
- [2] D. Azé, Caractérisation de la convergence au sens de Mosco en terme d'approximation inf-convolutives.
- [3] G. Beer, Convergence of linear functionals and their level sets.
- [4] G. Beer, A Polish topology for the closed subsets of a Polish Space.
- [5] G. Beer, Wijsman convergence of convex sets under renorming.
- [6] G. Beer, A. Lechicki, S. Levi and S. Naimpally, Distance junctionals and the suprema of hyperspace topologies.

- [7] G. Beer, R. Lucchetti Weak topologies for the closed subsets of a metrizable space.
- [8] J. Borwein and S. Fitzpatrick, Mosco convergence and the Kadec property.
- [9] P. Corradini, *Tesi di laurea*. Univ. di Milano, 1991.
- [10] B. Cornet, Topologies sur les fermés d'un espace métrique.
- [11] C. Hess. *Thèse d'Etat*. Montpellier, 1986.
- [12] S. Francaviglia, A. Lechicki, and S. Levi, *Quasi-uniformization of hyperspaces and the convergence of semi-continuous multifunctions*.
- [13] Y. Sontag, *Thèse d'Etat*. Université de Provence, 1982.
- [14] S. Sontag and C. Zalinescu, *Set convergences : an attempt of classification*.

Approximation and regularization of arbitrary sets in finite dimension

Joël Benoist

Université de Limoges, 87060 Limoges Cedex, France

Abstract

In this paper, we define the regularization of a set in R^n by using two kernels. More precisely, let (K_1, K_2) be a pair of two kernels (nonempty open subsets of R^n); then, for a nonempty closed subset $X \subset R^n$, we associate its regularizing family $(R_{\lambda,\mu}(X))_{0 < \mu < \lambda}$ defined by:

$$R_{\lambda,\mu}(X) = {}^c({}^c(X + \lambda K_1) + \mu K_2),$$

where $0 < \mu < \lambda$ be two real numbers and cA denotes the complement of A in R^n . When $K_1 = K_2$ is the open unit ball, we can easily build the regularizing family. When $K_1 = -K_2$ is a quadratic kernel and when X is the epigraph of a lower semicontinuous function, we retrieve the Lasry-Lions regularization (see [2]). Results are obtained in four directions:

1. general properties about $R_{\lambda,\mu}(X)$ which are issued of its definition ;
2. the convergence of the regularizing family when $(\lambda, \mu) \longrightarrow 0$;
3. the smoothness of the set $R_{\lambda,\mu}(X)$;
4. the asymptotic behaviour of Clarke's normal cone (see [1]) of the regularizing family.

References

- [1] J.-M. Lasry and P.-L. Lions, *A remark on regularization in Hilbert spaces*, Israel Journal of Mathematics, 55 (1986), pp. 257-266.
- [2] F. Clarke, *Optimization and nonsmooth Analysis*, New-York, John Wiley, 1983.

ASYMPTOTIC CONES, CLOSED IMAGES AND A THEOREM OF DIEUDONNE

E. Blum (Lima)

W. Oettli (Mannheim)

Asymptotic cones were introduced to describe the behaviour of convex sets "at infinity". If A is a convex, closed, nonempty subset of a real separated topological vector space E , then the asymptotic cone A_∞ of A is classically defined as

$$(1) \quad A_\infty := \bigcap_{\lambda > 0} \lambda(A-a),$$

when $a \in A$ is fixed arbitrarily. The definition is independent of the chosen $a \in A$. Dieudonné [3] has proved the following result :

Theorem. Let A, B be convex, closed, nonempty subsets of E . Let A be locally compact and $A_\infty \cap B_\infty = \{0\}$. Then $A-B$ is closed.

Dedieu [2] has generalized the definition of asymptotic cone for arbitrary subsets $A \subseteq E$ as follows :

$$(2) \quad A_\infty := \bigcap_{\varepsilon > 0} \overline{(0, \varepsilon) \cdot A}.$$

He extended Dieudonné's result to the case where A and B , instead of being convex, are radiating in $a \in A$ and $b \in B$ respectively. Dedieu's proof follows rather closely the proof of Dieudonné. This is possible, because if A is radiating in $a \in A$, then (1) still holds, as in the convex case.

Groinner [4] has carried over Dieudonné's result to convex multivalued mappings $T : E \rightrightarrows F$, where F is another topological vector space. Using Dieudonné's result he gave conditions for $T(A)$ to be closed, where A is a closed convex subset of E .

Here we want to prove the following result, which generalizes Dedieu's and Groinner's results :

Theorem. Let E, F be real topological vector spaces. Let $T : E \rightrightarrows F$ be a multivalued mapping. Let $\text{graph } T$ be closed. Let $A \subseteq E$ be closed, locally compact and radiating in $a \in A$. Let $\mathcal{M} := \{t \in E \setminus t \in A_\infty, (t, 0) \in (\text{graph } T)_\infty\}$. If $\mathcal{M} = \{0\}$, then $T(A)$ is closed.

Here A_∞ is to be understood in the sense of (2). If we set $T(X) := X - B$ we obtain Dedieu's result under weak assumptions. It should be noted that the requirement $\mathcal{M} = \{0\}$ above can be replaced by the requirement that $\mathcal{M}$ is a linear subspace and

$$A + \mathcal{M} \subseteq A, \text{ graph } T + (\mathcal{M} \times \{0\}) \subseteq \text{graph } T.$$

It is interesting to note that already Debreu in [1, p. 23] gave without proof the following result : If A, B are arbitrary closed subsets of $\mathbb{R}^n$ with $A_\infty \cap B_\infty = \{0\}$, then $A - B$ is closed.

References

- [1] G. Debreu, Theory of Value, Fifth printing, Yale University Press, New Haven 1973, Original Edition, 1959.
- [2] J.P. Dedieu, Cônes asymptotes d'ensembles non convexes, Bull. Soc. Math. France, Mémoire 60 (1979), 31-44.
- [3] J. Dieudonné, Sur la séparation des ensembles convexes, Math. Annales 163 (1966), 1-3.
- [4] J. Gwinne, Closed images of convex multivalued mappings in linear topological spaces with applications. J. Math. Anal. Appl. 60, (1977), 75-86.

EQUICOERCIVITE DE PROBLEMES VARIATIONNELS LE ROLE DES FONCTIONS DE RECESSION

G. BOUCHITTE (UTV Toulon)
en collaboration avec P.SUQUET (LMA Marseille)¹

Soit X un espace de Banach et τ une topologie rendant compactes les boules fermées de X . Etant données $F^n, F : X \rightarrow]-\infty, +\infty]$ des fonctionnelles convexes propres telles que :

$$\tau\text{-}\lim_{n \rightarrow \infty} F^n = F$$

et une forme linéaire τ -continue $L : X \rightarrow \mathbb{R}$, on se propose de caractériser l'ensemble des scalaires λ pour lesquels la suite $F_\lambda^n := F^n - \lambda L(\cdot)$ est équi-coercive dans X (ce qui assurera la convergence des infima associés). Lorsque cet ensemble J contient 0, il est montré que $J =]\lambda^-, \lambda^+[$ où :

$$\lambda^\pm = \text{Min } \{ F_\infty(u) ; u \in X, L(u) = \pm 1 \}$$

F_∞ étant la fonction de récession de F .

Diverses applications sont données en théorie de la plasticité, capillarité ..etc..

En liaison avec l'approximation des coefficients $\lambda^\pm$ à l'aide des problèmes approchés $\lambda_{n^\pm} = \inf \{ F_\infty^n(u) ; L(u) = \pm 1 \}$, on s'intéresse ensuite à la convergence des fonctions de récession F_∞^n vers F_∞ . En général l'égalité $\tau\text{-}\lim_{n \rightarrow \infty} F_\infty^n = F_\infty$ est fausse. Elle est vraie cependant dans le cas $X \subset Y$ avec Y Banach réflexif et injection compacte sous une hypothèse supplémentaire de compatibilité des domaines des fonctionnelles conjuguées sur Y^* :

$$w\text{-}\lim_{n \rightarrow \infty} \text{dom } (F^n)^* \subset \overline{\text{dom } F^*}$$

En application au problème des charges limites en plasticité, on obtient ainsi via l'inégalité de Deny-Lions un résultat d'homogénéisation en milieu incompressible.

¹ à paraître dans Séminaire EDP Collège de France 1991, Pitman

About proximal determinations of Huber's estimators

Mireille Bougeard

Université de Paris X, 92410 Ville d'Avray, France

and

Christian Michelot

Université de Paris I, 75634 Paris, France

Abstract

In recent years, *robustness* is a problem that has been given much attention in the statistical literature on the linear regression model. One reason of this interest is an increasing sensitivity of applied statisticians to potential deficiencies of least squares methods under the occurrence of outliers. Several robust alternatives have been considered. Among them, we find the class of *M*-estimators introduced by Huber [3]. *Huber's M-estimator*, proved to be the M-solution to a contaminated Gaussian model, is of particular interest. It consists in solving the following problem:

$$\text{Find } \hat{\beta} \in \operatorname{argmin} \varphi(\beta) = \sum_{i=1}^n \rho[(X\beta - y)_i] \quad (P)$$

where X is a given $n \times m$ matrix, $y \in \mathbb{R}^n$ and ρ is the cost function, depending on a given

tuning constant c , defined by

$$\rho(r) = \begin{cases} \frac{1}{2}r^2 & \text{if } |r| \leq c \\ c|r| - \frac{1}{2}c^2 & \text{otherwise} \end{cases}$$

Several iterative resolution procedures have already been investigated to find the *Huber M-estimator*.

Here, according to the closed connection between Huber's cost function ρ and the Moreau-Yosida c -approximate of the ℓ_1 cost function observed in [1], we propose to solve the optimization problem (P) by a proximal approach combined with duality whose ability in solving efficiently some closely related problems, namely location problems, has been proved [5][6].

In a first part of this paper we reformulate the Huber problem as a linearly constrained optimization problem. Then, thanks to duality (in the sense of Fenchel), we derive optimality conditions and we give several applications of these conditions (description of the entire set of optimal solutions, asymptotical results,...). We also show these optimality conditions can be solved by the *Partial Inverse Method* developed by Spingarn [7]. The algorithm, which can be viewed as a decomposition technique, gives very simple updating rules and permits parallel computations, what is of interest for large size data. As instance of the well known basic proximal algorithm, the procedure is known to be very stable and always globally convergent.

In a second part, we give a new equivalent formulation of the Huber problem that leads to a second duality scheme with decomposition properties in terms of "small residuals" and outliers. We show that this new formulation is also convenient to deal with duality and can be solved by the partial inverse method.

References

- [1] M.L. Bougeard, *Connection between statistical estimation criteria, lower C^2 functions and Moreau-Yosida approximates*, I.S.I. 47th Session, Vol.1, 159-160, 1989.
- [2] M.L. Bougeard and C. Michelot, *Collinearity in the linear model: statistical and numerical alternatives to least squares*, Journées "Systèmes de Référence Spatio-Temporels", Edit. N. Capitaine, 86-91, 1991.
- [3] P.J. Huber, *Robust estimation of a location parameter*, Ann. Math. Statist. 35, 73-101, 1964.
- [4] P.J. Huber, *Robust regression: asymptotics, conjectures and Monte Carlo*, Ann. Statist. 1, 799-821, 1973.
- [5] H. Idrissi, O. Lefebvre and C. Michelot, *Duality for constrained multifacility location problems with mixed norms and applications*, Annals of Oper. Res. 18, 71-92, 1989.
- [6] C. Michelot and O. Lefebvre, *A primal-dual algorithm for the Fermat-Weber problem involving mixed gauges*, Math. Prog. 39, 319-335, 1987.

- [7] J.E. Spingarn, *Partial inverse of a monotone operator*, Appl. Math. Optim. 10, 247-265, 1983.

A Eulogy on Nonconvexity

Arrigo Cellina

SISSA, 34014 Trieste, Italy

Abstract

There is some interest in problems of the Calculus of Variations and Optimization theory without the usual Convexity conditions.

This talk is mainly devoted in exploring and debating the reasons in favor of this approach to the problem.

Some recent existence results for problems involving the gradient will be discussed.

Convergence Issues in Penalty Methods. Linear Programming for Instance.

Roberto Cominetti and Jaime San Martín

Universidad de Chile, Casilla 170/3 Correo 3, Santiago, CHILE.

Abstract

Since the contributions of Kachiyan (1979) and Karmarkar (1984) on polynomial algorithms for linear programming, intensive research has been directed towards interior point methods for linear as well as nonlinear programming. A number of polynomial algorithms for LP and QP have emerged from this research, some of which have shown to be "competitive" with respect to more standard algorithms such as the simplex method. Recent advances have clarified the closed relationship between interior point methods and the classical (and somewhat forgotten) penalty methods.

In this lecture we present the analysis of trajectories associated to an exponential penalty function: the existence and convergence of these trajectories towards a "center" point of the optimal set, the exponentially fast rate of convergence towards this center, together with a fairly complete duality theory are all established under the sole and very weak assumption of boundedness of the solution set.

The analysis is presented in the simplest setting of LP, but some results of nonlinear nature are exposed as well. The computational efficiency of an algorithm based on exponential penalties, predicted by the fast convergence rate, have been confirmed by (limited) computational experience.

On the other hand, the (unexpected) good behavior of trajectories raise interesting questions concerning the limits to which one can push a satisfactory sensitivity analysis of nonlinear parametric programs in the absence of the usual strong second order hypothesis, linear independence of active gradients, etc.; and suggest that significant extensions are still possible in this area.

References

- [1] R. Cominetti & J-P. Dussault, *A stable exponential algorithm with superlinear convergence*, To appear, JOTA (1990).
- [2] R. Cominetti & J. San Martín, *Trajectory analysis for the exponential penalty method in linear programming*, Working Paper (1991).

Suprema of Wijsman topologies in normed spaces

Pietro Poggi Corradini

Via Cesariano 8, 20154 Milano

Abstract

Our work starts from a theorem of G. Beer which states that, on the hyperspace of closed and convex sets of a normed linear space X , the supremum of the Wijsman topologies ranging over the equivalent norms coincide with the Slice topology. In this context, we proved that, on the hyperspace of closed sets, it suffices to weaken a little the Slice topology to find a similar statement. That is to say, on the closed sets of X , the supremum of the Wijsman topologies made as above coincide with the Hit-and-Miss topology generated by the V^- with V open set of X , and the $(B^c)^{++}$ with B closed convex bounded and symmetric set of X . Here symmetric means that B is the translated of a set symmetric at the origin of X . This new topology always coincide with the Slice topology on the closed and convex sets, while on the closed sets this is true if and only if X is finite dimensional. Moreover, since a result of Borwein and Fitzpatrick states that, when X is reflexive, the supremum of the Wijsman is in fact a maximum, we examined the situation on the closed sets and found that in this case the supremum is realized if and only if X is finite dimensional.

References

- [1] G. Beer, *Wijsman convergence of convex sets under renorming*, preprint.
- [2] J. Borwein and S. Fitzpatrick, *Mosco convergence and the Kadec property*, Proc. Amer. Math. Soc. 106 (1989), 843-852.
- [3] C. Costantini, S. Levi and J. Zeminska, *Metriche che generano la stessa convergenza di Wijsman*, preprint.

A splitting property of the upper bounded-Hausdorff convergence

Camillo Costantini

Dip. Mat. Università di Milano, Via Saldini 50, 20133 Milano, Italy

Abstract

In the Hyperspace of a metric space (X, d) , a characterization of the upper bounded-Hausdorff convergence is provided. Precisely, we prove that a net $(A_i)_{i \in I}$ of closed subsets of X is bH^+ -convergent to a closed subset A if and only if for every $i \in I$ there exist closed subsets B_i, C_i of X such that:

$$1) A_i = B_i \cup C_i;$$

$$2) B_i \xrightarrow{H^+} A;$$

$$3) \forall x \in X: \lim_{i \in I} d(x, C_i) = +\infty \text{ (or, equivalently, } C_i \xrightarrow{w^+} \emptyset).$$

As an application, we obtain that if every A_i is connected and the set A is bounded, then the upper bounded-Hausdorff convergence of the A_i 's to A implies their upper Hausdorff convergence.

References

- [1] H. Attouch, R. Lucchetti, R.J.-B. Wets, *The topology of the ρ -Hausdorff distance*, Ann. Mat. Pura Appl.(4), **160** (1991), 303-320.
- [2] G. Beer, A. Di Concilio, *Uniform continuity on bounded sets and the Attouch-Wets topology*, Proc. Amer. Math. Soc., **112** (1991), 235-243.

Subdifferential Monotonicity as Characterization of Convex Functions *

Rafael Correa, Alejandro Jofré

Universidad de Chile, Facultad de Ciencias Físicas y Matemáticas,

Casilla 170-3 Correo 3, Santiago, Chile

and

Lionel Thibault

Université de Pau, Av. de l'Université, 64000, Pau, France

Abstract

It is known and easy to prove (Clarke (1983)) that the monotonicity of the Clarke subdifferential of a locally Lipschitz real valued function is equivalent to the convexity of this function. In order to prove the same result for a lower semicontinuous function $f : E \rightarrow \mathbf{R} \cup \{+\infty\}$ we have considered in Correa, Jofré and Thibault (1990) the Moreau-Yosida proximal approximation

$$f_\lambda(x) := \inf_{y \in E} \left[f(y) + \frac{1}{2\lambda} \|x - y\|^2 \right] \quad (1)$$

*This work was partially supported by Fondo Nacional de Ciencia y Tecnología, FONDECYT

since (under some general conditions) f_λ is locally Lipschitz and

$$f(x) = \sup_{\lambda > 0} f_\lambda(x).$$

Thus our procedure consisted in deriving the monotonicity of ∂f_λ from that of ∂f . This method required the reflexivity of the space E because it depended heavily on the fact that the above infimum is attained whenever the Frechet subdifferential $\partial^F f(x)$ of f at x is nonempty. This has been obtained by supposing (thanks to the reflexivity of E) that the norm of E is Kadec and by showing that (when $\partial^F f(x) \neq \emptyset$) there exists some minimizing sequence of (1) weakly converging to some point z and whose norms converge to $\|z\|$, which implies the strong convergence of the sequence. The same result was proved before by R.A. Poliquin (1988) for $E = \mathbb{R}^n$ with the help of his notion of quadratic conjugate function.

In this paper, by a completely different approach, we avoid the use of the Moreau-Yosida approximation f_λ in order to get the result for any Banach space E . In fact we prove that the monotonicity of any classical subdifferential ∂f of a lower semicontinuous function $f : E \rightarrow \mathbb{R} \cup \{+\infty\}$ defined on a Banach space E is equivalent to the convexity of this function f .

References

- [1] F. H. Clarke: *Optimization and nonsmooth analysis*, Wiley, New York, 1983.
- [2] R. Correa, A. Jofré and L. Thibault: Characterization of lower semicontinuous convex functions, Proc. Amer. Math. Soc. To appear.

Index multiplicatifs de convexité/concavité et applications

J.-P. Crouzeix et R. Kebbour

Université Blaise Pascal, Clermont Ferrand, France

Résumé

L'étude de la concavité/convexité d'un produit de fonctions séparables est faite à partir de la notion d'index de convexité/concavité dérivés de l'index de convexité introduite par Debreu-Koopmans et Crouzeix-Lindberg. Des applications sont données en économie et pour des fonctions potentielles introduites récemment en programmation linéaire.

Title to be announced

Ennio De Giorgi

Scuola Normale Superiore, 56126 Pisa, Italy

Stability of subdifferentials of non convex functionals

Robert Deville

Université de Franche Comté, 25030 Besancon Cedex, France

Abstract

We present some recent variational principles and we apply them to show various stability results of subdifferentials and superdifferentials of non convex functionals in infinite dimensions. These results are applied to the geometry of Banach spaces and to the study of Hamilton-Jacobi equations or of fully non linear second order partial differential equations in infinite dimensions.

On the Convergence of Efficient Sets

Dinh The Luc *

Institute of Mathematics, Hanoi

Roberto Lucchetti

University of Milano

Christian Malivert

Université de Limoges

Abstract

Let E be a normed space partially ordered by a convex cone C . The set of efficient points and of weakly efficient points of A are defined by

$$Min A = \{a \in A : a \in a' + C \text{ for some } a' \in A \text{ implies } a' \in a + C\}$$

$$WMin A = \{a \in A : \text{there is no } a' \in A \text{ with } a \in a' + int C\}$$

The question studied here is the dependence of $Min A$ and $WMin A$ on perturbations of A . The main results existing on this topic [1][2][4], deal with the Kuratovski-Painlevé limits. For instance, one can prove that if $\lim_{n \rightarrow \infty} A_n = A$ then $\limsup_{n \rightarrow \infty} WMin A_n \subset WMin A$. When E is finite dimensional and A is closed this result can be read as

*visiting the University of Limoges

: $\lim_{n \rightarrow \infty} d(A_n, x) = d(A, x)$, for all $x \in E$, implies $\liminf_{n \rightarrow \infty} d(WMin A_n, x) \geq d(WMin A, x)$ for all $x \in E$. Examples show that the same conclusion is not always valid in an infinite dimensional setting. Here we present two particular cases in which the result holds, either supposing that E is an Hilbert space with C a polyhedral cone, or with the condition that $\bigcup WMin A_n$ is relatively compact. Another interesting topic is to obtain an opposite inequality in the form $\limsup_{n \rightarrow \infty} d(Min A_n, x) \leq d(Min A, x)$ for all $x \in E$. For this type of conclusion we present sufficient conditions ensuring at the same time that $Min A$ is nonempty.

References

- [1] H. Attouch and H. Riahi, Stability results for Ekeland's e-variational principle and cone extremal solutions, To appear.
- [2] S.Dolecki and C.Malivert, Stability of efficient sets: Continuity of mobile polarities, Non linear Analysis, Theory, Methods and Applications Vol.12 **12** 1461-1486 (1988).
- [3] D.T. Luc, *Theory of Vector Optimization*, Lecture notes in Economics and Mathematical Systems, Springer-Verlag, 1989.
- [4] J.P. Penot and A. Sterna-Karwat, Parametrized multicriteria optimization, Continuity and closedness of optimal multifunctions, J. Math. Analysis Appl. **120** 150-168 (1986).

ON VARIATIONAL STABILITY IN COMPETITIVE ECONOMIES

Sjur Didrik Flåm

Bergen University, Department of Economics, 5008 Norway

ABSTRACT. We explore the variational stability of supply, demand and equilibria in perfectly competitive economies. The appropriate, and in fact, minimal limit notion is furnished by the concept of Kuratowski-Painlevé convergence together with the derived concepts of epi and hypo convergence. When technologies and preferences converge in such manners we show, subject to compactness assumptions, that equilibria of approximate economies cluster to those of the limiting economy.

Key words: Epi-convergence, hypo-convergence, slice topology, Hotelling's lemma, Shepard's lemma, excess demand, indirect utility, expenditure function, competitive equilibria.

1 INTRODUCTION.

Microeconomic theory deals with the behavior of firms and consumers interacting via markets. Typically one wants to understand (or predict) equilibrium demand and supply of these agents. The dominating paradigm is then constrained optimization: each firm (or consumer) is assumed to maximize its achievable profit (resp. utility) subject to constraints. As a rule, only one half of such optimization problems is visible for an outside observer: For a given, fixed price-vector p the objective of a profit-maximizing firm is a known linear function $\langle p, \cdot \rangle$. By contrast, its *technology*, that is, the set Y of all possible production plans y is often hard to describe. This situation is precisely reversed for a representative consumer: On one hand, an adequate description of his feasible set is readily given by the budget constraint $\langle p, x \rangle \leq b$, b denoting here his budget. On the other hand, his *utility function* u mapping the commodity space E into $[-\infty, \infty)$, is usually unknown, or at least, not made directly explicit.

This state of affairs prompts at least two questions. First, one regarding a *dual approach* (Diewert, 1982): Observing the *optimal choice correspondences*

$$p \rightarrow X(p, b), \quad p \rightarrow Y(p),$$

portraying demand and supply of an individual consumer and a single firm is it possible to reconstruct the underlying preference $u: E \rightarrow [-\infty, \infty)$ and the technology Y ?

Second, in terms of *sensitivity analysis* (also named comparative statics), knowing the two optimal value mappings, namely the *expenditure function*

$$p \rightarrow e(p) := \langle p, x \rangle, \quad x \in X(p, b)$$

and the *profit function*,

$$p \rightarrow \pi(p) := \langle p, y \rangle, \quad y \in Y(p)$$

may one predict changes $\Delta X(p,b)$, $\Delta Y(p)$ that result from price perturbations Δp ? Not surprisingly, good answers to these questions have been granted by the application of convex analysis. Some of these results are recalled in Prop. 3.1-2 below.

However, since economic models suffer from inaccuracies of various sorts, such answers are usually inexact. In fact, to provide estimates of competitive supply and demand is not well founded unless the correct, yet unknown economic model is stable in some appropriate sense. This observation motivates the present note to inquire: *What kind of convergence $E^n \rightarrow E$ between economies E^n, E ensures that competitive equilibria of the approximate economies E^n , $n=1,2,\dots$, cluster to those of the limit economy E ?*

At the level of technologies the convenient and natural notion $Y^n \rightarrow Y$ is that *Kuratowski-Painlevé convergence*. For preferences it turns out that the most suitable convergence mode $u^n \rightarrow u$ is that of *hypo-convergence*. All definitions are recalled in Section 2. As customary in microeconomics we focus first on *one* firm (Sect. 3) and *one* consumer (Sect. 4) the objective being to explore stability of individual demand and supply. Thereafter the results obtained for single agents are synthesized into a general equilibrium analysis (Sect. 5). To facilitate the exposition we shall assume, for the most part, that the commodity space E is finite-dimensional Euclidean. Sect. 6 points out generalizations.

The novelties of this paper are mostly in terms of applications; it opens up for variational analysis in competitive economic contexts, a branch so far not much developed.

REFERENCES

- C.D. Aliprantis and D.J. Brown, "Equilibria in markets with a Riesz space of commodities", *J. Math. Economics* II, 189-207 (1983).
- H. Attouch, "Convergences de fonctions convexes, des sous-différentiels et semi-groupes associés", *Comptes Rend. Acad. Sci. Paris* 284, 539-542 (1977).
- H. Attouch, *Variational Convergence for Functions and Operators*, Pitman, Boston (1984).
- H. Attouch and G. Beer, "On the convergence of subdifferentials of convex functions", WP, Montpellier (1991).
- D. Azé and M. Volle, "A stability result in quasi-convex programming", *J. of Optimization Th. and Applic.* vol. 67, no. 1, 175-184 (1990).
- G. Beer, "The slice topology: a viable alternative to Mosco convergence in nonreflexive spaces", WP, California State Univ., LA (1991).
- T.F. Bewley, "Existence of equilibria in economies with infinitely many commodities", *J. Econ. Theory* 4, 514-540 (1972).
- W.E. Diewert, "Duality approaches to microeconomic theory," in K.J. Arrow and M.D. Intriligator, *Handbook of Mathematical Economics*, North Holland, Amsterdam vol. II. Chapt. 14 (1982).
- G. Debreu, *The Theory of Value*, J. Wiley, New York (1959).
- J.P. Aubin and I. Ekeland, *Applied Nonlinear Analysis*, J. Wiley, N.York (1984).
- A. Mas-Colell, "The price equilibrium existence problem in topological vector lattices", *Econometrica* 54, 1039-1053 (1986).
- V. Mosco, "On the continuity of the Young-Fenchel transform", *J. Math. Anal. Applic.* 35, 518-535 (1971).
- R.T. Rockafellar, *Convex Analysis*, Princeton Univ. Press, New Jersey (1970).
- H. Varian, *Microeconomic Analysis*, 2.ed. Norton, New York (1984).
- M. Volle, "Convergence en niveaux et en epigraphs", *Notes aux Comptes Rendues de l'Académie des Sciences de Paris*, vol. 299, 295-298 (1984).
- R. Wijsman, "Convergence of sequences of convex sets, cones, and functions II", *Trans. Am. Math. Soc.* 123, 32-45 (1966).

Viability for Constrained Stochastic Differential Equations

Serge Gautier and Lionel Thibault

Université de Pau, 64000 Pau, France

Abstract

The existence of viable solutions of constrained stochastic differential equations (with Brownian motion) is established under a stochastic tangential condition. The approach is based on an appropriate notion of approximate solutions.

Multivalued strong law of large numbers in the Wijsman topology and the slice topology.

Christian HESS¹

In recent years, multivalued strong laws of large numbers (SLLN), especially for unbounded random sets, proved to be useful in applications to stochastic optimization, statistics and related fields. Results of such type were first provided by Artstein and Hart [ArH] for random sets (r. s.) with closed values in finite dimensional spaces. Later, Hiai [Hi] and Hess [He3, 4] independently proved similar results for random sets with values in an infinite dimensional Banach space. These multivalued SLLN were proved assuming that the set of all closed subsets of the Banach space X , denoted by $\mathcal{C}(X)$, is endowed with the Mosco topology (which reduces to the Kuratowski-Painlevé topology for finite dimensional spaces).

Although these SLLN provide useful probabilistic convergence properties, one may ask for other results involving more explicitly the norm of the space X . A natural and well-known topology for this purpose is the Wijsman topology, denoted by $\mathcal{T}_W$. It is the topology of pointwise convergence of distance functions. So, a natural question is : in a general (separable) Banach space, does the SLLN for random sets with values in $\mathcal{C}(X)$ hold when this set is endowed with $\mathcal{T}_W$? An affirmative answer to this question is provided by theorem 1 (A) below.

On the other hand, more recently, G. Beer [Be1, 2] introduced the "slice topology" on $\mathcal{C}(X)$, which is a natural extension of the Mosco topology. Indeed, Beer showed that, in a general Banach space, the slice topology, denoted by $\mathcal{T}_s$, is stronger than the Mosco topology, and that both topologies coincide if and only if X is reflexive. Further, on $\mathcal{C}(X)$ the slice topology is stronger than $\mathcal{T}_W$. Consequently, one may also ask if the multivalued SLLN holds in the slice topology. A positive answer to that question is furnished by theorem 1 (B).

Theorem 1 - A) Consider a separable Banach space X , an integrable r. s. Γ with values in $\mathcal{C}(X)$ and a sequence $(\Gamma_n)_{n \geq 1}$ of pairwise independent r. s. having the same distribution as Γ .

Then, there exists a negligible subset N such that, for any $\omega \in \Omega \setminus N$, one has

$$(1) \quad \bar{\text{co}} I(\Gamma, \mathcal{A}) = \mathcal{T} - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \Gamma_i(\omega)$$

where $\mathcal{T} = \mathcal{T}_W$, that is, for any $x \in X$,

$$d(x, \bar{\text{co}} I(\Gamma, \mathcal{A})) = \lim_{n \rightarrow \infty} d(x, \frac{1}{n} \sum_{i=1}^n \Gamma_i(\omega)).$$

B) Moreover, if X^* is strongly separable then (1) holds with $\mathcal{T} = \mathcal{T}_s$.

¹CEREMADE, Université Paris Dauphine, 75775 PARIS CEDEX 16, FRANCE

Théorème 2 (Application to integrands) - Consider an integrable convex lower semi-continuous normal integrand f defined on $\Omega \times X$ with values in $\bar{\mathbb{R}}$ and a sequence $(f_n)_{n \geq 1}$ of lsc normal integrands, pairwise independent with the same distribution as f . Also define the mean functional of f , namely the function F given by

$$F(x) := \int_{\Omega} f(\omega, x) d\omega \quad x \in X.$$

Then for almost all $\omega \in \Omega$, one has

$$(2) \quad F^{**} = \tau_s\text{-}\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n f_i(\omega, \cdot).$$

where F^{**} is the biconjugate of F . In (2) 'lim' means 'epigraphical limit'. More precisely, for almost all ω , the sequence of epigraphs

$$\text{epi} \left(\frac{1}{n} \sum_{i=1}^n f_i(\omega, \cdot) \right) \quad n \geq 1$$

τ_s -converges to $\text{epi } F^{**}$.

See [He5, 6] for the proofs and further discussions. The proof of theorem 1 (A) heavily relies upon specific properties on the distribution of random sets ([ArH] and [He1, 2]). Also note that results similar to theorem 2, but for other topologies on $\mathcal{C}(X)$, has been obtained by Attouch and Wets [AtW].

REFERENCES

- [ArH] Z. Artstein and S. Hart, Law of large numbers for random sets and allocation processes, Math. of Oper. Res., 6, p. 482-492 (1981)
- [AtW] H. Attouch and R. Wets, Epigraphical process : laws of large numbers for random lsc functions, preprint (Février 1991)
- [Be1] -----, Topologies on closed and closed convex sets and the Effrös measurability of set valued functions, Department of Mathematics, California State University, Los Angeles, CA 90032, Preprint (and : Séminaire d'Analyse Convexe de l'Université de Montpellier, Exposé n°3 (1991))
- [Be2] -----, The slice topology : a viable alternative to Mosco convergence in nonreflexive spaces, Department of Mathematics, California State University, Los Angeles, CA 90032, Preprint (and : Séminaire d'Analyse Convexe de l'Université de Montpellier, Exposé n°2 (1991))
- [He1] C. Hess, Loi de probabilité des ensembles aléatoires à valeurs fermées dans un espace de Banach séparable, C. R. Acad. Sc. Paris, t. 296 (1983)
- [He2] -----, Loi de Probabilité et indépendance pour des ensembles aléatoires à valeurs fermées dans un espace de Banach, Séminaire d'Analyse Convexe de l'Université de Montpellier, Exposé n°7 (1983)
- [He3] -----, Loi forte des grands nombres pour des ensembles aléatoires non bornés à valeurs dans un espace de Banach séparable, C. R. Acad. Sc. Paris, série I, t. 300, p.177-180 (1985)
- [He4] -----, Quelques théorèmes limites pour des ensembles aléatoires bornés ou non, Séminaire d'Analyse Convexe de l'Université de Montpellier, Exposé n°12 (1984)
- [He5] -----, On the distribution of random sets and the multivalued strong law of large numbers in the Wijsman topology, preprint (March 1992)
- [He6] -----, Multivalued strong law of large numbers in the slice topology, preprint (May 1992)

SUBDIFFERENTIAL CALCULUS WITHOUT "QUALIFICATION" CONDITIONS

J.-B. HIRIART-URRUTY

and

R. R. PHELPS

University Paul Sabatier
TOULOUSE

University of Washington
SEATTLE

Abstract. Calculus with subdifferentials of convex functions is important for dealing with variational problems ; if f is constructed from some other convex functions f_j , the problem is to compute (exactly) ∂f in terms of the ∂f_j 's. When the functions involved are extended-valued, some "qualification" conditions are usually required to get the desired formulas. For example, various conditions ensure that $\partial(f_1+f_2) = \partial f_1 + \partial f_2$, but these conditions are not always satisfied in variational problems. The aim of the present work is to derive a new set of calculus rules for subdifferentials of convex functions, without any qualification condition, substituting the information $(\partial_\epsilon f_j(x), \epsilon \in]0, \bar{\epsilon}])$ for $\partial f_j(x)$. The threshold $\bar{\epsilon} > 0$ can be taken as small as desired, the "zero" of your computer for example. Again the role of the ϵ -subdifferential $\partial_\epsilon f$ as a "smoothing effect" or "viscosity enlargement" of ∂f is emphasized. We give hereunder some snapshots of the calculus rules obtained.

(1) **Sum.** Suppose that f_1 and f_2 are proper lower semicontinuous convex functions ; then (with the bar denoting the closure in the weak* topology)

$$\partial(f_1+f_2)(x) = \bigcap_{0 < \epsilon \leq \bar{\epsilon}} [\overline{\partial_\epsilon f_1(x) + \partial_\epsilon f_2(x)}] \quad (\bar{\epsilon} > 0 \text{ arbitrary})$$

$$\left(= \lim_{\epsilon \downarrow 0} [\overline{\partial_\epsilon f_1(x) + \partial_\epsilon f_2(x)}] \right).$$

(2) **Inf-convolution.** If f_1 and f_2 are convex lower semicontinuous,

$$\partial(f_1 \square f_2)(x) = \bigcap_{0 < \varepsilon \leq \bar{\varepsilon}} \bigcup_{y \in E_\varepsilon(x)} [\partial_\varepsilon f_1(y) \cap \partial_\varepsilon f_2(x-y)],$$

where $E_\varepsilon(x) = \{y : f_1(y) + f_2(x-y) < (f_1 \square f_2)(x) + \varepsilon\}$.

(3) **Maximum.** If f_1 and f_2 are convex lower semicontinuous, if $f_1(x) = f_2(x)$,

$$\partial[\text{Max}(f_1, f_2)](x) = \bigcap_{0 < \varepsilon \leq \bar{\varepsilon}} \overline{\text{co}}[\partial_\varepsilon f_1(x) \cup \partial_\varepsilon f_2(x)] \quad (\text{A. BRONDSTED, 1972}).$$

Once some key-formulas are derived (like the one on the sum of functions), formulas concerning further operations are deduced by using the usual tricks and transformations in Convex Analysis.

The cases of marginal functions and differences of functions have been more thoroughly studied in AVIGNON (A. SEEGER, M. VOLLE, R. MOUSSAOUI) with applications in the area of Calculus of Variations.

Lipschitz Approximation of Lattice Valued Functions

VINCENT JALBY

Université Montpellier II
Laboratoire d'Analyse Convexe
34095 Montpellier Cedex 05

ABSTRACT

Let E be an order-complete Banach lattice with norm $\|\cdot\|$ and positive cone E_+ . We adjoin to E a greatest element $+\infty$ and we set $E^\bullet = E \cup \{+\infty\}$, $E_+^\bullet = E_+ \cup \{+\infty\}$. We introduce a new concept of lower semi-continuity for the functions ψ from a separable metric space (X, d) into $E^\bullet$: a function $\psi : X \rightarrow E^\bullet$ is *inf-continuous* at $x \in X$ if for every $e \in E$ with $e \leq \psi(x)$ and for every neighbourhood V of 0 in E , there exists a neighbourhood U of x such that $\inf \psi(U) \in e + V + E_+^\bullet$. This notion is stronger than the usual one (cf. [2]). Both coincide when E_+ has a non empty interior. We can now state the result of Lipschitz approximation.

THEOREM. *Let $\psi : X \rightarrow E^\bullet$ an inf-continuous proper function on X . Assume that there exist $a, b \in E_+$ and $x_0 \in X$ such that $\psi(x) \geq -a - d(x, x_0)b$, $\forall x \in X$. Then there exist $h \in E_+$ with $\|h\| \leq 1$ and $k_0 \in \mathbb{N}^*$ such that for all $k \geq k_0$, the functions $\psi^k : X \rightarrow E$, $x \mapsto \inf_{y \in X} \{\psi(y) + kd(x, y)h\}$ verify the following properties:*

- (1) $|\psi^k(x) - \psi^k(y)| \leq kd(x, y)h$, $\forall (x, y) \in X \times X$.
- (2) $\|\psi^k(x) - \psi^k(y)\| \leq kd(x, y)$, $\forall (x, y) \in X \times X$.
- (3) $\psi(x) = \sup_k \psi^k(x)$, $\forall x \in X$.
- (4) $\psi(x) = \lim_k \psi^k(x)$, $\forall x \in X$ such that $\psi(x) \in E$.

We point out that the element h is not unique and that it depends on ψ . Also, if ψ is such that there exists a sequence $(\psi_n)_n$ of functions verifying (1) to (4) then ψ is inf-continuous.

We use this approximation result to characterize an epi-like convergence of lattice valued functions: let $(\psi_n)_n$ a sequence of functions from X into $E^\bullet$ and define $\lim_p \psi_n = \sup_p \underline{\lim}_n \inf \psi_n(B(x, 1/p))$, $\lim_e \psi_n = \sup_p \overline{\lim}_n \inf \psi_n(B(x, 1/p))$. If

$\psi, (\psi_n)_n$ are finite inf-continuous functions minored by $-a-d(x, x_0)b$ and such that $\psi_n \leq \psi$, then one has $\text{lie } \psi_n(x) = \sup_k \underline{\lim}_n \psi_n^k(x)$, $\text{lse } \psi_n(x) = \sup_k \overline{\lim}_n \psi_n^k(x)$. It seems that the "good" definition of convergence is the following: a sequence $(\psi_n)_n$ **-epi-converges uniformly* on X to ψ_∞ if from every subsequence of $(\psi_n)_n$ one can extract another subsequence $(\psi_{n_p})_p$ such that $\text{lie } \psi_{n_p} = \text{lse } \psi_{n_p} = \psi_\infty$. This notion have many variational properties such that $\inf \psi_\infty \geq \overline{\lim}_p [\inf \psi_{n_p}]$ for a subsequence $(\psi_{n_p})_p$. Actually, the only relation between uniform *-epi-convergence and the convergence of the epigraphs is the following inclusion: $\text{Epi}(\text{lse } \psi_n) \subset \text{Li}(\text{Epi } \psi_n)$.

We also extend to lattice valued integrands the previous approximation result and prove a strong law of large numbers for this integrands.

REFERENCES

1. H. Attouch, "Variational convergence for functions and operators," Pitman, London, 1984.
2. J.M. Borwein, J.P. Penot & M. Théra, *Conjugate convex operators*, J. Math. Anal. Appl. 102 (1984), 399-414.

On B-subgradients and applications

Alejandro Jofre

Departamento de Ingenieria Matemática, Universidad de Chile

Casilla 170/3, Correo 3, Santiago, Chile

Abstract

The set of b -subgradients of a function f , $\partial_b f$, is a notion which generalizes the derivative, in the sense that, it is reduced to a singleton when the function f is differentiable. This subgradient set and the related normal cone N_b are always smaller than the Clarke generalized gradient and its correspondent normal cone. Recently, an important number of calculus rules (Michel & Penot, Treiman), optimality conditions in mathematical programming and optimal control, Proximal normal and b -subgradient formulae (Jofre & Thibault, Treiman) and Frechet (ϵ, r) -normal formula (Jofre & Thibault) have been proved using these concepts. In this talk, we show two consequences of the proximal b -normal formulae. Firstly, we give estimates to b -subgradient of the optimal value function $p(u)$

$$p(u) = \inf \{ f(x) : g(x) + u \in -K, x \in C \}$$

associated to the parameterized nonsmooth optimization problem

$$\begin{aligned} &\text{minimize } f(x) \text{ subject to} \\ &g(x) + u \in -K \\ &x \in C \end{aligned}$$

where $f : E \rightarrow \mathbf{R}$ and $g : E \rightarrow \mathbf{R}^p$ are locally Lipschitzian functions, C is a nonempty closed subset of a Banach space E , and K is a convex cone in $\mathbf{R}^p$. Our proof follows the ideas of Rockafellar and Clarke's works with a major difference, the set-valued map $\partial_b f(\cdot)$ is not upper semicontinuous even if f is locally Lipschitzian.

Secondly, we give a result on the b -normal cones, in the vein of Cornet-Rockafellar's theorem (1989), which coincide with a separation theorem when the sets are convex.

Semi-Infinite Optimization: Structure and Stability of the Feasible Set

H. Th. Jongen

RWTH Aachen, Germany

Abstract

This research was done in collaboration with F. Twilt and G.-W. Weber. The problem of the minimization of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ under finitely many equality constraints and perhaps infinitely many inequality constraints gives rise to a structural analysis of the feasible set $M[H, G] = \{x \in \mathbb{R}^n : H(x) = 0, G(x, y) \geq 0, y \in Y\}$ with compact Y in $\mathbb{R}^r$. An extension of the well-known *Mangasarian-Fromowitz* constraint qualification (EMFCQ) is introduced. The main result for compact $M[H, G]$ is the equivalence of the topological stability of the feasible set $M[H, G]$ and the validity of EMFCQ. As a byproduct, we obtain under EMFCQ that the feasible set admits local linearizations and also that $M[H, G]$ depends continuously of the pair (H, G) . Moreover, EMFCQ is shown to be satisfied generically.

Metric regularity and stability in semi-infinite optimization

Diethard Klatte

Institute for Operations Research

University of Zurich

CH - 8044 Zurich

Abstract

In the paper we study parametric nonlinear optimization problems of the type SIP(t):

$$\begin{aligned} f(x, t) &\longrightarrow \min_x \\ \text{subject to} \\ h_i(x, t) &= 0, \quad i \in I, \\ g(j, x, t) &\geq 0, \quad j \in K, \end{aligned}$$

where t is regarded as a parameter varying over some metric space T , $I := \{1, \dots, p\}$, K is a compact subset of $\mathbf{R}^s$, and $h_i : \mathbf{R}^s \times T \rightarrow \mathbf{R}$ and $g : \mathbf{R}^s \times \mathbf{R}^n \times T \rightarrow \mathbf{R}$ are continuous functions being C^1 w.r. to x resp. (j, x) .

Given $t^0 \in T$ and some solution x^0 to SIP(t^0) of interest (e.g., a local minimizer or a stationary solution in the Kuhn-Tucker sense), we are looking for conditions ensuring regularity of the constraint system near (x^0, t^0) and local stability of the solution. As a main regularity requirement on the constraints, we use the Extended Mangasarian Frolovitz Constraint Qualification (EMFCQ), cf. Jongen, Twilt and Weber [3]. First we show that EMFCQ at some feasible point z is necessary and sufficient for metric regularity

of the constraint system at z , cf. Henrion and Klatte [2]. This fact increases the number of equivalent characterizations of EMFCQ: In Henrion [1], equivalence between EMFCQ and some local epigraph representability of the constraint set was observed, and in [3], there is shown that the feasible set of a semi-infinite program is (topologically) stable in Jongen's sense if and only if EMFCQ holds on the whole set.

Then, assuming EMFCQ at a strict local minimizer x° of the initial semi-infinite problem (i.e., at t°), the continuity of some portion of the local minimizing set mapping at t° immediately follows, and, under more structure, even "upper Hoelder" continuity holds, cf. Klatte [4]. Using a certain extension of the Gauvin–Robinson result on local boundedness of the Lagrange multiplier set mapping under MFCQ to our situation, one may show the upper semicontinuity of some portion of the (Kuhn–Tucker) stationary solution set mapping.

Finally, we present strong stability results under the well-known reduction ansatz. This approach leads – even under restrictive smoothness and regularity conditions on the original data – to a $C^{1,1}$ optimization problem with finitely many constraints, and one may apply the stability properties known in this case, cf. Klatte [4].

References

- [1] R. Henrion. On constraint qualifications. *J. Optim. Theory Appl.* 72 (1992), 187-197.
- [2] R. Henrion and D. Klatte, *Metric regularity of the feasible set mapping in semi-infinite optimization*. Preprint Nr 92-5, Humboldt-Univ. Berlin, Fachbereich Mathematik, 1992.
- [3] H. Th. Jongen, F. Twilt and G.W. Weber, Semi-infinite optimization: Structure and stability of the feasible set. *J. Optim. Theory Appl.* 72 (1992) 529-552.
- [4] D. Klatte. Nonlinear optimization problems under data perturbations. In: W. Krabs and J. Zowe (Eds.), *Modern Methods of Optimization*, pp. 204-235, Springer-Verlag, 1992.

Convergence of diagonally stationary sequences in Convex Optimization

B. Lemaire & M.A. Bahraoui

Université des Sciences et Techniques du Languedoc

Place E. Bataillon, 34095 Montpellier Cedex 5

Let f be a closed proper convex function defined on a real normed vector space X . Denoting by $\partial f(x)$ the subdifferential of f at x and by $d_*(0, C)$ the distance, in the dual space X^* of X , from 0 to the subset C of X^* , we say that a sequence $\{x_n\}$ in X is **stationary** for f if it satisfies

$$\lim_{n \rightarrow +\infty} d_*(0, \partial f(x_n)) = 0$$

that is, for each $n \in \mathbb{N}$, x_n is determined along with a subgradient $x_n^* \in \partial f(x_n)$ such that $x_n^* \rightarrow 0$ strongly in X^* . Some infinite constructive processes for minimizing f generate such a sequence. Actually, in most situations (infinite dimension of X , constraints in f), such a process is not applied to f itself but to a **fixed** approximation of it.

The idea of diagonal processes, as introduced in [Boy 74], is to combine a basic process with a sequence $\{f^n\}$ of (closed proper convex) approximations of f **changing** the approximation at each step of the process. For stationary sequence generating processes this leads to generating what we call a **diagonally stationary sequence** (DSS) for $\{f^n\}$, that is, a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow +\infty} d_*(0, \partial f^n(x_n)) = 0$$

A natural question arises: under what conditions on $\{f^n\}$ with respect to f is the considered basic process stable in the sense that the diagonal process inherits the convergence properties of the basic one? Variational convergence theory has revealed powerful tools to study this question in the context of particular basic processes [Lem 88, Mou 89, Tos 90, A-L 91, Lem 91]. The purpose of the present work is to proceed on that way for general DSS. We first consider the case of **bounded** DSS with different kinds of convergence

of f^n to f . For **unbounded DSS** we introduce the notion of **equi well asymptotical behaviour** (of the sequence $\{f^n\}$) which extends the notion of **well asymptotical behaviour** introduced in [A-C 89] for a single convex function. If X is a (general) Banach space, this notion appears to be equivalent to a notion of **equi well posedness** closely related to the one introduced in [Zol 78], equivalence yet established in [Lem 92] for a single convex function between well asymptotical behaviour and well posedness in the (extended to nonuniqueness) sense of Thikhonov.

References

- [A-L 91] **Alart P. & Lemaire B.** Penalization in non-classical convex programming via variational convergence, *Math. Progr.* 51, 1991, 307-331.
- [A-C 89] **Auslender A. & Crouzeix J.P.** Well behaved asymptotical convex functions. in *Analyse non linéaire*, Gauthier- Villars, 1989, 101-122.
- [Boy 74] **Boyer R.** Quelques algorithmes diagonaux en optimisation convexe, Thèse de 3ème cycle, Université de Provence, 1974.
- [Lem 88] **Lemaire B.** Coupling optimization methods and variational convergence. in *Trends in Mathematical Optimization*, International Series of Numerical Mathematics, Vol 84(c), Birkhäuser Verlag, 1988, 163-179.
- [Lem 91] **Lemaire B.** About the convergence of the proximal method, 6th French-German Conference on Optimization, Lambrecht, june 1991.
- [Lem 92] **Lemaire B.** Bonne position, conditionnement, et bon comportement asymptotique. Séminaire d'Analyse Convexe, Montpellier, mars 1992.
- [Mou 89] **Mouallif K.** Convergence variationnelle et méthodes perturbées pour les problèmes d'optimisation et de point selle, Thèse d'Etat, Université de Liège, 1989.
- [Tos 90] **Tossings P.** Sur les zéros des opérateurs maximaux monotones et applications, Thèse de Doctorat, Université de Liège, 1990.
- [Zol 78] **Zolezzi T.** On Equiwellset Minimum Problems, *Appl. Math. Optim.* 4, 1978, 209-223.

SOME NEW RESULTS ON WIJSMAN TOPOLOGIES

S. LEVI (Milano)

Following the paper [BLLN], new results are appearing (see [Be] and [CLP]) that point towards the centrality of Wijsman topologies within the theory of set convergences.

It is therefore of interest to describe the conditions under which two equivalent metrics generate the same Wijsman convergence in the hyperspace.

$c_0(X)$ will denote the hyperspace of nonempty closed subsets of the metrizable space X .

We recall the following results from [BLLN] :

- the Vietoris topology on $c(X)$ is the supremum of the Wijsman topologies over all equivalent metrics on X ;
- the upper Hausdorff topology generated by a metric d on X is the supremum of the Wijsman topologies over all metrics that are uniformly equivalent to d .

and from [Be] :

- the slice topology on the closed convex subsets of a normed space is the supremum of the Wijsman topologies over all metrics generated by equivalent norms.

In the forthcoming paper [CLP] the following result is proved :

- Kuratowski convergence on $c(X)$ is the infimum - in the lattice of convergences - of the Wijsman topologies over all equivalent metrics on X .

As for equality of two Wijsman topologies we have the

Theorem [CLZ] : Let d and r equivalent metrics on X ; then the Wijsman topology generated by r is stronger on $c_0(X)$ than that generated by d if and only if for every x in X and $0 < e < a$ there exists $x_1, \dots, x_n$ in X and $0 < d_i < s_i$ ($i = 1, \dots, n$) with

$$B_e^d(x) \subset \bigcup_{i=1}^n B_{d_i}^r(x_i) \subset \bigcup_{i=1}^n B_{s_i}^r(x_i) \subset B_a^d(x)$$

where $B_e^d(x)$ is the d -open ball of center x and radius e .

As a corollary we obtain that on the hyperspace of a normed space two Wijsman topologies generated by equivalent non homothetic norms agree if and only if the space is finite dimensional.

[Be] G. Beer, "Wijsman convergence of convex sets under renorming", Preprint 1991.

[BLLN] G. Beer, A. Lechicki, S. Levi, S. Naimpally, "Distance functionals and suprema of hyperspace topologies". To appear in *Annali di Mat. pura ed appl.*

[CLP] C. Costantini, S. Levi, J. Pelant, "Infima of hyperspace topologies", Preprint 1992.

[CLZ] C. Constantini, S. Levi, J. Zeminski, "Metrics that generate the same hyperspace convergence", Preprint 1992.

WEAK TWO-LEVEL OPTIMIZATION PROBLEMS AND TYKHONOV REGULARIZATION

Pierre LORIDAN

Département de Mathématiques,
Laboratoire d'Analyse numérique, Univ. de Bourgogne
BP.138, 21004 Dijon, France

Jacqueline MORGAN

Dipartimento di Matematica e Applicazioni
Complesso di Monte San Angelo, Via Cintia
80126, Napoli, Italia

Let U and V be two finite dimensional euclidian spaces, X a non empty subset of U , K a multifunction from U to the nonempty subsets of V , f_1 and f_2 two functionals defined on $X \times V$ and valued in $\mathbb{R}$. We consider the following two-level optimization problem (with explicit constraints in the lower level problem) corresponding to a weak Stackelberg problem in which the set of solutions to the lower level is not a singleton :

$$\text{Find } \bar{x} \in X \text{ such that: } \sup_{y \in M_2(\bar{x})} f_1(\bar{x}, y) = \min_{x \in X} \sup_{y \in M_2(x)} f_1(x, y)$$

(S) where $M_2(x)$ is the set of optimal solutions to the lower level problem

$$P(x): \inf_{y \in K(x)} f_2(x, y)$$

When the problem (S) fail to have a solution, in order to obtain an approximation of (S), ϵ -regularizations have been considered in preceeding papers ([LO-MO.1],[LO-MO.2],[LO-MO.3],[LO-MO.4],[LI-MO],[MA-MO]). Here, in order to transform the problem (S) in a "better" one, different new regularizations are presented and more particularly a regularization of a Tykhonov type, which allows to substitute an ill-posed two-level problem by a quasi well-posed and approximating well-posed problem ([MO.]. Approximation results for the regularized problems and connections between the different approaches are given.

- [LO-MO.1] P.Loridan, J.Morgan, "New results on approximate solutions in two level optimization", Optimization, 20, 1989, 819-836.
- [LO-MO.2] P.Loridan, J.Morgan, " ϵ -regularized two-level optimization problems", Lecture Notes in Mathematics 1405, Springer Verlag, 1989, 99-113.
- [LO-MO.3] P.Loridan, J.Morgan, "Quasi convex lower level problem and applications in two-level optimization", Lecture Notes in Economic and Mathematical Systems, Springer-Verlag, n°345, 1990, 325-341.
- [LO-MO.4] P.Loridan, J.Morgan, "On strict ϵ -solution for a two level optimization problem", Operation Research Proceedings of the International Conference on Operation Research 90 in Vienna, G.Feichtinger, W.Buhler, F.J.Radermecher and P.Stahly eds., Springer Verlag, 1992, 165-172.
- [LI-MO] M.B.Lignola and J.Morgan, "Topological existence and stability for Stackelberg problems" Preprint n°3-1991, Pubblicazione dell' Istituto di Matematica dell' Universita' di Salerno.
- [MA-MO] L.Mallozzi, J.Morgan, " ϵ -mixed strategies for static continuous Stackelberg problem", to appear on J.O.T.A.
- [MO.] J.Morgan, "Constrained well-posed two-level optimization problems", Nonsmooth Optimization and related topics, Ettore Majorana International Sciences Series, Edited by F.Clarke, V.Demianov and F.Giannessi, Plenum Press, New-York, 1989, 307-326.

Hypertopologies: old and new approaches

R.Lucchetti
 Department of Mathematics
 University of Milano
 Via Saldini 50, 20133 Milano, Italy

Abstract

Two recent papers [BL1,SZ] deal with the description of the topologies on the subspace $CL(X)$ of the closed sets of a metrizable space X . They are presented as the weakest topologies making continuous certain families of geometric functionals defined on $CL(X)$. Not only this allows to unify the subject, but it also suggests the applications of general methods to get results in optimization, in presence of perturbations described by means of moving sets, as epigraphs, constrained sets etc. [BL2,LSS]. A new way of describing hypertopologies also outside the metric case is investigated in [LP].

References

- [BL1] G.Beer-R.Lucchetti: "*Weak topologies on the closed subsets of a metrizable space*" (to appear on Trans.Amer.Math.Soc.).
- [BL2] G.Beer-R.Lucchetti: paper in progress.
- [LP] R.Lucchetti-A.Pasquale: paper in progress.
- [LSS] R. Lucchetti-P.Shunmugaraj-Y. Sonntag: "*Recent hypertopologies and continuity of the value function and of the constrained level sets*" (submitted).
- [SZ] Y.Sonntag-C.Zalinescu: "*Set convergences:an attempt of classification*" (submitted).

Bifurcation for variational inequalities with constraints on the derivatives

Antonio Marino

Dipartimento di Matematica, Università di Pisa,

Via Buonarroti 2, 56127 Pisa, Italy

Abstract

There are several problems which can be framed in the following abstract scheme : to study, for a given real smooth function f on a manifold V , the equations

$$\operatorname{grad}_V f(u) = 0 \quad u \in V; \quad (1)$$

$$\mathcal{U}' = -\operatorname{grad}_V f(\mathcal{U}) \quad \mathcal{U} : I \rightarrow V \text{ (} I \text{ is an interval)}. \quad (2)$$

For the nonsmooth case, several important theories have been developed with various aims by Brézis, Clarke, Rockafellar, Temam, among others.

The theory of subdifferential and evolution equations for convex functions (and suitable perturbations of theirs), V being a convex set, provides a classical extension of (1) and (2), which allows to study problems of differential equations and inequalities of elliptic and parabolic type.

Nevertheless, several problems related to the above mentioned ones lead to consider some classes of nonconvex constraints V .

A typical example is provided by the study of Van Karman's Inequalities, which are related to the problem of the well-clamped plate, in presence of unilateral constraints on the displacement or on the curvature. To be more precise, given a bounded open subset

Ω of $\mathbf{R}^2$, a C^2 function $F_o : \bar{\Omega} \rightarrow \mathbf{R}$ and two functions (the obstacles) $\varphi_1, \varphi_2 : \Omega \rightarrow \mathbf{R}$ with $\varphi_1 \leq 0 \leq \varphi_2$, one considers the problem

$$(V.K.I.) \quad \begin{cases} \Delta^2 h + [u, u] = 0; \\ \int_{\Omega} (\Delta u \Delta(v - u) - [\lambda F_o + h, u](v - u)) \, dx dy \geq 0 \quad \forall v \in \mathbf{K}; \\ \lambda \in \mathbf{R}, h \in W_o^{2,2}(\Omega), u \in \mathbf{K}, u \neq 0 \end{cases}$$

(u represents the displacement of the plate and h the Airy stress function), having defined $[u, v] = u_{xx}v_{yy} - 2u_{xy}v_{xy} + u_{yy}v_{xx}$ and

$$\mathbf{K} = \mathbf{K}_1 = \{u \in W_o^{2,2}(\Omega) \mid \varphi_1 \leq u \leq \varphi_2\}$$

or

$$\mathbf{K} = \mathbf{K}_2 = \{u \in W_o^{2,2}(\Omega) \mid \varphi_1 \leq \text{eigenvalues of } H_u \leq \varphi_2\};$$

(H_u is the Hessian matrix of u).

To study the previous problem, we may introduce suitable definitions of "subdifferential" and "lower critical point" (extending the meaning of equation (1)). Now, introducing the functional $f : W_o^{2,2}(\Omega) \rightarrow \mathbf{R}$ defined by

$$f(u) = \int_{\Omega} \left(\frac{1}{2} (\Delta u)^2 + \frac{1}{4} ([\Delta^{-2}[u, u], u] u) \right) \, dx dy$$

and $\mathbf{M}_{\rho}$ defined by

$$\mathbf{M}_{\rho} = \{u \in W_o^{2,2}(\Omega) \mid \int_{\Omega} [F_o, u] u \, dx dy = \rho\}$$

the problem (V.K.I.) is reduced to finding the lower critical points of f on the constraint $\mathbf{V} = \mathbf{K} \cap \mathbf{M}_{\rho}$, as the parameter ρ varies.

The problem with assigned ρ has been studied for $\mathbf{K} = \mathbf{K}_1$, and some results are exposed in [2,4].

We wish to consider here the bifurcation problem for (V.K.I.), namely the relationship between "branches of solutions" (λ, h, u) of (V.K.I.), with ρ close to zero, and the nontrivial solutions of the corresponding "0-asymptotic" problem :

$$(V.K.I.)_{\infty} \quad \begin{cases} \int_{\Omega} (\Delta u \Delta(v - u) - [\lambda F_o, u](v - u)) \, dx dy \geq 0 \quad \forall v \in \mathbf{K}_{\infty}; \\ \lambda \in \mathbf{R}, u \in \mathbf{K}_{\infty}, u \neq 0 \end{cases}$$

where $\mathbf{K}_{\infty} = \overline{\bigcup_{t \in \mathbf{R}} t\mathbf{K}}$.

Notice that this problem is twofold:

I- If (λ, u) solves $(V.K.I.)_{\infty}$, does the "eigenvalue" λ give rise to bifurcating branches of solutions of (V.K.I.) (and *viceversa*)?

II) Do eigenvalues of $(V.K.I.)_\omega$ exist?

It is worth noticing that for the study of $(V.K.I.)$, in the case of assigned ρ and in the bifurcation case as well, the "curves of steepest descent" for f on $\mathbf{K} \cap \mathbf{M}_\rho$ (which extend the meaning of equation (2)) are used.

References

- [1] G. Còbanov, A. Marino, D. Scolozzi, Multiplicity of eigenvalues of the Laplace operator with respect to an obstacle and nontangency conditions, *Nonlinear Anal. The. Meth. Appl.* **15** (1990), 199-215.
- [2] G. Còbanov, D. Scolozzi, Equazioni di Von Karman rispetto a un ostacolo. Esistenza e molteplicità di soluzioni. (preprint), *Dip. Mat. Pisa*, 256(556), Pisa, 1990.
- [3] E. De Giorgi, A. Marino, M. Tosques, Problemi di evoluzione in spazi metrici e curve di massima pendenza, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (S)* **68** (1980), 180-187.
- [4] M. Degiovanni, A. Marino, Eigenvalue problems for Von Karman's equations with obstacle, in *Symmetry in Nature*, Quaderni S.N.S., volume in honour of L. A. Radicati di Brozolo, Pisa 1989.
- [5] M. Degiovanni, A. Marino, M. Tosques, Evolution equations with lack of convexity, *Nonlinear Anal. The. Meth. Appl.* (12) **9** (985), 1401-1443.
- [6] A. Marino, C. Saccon, M. Tosques, Curves of maximal slope and parabolic variational inequalities on nonconvex constraints, *Ann. Sc. Norm. Sup. Pisa*, **XVI** (1989), 281-330.

DUAL REPRESENTATION OF COOPERATIVE GAMES

J.-E. Martinez-Legaz

Departament d'Economia i d'Història Econòmica

Universitat Autònoma de Barcelona

Abstract.

One can associate to a cooperative game, defined by a set of players $N = \{1, 2, \dots, n\}$ and the characteristic function $v : 2^N \rightarrow \mathbb{R}$ (assumed to be non decreasing with respect to inclusion and to satisfy $v(\emptyset) = 0$), the function $\pi : \mathbb{R}_+^n \rightarrow \mathbb{R}$

given by $\pi(w) = \max_S \{v(S) - \sum_{i \in S} w_i\}$ for all $w = (w_1, \dots, w_n) \in \mathbb{R}_+^n$. Interpreting the

components of w as wages for the players, π associates to each possible set of wages the net profit an employer can get hiring some set of players according to these wages and then obtaining the value they generate by acting as a coalition. Clearly, π is a nonincreasing polyhedral convex function such that $\inf_w \pi(w) = 0$ and, at any $w \in \mathbb{R}_+^n$,

the subdifferential $\partial\pi(w)$ intersects the set $\{0, -1\}^n$. The function π provides exactly the same information as the characteristic function v , since one has $v(S) = \inf_w \{ \pi(w)$

$+ \sum_{i \in S} w_i \}$; it is worth noticing that, for this equality to hold, no assumption on v is

needed. The interpretation of this expression is the following : the value of a coalition is the total amount of agents (i.e., the players and the employer) are sure to obtain whatever the wages may be. If v is superadditive, the function π satisfies the condition

$\pi(w) \geq \inf_{w_{N \setminus S}} \pi(w_S, w_{N \setminus S}) + \inf_{w_S} \pi(w_S, w_{N \setminus S})$ for all $w \in \mathbb{R}_+^n$ and $S \subset N$, and

conversely. For arbitrary games, the core coincides with the set of vectors $x \in \mathbb{R}_+^n$

such that $\pi(x) = 0$ and $(-1, \dots, -1) \in \partial\pi(x)$.

ERGODIC THEORY AND THE CALCULUS OF VARIATION

E. CHABI

Faculté des Sciences
Université M.Y. Ismail
Meknès (Maroc)

G. MICHAILLE

Laboratoire d'Analyse Convexe
Département des Sciences Mathématiques
Université Montpellier II

Let $(\Sigma, \mathcal{C}, P)$ be a probability space, Ω a bounded regular open set in $\mathbb{R}^d$, $M(\mathbb{R}^d)$ the set of non negative regular Borel measure on $\mathbb{R}^d$ equipped with its Borel field $\mathcal{B}(\mathbb{R}^d)$. $M(\Omega)$ is the set of non negative bounded regular Borel measure on Ω .

A Random Borel measure is any map from Σ into $M(\mathbb{R}^d)$ which is $(\mathcal{C}, \mathcal{M})$ measurable, $\mathcal{M}$ denoting the trace of the product tribe of $\mathbb{R} \cup \{+\infty\}$ $\mathcal{B}(\mathbb{R}^d)$.

Given $(T_z)_{z \in S}$ a group of P -preserving transformation on Σ where S is any subgroup $k\mathbb{Z}^d$ of $(\mathbb{Z}^d, +)$, and μ a Random Borel measure satisfying, for every bounded set A in $\mathcal{B}(\mathbb{R}^d)$, the two following properties

$$\mu(\omega)(A+z) = \mu(T_{-z}\omega)(A)$$

$$\omega \rightarrow \mu(\omega)(A) \text{ belongs to } L^1(\Sigma, \mathcal{C}, P),$$

we define the sequence μ_n from Σ into $M(\Omega)$ by

$$\mu_n(\omega)(A) := \epsilon_n^d \mu(\omega) \left(\frac{1}{\epsilon_n} A \right)$$

where ϵ_n tends to 0^+ .

Using Ergodic Theory and more precisely, an additive ergodic theorem due to Nguyen Xuan Xanh and H. Zessin, we prove the following result

THEOREM 1.

(i) If a.s., $\{\mu_n(\omega); n \in \mathbb{N}\}$ is tight then a.s., $\mu_n(\omega)$ converges for the narrow topology towards $\theta(\omega) dx$ where

$$\theta(\omega) := \frac{1}{k^d} E^{\mathcal{F}} \mu(\cdot)([0, k]^d),$$

$E^{\mathcal{F}}$ denoting the conditional expectation operator with respect to the tribe $\mathcal{F} := \{E \in \mathcal{C}, T_z E = E \ \forall z \in S\}$;

(ii) if $(T_z)_{z \in S}$ is Ergodic, that is to say if $\mathcal{F}$ contains only sets of $\mathcal{C}$ with probability 0 or 1 then a.s., $\mu_n(\omega)$ converges for the narrow topology towards θ dx where

$$\theta := \frac{1}{k^d} E \mu(\cdot)([0, k]^d).$$

In the case where $\mu(\omega) = f(\omega, \cdot) dx$, $f(\omega, \cdot)$ belonging to $L^p_{loc}(\mathbb{R}^d)$, $1 \leq p \leq +\infty$, we get under suitable measurability hypothesis, the following stronger result.

THEOREM 2. Setting $f_n(\omega, x) := f(\omega, \frac{x}{\epsilon_n})$, we have

- (i) in the case $1 < p \leq +\infty$, a.s. $f_n(\omega, \cdot)$ converges towards $\frac{1}{k^d} E^{\mathcal{F}} \int_{[0, k]^d} f(\cdot, x) dx$ in $L^p(\Omega)$ weak (* weak if $p = +\infty$);
- (ii) in the case $p = 1$, when $(T_z)_{z \in S}$ is Ergodic, a.s. $f_n(\omega, \cdot)$ converges towards $\frac{1}{k^d} E \int_{[0, k]^d} f(\cdot, x) dx$ in $L^1(\Omega)$ weak.

These two results have been used to solve stochastic homogenization problems, in particular to construct suitable random test functions and random Radon measure in nonconvex stochastic homogenization and in a stochastic version of Darcy's Law.

REFERENCES

- H. ATTOUCH, Variational Convergence for functions and operators, Research Notes in Mathematics, Pitman, London (1984).
- J. P. AUBIN, H. FRANKOWSKA, Set Valued Analysis, Birkhäuser, (1990).
- C. CASTAING, M. VALADIER, Convex Analysis and measurable multifunctions, Lecture Notes in Mathematics, Vol 580, Springer (1977).
- U. KRENGEL, Ergodic Theorems, de Gruyter, Studies in Mathematics.
- K. MESSAOUDI, G. MICHAILLE, Stochastic Homogenization of Nonconvex Integral Functionals, Duality in the Convex Case, Séminaire d'Analyse Convexe, Montpellier 1991, Exposé n° 14.
- NGUYEN XUAN XANH, H. ZESSIN, Ergodic Theorems for Spatial Processes, Z. Wah. Verw. Gebiete, 48, 133-158 (1979).

**VARIATIONAL TOPOLOGICAL PROPERTIES OF
THE SPACE OF DIRICHLET FORMS**

Umberto MOSCO

Abstract.

We study compactness and closure properties of families of local and nonlocal Dirichlet forms with respect to suitable variational topologies.

Generalized second-order derivatives of convex functions in locally convex topological vector spaces

James Louis Ndoutoume *

Institut Africain d'Informatique B. P. 2263, Libreville, Gabon

and

Michel Théra

Université de Limoges, 87060 Limoges Cedex, France

Abstract

Generalized second-order derivatives introduced by T. Rockafellar in the finite dimensional setting are extended to convex functions defined on locally convex topological spaces. The main result which is obtained is the exhibition of a particular generalized Hessian when the function admits a generalized second derivative

References

[1] H. Attouch, *Variational convergences for functions and operators*. Pitman, London, 1984.

*The author would like to thank the Commission for Development and Education from which he obtained a grant to attend the meeting

- [2]J.-M. Borwein and D. Noll, Second order differentiability of convex functions: A. D. Alexandrov's Theorem in Hilbert spaces. Preprint, Dalhousie University, 1991.
- [3]C. N. Do, Generalized second-order derivatives of convex functions in reflexive Banach spaces. Preprint, 1991.
- [4]S. Navry, Formes quadratiques généralisées. C. R. Acad. Sci. Paris, **271**, série A. 1054-1057, 1970.
- [5] L. Ndoutoume, Calcul différentiel généralisé du second ordre. Publication **AVAMAC** de l'Université de Perpignan, 1987.
- [6]J. L. Ndoutoume, *Epi-convergence en calcul différentiel généralisé. Résultats de convergence et Approximation*. These de doctorat, Université de Perpignan, 1987.
- [7]R. T. Rockafellar, Maximal monotone relations and the second order derivative of nonsmooth functions. Ann. Inst. H. Poincaré, Analyse non-linéaire, **2**, 167-184, 1985.
- [8]R. T. Rockafellar, Second order conditions in mathematical programming obtained by the way of epiderivatives, Math. of Oper. Res. **14**, (3), 462-484, 1989.
- [9]R. T. Rockafellar, Generalized second derivatives of convex functions and saddle functions, Trans. Am. Math. Soc., **320**, 810-822, 1990.

Graphical convergence and generalized second derivatives for nonsmooth functions

Dominikus Noll

Universität Stuttgart, 7000 Stuttgart, F. R. G

Abstract

Second order epi derivatives for convex and nonsmooth functions, as introduced by R.T. Rockafellar around 1987, have emerged in an important role in nonsmooth analysis and optimization. Here we present some results obtained recently by D. Noll [3], [4], and J. Borwein and D. Noll [1]. In particular, we address the following questions:

- Give a refined analysis of the interrelation between second epi derivatives and the usual pointwise second derivatives.
- Give a geometric characterization of second order epi differentiability. A nonsmooth version of the Theorem of Meusnier.
- In the convex case, give a link between second epi derivatives and approximate second derivatives in the sense of J.B. Hiriart-Urruty. How to define the Dupin indicatrix using second epi derivatives?
- The infinite dimensional dilemma! What is the right notion of convergence for the second order difference quotients of non-convex functions in Hilbert space?

References

- [1] J.M. Borwein, D. Noll, 'Second order differentiability of convex functions in Banach spaces', Trans. Amer. Math. Soc., under press.
- [2] J.B. Hiriart-Urruty, A. Seeger, 'The second-order subdifferential and the Dupin indicatrices of a nondifferentiable convex function', Proc. London Math. Soc. 58 (1989), 351 - 365.

[3] D. Noll, 'Generalized second fundamental form for Lipschitzian hypersurfaces by way of second epi derivatives', *Bull. Can. Math.* under press.

[4] D. Noll, 'Second order differentiability of integral functionals on Sobolev spaces and L^2 -spaces', to appear.

Approximation by convex functions

Pier Luigi Papini

Dipartimento di Matematica, Piazza Porta S. Donato 5, 40127 Bologna, Italie

Abstract

Consider the following problem: let X be a space of real functions, endowed with a norm; for $f \in X$, consider the function(s) which best approximate f from the class of convex functions. Also, if C denotes the subset of X containing all continuous functions, let us consider the same problem by using as approximants the elements of the set $A \cap C$. The sets A and $A \cap C$ are cones; in case the norm of X is "good" and A (or $A \cap C$) is closed, then some obvious results spring from the general theory of approximation, while simple characterizations of best approximations are possible. But in different situations, some problems can arise, for example concerning convergence of sequences which best approximate f with approximating norms; in this context, many papers appeared recently. Here we try to survey some results of this type.

References

- [1] R. Huotari - D. Zwick, Approximation in the mean by convex functions, Numer. Funct. Anal. and Optimiz., 10, 489-498 (1989).
- [2] D.A. Legg - D.W. Townsend, The Polya-1 property for convex approximation, Approx. Th. Appl. 6, No.4, 9-17 (1990).
- [3] V.A. Ubhaya, L_p Approximation by quasi-convex and convex functions, J. Math. Anal. Appl., 139, 574-585 (1989).
- [4] D. Zwick, Best approximation by convex functions, Amer. Math. Monthly, 94, 528-534 (1987).
- [5] D. Zwick, Characterizing shape preserving L_1 -approximation, Proc. Amer. Math. Soc., 103, 1139-1146 (1988).

Convergent and divergent concepts from convergence theory and their uses in optimization theory

Jean-Paul Penot

Faculté des Sciences, Av. de l'Université 64000 Pau, France

Abstract

We endeavour to give a general account of the impact of convergence notions for optimization theory. Whether these tools can be considered as a convergent array of concepts or as a divergent and diverse bundle of tools is still a matter of taste or opinion. At least it cannot be denied that the number of topics in optimization theory for which notions of convergence become instrumental is increasing and covers a large extend of subjects: approximation of functions and of sets, optimality conditions, sensitivity analysis, well-posedness. . . A number of recent or less recent contributions of the author to this stream are reviewed. The appearance of "alien terms" in optimality conditions is pointed out as it corresponds to a well known phenomenon in homogenization theory.

References

- [1] H. Attouch, *Variational convergence for functions and operators*, Applicable Math. Series 129, Pitman, London, 1984.
- [2] H. Attouch and G. Beer, *On the convergence of subdifferentials of convex functions* Séminaire d'analyse convexe, Montpellier, vol. 21 (1991) (8) 1-11.

- [3] H. Attouch, J.L. Ndoutoume and M. Théra, *On the equivalence between epi-convergence of sequences of functions and graph-convergence of their derivatives in Banach spaces*, Séminaire d'analyse convexe, Montpellier, vol. 20 (1990) (9) 1-45.
- [4] H. Attouch and R. J.-B. Wets *Quantitative stability of variational systems : I. The epigraphical distance*, Trans. Amer. Math. Soc. 328 (2) (1991), 695-729.
- [5] D. Azé and J.-P. Penot, *Recent quantitative results about the convergence of convex sets and functions* in "Functional Analysis and Approximation", Bagni di Lucca, May 1988, P.L. Papini, ed. Pitagora, Bologna (1988).
- [6] D. Azé and J.-P. Penot, *Qualitative results about the convergence of convex sets and convex functions* in "Optimization and Nonlinear Analysis" workshop Haifa, March 1990, A. Ioffe, M. Marcus and S. Reich, ed., Pitman, London (1992).
- [7] D. Azé and J.-P. Penot, *Operations on convergent families of sets and functions*, Optimization 21, (1990) 4, 521-534.
- [8] G. Beer, *The slice topology : a viable alternative to Mosco convergence in nonreflexive spaces*, Séminaire d'analyse convexe, Montpellier, vol. 21 (1991) (3) 1-35.
- [9] G. Beer and R. Lucchetti, *The epi-distance topology: continuity and stability results with applications to convex optimization problems*, preprint 1990.
- [10] L. Contesse and J.-P. Penot *Continuity of the Fenchel correspondence and continuity of polarities*, J. Math. Anal. Appl. (156) (1991) , 305- 328.
- [11] B. Lemaire *Bons comportement et conditionnement linéaire* Journées d'analyse non-linéaire et optimisation, Avignon, Nov. 1991.
- [12] J.-P. Penot, *Penalization and regularization methods for nonsmooth analysis*, unpublished lecture, workshop on Nonlinear Analysis and Optimization, Louvain la Neuve, June 16-17, 1983.
- [13] J.-P. Penot, *Preservation of persistence and stability under intersections and operations*, preprint, Nov. 1986.
- [14] J.-P. Penot, *Topologies and convergences on the space of convex functions*, Nonlinear Analysis, Th., Methods Appl. 18 (1992).
- [15] J.-P. Penot, *The cosmic Hausdorff topology, the bounded Hausdorff topology and continuity of polarity*, Proc. Amer. Math. Soc. 113, (1) (1991), 275-285.
- [16] J.-P. Penot, *On the convergence of subdifferentials of convex functions*, preprint, submitted January 1991.
- [17] J.-P. Penot, *Second order derivatives : comparisons of two types of epi-derivatives*, Sixth French-German Conference on Optimization, Lambrecht, June 1991, D. Pallaschke ed., Springer Verlag.
- [18] J.-P. Penot, *Second order generalized derivatives : relationships with convergence notions*, in "Nonsmooth Analysis, Methods and Applications, Erice, June 1991", F. Giannessi ed., Springer Verlag.

Calculus of epi-derivatives and proto-derivatives

René Poliquin *

University of Alberta, Math.Dept., Edmonton Canada, T6G 2G1

Abstract

A calculus of first- and second-order epi-derivatives is presented for amenable and fully amenable functions (all functions are considered on a finite dimensional space). An amenable function is the composition of a lower semicontinuous proper convex function with a smooth mapping and a natural constraint qualification. A fully amenable function is an amenable function where the convex function is piecewise linear-quadratic and the smooth function is of class C^2 . The calculus rules have direct applications in optimization because most problems encountered in practice can be reformulated using fully amenable functions; in addition, first- and second-order optimality conditions can be derived using epi-derivatives. Finally, calculus rules for the proto-derivatives of subgradient mappings of fully amenable functions are presented. These calculus rules for proto-derivatives have applications in the sensitivity analysis of optimal solution mappings in parametric optimization.

References

- [1] R.A.Poliquin and R.T.Rockafellar. A calculus of epi-derivatives applicable to optimization. *Can. Jour. Math.* to appear.

*This work was supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC) under grant OGP 41983

[2] R.A.Poliquin and R.T.Rockafellar, Amenable functions in optimization, to appear in the proceedings of the International school of mathematics G. Stampacchia, 10th course: Nonsmooth Optimization: Methods and Applications.

Solving Nonsmooth equations via Generalized Jacobians and Iteration Functions

Liqun Qi *

School of Mathematics, University of New South Wales
Kensington, NSW 2032, Australia

Abstract

Systems of nonlinear equations defined by nondifferentiable, locally Lipschitzian functions arise from many different areas including partial differential equations, nonlinear complementarity, variational inequality, nonlinear programming and maximal monotone operator problems [2]. The generalized Jacobians of the functions defining these nonsmooth equations are set-valued. Superlinearly convergent Newton's methods for solving nonsmooth equations are constructed via generalized Jacobians [3] [4]. A characterization of superlinear convergence under the condition of semismoothness extends the classic results Dennis-Moré for smooth equations [2]. Broyden-type methods are also developed for solving nonsmooth equations [1]. Global convergent methods are established via iteration functions.

References

*This work is supported by the Australian Research Council. The author would like also thank the meeting organizers Hedy Attouch and Michel Théra for their help during the meeting.

- [1] X. Chen and L. Qi, A Broyden-like method for solving nonsmooth equations, in preparation.
- [2] J.S. Pang and L. Qi, Nonsmooth equations: Motivation and algorithms, to appear in: SIAM J. Optimization.
- [3] L. Qi, Convergence analysis of some algorithms for solving nonsmooth equations, to appear in: Mathematics of Operations Research.
- [4] L. Qi and J. Sun, A nonsmooth version of Newton's method, to appear in: Mathematical Programming.

Necessary and Sufficient Conditions for the Existence of Densely Defined Selections of Multivalued Mappings

Julian P. Revalski

Institute of Mathematics, Bulgarian Academy of Sciences,
Acad. G. Bonchev Street, Block 8, 1113 Sofia, Bulgaria

Abstract

Let $F: X \rightarrow Y$ be a multivalued mapping with non-empty images acting from the topological space X into the topological space Y (i.e. for each $x \in X$ the value $F(x)$ is a non-empty subset of Y). We give here sufficient (and necessary) conditions for the existence of an upper semicontinuous and non-empty-compact-valued (usco) mapping $G: X_1 \rightarrow Y$, where X_1 is a dense G_δ -subset of X and G is a selection of F . In contrast to what is widely accepted under "*a selection of F on X_1* " we understand here a set-valued mapping $G: X_1 \rightarrow Y$ such that $\emptyset \neq G(x) \subset F(x)$ for every $x \in X_1$. In some particular cases the selection G coincides with the restriction of F on X_1 . If the range space Y is completely metrizable, then G can be considered to be single-valued.

More precisely, we call F *lower demicontinuous* (ldc) in X if for every open V in Y , the interior of the closure of the set $F^{-1}(V) := \{x \in X: F(x) \cap V \neq \emptyset\}$ is dense in the closure of $F^{-1}(V)$, i.e. $\text{IntCl}(F^{-1}(V))$ is dense in $\text{Cl}(F^{-1}(V))$. The class of ldc mappings contains for example all lower semicontinuous mappings and all minimal usco mappings. Several results are proved asserting that a given ldc mapping F admits an usco selection defined on a dense subset of X . A typical theorem reads as follows:

Let $F: X \rightarrow Y$ be an open ldc mapping with closed graph, where X is a Baire space and Y contains a dense Čech complete subspace Y_1 . Then there exist a dense G_δ -subset X_1 of X and an usco mapping $G: X_1 \rightarrow Y_1$ which is a selection of F . If, in addition, Y_1 is completely metrizable, then G can be considered to be single-valued.

In fact, the existence of such kind of selections for every mapping from the above class is a characterization of the fact that the space Y contains a dense Čech complete subspace or a dense completely metrizable subspace.

The above result generalizes some of the results in the recent paper of E. Michael [M].

We describe also situations in which the mapping F itself (not merely a selection of it) is usco at the points of a dense G_δ -subset of X . Here a typical result says that, if $F: X \rightarrow Y$ has a closed graph, X is a Baire space, Y is Čech complete and for every open $V \subset Y$ the interior of the set $\{x \in X: \text{cl } F(x) \subset V\}$ is dense in $F^{-1}(V)$, then there exists a dense G_δ -subset X_1 of X at the points of which F is usco. If, in addition, Y is completely metrizable then F is single-valued at the points of X_1 .

These results are used to obtain some new as well as known results like:

- to get a new version of the classical Lavrentieff theorem concerning the extension of a densely defined homeomorphism to a subset containing a dense G_δ -subset of the domain space;
- to prove that a given convex and continuous function is differentiable at the points of some dense G_δ -subset of its domain ([Ph]);
- to obtain generic results about well-posed optimization problems ([LP], [ČK1], [ČK2], [ČKR]);
- to derive results about generic non-multivaluedness of metric projections and antiprojections ([Zh]);

References

- [ČK1] M.M. Čoban and P.S. Kenderov, *Dense Gâteaux differentiability of the sup-norm in $C(T)$ and the topological properties of T* , Compt. rend. Acad. bulg. Sci. **38**(1985), 1603-1604.
- [ČK2] M.M. Čoban and P.S. Kenderov, *Generic Gâteaux differentiability of convex functionals in $C(T)$ and the topological properties of T* , Math. and Education in Math., Proc. of the XV-th Spring Conf. of the Union of Bulg. Mathematicians, 1986, pp. 141-149.
- [ČKR] M.M. Čoban, P.S. Kenderov and J.P. Revalski, *Generic well-posedness of optimization problems in topological spaces*, Mathematika, **36**(1989), 301-324.
- [LP] R. Lucchetti and F. Patrone, *Sulla densità e genericità di alcuni problemi di minimo ben posti*, Boll. U.M.I. **15-B**(1978), 225-240.
- [M] E. Michael, *Almost complete spaces, hypercomplete spaces and related mapping theorems*, Topology Appl. **41**.n.1-2,(1991), 113-130.
- [Ph] R.R. Phelps, *Convex Functions, Monotone Operators and Differentiability*, Lect. Notes in Math., #1364, Springer Verlag, Berlin, 1989.
- [Zh] N.V. Zhivkov, *Continuity and non-multivaluedness properties of metric projections and antiprojections*, Serdica Bulg. Math. Publ. **8**(1982), 378-385.

TOPOLOGICAL DEGREE FOR MAXIMAL MONOTONE OPERATORS

Hassan RIAHI

Université CADI AYYAD, Faculté des Sciences I, B.P. S. 15, Marrakech, MAROC

Abstract

The generalized degree theory is a replacement for the Brouwer and Leray-Schauder degrees. The purpose of this work is two-fold. One goal is to show how Browder's degree, given for operators of class (S_+) , see [4], can be extended to the case of maximal monotone operators by relying on generalized Yosida approximations. Particular attention is paid to the normalization and invariance under homotopies for the topological degree we define. This allows us to extend some recent results of Attouch, Penot and Riahi [2] about the continuation method for solutions of parametrized monotone nonlinear equations by withdrawing the compactness assumptions. It is also possible that our definition, by relying on subdifferentials, could be used to establish definitions of topological degree for real convex functions and convex-concave saddle bifunctions.

1- Let be given a real reflexive Banach space X with the topological dual space X^* , such that X and X^* are locally uniformly convex. This implies that the duality mapping J of X is a homeomorphism between X and X^* .

In the sequel we will identify an operator $A : X \rightrightarrows X^*$ with its graph in $X \times X^*$. An operator A is said to be monotone if for any $(x_1, y_1), (x_2, y_2) \in A$, one has $\langle y_1 - y_2, x_1 - x_2 \rangle \geq 0$. A is maximal monotone if it is maximal in the family of monotone operators in $X \times X^*$, ordered by inclusion.

The Yosida approximation is given by $A_\lambda x = (A^{-1} + \lambda J^{-1})^{-1}x$. An extension, called generalized Yosida approximation, we'll use is given by $A^\lambda = A_\lambda + \lambda J$. A^λ is of class (S_+) .

Theorem 1. If a family of maximal monotone operators $\{A_t \subset X \times X^* ; t \in T\}$ is graph-continuous (given $t_i \rightarrow t$ in T one has $\text{Limsup}_{i \in I} A_{t_i} \subset A_t \subset \text{Liminf}_{i \in I} A_{t_i}$) then for any open bounded subset $\Omega \subset X$, the family $\{(A_t)^\lambda : \bar{\Omega} \rightarrow X^* ; t \in T, \lambda > 0\}$ is of class (S_+) .

2- Now, we are in a position to define the topological degree for a maximal monotone operator A over Ω at zero by the formula :

$$\deg(A, \Omega, 0) = \lim_{\lambda \downarrow 0} d(A^\lambda, \Omega, 0). \quad (1)$$

where d denotes the Browder's degree for operators of class (S_+) , we refer to [4] for more details. We verify that in the definition above, the degree function $\deg$ is independent of $\lambda > 0$ for λ sufficiently small. The following Theorem summarizes familiar properties of degree theory for maximal monotone mappings.

Theorem 2. Let Ω be an open bounded subset, $A \subset X \times X^*$ be maximal monotone. Then we have :

- (i) $\deg(J, \Omega, 0) = 1$, provided $0 \in \Omega$;

- (ii) $\deg(A, \Omega, 0) = 0$ whenever $0 \notin A(\bar{\Omega})$;

- (iii) if the homotopy of maximal monotone operators $\{A_t \subset X \times X^*; t \in T\}$ is graph continuous and satisfies $0 \notin \cup \{A_t(\partial\Omega); t \in T\}$, then $\deg(A_t, \Omega, 0)$ is independent of t in T ;

- (iv) if Ω_1 and Ω_2 are two disjoint subsets of Ω such that $0 \notin A(\bar{\Omega} \setminus \Omega_1 \cup \Omega_2)$, then

$$\deg(A, \Omega, 0) = \deg(A, \Omega_1, 0) + \deg(A, \Omega_2, 0).$$

- (v) On the family of maximal monotone operators, there exists one and only one degree function, with the invariance under graph-continuous homotopies.

As a consequence of this theorem we shall sharpen and extend [2] results for Hilbert to reflexive spaces, as well as ruling out the compactness conditions.

Corollary 3. Suppose that assumptions of the preceding theorem (iii) hold and the following condition is satisfied : for some $t_0 \in T$ one has $\deg(A_{t_0}, \Omega, 0) \neq 0$.

Then for each t in T one has $A_t^{-1}(0)$ is nonempty and contained in Ω .

3- In this brief section we apply the results of section 2 to convex functions and convex-concave saddle bifunctions.

A- Since the subdifferential $\partial\phi$ of a proper lsc function ϕ is a maximal monotone operator, see [1, 5] one can define the degree at 0 relatively to an open bounded subset of X by :

$$\deg(\phi, \Omega, 0) = \deg(\partial\phi, \Omega, 0). \quad (2)$$

Proposition 4. a) For a proper convex lsc function ϕ , $\deg(\phi, \Omega, 0) \neq 0$ and $M(\phi) \cap \partial\Omega = \emptyset$ implies that $\emptyset \neq M(\phi) \subset \Omega$, where $M(\phi)$ denotes the minimum set of the function ϕ .

b) Let $\{\phi_t; t \in T\}$ be a Mosco-epicontinuous family of proper convex lsc functions such that $M(\phi_t) \cap \partial\Omega = \emptyset \quad \forall t \in T$ and $\deg(\phi_{t_0}, \Omega, 0) \neq 0$ for some $t_0 \in T$.

Then for each t in T one has $\emptyset \neq M(\phi_t) \subset \Omega$.

B- For a closed convex-concave saddle bifunction $F: X \times Y \rightarrow \bar{\mathbb{R}}$ which is convex lsc (resp. concave usc) with respect to the variable x in X (resp. y in Y), the operator $A = \partial_1 F x (-\partial_2 F)$, where $\partial_1 F$ and $\partial_2 F$ denote the partial subdifferentials of the convex functions $F(\cdot, y)$ and $-F(x, \cdot)$, see [6], is maximal monotone.

The degree of F at (x, y) is defined as follows $\deg(F, \Omega, (x, y)) = \deg(A, \Omega, (x, y))$. With the above definition one can state the analogue of Proposition 4 for bifunctions:

Proposition 5. a) $\deg(F, \Omega, 0) \neq 0$ implies that $S(F) = \{\text{saddle points of } F \text{ in } X \times Y\} \cap \Omega \neq \emptyset$.
b) Let $\{F_t; t \in T\}$ be a Mosco-epi/hypocontinuous homotopy such that $\deg(F_{t_0}, \Omega, 0) \neq 0$ for some $t_0 \in T$ and $S(F_t) \cap \partial\Omega = \emptyset$ for each t in T . Then for every $t \in T$ one has $\emptyset \neq S(F_t) \subset \Omega$.

SELECTED REFERENCES

- [1] H. Attouch, Variational convergence for functions and operators, Appl. Math. Series Pitman, London, 1984.
- [2] H. Attouch, J.-P. Penot and H. Riahi, The continuation methods and variational convergence, Fixed point theory and applications, B. Baillon, M. Théra, ed., Research Notes in Math. Pitman, London, (1991), 9-32.
- [3] F.E. Browder, Nonlinear operators and nonlinear equations of evolution in Banach spaces, Proc. Symp. Pura Math., 18, 2, AMS Providence, 1976.
- [4] F.E. Browder, Degree of mapping for nonlinear mappings of monotone type; strongly nonlinear mappings, Proc. Nat. Acad. Sci. USA, 89, (1983), 2408-2409.
- [5] R. Phelps, Convex functions, monotone operators, and differentiability, Lecture Notes in Maths. # 1364, Springer-Verlag, Berlin, 1989.
- [6] R.T. Rockafellar, Monotone operators associated with saddle functions and minimax problem, In Nonlinear Func. Anal., F.E. Browder ed., Proc. Pure Math., 18, (1970), 241-250.
- [7] E. Zeidler, Nonlinear functional analysis and its applications, II B, Nonlinear Monotone operators, Springer-Verlag, New York, 1990.

Some Topological Min-Max Theorems via an Alternative Principle for Multifunctions

Biago Ricceri

Universita di Catania, Italy

Abstract

Let X, Y be two non-empty sets and let f be a real function defined on $X \times Y$. We are interested in the classical problem of finding suitable conditions under which the equality

$$(1) \quad \sup_{y \in Y} \inf_{x \in X} f(x, y) = \inf_{x \in X} \sup_{y \in Y} f(x, y)$$

does hold.

In this talk we intend to discuss some of our recent results on this subject. Here is a sample:

THEOREM - *Let X, Y be topological spaces, with Y connected and admitting a continuous bijection onto $[0, 1]$. Assume that, for each $\lambda > \sup_{y \in Y} \inf_{x \in X} f(x, y)$, $x_0 \in X$, $y_0 \in Y$, the sets*

$$\{x \in X : f(x, y_0) \leq \lambda\}$$

and

$$\{y \in Y : f(x_0, y) > \lambda\}$$

are connected. In addition, assume that at least one of the following three sets of conditions is satisfied:

(h₁) $f(x, \cdot)$ is upper semicontinuous in Y for each $x \in X$, and $f(\cdot, y)$ is lower semicontinuous in X for each $y \in Y$;

(h_2) Y is compact, and f is upper semicontinuous in $X \times Y$;

(h_3) X is compact, and f is lower semicontinuous in $X \times Y$.

Under such hypotheses, equality (1) does hold.

Just one remark. Namely, the condition that Y must admit some continuous bijection onto $[0,1]$ is, of course, very restrictive. This is, in practice, the necessary price one has to pay for the generality of the other assumptions. In this connection, consider that, almost every known topological mini-max theorem, ensuring the validity of (1), contains at least the following assumptions: the space X is compact and, for each $\lambda > \sup_{y \in Y} \inf_{x \in X} f(x, y)$ and each non-empty finite set $A \subseteq Y$, the set

$$\bigcap_{y \in A} \{x \in X : f(x, y) \leq \lambda\}$$

is connected (see [1]-[8]).

The fact that the above theorem can badly fail even when Y is only a "little" out of the considered class, is clearly shown by the following

EXAMPLE - Take:

$$X = Y = \{(t, u) \in \mathbf{R}^2 : t^2 + u^2 = 1\}$$

and, for each $(t, u), (v, z) \in X$,

$$f(t, u, v, z) = tv + uz.$$

In this case, of course, we have

$$-1 = \sup_{y \in Y} \inf_{x \in X} f(x, y) < \inf_{x \in X} \sup_{y \in Y} f(x, y) = 1$$

while, except that on Y , each other assumption of the theorem is satisfied.

References

- [1] M.A.GERAGHTY and B.-L.LIN, *Topological minimax theorems*, Proc.Amer.Math.Soc., **91** (1984), 377-380.
- [2] C.HORVATH, *Quelques théorèmes en théorie des mini-max*, C.R.Acad.Sci.Paris, Série I, **310** (1990), 269-272.
- [3] J.KINDLER, *On a minimax theorem of Terkelsen's*, Arch.Math., **55** (1990), 573-583.
- [4] H.KÖNIG, *A general minimax theorem based on connectedness*, Arch.Math., to appear.
- [5] H.KÖNIG and F.ZARTMANN, *New versions of the minimax theorem*, preprint.
- [6] F.TERKELSEN, *Some minimax theorems*, Math.Scand., **31** (1972), 405-413.
- [7] H.TUY, *On a general minimax theorem*, Soviet Math.Dokl., **15** (1974), 1689-1693.
- [8] WU WEN-TSÜN, *A remark on the fundamental theorem in the theory of games*, Sci.Rec.(New Ser.), **3** (1959), 229-233.

Graph Convergence in Mathematical Programming

Stephen M. Robinson

University of Wisconsin-Madison, Madison, WI 53706, USA

Abstract

Methods using graph convergence, particularly the proto-derivatives introduced by R.T. Rockafellar [3], seem to be the most natural tools available for obtaining certain optimality conditions in mathematical programming (see, e.g., [4]). On the other hand, once the optimality conditions are written, the analysis of solutions to those conditions generally proceeds using very different tools, often some variety of implicit-function theorem.

For such analysis the formalism of *normal maps* [2] leads to equations involving single-valued, though generally nonsmooth, functions. For example, the theorem of [2] together with the functional form first introduced by Kojima [1] permits one to conduct a complete local analysis in very much the same way as one would do for a system of nonlinear equations. Noteworthy in this analysis is the complete absence of any graph convergence methodology.

In this lecture we will investigate connections between the graph convergence methodology for obtaining optimality conditions and the analytical methods for exploiting the mathematical structure of those solutions once they are obtained. We will describe the normal-map approach and ask whether any natural relationships exist between the normal map derived from the optimality conditions and the graph-convergence techniques that lead to those conditions.

References

- [1] M. Kojima, "Strongly stable stationary solutions in nonlinear programs," in: S. M. Robinson, ed., *Analysis and Computation of Fixed Points*, Academic Press, New York, 1980, pp. 93-138.
- [2] S. M. Robinson, "An implicit-function theorem for a class of nonsmooth functions," *Math. Operations Res.* **16** (1991) 292-309.
- [3] R.T. Rockafellar, "First and second-order epi-differentiability in nonlinear programming," *Trans. Amer. Math. Soc.* **307** (1988) 75-108.
- [4] R.T. Rockafellar, "Second-order optimality conditions in nonlinear programming obtained by way of pseudo-derivatives," *Math. Operations Res.* **14** (1989) 462-484.

Proto-differentiability of solution mappings in optimization

Terry Rockafellar

University of Washington, Seattle, WA 98195, U.S.A.

Abstract

The most commonly used model for sensitivity analysis of solution mappings in parameterized problems of optimization is the generalized equation model of Robinson, which corresponds closely to optimality conditions in the sense of variational inequalities. Much of the work on generalized equations has centered on obtaining conditions under which the solution is unique and depends Lipschitz continuously on the parameters, but one-sided differentiability of the single-valued mapping obtained under such circumstances has been studied as well. In contrast, proto-differentiability properties can be developed even when the solution mapping is multi-valued. One-sided differentiability is then simply the case where the mapping does happen to be single-valued and locally Lipschitz, which in finite-dimensional spaces can be characterized through Mordukhovich's criterion on coderivatives. Furthermore, proto-differentiability also is an effective tool in broader models in sensitivity analysis formulated in terms of subgradient mappings instead of generalized equations. In that setting it relates to the calculation of perturbations through the solution of auxiliary problems of optimization involving second-order epi-derivatives of functions in the original problem.

APPROXIMATE EULER-LAGRANGE INCLUSION, APPROXIMATE TRANSVERSALITY
CONDITION, AND SENSITIVITY ANALYSIS OF CONVEX PARAMETRIC PROBLEMS
OF CALCULUS OF VARIATIONS.

Alberto SEEGER
University of Avignon
Department of Mathematics
33, rue Louis Pasteur
84000 Avignon, France

ABSTRACT. We study the first-order behaviour of the optimal value function associated to a convex parametric problem of calculus of variations. An important feature of this paper is that we do not assume the existence of optimal trajectories for the unperturbed problem. The concepts of approximate Euler-Lagrange inclusion and approximate transversality condition are key ingredients in the writing of our sensitivity results.

1991 Mathematics Subject Classification : 49 N 99, 90 C 31

Key words : Euler-Lagrange inclusion, transversality condition, approximate subdifferential, sensitivity analysis.

References :

- [1] M.MOUSSAOUI and A.SEEGER. Sensitivity analysis of optimal value functions of convex parametric programs with possibly empty solution sets. Preprint, University of Avignon, March 1992.
- [2] A.SEEGER. Approximate Euler-Lagrange inclusion, approximate transversality condition, and sensitivity analysis of convex parametric problems of calculus of variations. Preprint, University of Avignon, May 1992.

Swimming below icebergs, and the maximal monotonicity of subdifferentials

Stephen Simons

University of California Santa Barbara, CA, USA

Abstract

Let E be a Banach space, $\phi : E \rightarrow \mathbb{R} \cup \{\infty\}$ be proper, convex and lower semicontinuous and Pr_2 be the projection map from $E \times \mathbb{R}$ to $\mathbb{R}$. If B is a nonempty subset of $E \times \mathbb{R}$ such that $(x, \lambda) \in B \implies \lambda \leq \phi(x)$ and $\sup \text{Pr}_2(B) > \phi(E)$ then we write

$$[B \downarrow \phi] := \sup_{(x, \lambda) \in B, \phi(y) < \lambda} \frac{\lambda - \phi(y)}{\|x - y\|}$$

If B is a rock and ϕ is the bottom of an iceberg, $[B \downarrow \phi]$ tells you at what slope you have to swim down starting from an arbitrary point on the rock and still be sure that you will not hit the iceberg. There are significant situations in which $0 < [B \downarrow \phi] < \infty$. We can deduce from this a number of results on the existence of subtangents to ϕ satisfying various conditions, with sharp lower bounds on the slopes of the subtangents. One of them improves a recent result of Beer on separating subtangents, and another improves a recent result of Beer on separating subtangents, and another improves a recent result of Attouch and Beer on the approximation of conjugate functions. Our results also lead to generalizations of Rockafellar's fundamental result that $\partial\phi$ is *maximal monotone*, that is to say,

if: $q \in E$, $a \in E'$ and $\forall (z, b) \in \partial\phi$, $\langle q - z, a - b \rangle \geq 0$ then $(q, a) \in \partial\phi$.

For instance, we can prove that : if Q is a nonempty weakly compact convex subset of E , $a \in E'$ and,

$$\forall (z, b) \in \partial\phi, \exists q \in Q \text{ such that } \langle q - z, a - b \rangle \geq 0$$

then

$$\exists q \in Q \text{ such that } (q, a) \in \partial\phi.$$

Dually, we can prove that : if A is a nonempty weak* compact convex subset of E' , $q \in E$ and,

$$\forall (z, b) \in \partial\phi, \exists q \in Q \text{ such that } \langle q - z, a - b \rangle \geq 0$$

then

$$\exists a \in Q \text{ such that } (q, a) \in \partial\phi.$$

Since some of these result are quite technical, in the talk we will simply show how the techniques can be used to give a very short proof that $\partial\phi$ is maximal monotone,

Set convergences: a survey and a classification

Yves Sonntag

Université de Provence, case X, 13331-Marseille Cedex 3, France

and

Constantin Zălinescu

University of Iași, Department of Mathematics, 6600-Iași, România

Abstract

Let (E, d) be a metric space and $\mathcal{F}$ be the class of nonempty and closed subsets of E . By a *set convergence* we mean a procedure that associates an element $A \in \mathcal{F}$ to a net $(A_i)_{i \in I} \subset \mathcal{F}$. The number of these convergences continues to increase: there are more than 35 (and, of course, the same number of epigraphical convergences for nets of functions $f_i: E \rightarrow \overline{\mathbb{R}}$ and graphical convergences for nets of operators $\Phi_i: E \rightarrow E$). This proliferation is justified by the theory and by the applications, but the theory becomes more and more complicated and the perplexity of the potential user is comprehensible ... The difficulty comes from the disparate aspect of definitions that does not make easy a global view of convergences: having in mind all the results concerning the relative fineness between them causes already problems. A solution consists to modify the original definitions of convergences using only few mathematical notions that can be easily compared (inequalities, inclusions for example).

We proposed the following method in [SZ]: each convergence corresponds to a semi-metric space defined by a couple formed by a family $\mathcal{X} \subset \mathcal{F}$ (the *class* of the convergence) and a function $f: \mathcal{F} \times \mathcal{X} \rightarrow \mathbb{R}$ or $g: E \times \mathcal{F} \rightarrow \mathbb{R}$ (the *type* of the convergence). The semi-metric space is defined by the semi-metrics

$$(f_X)_{X \in \mathfrak{X}}, \text{ where } f_X(A, B) = |f(A, X) - f(B, X)| \text{ for } A, B \in \mathfrak{F},$$

or

$$(g_X)_{X \in \mathfrak{X}}, \text{ where } g_X(A, B) = \sup_{x \in X} |g(x, A) - g(x, B)| \text{ for } A, B \in \mathfrak{F}.$$

It is also possible to take several families of such semi-metrics. The natural choices $f(A, X) = d(A, X)$ (the type p) and $g(x, A) = d(x, A)$ (the type q) give the possibility to *redefine* the most part of known convergences. One can use results of G. Beer (for type p), of B. Cornet (for type q) and of the authors. The convergences can be written simply:

$$\mathfrak{X}(f)\text{-}\lim(A_i) = A \text{ iff } \lim f(A_i, X) = f(A, X) \quad \forall X \in \mathfrak{X},$$

$$\mathfrak{X}(g)\text{-}\lim(A_i) = A \text{ iff } \lim[\sup_{x \in X} |g(x, A_i) - g(x, A)|] = 0 \quad \forall X \in \mathfrak{X}.$$

The classification of convergences becomes now very simple and is based on the inequality $p_X \leq q_X$ and on evident inclusions of classes (compact sets $\subset$ closed bounded sets $\subset \mathfrak{F}$, etc ...).

Remark that there are many convergences of type p, but very few, till now, of type q: Hausdorff, Attouch-Wets and Wijsman.

Of course, one can define other types taking for f and g other canonical functions associated to (E, d) : for example the Chebyshev radius, the Hausdorff excess function, etc...

G. Beer and R. Lucchetti developed the study of types corresponding to the Hausdorff excess function, giving so a good framework for fine and very fine convergences.

The papers of [SZ] and [BL] (and some others), which represent the base for our talk, give a clear and coherent vision of the majority of set convergences. We hope that this presentation will facilitate the work of ès-convergences students and the task of users (optimization, multivalued functions, probability, etc...) in choosing the adequate convergence(s) for their own study.

References

[BL] G. Beer and R. Lucchetti, *Weak topologies on the closed subsets of a metrizable space*, to appear in Trans. Amer. Math. Soc.

[SZ] Y. Sonntag and C. Zălinescu, *Set convergences. An attempt of classification*, to appear in Trans. Amer. Math. Soc.

A new Topology for the Solid Sets of a Topological Space

Anna Torre

Dip. Matematica Università di Pavia

Pavia ITALIA

Abstract

In the paper [LTW] a new hypertopology τ_r is defined with the aim of identifying classes of sets where narrow convergence of a sequence of probability measures implies uniform convergence (see also [BT, LSW] and [SW] for applications).

The result can be expressed in this way: in any τ_r -compact class $\mathcal{A}$ of sets which is contained in the family of the continuity sets of a probability P , narrow (i.e. pointwise) convergence of P^n to P implies uniform convergence of P^n to P on the class $\mathcal{A}$. Also, it turns out that the condition of A being a P -continuity set (i.e $P(\partial A) = 0$) is necessary as well as sufficient for the function $P : (CL(X), \tau_r) \longrightarrow [0, 1]$ being continuous at A . The paper investigates the topological properties of τ_r on the subset of the sets that are the closure of their interior. In particular, it is shown that the hyperspace is metrizable if and only if the space X is separable and locally compact.

References

- [Bi] P. Billingsley, Convergence of probability measures, J. Wiley & Sons, N.Y. 1968.

[LTW] R. Lucchetti, G. Salinetti and R.J-B. Wets, A new topology for the solid sets of a topological space, submitted.

[SW] G. Salinetti and R. J-B. Wets, A Glivenko-Cantelli type theorem: an application of the convergence theory of stochastic suprema, to appear on Annals of Operation Research.

Strong convergence implied by weak in L^1

Michel Valadier

(in collaboration with Allal Amrani and Charles Castaing)
Université Montpellier II, 34095 Montpellier Cedex 5, France

Abstract

The Visintin's theorem and its extension by E.J. Balder have been extended to infinite dimensional Banach spaces by several authors. The best conclusion is obtained when some strong compactness is assumed. But the weak topology is also considered. Some of the results can be proved without Young measures. Then truncations are essential parts of proofs and are related to some extensions of the tightness condition.

References

- Amrani, A.
[A] "Lemme de Fatou pour l'intégrale de Pettis. Extrémalité et convergence dans l'espace des fonctions intégrables," Thèse de Doctorat. Montpellier, 1991.
- Amrani, A., C. Castaing & M. Valadier
[ACV1] Convergence forte dans L^1 impliquée par la convergence faible. Méthodes de troncature, *C. R. Acad. Sci. Paris Sér. I* **314** (1992), 37-40.
[ACV2] Méthodes de troncature appliquées à des problèmes de convergence faible ou forte dans L^1 , *Arch. Rational Mech. Anal.* **117** (1992), 167-191.
- Balder, E.J.
[Ba1] On weak convergence implying strong convergence in L_1 -spaces, *Bull. Austral. Math. Soc.* **33** (1986), 363-368.
[Ba2] On equivalence of strong and weak convergence in L_1 -spaces under extreme point conditions, *Israel J. Math.* **75** (1991), 21-47.
[Ba3] "From weak to strong L_1 -convergence by an oscillation restriction criterion of BMO type," University of Utrecht, Department of Mathematics, Preprint, may 1991 (10 pages).
[Ba4] On weak convergence implying strong convergence under extremal conditions, *J. Math. Anal. Appl.* **163** (1992), 147-156.
- Benabdellah, H.
[Be] "Contribution aux problèmes de convergence fort-faible, à la géométrie des espaces de Banach et aux inclusions différentielles," Thèse de Doctorat. Montpellier, 1991.
- Castaing, C.
[C1] Convergence faible et sections extrémales, *Sém. Anal. Convexe Montpellier* (1988), exposé 2 (18 pages).

- [C2] Sur la décomposition de Slaby. Applications aux problèmes de convergence en probabilités, économie mathématique, théorie du contrôle, minimisation, *Sém. Anal. Convexe Montpellier* (1989), exposé 3 (35 pages).
- Girardi, M.
- [G1] Compactness in L_1 , Dunford-Pettis operators, geometry of Banach spaces, *Proc. Amer. Math. Soc.* **111** (1991), 767-777.
- [G2] Weak vs. norm compactness in L_1 : the Bocce criterion, *Studia Math.* **98** (1991), 95-97.
- Rzeżuchowski, T.
- [R1] Strong convergence of selections implied by weak, *Bull. Austral. Math. Soc.* **39** (1989), 201-214.
- [R2] "Impact of dentability on weak convergence in L^1 ," Technical University of Warsaw, Institute of Mathematics, 1990 (12 pages).
- Valadier, M.
- [Va1] Différents cas où, grâce à une propriété d'extrémalité, une suite de fonctions intégrables faiblement convergente converge fortement, *Sém. Anal. Convexe Montpellier* (1989), exposé 5 (20 pages).
- [Va2] Young measures, in "Methods of Nonconvex Analysis" (ed. A. Cellina), Lecture Notes in Math. 1446, Springer Verlag, Berlin, 1990, pp. 152-188.
- [Va3] Oscillations et compacité forte dans L^1 , *Sém. Anal. Convexe Montpellier* (1991), exposé 7 (10 pages).
- Visintin, A.
- [Vi] Strong convergence results related to strict convexity, *Comm. Partial Differential Equations* **9** (1984), 439-466.

Remarks about the Infimum of Hausdorff Metric Topologies

Paolo Vitolo

Universita' della Basilicata, 85100, Potenza, Italy

Abstract

We consider a metrisable topological space X and denote by D the set of all compatible metrics on X . For every d in D , the upper and lower Hausdorff topologies - $H(d,+)$ and $H(d,-)$, respectively - are defined on the collection $C(X)$ of all closed subsets of X . (See [1]). Let P be the infimum of the topologies of the form $H(d,+)$, where d runs over D , and M the infimum of the topologies of the form $H(d,-)$. We show that M coincides with the lower Vietoris topology, if and only if X separable (compare with [3], cor. 6). We also show that (for locally compact X) P coincides with the co-compact topology (see [2]) if and only if X is compact.

References

- [1] E. C. Klein, A. Thompson, Theory of Correspondences, Wiley 1984.
- [2] S. Francaviglia, A. Lechicki, S. Levi, Quasi-Uniformization of Hyperspaces and Covergence of Nets of Semicontionuous Multifunctions, J. Math. Anal. Appl. 112 (1985), 347-370.
- [3] S. Levi, R. Lucchetti, J. Pelant, On the infimum of the Hausdorff and Vietoris Topologies, to appear on Proc. Amer. Math. Soc.
- [4] C. Costantini, P. Vitolo, Some Remarks about the infimum of Hausdorff Metric Topologies, preprint 1992.

**THE USE OF MONOTONE NORMS
IN CONVEX ANALYSIS**

Michel VOLLE
Université d'Avignon

Abstract. We introduce some new binary operations for convex sets and convex functions. These operations provide a general framework for dealing with the calculus of epigraphs, polar sets, and Fenchel conjugates.

1991 Mathematics Subject Classification : 90 C 25

Key words : monotone norms, convex duality, epigraphical analysis.

References

- [1] R.T.Rockafellar, Convex Analysis, Princeton University Press, 1969.
- [2] H.Attouch-R.Wets, Epigraphical analysis, analyse non linéaire, Gauthier-Villars, 1989.
- [3] A.Seeger, Direct and inverse addition in Convex Analysis and Applications, J.Math. Anal. Appl., 1990.

Convergence of Integral Functionals

Roger J-B Wets *

Mathematics, University of California, Davis, CA 95616, U.S.A.

Abstract

Let $(T, \mathcal{A}, \mu)$ be a measure space with σ -additive, nonnegative measure μ , and $\mathcal{X}$ a Banach subspace of $\mathcal{M}(T; \mathbb{R}^n)$, the space of measurable functions from T to $\mathbb{R}^n$. One refers to a bivariate function $f : T \times \mathcal{X} \rightarrow \overline{\mathbb{R}}$ as an *integrand* if for all $x \in \mathcal{M}(T; \mathbb{R}^n)$, the function $t \mapsto f(t, x(t))$ is $\mathcal{A}$ -measurable. We are interested in *integral functionals* of the type:

$$F : \mathcal{X} \rightarrow \overline{\mathbb{R}}, \quad F(x) := \int_T f(t, x(t)) \mu(dt).$$

The need for approximation techniques to deal with integral functionals that arise in the calculus of variations, stochastic optimization and other variational problems (pde), provided at least initially, the major motivation for the development of the theory of *epi-convergence* and its extensions (epi/hypo-convergence, Γ -convergence, etc.). This lecture will try to take stock of the progress made during the last decade or so in dealing with the convergence of integral functionals.

More specifically, let $\{f^\nu : T \times \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \nu \in \mathbb{N}\}$ be a sequence of integrands, $\{\mu^\nu : \mathcal{A} \rightarrow \mathbb{R}_+, \nu \in \mathbb{N}\}$ a sequence of measures, and F^ν the associated sequence of integral functionals

$$F^\nu(x) := \int_T f^\nu(t, x(t)) \mu^\nu(dt), \quad \nu \in \mathbb{N}.$$

Assuming that the f^ν and μ^ν approximate f and μ in some sense, what can be said about the convergence of the integral functionals F^ν to F ? In particular, when can we guarantee the epi-convergence of the F^ν to F ?

*Supported in part by a grant of the National Science Foundation

References

- [1] Zvi Artstein and Roger J-B Wets, "Stability of stochastic programs with possibly discontinuous objective functions," Weizmann Institute, Rehovot, 1992
- [2] Zvi Artstein and Roger J-B Wets, "Approximating the integral of a multifunction," *J. Multivariate Analysis*, 24 (1988), 285-308.
- [3] Hedy Attouch, "Famille d'opérateurs maximaux monotones et mesurabilité," *Annali di Matematica Pura ed Applicata*, 120 (1979), 35-111.
- [4] Hedy Attouch and Roger J-B Wets, "Epigraphical processes: laws of large numbers for random lsc functions," manuscript University of California, Davis, 1991.
- [5] Giuseppe Buttazzo and Gianni Dal Maso, " Γ - Limits of integral functionals," Technical Report, Scuola Normale Superiore Pisa, 1979.
- [6] Gianni Dal Maso and Umberto Mosco, "Wiener's criterion and Γ -convergence," *Applied Mathematics and Optimization*, 15 (1987), 15-63.
- [7] Ennio De Giorgi, Gianni Dal Maso and Placido Longo, " Γ - Limiti de ostacoli," Technical Report, Scuola Normale Superiore Pisa, 1980.
- [8] Ennio De Giorgi, "Generalized limits in calculus of variations," in *Topics in Functional Analysis 1980-1981*, F. Strocchi, E. Zarantonello, E. De Giorgi, G. Dal Maso and L. Modica (eds.), Tecnoprint, Bologna, 1981.
- [9] Jitka Dupačová and Roger J-B Wets, "Asymptotic behavior of statistical estimators and of optimal solutions for stochastic optimization problems" *The Annals of Statistics*, 16 (1988), 1517-1549.
- [10] Christian Hess, "Epi-convergence of sequences of normal integrands and strong consistency of the maximum likelihood estimator," *Cahiers mathématiques de la décision* no. 9121, Céremade, Paris, 1991.
- [11] Alan J. King and Roger J-B Wets, "Epi-consistency of convex stochastic programs", *Stochastics and Stochastics Reports*, 34 (1990), 83-92.
- [12] Roberto Lucchetti and Roger J-B Wets, "Convergence of minima of integral functionals, with applications to optimal control and stochastic optimization," *Statistics and Decisions*, 11 (1993), to appear.

ON THE CONVERGENCE OF THE GENERALIZED GRADIENTS

T. Zolezzi.

Dipartimento di Matematica , Universita' di Genova, via L.B.Alberti
4. 16132 Genova (Italy). e-mail : Zolezzi at IGECUNIV. BITNET.

EXTENDED ABSTRACT for the conference " Convergences en
Analyse Multivoque et Unilatérale " ; Marseille-Luminy , June
1992.

We consider a real Banach space E , and sequences of locally
Lipschitz functions

$$f, f_n : E \rightarrow (-\infty, +\infty), n = 1, 2, 3, \dots$$

We address the problem, under which conditions on the
convergence of f_n towards f , it is possible to obtain some
form of convergence of the sequence of their Clarke's generalized
gradients

$$\partial f_n \rightarrow \partial f .$$

As an application (and a motivation), we mention the stability
of the Clarke multiplier theorem [1] , which is connected to the
stability under perturbations of sensitivity estimates for
constrained optimization problems under (in)equality constraints.

If each term f_n of the sequence is a proper, convex, lower
semicontinuous function, the problem above has been thoroughly
investigated. Necessary and sufficient conditions, linking epi -
convergence of f_n toward f with the Kuratowski convergence
of the graphs of the generalized gradients are known : see [2] ,
[3], [4] , [5] . The bounded Hausdorff convergence of f_n is also
related to the convergence of the subgradients, see [6].

If E is a finite - dimensional space , sufficient conditions are known yielding

$$(1) \quad \limsup \partial f_n (y) \subset \partial f(x), \text{ as } n \rightarrow +\infty \text{ and } y \rightarrow x,$$

for every $x \in E$, again under epi - convergence of f_n to f , see [7].

If E is a Hilbert space , and each f_n is (p,q) -convex, then results linking epi-convergence of the functions with various Kuratowski convergences of their lower semigradients are proved in [8].

If E is infinite - dimensional and fulfills some regularity conditions, it is possible to extend (1) under epi-convergence of the sequence f_n . No convexity is required, and an equi - lower - semidifferentiability condition is imposed (as in [7]) on the sequence f_n . The proof makes use of convergence results for lower semigradients of lower semicontinuous functions.

This allows us to pass to the limit in the Clarke's multipliers rule in the infinite - dimensional setting, when data perturbations are present. In this way we obtain sufficient conditions for the stable behavior of the multipliers, under epi - convergence conditions.

Further applications to the behavior of the generalized gradients of integral functionals are also possible using the above result.

REFERENCES

- [1] F.H.CLARKE. A new approach to Lagrange multipliers. Math. Oper. Res. 1 , 1976.
- [2] L.BOCCARDO - P.MARCELLINI . Sulla convergenza delle soluzioni di disequazioni variazionali . Ann. Mat. Pura Appl. 110, 1976.

- [3] M.MATZEU . Su un tipo di continuita' dell'operatore subdifferenziale. Boll. Un. Mat. Ital. 14 B, 1977.
- [4] H.ATTOUCH . Convergence de fonctions convexes, des sous-différentielles et de semi-groupes associés. C.R.Acad.Sci.Paris 284 , 1977.
- [5] H.ATTOUCH . Variational convergences for functions and operators. Pitman, 1984 .
- [6] H.ATTOUCH- J.L.NDOUTOUME - M.THERA . Epigraphical convergence of functions and convergence of their derivatives in Banach spaces. Preprint , 1991.
- [7] T.ZOLEZZI . Continuity of generalized gradients and multipliers under perturbations . Math. Oper. Res. 10 , 1985.
- [8] M.DEGIOVANNI - A.MARINO - M.TOSQUES . Evolution equations associated with (p,q) - convex and (p,q) - monotone operators. Ric. Mat. 33 , 1984.

Adresses des participants

Samir Adly
Département de Mathématiques
Faculté des Sciences
123, Avenue A. Thomas
87060 Limoges Cedex

Erick Balder
University of Utrecht
P.O. Box 80. 010
NL 3508 TA Utrecht
Pays-Bas

Zvi Artstein
Department of Theoretical Mathematics
The Weizman Institute
Rehovot
76100 Israel

Gerald Beer
Department of Mathematics
California State University
5151 State University Drive
Los Angeles Ca 9003 U.S.A

Hedy Attouch
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Joël Benoist
Département de Mathématiques
Faculté des Sciences
123, avenue A. Thomas
87060 Limoges Cedex

Allred Auslender
Département de Mathématiques Appliquées
Université Blaise Pascal
B.P. 45
63170 Aubière Cedex

Guy Bouchitté
Mathématiques
Université de Toulon et du Var
Avenue de l'Université
B.P. 13 83957 La Garde Cedex

Dominique Azé
Département de Mathématiques
Université de Perpignan
Avenue de Villeneuve
66025 Perpignan Cedex

Mireille Bougeard
Observatoire de Paris
61, avenue de l'Observatoire
75014 Paris

M. A. Bahraoui
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

P. Brandi
Università di Perugia
Dipartimento di Matematica
via Vanvitelli 1
20130 Perugia Italie

Jean-Bernard Barillon
College Sisley
93450 L'île Saint-Denis

Arigo Cellina
Mathematics
Sissa
Strada Castiera 11
34014 Trieste Italie

Zaki Chbani
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Jean-Pierre Crouzeix
Département de Mathématiques Appl.
Université Blaise Pascal
B.P. 45
63170 Aubiere Cedex

Roberto Cominetti
Matematicas
Universidad de Chile
Castilla 170 Coreo 3
Santiago Chili

Alain Damlamian
Centre de Mathématiques Appliquées
Ecole Polytechnique
91128 Palaiseau Cedex

Pietro, Poggi Corradini
Università di Milano
Dipartimento di Matematica
via C. Saldini 50
20130 Milano Italie

Ennio De Giorgi
Scuola Normale Superiore
Piazza dei Cavalieri 7
56126 Pisa Italie

Raphael Correa
Matematicas
Universidad de Chile
Castilla 170 Coreo 3
Santiago Chili

Marco De Giovanni
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

J.-N. Corvelec
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

Robert Deville
Département de Mathématiques
Faculté des Sciences
Route de Gray, La Bouloie
25030 Besançon Cedex

Camillo Costantini
Università di Milano
Dipartimento di Matematica
via C. Saldini 50
20130 Milano Italie

Seghir Driss
Laboratoire d'Analyse Numérique
Université Paul Sabatier
118, route de Narbonne
31062 Toulouse Cedex

Philippe Coutat
CERMSEM
Université Paris I
90, Rue de Tolbrac
75634 Paris Cedex 13

Rachid Eilaya
Laboratoire d'Analyse Numérique
Université Paul Sabatier
118, route de Narbonne
31062 Toulouse Cedex

Sjur D. Flam
Department of Economics
University of Bergen
5008 Bergen
Norvège

Alejandro Jofré
Matematicas
Universidad de Chile
Castilla 170 Correo 3
Santiago Chili

Gilles Godefroy
Université Paris 6
Boite 186
4 place Jussieu
75252 Paris Cedex 05

Bert Jongen
Department of Mathematics
RWTH
Aachen Germany

Jean Guillerme
Département de Mathématiques
Faculté des Sciences
123, avenue A. Thomas
87060 Limoges Cedex

Diethard Klatte
Inst. für Oper. Research
der Universität Zurich
Moussonstrasse 15
CH-8044 Zürich Suisse

Christian Hess
Ceremade
Université Paris Dauphine
Pl. Mal. de Lattre de Tassigny
75775 Paris Cedex 16

Jaafar Lahrache
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Jean-Baptiste Hiriart-Urruty
Laboratoire d'Analyse Numérique
Université Paul Sabatier
118, route de Narbonne
31062 Toulouse Cedex

Alec Lechicki
Harbstrasse 43
D-8510 Fürth
v

Vincent Jalby
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Bernard Lemaire
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Robert Janin
Département de Mathématiques
Université de Poitiers
Avenue du Recteur Pineau
86000 Poitiers

Sandro Levi
Università di Milano
Dipartimento di Matematica
via C. Saldini 50
20130 Milano Italie

Pierre Loridan
Département de Mathématiques
Faculté des Sciences Mirande
21004 Dijon Cedex

Gérard Michaille
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Dinh The Luc
Département de Mathématiques
Faculté des Sciences
123, Avenue A. Thomas
87060 Limoges Cedex

Christian Michelot
CERMSEM
Université Paris I
90, Rue de Tolbrac
75634 Paris Cedex 13

Roberto Lucchetti
Università di Milano
Dipartimento di Matematica
via C. Saldini 50
20130 Milano Italia

Jacqueline Morgan
Dipartimento di Matematica
Università degli Studi di Napoli
Via Mezzocannone 8
80134 Napoli Italia

Mongi Mabrouk
Département de Mathématiques
Faculté des Sciences
Route de Gray, La Bouloie
25030 Besançon Cedex

Umberto Mosco
Dipartimento di Matematica
Università degli Studi di Roma
Piazza A. Jdo Morro, 2
00185 Roma Italia

Christian Malivert
Département de Mathématiques
Faculté des Sciences
123, Avenue A. Thomas
87060 Limoges Cedex

Abdelattif Moudafi
Département de Mathématiques Appli.
Université Blaise Pascal
B.P. 45
63170 Aubiere Cedex

Antonio Marino
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italia

Mohammed Moussaoui
Département des Mathématiques
Faculté des Sciences
33 rue Louis Pasteur
84000 Avignon

Juan-Enrique Martinez-Legaz
Dept. of Mat. Apl. y Anal.
Universitat de Barcelona
Gran Via, 585
08071 Barcelona

Josef Myjak
Dipartimento di Matematica
Facoltà di Scienza
Via Roma 33
67100 L'Aquila Italia

James-Louis Ndoutoume
Institut Africain d'Informatique
Libreville
Gabon

Jean-Paul. Penot
Département de Mathématiques
Faculté de Sciences de Pau
Av. Philippon
64000 Pau Cedex

V. Hien Nguyen
Département de Mathématiques
Universités Notre-Dame de la Paix
Rempart de la Vierge
B-5000 Namur Belgique

Robert Phelps
Department of Mathematics
University of Washington
Seattle WA 98195 U.S.A

Dominikus Noll
Mathematisches Institut B
Universität Stuttgart
Plattenwaldring 57
7000 Stuttgart R.F.A

Angela Pistoia
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

Werner Oetli
Lehrstuhl für Mathematik VII
Universität Mannheim
Senumargebaude A5
6800 Mannheim 1 R.F.A

René Poliquin
Department of Mathematics
University of Alberta
Edmonton Alberta
T6G-2G1 Canada

Pier-Luigi Papini
Dipartimento di Matematica
Piazza di Porta S. Donato 5
40127 Bologna Italie

Liqun Qi
Department of Applied Mathematics
University of New South Wales
Kensington NSW
Australie

Angela Pasquale
Department of Mathematics
University of Washington
Seattle WA 98195 U.S.A

A. Ramouni
Département de Mathématiques
Université de Perpignan
Avenue de Villeneuve
66025 Perpignan Cedex

Donato Passaro
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

Julian Revalski
Institute of Mathematics
Bulgarian Academy of Sciences
G. Donchev, 8
1113 Sofia Bulgarie

Hassan Riahi
Département de Mathématiques
Université Cadi AYYAD
Faculté des Sciences
BP S 15 Marrakech Maroc

Stephen Simons
Mathematics
UC Santa Barbara
Santa Barbara Ca 93106 U.S.A

Biagio Ricceri
Dipartimento di Matematica
Viale A. Doria, 6-1
95125 Catania Italie

Yves Sonntag
Département de Mathématiques
Université de Provence
3, place Victor Hugo
13331 Marseille Cedex

Steve Robinson
Department of Industrial Engineering
513 University Avenue
Madison Wisconsin 53706 U.S.A

Michel Théra
Département de Mathématiques
Faculté des Sciences
123, Avenue J. Thomas
87060 Limoges Cedex

Terry Rockafellar
Department of Mathematics
University of Washington
Seattle WA 98195 U.S.A

Lionel Thibault
Département de Mathématiques
Université de Pau
Avenue de l'Université
64000 Pau Cedex

Claudio Saccon
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

Anna Torre
Università di Pavia
Dipartimento di Matematica
Pavia Italie

Anna Salvadori
Università di Perugia
Dipartimento di Matematica
via Vanvitelli 1
20130 Perugia Italie

Claudio Tosques
Dipartimento di Matematica
Università di Pisa
56100 Pisa Italie

Alberto Seeger
Département des Mathématiques
Faculté des Sciences
33 rue Louis Pasteur
84000 Avignon

Patricia Tossings
Institut de Mathématiques
Université de Liège
15, avenue des Tilleuls
B-4000 Liège Belgique

Annick Truffert
Département de Mathématiques
Université de Perpignan
Avenue de Villeneuve
66025 Perpignan Cedex

Michel Valadier
Département de Mathématiques
Université du Languedoc
Place E. Bataillon
34060 Montpellier Cedex

Paolo Vitolo
Università della Basilicata
Dipartimento di Matematica
85100 Potenza Italie

Michel Volle
Département des Mathématiques
Faculté des Sciences
33 rue Louis Pasteur
84000 Avignon

Roger Wets
2121 Pinhurst
El Cerrito Ca 94530
U.S.A

Constantin Zălinescu
Département de Mathématiques
Université de Provence
3, place Victor Hugo
13331 Marseille Cedex

Tullio Zolezzi
Istituto di Matematica
Università di Genova
Via Leon Battista Alberti 4
16132 Genova Italie