



dated 12-1-1988. It is for "MORRIS" including the time for "Training instructions, searching existing data sources, reviewing the Collection of information, long comments regarding this Bureau estimate or any other aspect of this Bureau's: A. Information Management Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Management and Support, 0 userswork reduction project (1275-0188), Arlington, DC 22202).

3. REPORT TYPE AND DATES COVERED
Technical Report

**Grant
#N00014-91-J-1043
R&T Code 3135014
Harold E. Guard,
Program Officer**

Esther Spaltenstein, Pilar Palma, Kristina A. Kreutzer, Christopher A. Willoughby, William M. Davis, and Stephen L. Buchwald

**Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139**

94-35064



**Office of Naval Research
Department of the Navy
Chemistry Division, ONR 331
800 North Quincy Street
Arlington, VA 22217-5660**

Technical Report No. 7

Prepared for Publication in the Journal of the American Chemical Society

Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release and sale, distribution unlimited.

12b. DISTRIBUTION CODE

The synthesis and X-ray structure determination of the novel complex, $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, **4** is reported. The spectroscopic and structural data indicate that **4** is best described as being a hybrid between a titanium(IV)(silyl)hydride complex and a titanium(II) silane complex. Complex **4** reacts reversibly with N_2 to form the bridging dinitrogen complex **5** and, in the presence of norbornene, **4** catalytically dimerizes Ph_2SiH_2 in quantitative yield.

silane complex, titanium, norbornene

19. NUMBER OF PAGES	10
---------------------	----

VARIABLES

Unclassified

Unclassified

Unclassified

UL

94 11 14 010

PREPARATION AND X-RAY STRUCTURE OF $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$

Esther Spaltenstein, Pilar Palma, Kristina A. Kreutzer, Christopher A. Willoughby, William M. Davis and Stephen L. Buchwald*

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, MA 02139

Abstract: The synthesis and X-ray structure determination of the novel complex, $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, **4** is reported. The spectroscopic and structural data indicate that **4** is best described as being a hybrid between a titanium(IV)(silyl)hydride complex and a titanium(II) silane complex. Complex **4** reacts reversibly with N_2 to form the bridging dinitrogen complex **5** and, in the presence of norbornene, **4** catalytically dimerizes Ph_2SiH_2 in quantitative yield.

REVISED

Title		X	
Author			
By			
Date			
A.			
Dist			
A-1			

OFFICE OF NAVAL RESEARCH

GRANT #N00014-91-J-1043

**R&T Code 3135014
Harold E. Guard, Program Officer
Chemistry Division**

Technical Report No. 7

**Preparation and X-Ray Structure
of $\text{Cp}_2\text{Ti}(\text{PH}_2\text{SiH}_2)(\text{PMe}_3)$**

by

**Esther Spaltenstein, Pilar Palma, Kristina A. Kreutzer,
Christopher A. Willoughby, William M. Davis and Stephen L. Buchwald**

**Prepared for Publication
in the
Journal of the American Chemical Society**


**Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139**

October 18, 1994

**Reproduction in whole or in part is permitted for any purpose of the United States
Government**


Transition metal(silyl)hydride complexes have been proposed as intermediates in the catalytic hydrosilylation of olefins¹ and the polymerization of organosilanes.^{2,4,6} Several of these complexes have been prepared and studied in detail.³ Harrod and coworkers have isolated and characterized several dinuclear⁴ and mononuclear⁵ titanocene complexes from the reaction of dimethyltitanocene and phenylsilane (Figure 1). The compound $\text{Cp}_2\text{Ti}(\mu\text{-Si}(\text{H})_2\text{Ph})(\mu\text{-H})\text{TiCp}_2$ **2**, is of particular interest and can be viewed as being composed of a titanocene(hydrido)silyl complex and a titanocene. It has been suggested that the titanocene silyl hydride monomer is an active participant in silane polymerization.⁶

Figure 1 Representative Titanium(silyl)hydride Complexes




The bonding in transition metal(silyl)hydride complexes has classically been described by either of two limiting structures, **A** or **C** (Figure 2). However, there are many complexes that cannot be accurately described by either **A** or **C** and are thus best represented by an intermediate structure such as **B**.^{3a} We have previously reported the synthesis of some group 4 silyl hydride complexes, $\text{Cp}_2\text{M}(\text{H})(\text{SiR}_3)(\text{PMe}_3)$, $\text{M}=\text{Zr}, \text{Hf}$.⁷ The physical data for these complexes indicate that they are true metal(silyl)hydride complexes (structure **A**). In this paper we report the preparation and characterization of a novel titanocene complex, $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, which, in our view, has a hybrid structure like **B**.

Figure 2 Bonding in Transition Metal(silyl)hydride Complexes



Treatment of a benzene solution of $\text{Cp}_2\text{Ti}(\text{PMe}_3)_2$ ⁸ with diphenylsilane followed by addition of cold hexanes caused precipitation of an air and moisture sensitive yellow solid identified as $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, **4** (eq. 1). Compound **4** was obtained in 63 % isolated yield after washing with hexane. Complex **4** is unstable in solution and decomposes to the dinuclear species $\{(\mu-(\eta^1:\eta^5\text{C}_5\text{H}_4))(\text{Cp})\text{Ti}(\text{PMe}_3)\}_2$ ⁹ and Ph_2SiH_2 . However, in the presence of excess Ph_2SiH_2 , benzene solutions of **4** are stable for several days.




The ^1H NMR spectrum (300 MHz, C_6D_6) of **4** at room temperature exhibits a broad resonance at 6.36 ppm assigned to the silicon hydride and a broad doublet at -3.93 ppm assigned to the titanium hydride. At -55 °C in THF-d_8 (500 MHz) these resonances are completely resolved, the Si-H resonance is located at 6.03 ppm as a doublet of doublets ($J_{\text{HH}} = 11$ Hz, $J_{\text{HP}} = 4$ Hz) while the Ti-H signal is observed at -4.14 ppm, also as a doublet of doublets ($J_{\text{HH}} = 11$ Hz, $J_{\text{HP}} = 68$ Hz). The Ti-H resonance is 3.87 ppm up field of that for the corresponding zirconium complex,⁷ reflecting the more electron rich character of the metal in **4**. In comparison, the Ti-H resonance for the recently reported titanium(IV) hydride, $(^t\text{Bu}_3\text{SiO})_2(^t\text{Bu}_3\text{SiNH})\text{TiH}$, is at 8.62 ppm.¹⁰

The ^{29}Si NMR spectrum of complex **4** (99 MHz, THF-d_8 , -50 °C) shows a doublet of doublets indicating coupling of the silicon to two hydrogens¹¹ ($J_{\text{SiH}_1} = 28$ Hz, $J_{\text{SiH}_2} = 161$ Hz). The large silicon hydrogen coupling constant of 161 Hz is consistent with those found in normal silanes (150 - 200 Hz)^{3a} while the smaller value of 28 Hz is slightly larger than values typically observed for silyl-hydride complexes in which there is no interaction between Si and H ($J_{\text{SiH}} < 20$ Hz).^{3a} This suggests the possibility of a 3 center interaction between the titanium hydride, the titanium and the silicon atom. Consistent with this hypothesis is the large H-H coupling constant of 11 Hz. For the corresponding zirconium analog, $\text{Cp}_2\text{Zr}(\text{H})(\text{SiPh}_2\text{H})(\text{PMe}_3)$,^{1d,7} in which a 3 center interaction is not present, the H-H coupling

constant is 3 Hz. The IR spectrum displays a weak Ti-H stretch at 1508 cm^{-1} ($\nu\text{ Ti-D} = 1042\text{ cm}^{-1}$). This value is slightly lower than that reported for the only other known titanocene (IV) hydride, $\text{Cp}_2^*\text{TiH}_2$ ($\nu\text{ Ti-H} = 1560\text{ cm}^{-1}$)¹² and slightly higher than the value reported for the titanocene (III) hydride $[\text{Cp}^*\{\text{C}_5\text{Me}_4\text{CH}_2(\text{C}_5\text{H}_3\text{N})\}\text{TiH}]$ ($\nu\text{ Ti-H} = 1475\text{ cm}^{-1}$).¹³ The IR stretch for $(^t\text{Bu}_3\text{SiO})_2(^t\text{Bu}_3\text{SiNH})\text{TiH}$ is observed at 1645 cm^{-1} .¹⁰

The silane moiety in complex **4** is only weakly bound and is readily displaced by dinitrogen according to equation 2, leading to formation of the previously reported dinitrogen complex **5**.¹⁴ Under vacuum, nitrogen can be removed to regenerate **4**.



In order to achieve a more complete bonding picture, complex **4** was examined by X-ray crystallography. Importantly, the hydrogen atoms were located and anisotropically refined. The ORTEP drawing of complex **4** is shown in Figure 3, selected bond distances and angles are listed in Table 1.

Figure 3 ORTEP drawing of $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, **4**

Table 1 Some Selected Bond Distances and Angles for $\text{Cp}_2\text{Ti}(\text{Ph}_2\text{SiH}_2)(\text{PMe}_3)$, **4**.

Bond Distances, Å		Bond Angles, deg	
Ti-H ₁	1.81 (5)	Si-Ti-H ₁	44 (2)
Ti-Si	2.597 (2)	Si-Ti-H ₂	112 (2)
Ti-P	2.550 (2)	Si-Ti-P	111.78 (7)
Si-H ₁	1.69 (5)	H ₁ -Si-H ₂	156 (2)
Si-H ₂	1.56 (5)	C ₂₀ -Si-H ₁	92 (2)
		C ₂₀ -Si-H ₂	100 (2)
		C ₉ -Si-H ₁	94 (2)
		C ₉ -Si-H ₂	103 (2)
		C ₉ -Si-C ₂₀	105.4 (3)
		Ti-Si-C ₉	118.2 (2)

The Ti-H bond of 1.81 Å is 0.11 Å longer than that of a terminal Ti(III)-H bond in $[\text{Cp}^*(\text{C}_5\text{Me}_4\text{CH}_2(\text{C}_5\text{H}_3\text{N}))\text{TiH}]$ (Ti-H = 1.70 Å)¹³ but well within the normal range of 1.66 to 1.97 Å found for bridging titanium hydrides.^{4,15} The Ti-P distance (2.550 Å) is in good agreement with that of the dinitrogen complex **5** (2.539 Å)¹⁴ and 0.1 shorter than that of complex **3** (Figure 1, Ti-P = 2.651 Å)^{5a} (the longer bond in this complex may be due to greater steric crowding of the more bulky PEt_3 ligand). The Ti-Si distance of 2.597 Å is similar to that of other titanocene silyl derivatives which are typically between 2.60 and 2.63 Å.^{4,5}

A most interesting feature of the structure is the geometry of the silicon atom. This geometry is best described as a distorted trigonal bipyramid with the two hydrogen atoms occupying the apical positions and Ti, C₂₀ and C₉ defining the equatorial plane. The silicon atom is displaced from the equatorial plane toward H₂ with C₂₀-Si-H₂ and C₉-Si-H₂ being 100° and 103° respectively. The Si-H₂ distance of 1.56 Å is 0.09 Å longer than the terminal Si-H distance in the related complex **2** (Figure 1, Si-H = 1.47 Å),⁴ likely due the trans influence of the H₁ ligand. The Si-H₁ distance of 1.69 Å is substantially (15 %) longer than the typical Si-H distance (1.47 Å) but still short enough for an interaction with the silicon atom.^{3a}

The Si-Ti-H₁ angle of 44° is 10° smaller than that of the related zirconium complex $\text{Cp}_2\text{Zr}(\text{H})(\text{SiPh}_3)(\text{PMe}_3)$ ⁷ while the Si-Ti-P angle of 111.78° is roughly the same as that found in $\text{Cp}_2\text{Zr}(\text{H})(\text{SiPh}_3)(\text{PMe}_3)$ (Si-Zr-P = 112.68°). The similarity of the Si-M-P angle in these two complexes indicates that the smaller Si-Ti-H₁ angle is not due to steric crowding resulting from the smaller size of the titanium atom. The smaller Si-Ti-H₁ angle thus reflects a bonding interaction between silicon and H₁.

Since complexes similar to **4** have been implicated in the catalytic polymerization of organosilanes,⁶ we investigated the use of complex **4** as a catalyst for this reaction. When a solution of **4** is treated with excess Ph_2SiH_2 , no reaction is observed. However, in the presence of norbornene, complex **4** catalytically and quantitatively converts Ph_2SiH_2 to the dimer, $[\text{Ph}_2\text{Si}(\text{H})]_2$. This reaction is complete within 1 hour at 45 °C and no induction period is observed. Furthermore, no olefin hydrosilylation^{1d, 6} is observed.

In summary, we have prepared and characterized the first adduct of $\text{Cp}_2\text{Ti}(\text{PMe}_3)$ and diphenyl silane. Both the spectroscopic and structural data suggest that there is a 3 center interaction between the silicon, the titanium and the hydride ligand. The bonding in complex **4** is best described by the intermediate structure **B** (Figure 2). While not totally unambiguous, the low Si-H₁ coupling constant as well as the structural data (short Ti-Si distance and long Si-H₁ distance) indicate that **4** has more character of a metal(silyl)hydride complex (structure **A**) than of a metal silane complex (structure **C**).

Acknowledgment: This work was supported, in part, by the Office of Naval Research, the National Science Foundation, W. R. Grace and the donors of the Petroleum Research Fund, administered by the American Chemical Society to whom we are grateful. ES thanks the Ford Foundation for a fellowship. PP acknowledges the Spanish Ministry of Education and Science for a fellowship. CAW acknowledges support as an ACS Organic Division Predoctoral Fellow (sponsored by the Monsanto Company). SLB acknowledges additional support as an Alfred P. Sloan Foundation Fellow and a Camille & Henry Dreyfus Foundation Teacher Scholar.

Supplementary Material Available: Detailed experimental procedures for the preparation and spectroscopic characterization of Complex **4**, along with tables of bond distances, bond angles and final positional and thermal parameters for **4** (32 pages).

References:

1. a) Chalk, A. J.; Harrod, J. F.; *J. Am. Chem. Soc.* **1965**, *187*, 16. b) Schroeder, M. A.; Wrighton, M. S. *J. Am. Chem. Soc.* **1981**, *103*, 975. c) Brookhart, M.; Grant, B. E. *J. Am. Chem. Soc.* **1993**, *115*, 2151. d) Takahashi, T.; Hasegawa, M.; Suzuki, N. Saburi, M.; Rousset, C. J.; Fanwick, P. E.; Negishi, E. *J. Am. Chem. Soc.* **1991**, *113*, 8564.
2. Harrod, J. F. in *Inorganic and Organometallic Polymers*; ACS Symposium Series 360; Zeldin, M.; Wynne, K. J.; Allcock, H. R.; Eds; American Chemical Society: Washington DC, 1988, Chapter 7. b) Harrod, J. F.; Zeigler, T.; Tschinke, V. *Organometallics* **1990**, *9*, 897. c) Woo, H. G.; Tilley, T. D. *J. Am. Chem. Soc.* **1989**, *111*, 8043. d) Woo, H. G.; Tilley, T. D. *J. Am. Chem. Soc.* **1989**, *111*, 3757. e) Chang, L. S.; Corey, J. Y. *Organometallics*, **1989**, *8*, 1885. f) Campbell, W. H.; Hilty, T. K.; Yurga, L. *Organometallics*, **1989**, *8*, 2615. g) Aitken, C. T.; Harrod, J. F.; Samuel, E. *J. Organomet Chem.* **1985**, *279*, C11. h) Aitken, C. T.; Harrod, J. F.; Samuel, E. *Can. J. Chem.* **1986**, *64*, 1677.
3. a) Schubert, U. in "Advances in Organometallic Chemistry," vol. 30, p 151, 1990. b) Tilley, T. D. in "Chemistry of Organic Silicon Compounds," S. Patai and Z. Rappoport, Ed., p. 1415, 1989.
4. Aitken, C. T.; Harrod, J. F.; Samuel, E. *J. Am. Chem. Soc.* **1986**, *108*, 4059.
5. a) Samuel, E.; Mu, Y.; Harrod, J. F.; Dromzee, Y.; Jeannin, Y. *J. Am. Chem. Soc.* **1990**, *112*, 3435. b) Woo, H. G.; Harrod, J. F.; Henique, J.; Samuel, E. *Organometallics*, **1993**, *12*, 2883.
6. Harrod, J. F.; Yun, S. S. *Organometallics*, **1987**, *6*, 1381.
7. Kreutzer, K. A.; Fisher, R. A.; Davis, W. M.; Spaltenstein, E.; Buchwald, S. L. *Organometallics*, **1991**, *10*, 4031.
8. Kool, L. B.; Rausch, M. D.; Alt, H. G.; Thewalt, U. *J. Organomet. Chem.* **1987**, *37*.
9. Kool, L. B.; Rausch, M. D.; Alt, H. G.; Huberhold, M.; Thewalt, U.; Honolt, B. *J. Organomet. Chem.* **1986**, *27*.
10. Bennet, J. L.; Wolcznaski, P. T. *J. Am. Chem. Soc.* **1994**, *116*, 2179.
11. The $\{^1\text{H}\}^{29}\text{Si}$ spectrum shows a single resonance indicating no Si-P coupling.

12. a) Bercaw, J. E.; Brintzinger, H. H. *J. Am. Chem Soc.* **1971**, *93*, 2046. b) Bercaw, J. E.; Marvich, R. H.; Bell, L. G.; Brintzinger, H. H. *J. Am. Chem. Soc.* **1972**, *94*, 1219.
13. Pattiasina, J. W.; Bolhuis, F.; Teuben, J. H. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 330.
14. Berry, D. H.; Procopio, L. J.; Carrol, P. J. *Organometallics*, **1988**, *7*, 570.
15. a) Troyanov, S. I.; Varga, V.; Mach, K. *J. Chem. Soc. Chem Commun.* **1993**, 1174. b) Melmed, K. M.; Coucouvanis, D.; Lippard, S. J. *Inorg. Chem.* **1973**, *12*, 232. c) Guggenberger, L. J.; Tebbe, F. N. *J. Am. Chem. Soc.* **1973**, *95*, 7870.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

**Office of Naval Research (1)*
Chemistry Division, ONR 331
800 North Quincy Street
Arlington, Virginia 22217-5660**

**Dr. Richard W. Drisko (1)
Naval Facilities & Engineering
Service Center
Code L52
Port Hueneme, CA 93043**

**Defense Technical Information Center (2)
Building 5, Cameron Station
Alexandria, VA 22314**

**Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198**

**Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5320**

**Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000**

**Dr. John Fischer, Director (1)
Chemistry Division, C0235
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001**

**Dr. Peter Seligman (1)
Naval Command, Control and
Ocean Surveillance Center
RDT&E Division
San Diego, CA 92152-5000**

* Number of copies to forward