# **MU-ESTIMATING AND SMOOTHING**

#### **Running Title:** $\mu$ -estimation

#### Z.J. Liu and C.R. Rao

Technical Report 99-01

March 1999

Center for Multivariate Analysis 417 Thomas Building Penn State University University Park, PA 16802 19990621 003

Research work of authors was supported by the Army Research Office under Grant DAAHO4-96-1-0082. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

•

1

11

# . . . SF 298 MASTER COPY KEEP THIS COPY FOR REPRODUCTION PURPOSES

REPORT DOCUMENTATION PAGE			Fo	Form Approved	
Public recording burden for this collection of info gathering and maintaining the data needed, and collection of information including suggestions in succession, Science 204, Ariundron, VA 22207	mation is estimated to average 1 hour per completing and reviewing the collection of or reducing this burden, to Washington He 4402, and In the Office or Macagement a	response, including the time for revie information. Send comment regardini adduarters Services. Directorate for divugate Paseework Reduction Pro-	wing instructions. s g this burden estim information Operat	searching existing data sources. lates or any other aspect of this lons and Reports. 1215 Jefferson Nathorithm OC 20503	
1 AGENCY USE ONLY (Leave Diank)	2. REPORT DATE	3. REPORT TYPE	AND DATES		
4 TITLE AND SUBTITLE	5-20-99	fiechnical -	5. FUNDIN		
Mu-estimating and Smoothing			DAAF	104-96-1-0082	
6. AUTHOR(S)					
Z.J. Liu and C.R. Rao					
7 PERFORMING ORGANIZATION NAMES(S) AND ADDRESS(ES)			8. PERFO	RMING ORGANIZATION	
Center for Multivariate Analysis 417 Thomas Building			REPOR		
Department of Statistics					
Penn State University University Park, PA 16802			99	9-01	
9 SPONSCRING, MONITORING AGENCY NAME(S, AND ADDRESS(ES)			10. SPON AGEN	NSORING / MONITORING NCY REPORT NUMBER	
U.S. Army Research Office					
P.O. Box 12211 Research Triangle Park. NC 27709-2211			ARO:	35518.49-MA	
11. SUPPLEMENTARY NOTES					
The views, opinions and/or f an official Department of the	indings contained in this r Army position, policy or	eport are those of the a decision, unless so des	uthor(s) and ignated by	d should not be construed as other documentation.	
12a. DISTRIBUTION / AVAILABILITY	STATEMENT		12 b. DIS	TRIBUTION CODE	
Approved for public release: distribution unlimited.					
13. ABSTRACT In the traduunder estima a discrepancy random varia as multivaria minimizes the are independ studied the e estimation (of function of $\theta_{1}$ as in their co DISM with a allows up to computation 14. SUBJECT TERMS Asympton equation, kernel, multi-	itional M-estimation the tion is the value of $\theta$ way measure (DISM) $\delta(x)$ ble X. Such a setting the median defined by C expectation of a DISM ent copies of the under stimation of such parameter or $\mu$ -estimation for com- some complexities arise imputation. In such a con- develop an elegant app of estimations.	heory developed by $\theta$ which minimizes the $\theta$ , $\theta$ ) which is a funct does not cover the en- operator of the type $\delta(X_1,,,,,,,, .$	Huber (19 expectation ion of $\theta$ a stimation .990), as the , $X_m, \theta$ ) ble X. A associated DISM is the perties of new methor of asympt the state of the second construction of the second of the second construction of the second constr	64), the parameter on of what is called and the underlying of parameters such he value of $\theta$ which where $X_1, \ldots, X_m$ recones et al (1994) M-estimation MU- not a differentiable d of smoothing the een that smoothing otic properties and 15. NUMBER IF PAGES 21 16. PRICE CODE	
U-statistic	18. SECURITY CLASSIFICATIO	ON 19. SECURITY CLASS	SIFICATION	20. LIMITATION OF ABSTRACT	
CR REPORT	OF THIS PAGE	OF ABSTRACT		1 11	
UNCLASSIFIED	UNCLASSIFIED	UNCLASSI	FIED		
NSN 7540-01-280-5500				Standard Porti 230 (Nev. 2-8	

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. 239-18

# **MU-estimation and Smoothing**

• ,

(Running Title:  $\mu$ -estimation)

by

Z. J. Liu and C. R. Rao

Department of Mathematics and Statistics, Mississippi State University, Mississippi

State, MS 39762

Department of Statistics, Penn State University, University Park, PA 16802

## Abstract

In the traditional M-estimation theory developed by Huber (1964), the parameter under estimation is the value of  $\theta$  which minimizes the expectation of what is called a discrepancy measure (DISM)  $\delta(x,\theta)$  which is a function of  $\theta$  and the underlying random variable X. Such a setting does not cover the estimation of parameters such

Key Words and Phrases: Asymptotic, data depth, discrepancy, estimating equation, kernel, multivariate median, M-estimation, smooth, U-statistic.

AMS 1991 Subject Classifications. Primary: 62F35; Secondary: 62F12, 62H12.

as multivariate median defined by Oja (1983) and Liu (1990), as the value of  $\theta$  which minimizes the expectation of a DISM of the type  $\delta(X_1, ..., X_m, \theta)$  where  $X_1, ..., X_m$ are independent copies of the underlying random variable X. Arcones et al (1994) studied the estimation of such parameters. We call the associated M-estimation MUestimation (or  $\mu$ -estimation for convenience). When a DISM is not a differentiable function of  $\theta$ , some complexities arise in studying the properties of estimations as well as in their computation. In such a case, we introduce a new method of smoothing the DISM with a kernel function and using it in estimation. It is seen that smoothing allows us to develop an elegant approach to the study of asymptotic properties and computation of estimations.

•••

#### 1 Introduction

Let us consider a parameter  $\theta_F \in \mathbb{R}^q$  of a *p*-variate distribution F that

$$\theta_F = \arg\min\left\{\Delta(\mu, F)\right\},\tag{1}$$

where

$$\Delta(\theta, F) = E_F \delta(X_1, \dots X_m, \mu), \tag{2}$$

where  $E_F(.)$  denotes the expectation with respect to F,  $\delta(x_1, ..., x_m, \theta) : R^{mp} \times R^q \to R$ is a symmetric function of  $x_1, ..., x_m$  for every fixed  $\theta$ , called discrepancy measure (DISM), and  $X_1, ..., X_m$  are independent copies of a random variable X distributed as F. Let  $X_1, ..., X_n$  be a sample of n independent observations on  $X \sim F$ , and define  $F_n$  as the empirical distribution function based on  $X_1, ..., X_n$ . The  $\mu$ -estimate of  $\theta = (\theta_1, ..., \theta_q)$  is defined as

$$\hat{\theta} = \arg\min\{\Delta(\theta, F_n)\},\tag{3}$$

where we define

$$\Delta(\theta, F_n) = \binom{n}{m}^{-1} \sum_{1 \le i_1 < i_2 \dots < i_m \le n} \delta(X_{i_1}, \dots, X_{i_m}, \theta).$$
(4)

If (4) is differentiable, then  $\hat{\theta}$ , an estimator of  $\theta$ , is a solution of

$$D\Delta(\theta, F_n) = 0, \tag{5}$$

where D is the differential operator

$$D = \left(\frac{\partial}{\partial \mu_i}\right)_{1 \times q}, \quad D^2 = \left(\frac{\partial^2}{\partial \mu_i \partial \mu_j}\right)_{q \times q}.$$
 (6)

If the second derivative exists, we could use an iterative procedure like Newton-Raphson to compute the solution of (5). Further the asymptotic properties of the estimator can be studied without heavy assumptions. If the DISM is not differentiable, as in the case of median estimations of Oja (1983) and Liu (1990), then the problems of computing the  $\mu$ -estimate and deriving its asymptotic properties need complex approaches. To overcome the possible difficulties, we propose to smooth the DISM using a differentiable kernel function, which results in a differentiable function. We work with such a smoothed DISM and derive the asymptotic properties of the resulting  $\mu$ -estimators.

Let  $k(u): \mathbb{R}^q \to \mathbb{R}$  be a differentiable function, called a kernel function, such that

$$\int_{R^q} k(u)du = 1. \tag{7}$$

Using such a k(u), we obtain the smoothed version of the DISM by convolution

$$\delta_n(x_1, ..., x_m, \theta) = \int_{R^q} \delta(x_1, ..., x_m, \mu - h_n u) k(u) du,$$
(8)

where  $h_n$  is a bandwidth such that  $\{h_n, n = 1, 2, ...\}$  decreasing to zero as n tends to infinite.. The derivative of  $\delta_n$  is

$$D\delta_n(x_1,...,x_m,\theta) = \frac{1}{h_n^q} \int_{R^q} \delta(x_1,...,x_m,u) Dk\left(\frac{\theta-u}{h_n}\right) du.$$
(9)

Let

$$\Delta_n(\theta, F_n) = \binom{n}{m}^{-1} \sum_{1 \le i_1 < i_2 \dots < i_m \le n} \delta_n(X_{i_1}, \dots, X_{i_m}, \theta).$$
(10)

Then the  $\mu$ -estimator obtained from the smoothed DISM is a solution of the equation

$$D\Delta_n(\theta, F_n) = 0. \tag{11}$$

Since we have the estimating equation in an explicit form, we can use standard computational techniques to obtain the estimator  $\hat{\theta}$ . We now investigate the asymptotic stochastic properties of the smoothed  $\mu$ -estimator. We establish the consistency, asymptotic normality and asymptotic representation of  $\hat{\theta}$  in Section 3 of the paper. A discussion of smoothed  $\mu$ -estimator of the median when p = 1 and m = 1, and multivariate median of Liu (1990) is given in Section 4, and the results are compared with those obtained by Arcones et al (1994) using the unsmoothed DISM.

For convenience and easy understanding of the paper, we list some symbols and statements in the following:

- $\{X_i, i = 1, ..., n\}$  is an *i.i.d.* sample from a *p*-variate distribution *F*. f(x) denotes the density function, and  $F_n$  the empirical distribution.
- ||.|| denotes the Euclidean norm. C denote a positive number which may take different values in different places. Denote a<sub>n</sub> = n<sup>-1</sup> log log n. A matrix may be denoted as O(.) (or o(.)) if its norm is O(.) (or o(.)). E(.) and cov(.) denote mean and variance-covariance matrix respectively.
- Conditions for changing the order between integrals, between derivatives, and between integral and derivative are assumed to be satisfied.
- $k(u): \mathbb{R}^q \to \mathbb{R}$  is a function, called a kernel function, such that  $\int_{\mathbb{R}^p} k(u) du = 1$ .  $\{h_n, n = 1, 2...\}$  is a sequence of positive numbers called bandwidth.
- $\delta(x,\theta) = \delta(x_1,...,x_m,\theta) : R^{mp} \times R^q \to R$  is a function called a DISM (discrepancy measure) which is symmetric with respect to permutations of  $(x_1,...,x_m)$ .

A kernel smoothed DISM is

· ·

$$\delta_n(x,\theta) = \int_{R^q} \delta(x_1,...,x_m,\theta - h_n u) k(u) du.$$

• Denote  $A_{i,j}$  the element in the *i*th row and *j*th column of matrix A. A' denotes the transpose of matrix A. And

$$\begin{split} D &= \left(\frac{\partial}{\partial \theta_i}\right)_{1\times q}, \quad D^2 = D \otimes D', \quad D^3 = D \otimes \left(D \otimes D'\right), \\ D_i &= \frac{\partial}{\partial \theta_i}, \quad D_{i,j}^2 = \frac{\partial^2}{\partial \theta_i \partial \theta_j}, \quad D_{i,j,l}^3 = \frac{\partial^3}{\partial \theta_i \partial \theta_j \partial \theta_l}, \\ k_1(u) &= Dk(u), \quad k_2 = D^2 k(u), \\ \Delta(\theta, F) &= E_F \delta(X_1, \dots, X_m, \theta), \quad \Delta(\theta, F_n) = \binom{n}{m}^{-1} \sum_{1 \leq i_1 < i_2 \dots < i_m \leq n} \delta(X_{i_1}, \dots, X_{i_m}, \theta), \\ \Delta_n(\theta, F) &= E_F \delta_n(X_1, \dots, X_k, \theta), \quad \Delta_n(\theta, F_n) = \binom{n}{m}^{-1} \sum_{1 \leq i_1 < i_2 \dots < i_m \leq n} \delta_n(X_{i_1}, \dots, X_{i_m}, \theta), \\ G(\theta, F) &= D\Delta(\theta, F), \quad G(\theta, F_n) = D\Delta(\theta, F_n), \\ G_n(\theta, F) &= D\Delta_n(\theta, F), \quad G_n(\theta, F_n) = D\Delta_n(\theta, F_n). \\ S(\theta, F) &= D^2\Delta(\theta, F), \quad S(0, F) = D^2\Delta(\theta, F)|_{\theta=0}. \\ \Sigma &= cov \left(E \left(D\delta(X_1, \dots, X_m, 0) \mid X_1\right)\right), \quad \Sigma_n = cov \left(E \left(D\delta_n(X_1, \dots, X_m, 0) \mid X_1\right)\right) \end{split}$$

# 2 Existence and consistency

Under some regularity conditions, the strong law of large numbers of U-statistic implies that  $G_n(\theta, F_n)$  will converge to  $G(\theta, F)$  for every fixed  $\theta$ , and therefore when  $G(\theta, F) = 0$  is solvable, the existence of solution of (11) depends on the uniform closeness of  $G_n(\theta, F_n)$  and  $G(\theta, F)$ .

Theorem 1 Assume:

A1.  $\Delta(\theta, F)$  has a unique minimum value, reached at  $\theta = \theta_0$ . Without loss of generality, assume  $\theta_0 = 0$ .  $G(\theta, F)$  and  $S(\theta, F)$  exist for  $\theta \in B$ , an open neighborhood of 0, and  $S(\theta, F)$  is continuous and positive definite for every  $\theta \in B$ .

A2. the kernel function chosen has a compact support, is three times differentiable.  $D^2\delta_n(x_1,...,x_n,\theta)$  is continuous and has full rank for every  $\theta \in B$ .

A3. the bandwidth sequence is such that

$$\lim_{n \to \infty} h_n = 0 \tag{12}$$

A4.

$$\lim_{n \to \infty} \sup_{\theta \in B} \|G_n(\theta, F_n) - G_n(\theta, F)\| = 0. \quad w.p.1.$$
(13)

Then for large n, (11) has a unique solution  $\hat{\theta}$ , w.p.1 and

$$\lim_{n \to \infty} \hat{\theta} = 0, \quad w.p.1.$$
(14)

Remark 1 (1) It would have been more satisfactory if the conditions under which A4 holds could be stated. We hope to study this problem in a future communication. It may be easy to verify A4 in particular cases.

(2) When the DISM  $\delta$  is twice differentiable, no smoothing is needed. In such a case the bandwidth can be chosen to be 0.

*Proof.* We prove this theorem for q = 2 and  $h_n \neq 0$ . Similarly, one can prove the theorem for other cases. Note that

$$G_n(\theta, F) = \int_{\mathbb{R}^p} G(\theta - h_n u, F) k(u) du.$$
(15)

Then by conditions A2, A3, A4 and (15) we have

$$\lim_{n \to \infty} \sup_{\theta \in B} \|G_n(\theta, F) - G(\theta, F)\| = 0.$$
(16)

By condition A1 and the implicit function theorem, there exist continuous curves  $C_1(t) = (t, c_1(t))$  and  $C_2(t) = (t, c_2(t))$  such that

$$C_i(t) \subset B, \quad C_i(0) = (0,0), \quad G(C_i(t),F) = 0, \quad i = 1,2.$$
 (17)

By condition A4 and (16), it is obvious that

$$\lim_{n \to \infty} \sup_{\theta \in B} \|G_n(\theta, F_n) - G(\theta, F)\| = 0, \quad w.p.1.$$
(18)

By condition A2, (18), and the implicit function theorem, for every  $\varepsilon > 0$  and large n, there exist continuous curves  $C'_1(t) = (t, c'_1(t))$  and  $C'_2(t) = (t, c'_2(t))$  such that

$$C'_{i}(t) \subset B, \quad G_{n}(C'_{i}(t), F_{n}) = 0, \quad i = 1, 2,$$
(19)

 $\operatorname{and}$ 

$$C_i(t) - \varepsilon < C'_i(t) < C_i(t) + \varepsilon, \quad w.p.1, \quad i = 1.2.$$
(20)

Condition A1 implies that  $C_1(t)$  and  $C_2(t)$  cross with each other at a unique point which is (0,0). (20) implies that  $C'_1(t)$  and  $C'_2(t)$  cross with each other at a unique point in B when  $\varepsilon$  is small enough. This cross point is  $\hat{\theta}$ . Let  $\varepsilon \to 0$ , which requires  $n \to \infty$ . Then  $\hat{\theta} \to 0$ , w.p.1, which establish the desired result.

# 3 Asymptotic Normality and Asymptotic Representation

We first consider a  $\mu$ -estimator with a twice differentiable discrepancy function. Since we proved the existence and consistency of  $\mu$ -estimates under certain conditions, we assume  $\hat{\theta}$  exists and is consistent in the following.

**Theorem 2** Assume that the DISM  $\delta$  is twice differentiable with respect to  $\theta$ , and  $\hat{\theta}$  exists and is strongly consistent. Further assume:

B1. Both  $D\delta(X_1, ..., X_m, \theta)$  and  $D^2\delta(X_1, ..., X_m, \theta)$  have second order moments for every  $\theta \in B$ .

B2.  $G(0, F_n)$  is not a degenerate U-statistic, i.e.  $\Sigma$  is positive definite.

Then

$$\hat{\theta} = -S^{-1}(0, F)G(0, F_n) + O(a_n), \quad w.p.1,$$
(21)

and

$$n^{\frac{1}{2}}S(0,F)\widehat{\theta} \to N(0,m^{2}\Sigma), \quad in \ dist., \ as \ n \to \infty.$$
 (22)

*Proof.* By Taylor's theorem and condition B1

$$G(0, F_n) = G(0, F_n) - G(\hat{\theta}, F_n)$$

$$= -DG(0, F_n)\hat{\theta} + O(||\hat{\theta}||^2), \quad w.p.1.$$

$$(23)$$

By the law of iterated logarithm for U-statistics, we have

$$G(0, F_n) = O(a_n^{\frac{1}{2}}), \quad w.p.1,$$
 (24)

$$DG(0, F_n) = S(0, F) + O\left(a_n^{\frac{1}{2}}\right), \quad w.p.1.$$
 (25)

Therefore

$$\hat{\theta} = -S^{-1}(0, F)G(0, F_n) + O(a_n^{\frac{1}{2}}||\hat{\theta}||) + O\left(||\hat{\theta}||^2\right), \quad w.p.1.$$
(26)

(21) is proved. Since  $G(0, F_n)$  is a non-degenerated U-statistic, by Hoeffding's (1948) central limit theorem for U-statistics, (22) is true.

The following results are on smoothed estimates, i.e. those obtained by a smoothed DISM.

# **Theorem 3** Assume $\hat{\theta}$ exists and is strongly consistent and

C1. There exist  $\alpha > 0$  such that

$$cov\left(D^{2}\delta_{n}(X_{1},..,X_{m},0)\right) = O(h_{n}^{-\alpha}), \text{ therefore}$$

$$cov\left(D^{3}\delta_{n}(X_{1},...,X_{m},0)\right) = O(h_{n}^{-\alpha-2}).$$

$$(27)$$

C2. There is an integer l > 2 such that C4 holds,  $\Delta(\theta, F)$  is l times continuously differentiable, and  $\lim_{n\to\infty} \Sigma_n = \Sigma$  exists and is positive definite.

C3. k(u) has a compact support, is twice differentiable, and there is integer  $r \ge 1$  such that C4 holds and

$$\int_{R^q} u_1^{l_1} \dots u_m^{l_m} k(u) du = 0, \quad \text{for } l_j \ge 0 \text{ and } 0 < \sum_{j=1}^m l_j \le r \le l-2,$$
(28)

where  $u = (u_1, ..., u_m)$ . C4. Choose  $h_n = O\left(a_n^{\frac{1}{2r+2}}\right)$ ..

Then

٠

ς.

$$\widehat{\theta} = -S^{-1}(0, F)G_n(0, F_n) + O\left(a_n^{\frac{4r+\alpha}{4r+2\alpha}}\right), \quad w.p.1,$$
(29)

and

$$n^{\frac{1}{2}}S(0,F)\left(\widehat{\theta} - E\left(D\delta_n(X_1,...,X_k,0)\right)\right) \to N(0,k^2\Sigma), \text{ in dist. as } n \to \infty.$$
(30)

**Remark 2** 1. In condition C1, the value of  $\alpha$  depends on the smoothness of the underlying density and the DISM. In fact  $0 \le \alpha \le 4$ .

2. In the worst case where  $\alpha = 4$ , (29) becomes

$$\widehat{\theta} = -S^{-1}(0, F)G_n(0, F_n) + o\left(a_n^{\frac{r+1}{r+4}}\right), \ w.p.1.$$
(31)

The remainder can be as close to  $O(a_n)$  as possible if r is large enough. In the best case where  $\alpha = 0$ , (29) becomes

$$\hat{\theta} = -S^{-1}(0, F)G_n(0, F_n) + O(a_n), \ w.p.1.$$
(32)

3. In fact

$$EG_n(0, F_n) = \int_{R^q} D\Delta(-h_n u, F) k(u) du = O(h_n^{r+1}).$$
(33)

Condition C4 implies

$$\lim_{n \to \infty} n^{\frac{1}{2}} E G_n(0, F_n) = 0.$$
(34)

If we replace condition C4 with  $\lim_{n\to\infty} n^{1/2} h_n^{r+1}$  exists and is greater than 0, then  $\hat{\theta}$  converges to a non-central normal distribution.

Proof. By Taylor's theorem, we have

$$G_n(0, F_n) = G_n(0, F_n) - G_n(\widehat{\theta}, F_n)$$
(35)

$$= -D'G_n(0,F_n)\widehat{\theta} + O\left(||\widehat{\theta}||^2 \left\| D^3 \Delta_n(0,F_n) \right\|\right)$$

By condition C1 and the law of iterated logarithms for U-statistics, we have

$$D'G_n(0,F_n) = D'G_n(0,F) + O\left(h_n^{-\frac{\alpha}{2}}a_n^{\frac{1}{2}}\right), \quad w.p.1.$$
(36)

 $\operatorname{and}$ 

٠

...

$$D^{3}\Delta_{n}(0,F_{n}) = D^{3}\Delta_{n}(0,F) + O\left(h_{n}^{-\frac{\alpha}{2}-1}a_{n}^{\frac{1}{2}}\right), \quad w.p.1.$$
(37)

Further, by condition C3

$$D'G_{n}(0,F) = ED^{2} \int_{R^{q}} \delta(X_{1},...,X_{k},\theta - h_{n}u)k(u)du|_{\theta=0}$$
(38)  
$$= D^{2} \int_{R^{q}} \Delta(\theta - h_{n}u,F)k(u)du|_{\theta=0} = \int_{R^{q}} D^{2}\Delta(-h_{n}u,F)k(u)du$$
$$= S(0,F) + o(h_{n}^{r}).$$

Consequently

$$G_n(0,F_n) = -S(0,F)\widehat{\theta} + o(h_n^r ||\widehat{\theta}||) + O\left(h_n^{-\frac{\alpha}{2}} a_n^{\frac{1}{2}} ||\widehat{\theta}||\right)$$
(39)

$$+O(||\widehat{\theta}||^2)+O\left(h_n^{-\frac{\alpha}{2}-1}a_n^{\frac{1}{2}}||\widehat{\theta}||^2\right), \quad w.p.1.$$

By the law of iterated logarithms, we have

$$S(0,F)\widehat{\theta} = -EG_n(0,F_n) + O\left(a_n^{\frac{1}{2}}\right), \ w.p.1.$$

$$\tag{40}$$

On the other hand

٠

· •

$$EG_{n}(0, F_{n}) = D \int_{R^{q}} E\delta(X_{1}, ..., X_{m}, \theta - h_{n}u)k(u)du|_{\theta=0}$$
(41)  
$$= \int_{R^{q}} D\Delta(-h_{n}u, F)k(u)du = O(h_{n}^{r+1}).$$

Therefore

$$\hat{\theta} = O(h_n^{r+1}) + O\left(a_n^{\frac{1}{2}}\right) = O(a_n^{\frac{1}{2}}), \quad w.p.1.$$
 (42)

(29) is proved. By central limit theorem of U-statistics, we have

$$n^{\frac{1}{2}} \Sigma_n^{-\frac{1}{2}} \left( G_n(0, F_n) - EG_n(0, F_n) \right) \to N(0, m^2 I_{q \times q}), \text{ in dist.}$$
(43)

(30) is proved.

# 4 Applications

We apply the asymptotic theory of  $\mu$ -estimation established in previous sections to the median where p = 1, m = 1, and to the median of Liu (1994) using a smoothed version of the DISM.

## 4.1 Smoothed median estimator

It is well known that the  $L_1$ -norm estimator that minimizes

$$\sum_{i=1}^{n} |X_i - \theta| \tag{44}$$

is a sample median. Asymptotic properties of a sample median have been well studied. We use this example to demonstrate how well the kernel smoothing method performs. In this case, the observations are *i.i.d.* univariate random variables and m = 1. Without loss of generality, let us assume that the median is 0. Note that the smoothed discrepancy function becomes

$$\delta_n(X,\theta) = \int_R |X-\theta+h_n u| k(u) du.$$
(45)

Denote  $K(x) = \int_{-\infty}^{x} k(x) dx$ . Then we have

$$D\delta_n(X,\theta) = 2K\left(\frac{\theta-X}{h_n}\right) - 1,$$
(46)

and

$$G_n(\theta, F_n) = \sum_{i=1}^n \left( 2K\left(\frac{\theta - X_i}{h_n}\right) - 1 \right).$$
(47)

#### **Theorem 4** Assume that

D1. f(0) > 0 and f(x) is differentiable in a neighborhood of 0.

D2. k(u) is twice differentiable and has a compact support, and

$$\int_{R} uk(u)du = 0.$$
(48)

D3. Choose the bandwidth as

$$h_n = O\left(n^{-\frac{1}{3}} (\log \log n)^{\frac{1}{3}}\right).$$
(49)

Then

.

. .

1.  $\hat{\theta}$  is strongly consistent, i.e.

$$\lim_{n \to \infty} \hat{\theta} = 0, \ w.p.1.$$
<sup>(50)</sup>

2.  $\hat{\theta}$  is asymptotically normal, i.e.

$$2f(0)n^{\frac{1}{2}}\widehat{\theta} \to N(0,1), \text{ in dist., as } n \to \infty.$$
 (51)

3.  $\hat{\theta}$  can be represented as

$$\widehat{\theta} = \frac{1}{2f(0)n} \sum_{i=1}^{n} \left( 1 - 2K\left(\frac{-X_i}{h_n}\right) \right) + O\left(n^{-\frac{5}{6}} (\log\log n)^{\frac{5}{6}}\right), \ w.p.1.$$
(52)

*Proof:* We need only to verify conditions in previous theorems. We first verify condition A4 in theorem 1. Other conditions are easy to be verified. Note that

$$\sup |G_n(\theta, F_n) - G_n(\theta, F)|$$

$$= 2 \sup \left| \frac{1}{n} \sum_{i=1}^n K\left(\frac{\theta - X_i}{h_n}\right) - \int_R K\left(\frac{\theta - x}{h_n}\right) f(x) dx \right|$$

$$= 2 \sup \left| \int_R \left(F_n(\theta - h_n x) - F(\theta - h_n x)\right) k(x) dx \right|$$

$$\leq C \sup_x |F_n(x) - F(x)| \leq Cn^{-\frac{1}{2}}, \ w.p.1.$$
(53)

The last inequality is from Dvoretzky-Kiefer-Wolfowitz theorem (1956) on Kolmogorov-Simirnov distance (Serfling, 1980). By Theorem 1, (50) is true. Note that

$$S(0,F) = \frac{d^2}{d\theta^2} E|X_1 - \theta||_{\theta=0}$$

$$= \frac{d^2}{d\theta^2} \left( \int_{x>\theta} (x-\theta)f(x)dx - \int_{x<\theta} (x-\theta)f(x)dx \right)|_{\theta=0}$$

$$= \frac{d}{d\theta} \left( 2F(\theta) - 1 \right)|_{\theta=0} = 2f(0),$$
(54)

and

٠

. .

$$\Sigma_{n} = var\left(D\delta_{n}(X_{1},0)\right) = 4var\left(K\left(\frac{-X_{1}}{h_{n}}\right)\right)$$

$$= 4\int_{R}K^{2}\left(\frac{-x}{h_{n}}\right)f(x)dx - 4\left(\int_{R}K\left(\frac{-x}{h_{n}}\right)f(x)dx\right)^{2}$$

$$= 8\int_{R}F(-h_{n}u)K(u)k(u)du - 4\left(\int_{R}F(-h_{n}u)k(u)du\right)^{2}$$

$$\to 4F(0)\int_{R}2K(u)k(u)du - 4F^{2}(0) = 1, \text{ as } n \to \infty.$$
(55)

All other conditions in Theorem 3 are satisfied and note that  $\alpha = 1$ , and r = 1. Then (51) and (52) follow theorem 3.

**Remark 3** 1. Bahadur (1966) and Kiefer (1967) found an asymptotic representation of a sample quantile. The Bahadur-Kiefer representation of a sample median is given as

$$\xi_n = \xi + \frac{1}{2f(\xi)} \left( 1 - 2F_n(\xi) \right) + O\left( n^{-\frac{3}{4}} (\log \log n)^{\frac{3}{4}} \right), \ w.p.1.$$
(56)

The remainder rate cannot be improved. However, under the same conditions as those of Bahadur-Kiefer representation, a smoothed median estimator can have a representation with a remainder rate  $n^{-5/6}(\log \log n)^{5/6}$ . If f(x) is r times differentiable, the remainder rate can be  $n^{-(4r+1)/(4r+2)}(\log \log n)^{(4r+1)/(4r+2)}$ .

2. The smoothed median estimate provides an alternative computing method for  $L_1$ -norm locations and linear models. We will carry on comparison study on computation of  $L_1$ -norm for linear programing based methods and the smoothed estimation method in future communication.

#### 4.2 Simplicial median estimator

By the notion of data depth (Liu, 1990), the deepest point which is called simplicial median of a multivariate distribution is a generalization of the median to a multivariate distribution. Estimating of a simplicial median falls within the  $\mu$ estimation framework. Let  $A(x) = A(x_1, ..., x_{p+1})$  be a closed simplex in  $\mathbb{R}^p$  with vertices  $x_1, ..., x_{p+1}$ , and define

$$I_B(y) = \begin{cases} 1 & if \quad y \in B \\ 0 & if \quad y \notin B \end{cases}$$
(57)

A simplicial median DISM is

$$\delta(x,\theta) = -I_{A(x)}(\theta). \tag{58}$$

Applying kernel smooth method, we have a smoothed DISM.

$$\delta_n(x,\theta) = \int_{R^p} -I_{A(x)}(\theta - h_n u)k(u)du.$$
(59)

# Theorem 5 Assume:

٠

- E1. F(x) has a unique simplicial median which is assumed to be 0 for convenience.
- f(x) is differentiable, and S(0, F) is positive definite.
  - E2. k(u) has a compact support, twice differentiable and

$$\int_{R^p} uk(u)du = 0 \tag{60}$$

E3. 
$$h_n = O\left(n^{-\frac{1}{4}} (\log \log n)^{\frac{1}{4}}\right).$$

Then:

1.  $\hat{\theta}$  is strongly consistent, i.e.

$$\lim_{n \to \infty} \hat{\theta} = 0, \quad w.p.1. \tag{61}$$

2.  $\hat{\theta}$  can be represented as

$$\widehat{\theta} = S^{-1}(0, F)G_n(0, F_n) + O\left(n^{-\frac{3}{4}} \left(\log\log n\right)^{\frac{3}{4}}\right), \quad w.p.1.$$
(62)

3.  $\hat{\theta}$  is asymptotically normal, i.e.

$$n^{\frac{1}{2}}S(0,F)\widehat{\theta} \to N\left(0,(p+1)^{2}\Sigma\right), \quad in \ dist., \quad n \to \infty,$$
(63)

where  $\Sigma$  is the variance-covariance matrix of  $D(E(\delta(X,0)|X_1)$ 

**Remark 4** The asymptotic distribution we obtained is the same as Arcones et al's (1994) result. Our conditions are much milder, simpler, and easy to verify.

*Proof.* We need to verify condition A1 in Theorem 1 and find  $\alpha$  and r for the smoothed simplicial median estimator. By the conditions in the theorem, we know that r = 1. Note that

$$E\left(D_{i_{1},j_{1}}^{2}\delta_{n}(X,0)D_{i_{2},j_{2}}^{2}\delta_{n}(X,0)\right)$$

$$=h_{n}^{-4}\int_{R^{p}}\int_{R^{p}}\Pr\left(-h_{n}u,-h_{n}v\in A(X)\right)D_{i_{1},j_{1}}^{2}k(u)D_{i_{2},j_{2}}^{2}k(v)dudv=O(h_{n}^{-2}),$$
(64)

 $\operatorname{and}$ 

٠

• •

$$E\left(D_{i_1,j_1,l_1}^3\delta_n(X,0)D_{i_2,j_2,l_2}^3\delta_n(X,0)\right)$$
(65)

$$=h_n^{-6}\int_{\mathbb{R}^p}\int_{\mathbb{R}^p}\Pr\left(-h_nu,-h_nv\in A(X)\right)D^3_{i_1,j_1,l_1}k(u)D^3_{i_2,j_2,l_2}k(v)dudv=O(h_n^{-4}).$$

Therefore  $\alpha = 2$ . It is easy to show that

$$\lim_{n \to \infty} \sup_{x} |F_n(x) - F(x)| = 0, \quad w.p.1,$$
(66)

as in the multivariate case. It is seen that

$$\sup \|G_n(\theta, F_n) - G_n(\theta, F)\|$$

$$\leq C \sup \left\| \int I\left(\theta \in A(X)\right) d\underbrace{(F_n \times \dots \times F_n)}_{p+1} - \int I\left(\theta \in A(X)\right) d\underbrace{(F \times \dots \times F)}_{p+1} \right\|$$

$$\to 0, \quad w.p.1, \quad n \to \infty.$$
(67)

Other conditions in Theorem 3 are easy to verify. This theorem is proved.

## References

. .

Arcones, M. A, Chen, Z. and Gine, E (1994). Estimations related to U-processes with application to multivariate median: Asymptotic normality. *Ann. Statist.*, 22, 1460-1477.

Bahadur, R. R. (1966). A note on quantiles in large samples. Ann. Math. Statist.,37, 577-580.

Dvoretzky, A., Kiefer, J. and Wolfowitz, J. (1956). Asymptotic minimax character of the sample distribution function of the multinomial estimator. Ann. Math. Statist., 27, 642-669.

Hoeffding, W. (1948). A class of statistics with asymptotically normal distribution. Ann. Math. Statist., 19, 293-325.

Huber, P. J. (1964). Robust estimation of a location parameter. Ann. Math. Statist., 43, 1449-1458.

Kiefer, J. (1967). On Bahadur's representation of sample quantiles. Ann. Math. Statist., 38, 1323-1342.

Liu, R. Y. (1990). On a notion of data depth based on random simplicies. Ann. Statist., 18, 405-414. Oja, H. (1983). Descriptive statistics for multivariate distribution.. Statist. and Prob. Letters, 1, 327-333.

Serfling, R. J. (1980). Approximation theorems for mathematical statistics. Wiley, New York