

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 8-17-00	3. REPORT TYPE AND DATES COVERED Final		
4. TITLE AND SUBTITLE Quantum Optimization		5. FUNDING NUMBERS DAAH-04-96-1-0353		
6. AUTHOR(S) Griff Bilbro				
7. PERFORMING ORGANIZATION NAMES(S) AND ADDRESS(ES) North Carolina State University Raleigh, NC 27695-7514		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 36174.1-PH		
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.		12 b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) I proposed to use ideas from classical stochastic search algorithms to design a quantum computing algorithm for optimization which would tolerate quantum decoherence. I applied ideas from classical annealing algorithms to quantum optimization. I applied the Redfield theory for spin systems to model decoherence in a quantum computer. I designed a simple quantum computing cell which could be analyzed in terms of the Redfield equation. And I showed how a large number of such cells could be connected to perform a quantum version of simulated annealing for combinatorial optimization. Unfortunately, I did not succeed in effectively and correctly applying the proposed quantum annealing computer to the <i>Min Cut</i> combinatorial optimization problem I chose to attack.				
14. SUBJECT TERMS simulated annealing stochastic search quantum computing			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

Final Progress Report on Quantum Optimization

Griff L. Bilbro

Department of Electrical and Computer Engineering
North Carolina State University, Raleigh, NC

August 17, 2000

1 Statement of problem

I proposed to attack the problem of extreme sensitivity of quantum computing to decoherence by designing a new kind of quantum computer that could tolerate some decoherence by exploiting ideas from classical iterative optimization algorithms (such as steepest descent) and from stochastic search algorithms (such as simulated annealing).

QC depends delicately on the coherent superposition of so many quantum states that any computation may be too sensitive to imperfections in the implementation of program steps and maybe too sensitive to interactions with the external environment.

However some classical algorithms such as stochastic descent algorithms tolerate roundoff errors and even a finite number of large disturbances, since they are designed to gradually reduce any current error regardless whether its origin is poor initialization or perturbations in each step.

2 Results

2.1 Quantum version of stochastic search algorithm

I applied ideas from classical annealing algorithms[1, 2] to quantum computation of a minimum for the first time as follows.

Define $dU = \prod_m U_m$, so that for some H_s ,

$$dU = e^{-iH_s dt},$$

then execute the following algorithm:

REPEAT

Apply sequence of $U_m = U(i_m, j_m, k_m, \theta_m)$
 3 bits at a time, with each $|\theta_m| \ll \pi$,
 Interacting with environment

UNTIL measurement

Execution obeys $i\hbar \dot{\psi}_s = H_s(t)\psi_s$, so that the state of the computation is

$$\psi_s(t) = e^{iH_s t} \psi_s(0).$$

The program is H_s .

2.2 Redfield theory to analyze decoherence

I applied the Redfield theory for spin systems[3] to model decoherence in a quantum computer, for the first time as follows. The density matrix

$$\rho(t) = \sum_x p_x |x\rangle \langle x|$$

in the interaction representation

$$\rho^I \equiv e^{i(H_s + H_b)t/\hbar} \rho e^{-i(H_s + H_b)t/\hbar}$$

obeys the linear equations

$$\begin{aligned} \frac{d}{dt} \rho_{s\alpha\alpha'}^I(t) &= \\ &- \sum_{\beta\beta'} e^{\frac{i}{\hbar}(\epsilon_\alpha - \epsilon_\beta - \epsilon_{\alpha'} + \epsilon_{\beta'})} \\ &R_{\alpha\alpha'\beta\beta'} (\rho_{s\beta\beta'}^I - \langle \rho_{s\beta\beta'}^I \rangle_T) \end{aligned}$$




Figure 1: Quantum Shift Register

for finite T [4, p174].

2.3 Quantum computational cell

I designed a simple quantum computing cell which could be analyzed in terms of the Redfield equation as shown in Figure 1.

2.4 Quantum architecture

And I showed how a large number of such cells could be connected to perform a quantum version of simulated annealing for combinatorial optimization as shown in Figure 2.

Unfortunately, I did not succeed in effectively and correctly applying the proposed quantum annealing computer to the *Min Cut* combinatorial optimization problem I chose to attack.




Figure 2: Architecture of a Quantum Optimizer

References

- [1] S. Kirkpatrick, C. Gelatt, and M. Vecchi. Optimization by simulated annealing. *Science*, 220(4598):671–680, May 1983.
- [2] Griff L. Bilbro. Fast stochastic global optimization. *IEEE Trans. Systems, Man, and Cybernetics*, 24(4):685–689, 1994.
- [3] A. G. Redfield. *Adv. Magn. Reson.*, 1(1), 1966.
- [4] C. P. Slichter. *Principles of Magnetic Resonance*. Springer-Verlag, 2 edition, 1978.