AD_____

Award Number: W81XWH-05-1-0494

TITLE: Tumor Suppression by BRCA-1: A Critical Role at DNA Replication Forks

PRINCIPAL INVESTIGATOR: Jean Gautier, Ph.D.

CONTRACTING ORGANIZATION: Columbia University New York, NY 10032

REPORT DATE: October 2006

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188
Public reporting burden for this data needed, and completing a this burden to Department of D 4302. Respondents should be valid OMB control number. PL	collection of information is estii nd reviewing this collection of in efense, Washington Headquart aware that notwithstanding any EASE DO NOT RETURN YOU	nated to average 1 hour per resp nformation. Send comments rega ers Services, Directorate for Infor or other provision of law, no perso R FORM TO THE ABOVE ADDI	onse, including the time for revie arding this burden estimate or an rmation Operations and Reports (n shall be subject to any penalty f RESS.	wing instructions, seard y other aspect of this co (0704-0188), 1215 Jeff or failing to comply with	ching existing data sources, gathering and maintaining the ollection of information, including suggestions for reducing prson Davis Highway, Suite 1204, Arlington, VA 22202- n a collection of information if it does not display a currently
1. REPORT DATE 01-10-2006		2. REPORT TYPE		3. I 15	DATES COVERED Sen 2005 – 14 Sen 2006
4. TITLE AND SUBTIT	LE	i indi		5a.	CONTRACT NUMBER
Tumor Suppressio	n by BRCA-1:A Cri	tical Role at DNA R	eplication Forks	5b. VV8 5c.	GRANT NUMBER 31XWH-05-1-0494 PROGRAM ELEMENT NUMBER
6. AUTHOR(S)				50.	PROJECT NUMBER
Jean Gautier, Ph.[D.			5e.	TASK NUMBER
				5f.	WORK UNIT NUMBER
 F. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 				8. F	PERFORMING ORGANIZATION REPORT
Columbia Universi New York, NY 100	ty 32				IUMBER
9. SPONSORING / MO U.S. Army Medica Fort Detrick, Maryl	NITORING AGENCY N Research and Ma and 21702-5012	IAME(S) AND ADDRES teriel Command	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)
				11.	SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES Original contains colored plates: ALL DTIC reproductions will be in black and white.					
14. ABSTRACT					
BRCA1 is the major identical phenotype cellular phenotype during DNA replica laevis eggs that su response. Our key demonstrate that t BRCA1/BARD1 loa relocalized within o	or breast cancer su es suggesting that s consistent with a ation supporting DN pport: 1. Semi-con accomplishments he complex assem ading to chromatin chromatin compartr	sceptibility gene. It f these proteins funct DNA replication def IA transactions at re servative, cell-cycle were to generate sp bles to chromatin in does not dramatical nents.	forms heterodimers ion primarily as a co ect. We wished to te eplication forks. We a regulated DNA repl pecific antibodies ag a DNA replication-o lly increases followin	with BARD1. I omplex. BRCA est the hypoth are using cell- ication; 2. Ma ainst Xenopus lependent ma ng DNA dama	nactivation of either gene results in 1 deficiencies are associated with esis that BRCA1/BARD1 function free extracts derived from Xenopus ny facets of the DNA damage BARD1 and BRCA1. We also nner. Finally, we show that ge, suggesting that it might be
15. SUBJECT TERMS Tumor suppressor, genome stability, DNA replication					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	UU	7	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Cover1
SF 2982
Introduction4
Body4
Key Research Accomplishments6
Reportable Outcomes6
Conclusions6
References6

ABSTRACT

BRCA1 is the major breast cancer susceptibility gene. It forms heterodimers with BARD1. Inactivation of either gene results in identical phenotypes suggesting that these proteins function primarily as a complex. BRCA1 deficiencies are associated with cellular phenotypes consistent with a DNA replication defect. We wished to test the hypothesis that BRCA1/BARD1 function during DNA replication supporting DNA transactions at replication forks. We are using cell-free extracts derived from Xenopus laevis eggs that support: 1. Semi-conservative, cell-cycle regulated DNA replication; 2. Many facets of the DNA damage response. Our key accomplishments were to generate specific antibodies against Xenopus BARD1 and BRCA1. We also demonstrate that the complex assembles to chromatin in a DNA replication-dependent manner. Finally, we show that BRCA1/BARD1 loading to chromatin does not dramatically increases following DNA damage, suggesting that it might be relocalized within chromatin compartments.

INTRODUCTION

Mutations in the BRCA1 tumor suppressor gene occur in about 40% of familial cases of breast cancer. Carriers of BRCA1 germline mutations have a lifetime risk of 50-85% for breast cancer development. In addition, sporadic breast cancer patients often show reduced BRCA1 expression (Yoshida and Miki, 2004). The BRCA1 protein is expressed in most proliferating cells and was shown to be involved in various cellular processes including DNA repair, transcriptional regulation, chromatin remodeling and cell cycle checkpoint regulation. The BRCA1 protein contains two known amino acid motifs: a RING domain at the N terminus and two tandem copies of the BRCT domain at the C terminus. The BRCA1 RING domain forms a heterodimeric complex with another RING domain protein, BARD1. BARD1 is the major BRCA1 interacting protein in the cell. In knock-out mice, BRCA1 and BARD1 have very similar phenotypes and in Xenopus the two proteins are co-regulaed both at the mRNA and protein level, suggesting that most of BRCA1 functions might require BARD1 (Joukov et al., 2001; McCarthy et al., 2003). Indeed, BRCA1/BARD1 interaction was shown to be required for their proper localization, E3 ligase activity, DNA binding and protein stability. Following DNA damage, the BRCA1/BARD1 complex localizes to nuclear foci along with other proteins that are involved in DNA repair including Rad51 and the Mre11/Rad50/Nbs1 (MRN) complex. Both BRCA1 and BARD1 are phosphorylated in response to exogenous DNA damage. BRCA1 is phosphorylated by ATM in response to DSBs (Cortez et al., 1999) and by ATR in response to replication inhibition (Tibbetts et al., 2000) this may suggest a role for BRCA1/BARD1 complex in DNA repair following replication stress. In addition BRCA1/BARD1 nuclear foci are observed in untreated S-phase cells, supporting a role for BRCA1 and BARD1 during the DNA replication process.

DNA replication is a complex mechanism that requires the coordinated interplay of multiple factors. Many of which are involved in DNA transactions that maintain the integrity of the genome. The replication fork is a structure with the potential to trigger a DNA damage response. DNA replication fork stalling or collapse has been proposed to lead to DSBs formation and was shown to be highly reliant on homologous-recombination. Taken together, it is possible that BRCA1/BARD1 complex exerts its critical role as tumor suppressor at the replication fork where it stabilizes the fork and facilitates DNA transactions triggered by DNA damage. Our objectives are 1) to establish that BRCA1/BARD1 is present at the replication fork; 2) to determine if DNA lesion or fork stalling enhance BRCA1/BARD1 recruitment at the replication fork; 3) to assess the consequences of complete loss of BRCA1/BARD1 on fork progression and stalling.

BODY

Cell-free systems derived from Xenopus eggs can recapitulate cell cycle progression, DNA replication, mitosis, checkpoint response and DNA repair in a test tube (Costanzo et al., 2004). In our lab we have raised antibodies against xBRCA1 and xBARD1. However, only BARD1 antibodies were sensitive enough to detect a 100 kD band in Xenopus extracts as was previously described. As BRCA1/BARD1 is mainly found as a heterodimer in Xenopus extracts we focused on BARD1.



Figure 1. BARD1 associates with chromatin in a replication dependent manner.

In low-speed-supernatant (LSS) extract BARD1 is specifically associated with chromatin (Fig. 1 A). In the absence of Calcium, LSS extracts are arrested at the M phase of the cell cycle. Upon Calcium addition, that mimics fertilization, DNA templates can undergo cell-cycle regulated. semi-conservative DNA replication (Blow and Laskey, 1986). Under this condition a specific 100 kD corresponding to BARD1 chromatin association can be detected. Moreover, in the presence of Geminin, that binds to Cdt1 and inhibit MCM loading onto the

chromatin and replication initiation, BARD1 chromatin association was markedly reduced. When using Aphidicolin, that is a DNA polymerase inhibitor that blocks replication elongation, BARD1 chromatin association was not changed. This may indicate that BARD1 loades onto chromatin during the initiation steps of DNA replication.

In order to verify these results we have used a different extract preparation method term HSS (high speed supernatant) and NPE (nucleoplasmic extract). In this system DNA replication is a twostep reaction, where pre-replication complexes (pre-RC) assembly on the DNA template is physically and temporarily dissociated from origin firing. In the first step, pre-RC assembled in membrane-free cytosol (HSS) and this is followed by the addition of NPE that provides the regulatory activities required for firing including Cdk2/cyclinE and Ccd7/dbf4 (Walter et al., 1998). In HSS alone, no BARD1 chromatin association can be detected. As a loading control we have used xMCM6 antibodies. Upon addition of NPE, Cdc45 is loaded and replication can initiate, under these conditions, BARD1 is associated with the chromatin. This binding is not affected by aphidicolin that blocks the later replication initiation step, but is markedly reduced when using geminin that blocks pre-RC formation. In NPE alone that does not support DNA replication BARD1 chromatin binding could not be detected. Thus, we can conclude that BARD1 associates with chromatin during the replication initiation.



Figure 2. BARD1 loading requires pre-RC assembly. Left panel. Schematic of pre-RC assembly showing the steps that are inhibited by geminin and by aphidicolin.

Right panel. Chromatin loading of BARD1 following the inhibition of pre-RC assembly (Geminin) or DNA replication elongation (aphidicolin). In response to DNA damage, BRCA1/BARD1 localizes to nuclear foci containing other DNA repair proteins. Next we looked at BARD1 chromatin association in response to different DNA damaging agents. In response to Mitomycin C (150mM) DNA cross linking agent, BARD1 chromatin association did not change. However, and Cisplatin (25mM) which is also a DNA cross-linking agent completely abolished BARD1 chromatin association. This is probably due to high dose as indicated by the lack of MCM6 control loading that indicate that there was no replication. In response to Etoposide (100mM) treatment that inhibit topoisomerase II and induce DNA breaks and inhibit DNA synthesis BARD1 chromatin association did not change. Finally, Camptothecin (50mM) that inhibit topoisomerase I and increased DNA cleavage also did not change BARD1 chromatin association. Altogether, we can conclude the no increase in BARD1 chromatin association can be detected following DNA damage.



Figure 3: Chromatin-bound BARD1 following DNA damage.

BARD1 loading onto chromatin is not increased following treatments with mitomycin C, a cross-linking agent, etoposide, a DNA topoisomerase II inhibitor or Camptothecin, a DNA topoisomerase I inhibitor

KEY RESEARCH ACCOMPLISHMENTS

- To generate specific antibodies against Xenopus BARD1 and BRCA1.
- To demonstrate that the BRCA1/BARD1complex assembles to chromatin in a DNA replication-dependent manner.
- To show that BRCA1/BARD1 loading to chromatin does not dramatically increases following DNA damage, suggesting that it might be relocalized within chromatin compartments.

REPORTABLE OUTCOME

An abstract will be presented at the Keystone meeting on DNA recombination, January 2006

CONCLUSIONS

We have made significant progress by demonstrating the pre-Replicative-dependent loading of BARD1 onto chromatin. This is a critical accomplishment that is in strong support of our original hypothesis that BRCA1 is part of the replication machinery. It will serve as a base for future efforts. Furthermore, our preliminary data suggests that the BRCA1/BARD1 complex might not be enriched on chromatin following DNA damage but instead be recruited from its replication-dependent localization on chromatin to damaged sites.

REFERENCES

Blow, J. J., and Laskey, R. A. (1986). Initiation of DNA replication in nuclei and purified DNA by a cell-free extract of Xenopus eggs. Cell 47, 577-587.

Cortez, D., Wang, Y., Qin, J., and Elledge, S. J. (1999). Requirement of ATM-dependent phosphorylation of brca1 in the DNA damage response to double-strand breaks. Science 286, 1162-1166.

Costanzo, V., Robertson, K., and Gautier, J. (2004). Xenopus cell-free extracts to study the DNA damage response. Methods Mol Biol *280*, 213-227.

Joukov, V., Chen, J., Fox, E. A., Green, J. B., and Livingston, D. M. (2001). Functional communication between endogenous BRCA1 and its partner, BARD1, during Xenopus laevis development. Proc Natl Acad Sci U S A *98*, 12078-12083.

McCarthy, E. E., Celebi, J. T., Baer, R., and Ludwig, T. (2003). Loss of Bard1, the heterodimeric partner of the Brca1 tumor suppressor, results in early embryonic lethality and chromosomal instability. Mol Cell Biol *23*, 5056-5063.

Tibbetts, R. S., Cortez, D., Brumbaugh, K. M., Scully, R., Livingston, D., Elledge, S. J., and Abraham, R. T. (2000). Functional interactions between BRCA1 and the checkpoint kinase ATR during genotoxic stress. Genes Dev *14*, 2989-3002.

Walter, J., Sun, L., and Newport, J. (1998). Regulated chromosomal DNA replication in the absence of a nucleus. Mol Cell *1*, 519-529.

Yoshida, K., and Miki, Y. (2004). Role of BRCA1 and BRCA2 as regulators of DNA repair, transcription, and cell cycle in response to DNA damage. Cancer Sci *95*, 866-871.