

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 04-06-2008		2. REPORT TYPE ONR Report-Final		3. DATES COVERED (From - To) 30-DEC-2002 thru 31-DEC-2007	
4. TITLE AND SUBTITLE Accelerating Electronic Tag Development for Tracking Free-Ranging Marine Animals at Sea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER N000143-03-1-0325	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Barbara A . Block				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Board of Trustees of the Leland Stanford Junior University, Office of Sponsored Research, 340 Panama Street, Stanford CA 94305-6203				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Stanford University, Office of Sponsored Research, 340 Panama Street, Stanford CA 94305-6203				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This project developed, tested, and deployed archival and satellite tags for the Tagging of Pacific Pelagics (TOPP). Our efforts centered both on improving existing technologies and on developing new tools. These new tools have provided TOPP the tools necessary to address fundamental questions in biological oceanography concerning the distribution and critical habitats of pelagic organisms. We developed a GPS tag using Fastloc technology making it possible to study animal movements on a scale of 10-50 meters. The GPS tag design developed under this NOPP award is now commercially available from the Sea Mammal Research Unit, Wildlife Computers Inc (Redmond WA) and Sirtrack Ltd New Zealand. We developed a CTD tag that is now commercially available from the Sea Mammal Research unit. This tags is accurate to ± 5 dBar, $\pm 0.001^\circ\text{C}$, and ± 0.003 mS/cm. These tags have made possible a series of international collaborations SEAOS and the IPY (International Polar Year) initiative MEOP (Marine Mammals as Explorers of the Ocean Pole to Pole).					
15. SUBJECT TERMS Lotek archival tag pop-up satellite archival (PAT) blue shark, mako shark, salmon shark, white sharks.)SPOT (Satellite Position Only Tags or Temperature Transmitting) marine organisms pelagic predator North Pacific ecosystems marine mammals, tunas,					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Barbara Block
U	U	U	UU	1	19b. TELEPHONE NUMBER (Include area code) 831-655-6236

Dr. Barbara A. Block

bblock@stanford.edu

THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY

Biology

120 Ocean View Blvd

Pacific Grove, CA

93950

Phone: (831) 655-6236

Fax: (831) 375-0793

Website URL: www.topp Census.org

Project Title: Accelerating Electronic Tag Development for Tracking Free-Ranging Marine Animals at Sea

ONR Award No: N000140310325

Organization Award No: 26306

Final Report

Award Period: December 30, 2002 - August 30, 2007

LONG-TERM GOALS

The objective of this proposal is to complete the development, testing, and deployment of archival and satellite tags for the Tagging of Pacific Pelagics (TOPP) pilot program of the Census of Marine Life. Efforts have centered both on improving existing technologies and on developing new tools that are now allowing us to address more complex questions about the animals and their environment. First, electronic tags are providing TOPP with the tools necessary to address fundamental questions in biological oceanography concerning the distribution and critical habitats of pelagic organisms. This information is being used to describe the movements and behaviors of marine vertebrates and large squid in the North Pacific and to identify hotspots and migratory corridors. Second, with the new tags developed under this effort we are now using the tag-bearing animals as autonomous ocean profilers and we are providing oceanographic data to the growing global databases. The vertical and horizontal movements of tagged marine vertebrates allow sampling of more remote, traditionally under sampled areas, as well as providing unprecedented temporal and spatial coverage of the North Pacific. Not only are these data of great value to oceanographers but when the biological and physical data are merged, we can obtain an "organism-eye" view of how marine animals from several trophic levels use distinct oceanic regions.

OBJECTIVES

To accelerate the electronic tags available to track marine animal movements. The specific objectives were to: Develop a Conductivity Temperature depth tag (CTD), develop a GPS tag, further test and refine the archival and satellite tags currently available. This included testing and modification of tags built by archival and satellite tags from Wildlife Computers and archival tags from Lotek Wireless. Finally our goal was to test these tags and further the ability to use animals to acquire oceanographic data.

APPROACH

To accelerate tag technology, TOPP scientists have been working with engineers at Wildlife Computers (USA), Wildtrack (U.K), Sirtrack (New Zealand) Lotek Technology (Canada), and the Sea Mammal Research Unit (U.K.). New tags have been tested on elephant seals, salmon sharks and

20080707 037

bluefin tuna, all proven test platforms for tag deployment. Tags and algorithms that have been developed include Conductivity – Temperature – Depth (CTD), GPS, and single and dual light wavelength tags for estimating primary productivity. For marine mammal deployments, tags were initially tested through a series of 2-5 day elephant seal translocations experiments followed by longer deployments on seals during their post-breeding or post-molt migrations. Prior to deployment and after recovery, each tag is rigorously calibrated to determine instrument drift. Tags that were tested include a number of archival tags, pop-up satellite tags, single position only tags, and Satellite Relay Data Loggers (SRDLs), CTD and GPS tags. Tag performance is carefully evaluated by assessing the durability under field conditions, effectiveness of attachment on the animal, battery performance, accuracy, and responsiveness of tags to measure temperature, depth, light, and other variables. For fish and shark deployments, the focus is on solving algorithm problems and increasing the longevity of deployment duration without tag failure.

WORK COMPLETED

By working internationally as a team (TOPP, SMRU, Lotek, Wildlife Computers) we have used the support from NOPP to bring to fruition GPS tags for increasing the accuracy of marine mammal positions, CTD tags that provide salinity data along with temperature profiles, single position only tags for monitoring five species of shark (SPOT), archival tags for electronic tagging of tuna and seabirds, and algorithms that provide a measurement for in vitro chlorophyll in the ocean (Teo 2006). Using Fast-GPS tags, it is now possible to obtain animal locations to within 10m with a corresponding water temperature profile that has a resolution of 0.05°C and an accuracy of $\pm 0.1^\circ\text{C}$. The tests in 2006 and 2007 with seals and sea lions demonstrated that GPS tags allow identification of water column features significantly greater than ARGOS and at a much higher resolution than satellite remote sensing. We are now exploring incorporating Application-Specific Integrated Circuit (ASIC) technology to reduce the cost and size of the tags. ASIC chips in the tuna light-based geolocation tags would allow us to reduce the size of the tags and extend TOPP technology to coastal species such as salmon or smaller seabird species. This would be a significant advance for the biologging community.

Advances in electronic tag technology have provided TOPP with the tools necessary to describe the movements and behaviors of marine vertebrates and large squid in the North Pacific and to identify hotspots and migratory corridors. The tag-bearing animals continue to serve as autonomous ocean profilers providing key oceanographic data to the growing global databases. The vertical and horizontal movements of tagged marine vertebrates have provided the sampling of more remote, traditionally undersampled oceanic areas, as well as providing unprecedented temporal and spatial coverage of the North Pacific. To date, over 2,500 tags have been deployed and several million ocean observations have been documented. Efforts to database the entire TOPP data set are on-going and new collaborations with JPL-NASA for data assimilation into ocean general circulation models have been established.

RESULTS

GPS tag: Development of a GPS tag using *Fastloc* technology has made it possible to study animal movements on a scale of 10-50 meters compared to the 1-10 kilometers possible with ARGOS satellite tags (Figure 1). The development of a GPS system that could work with marine animals was a major goal of this grant as the achievement of this precision would allow measurements of animal movements relative to the mesoscale features, and would significantly improve the quality of the physical oceanographic data collected by the animals. We supported engineer Ed Bryant of Wildtrack Telemetry Systems Ltd (Leeds, England) to develop a functional prototype of a small GPS system that




Figure 1. Comparison of an ARGOS track in red to a GPS track in blue for a female California sea lion foraging in the Southern California Bight.

can operate within the narrow bandwidth confines of the ARGOS system. This system is called *Fastloc* and uses a novel intermediate solution that couples brief satellite reception with limited onboard processing to reduce the memory required to store or transmit the location. *Fastloc* first acquires a 10-20 ms snapshot of GPS satellite signals (stage 1) and then uses a digital signal processor (DSP) chip to identify the observed satellites and their pseudo-ranges (Stage II). DSP calculations take about 5 s, but this can be carried out when an animal is underwater. The resulting processed data are only 29 bytes long and so *Fastloc* data can be relayed via the ARGOS system or it can be archived for later retrieval if and when the tag is recovered. Final stage 3 *Fastloc* location determinations are calculated in the lab using GPS constellation orbitography data from archives available on line (e.g.). Satellite-linked GPS tags from

Sea Mammal Research Unit (SMRU), Sirtrack, and Wildlife Computers have been tested on 40 elephant seals, 26 California and 5 Galapagos sea lions, 5 Australian fur seals and 13 Cape fur seal. Further, using GPS tags we tested the quality of locations generated by the new "template math" algorithm developed by Lotek Ltd. GPS and Lotek archival tags were simultaneously deployed on 10 female California sea lions on San Nicolas Island. All tags were recovered and we are currently comparing the quality of the "template math" algorithm for light based geolocation to the high quality locations generated by the *Fastloc* GPS tags. Template math has the potential to reduce the error of light based geolocation to within 10 to 30 km.

CTD tag: The CTD tag is now commercially available from the Sea Mammal Research unit. The CTD-SRDL tag has operational characteristics of -5 to +35 °C with an accuracy of 0.005 °C and a resolution of 0.001 °C. Following successful tests with CTD tags, long-term deployments were conducted on northern and southern elephant seals and sea lions. In the spring of 2005 and winter and spring of 2006, 16 CTD tags, which relay near real-time data to Argos satellites, were deployed on elephant seals at Año Nuevo, California (Figures 2 & 3). From these deployments (excluding spring 2005), 20,190 dives, 7,150 filtered Argos locations, and 1,826 CTD casts were obtained. The overall




Figure 2. The inset at lower left shows an elephant seal with the CTD tag. The two images show the coverage across the North Pacific from only 7 CTD deployments on female elephant seals from the Año Nuevo rookery.

performance of the tags was good but several earlier models malfunctioned and post recovery calibrations on others were anomalous. Upon recovery of the tags that malfunctioned we were able to discover that the CTD casing had leaked. Given this information the Valeport LTD, the CTD manufacturer, undertook a complete redesign of the CTD head. The newly redesigned tags have been deployed on southern and northern elephant seals, crabeater, Weddell, grey, and hooded seals as part of the SEAOS and MEOP programs. In addition to the deployments on northern elephant seals, 19 CTD tags were also deployed on southern elephant seals from the South Shetland Islands in the Antarctic Peninsula with funding from the National Undersea Research Program, NOAA (7 tags in 2005) and Office of Polar Programs NSF (12 tags in 2006).

Archival tags: Between June 1, 2005 and May 31, 2006, three hundred and forty-six Lotek archival tags were deployed. Seventy-six tags were deployed on bluefin, 168 tags deployed on yellowfin, and 102 tags deployed on albacore tuna off the coast of California, USA and Baja California (Figure 4). Recovery of the current D-series of tags has indicated that algorithms to correct passivation problems employed have been useful for increasing the longevity of the track missions. From August 2002 until June 2006 a total of eight hundred and sixty-six Lotek archival tags were deployed on bluefin, yellowfin, and albacore tuna off the coast of California, USA and Baja California. Increased sampling intervals were tested (4 - 20 seconds) for comparison to prior deployments and increased oceanographic sampling. The data recovered from recaptured tunas (mean of 320 days) will allow for improved chlorophyll a estimates and water column temperature profiles. From previous deployments, we have recovered nearly 50% of deployed tags from bluefin (158) and yellowfin (133) tuna. Lotek Wireless 2400 archival tags were deployed on 60 albatrosses and 49 shearwaters for durations up to 313 days. These tags revealed post breeding movements and diving behavior of the birds as well as SSTs for a significant portion of the North and South Pacific. Locations of dives for shearwaters reveal strong




Fig 3. Elephant track showing temperature profile as the animal moves through the eddy.

relationships with remotely sensed primary productivity data. In fall 2005, we obtained the first tag recoveries from migrating sooty shearwaters equipped with archival geolocation tags during the breeding period in January 2005 at two breeding colonies in New Zealand. Upon completion of breeding in New Zealand, each bird migrated to one of three destinations in the North Pacific: 1)




Figure 5. Tracks of sooty shearwaters tagged at New Zealand. From Shaffer *et al.* 2006 PNAS.

western Pacific, 2) Alaska, and 3) Coastal California and Mexico (Figure 5). Sooty shearwaters spent nearly half the year (May – late September) at these over-wintering zones in the northern hemisphere. Shearwaters traveled more than 60,000 km roundtrip, which is one of the longest animal migrations yet recorded. This strategy of breeding in New Zealand during the austral summer followed by a migration to the northern hemisphere during the boreal summer allows sooty shearwaters to enjoy an endless summer while exploiting oceanic resources on a global scale (Shaffer *et al.* 2006).

deployed on blue, mako, salmon and white sharks. To date more than 70% of the tags deployed on all species have reported and tag performance in general has been excellent. The fast responding thermistor now allows for more accurate measurement of the water column profile and sea surface temperature (SST), which will improve SST-based latitude estimates. Over 5,000 temperature and depth profiles have been gathered in the North Pacific from shark data sets. Over the past year the white shark team has deployed 30 tags on 29 adult white sharks. One individual was tagged with two PAT tags, one long-term deployment tag and one short-term, high resolution tag. Thus far 5 tags have released including the high resolution tag. One tag released prematurely after only 6 weeks. The remaining four PAT tags surfaced in the mid Pacific area between California and Hawaii. All tags have provided high quality data. The remaining deployed tags have been silent for over seven months, which is very promising for attaining data on white shark movements during spring and summer, a season for which previous data are sparse.

PAT: Between June 1, 2005 and May 31, 2006 a total of (68) pop-up satellite archival (PAT) tags were

SPOT: Between June 1, 2005 and May 31, 2006 a total of (49) SPOT (Satellite Position Only Tags or Temperature SPOT: Between June 1, 2005 and May 31, 2006 a total of (49) SPOT (Satellite Position Only Tags or Temperature Transmitting) tags were deployed on blue, mako, and salmon sharks. Sixteen of the 25 tags deployed on salmon sharks and 1 of 9 tags on mako sharks were still transmitting on 31 May 2006, after nearly one year. We continue to receive transmissions from 1 tag deployed in 2003 and 8 tags deployed in 2004 on salmon sharks. SPOT tags have provided a phenomenal amount of data for both the salmon (Figure 2) and mako sharks. The SPOT tags on blue sharks have not performed as well and currently only 2 of 15 tags are transmitting after 9 months. We continue to use SPOT tags for our colony based research on albatrosses, which now totals 233 individuals tracked.




Fig. 4. A) Geolocation estimated for albacore (white), yellowfin (yellow) and bluefin (blue). B) The depth and temperature of bluefin, b) yellowfin and c) albacore for one day during the summer when all species are in the same regions off Northern Baja, Mexico.

TOPP's animal oceanographers include tracks from 32 elephant seals that generated 9,697 CTD profiles along the CCS and the NEP (Figure 2 & 3). These CTD profiles are augmented with 109,405 temperature profiles from 93 elephant seal, 1,781 temperature and depth profiles from California sea lions, and 200,000 temperature profiles from 130 bluefin tuna that heavily sample the California Current ecosystem in regions where there are few ship or AUV observations. Two hundred TOPP tagged sharks have provided over 16,000 profiles throughout the North Pacific. Albatrosses have collected over 10,000 SST measurements from across the entire Pacific in one summer season. Together these data provide important physical oceanographic input to ocean observation and essential habitat models.

IMPACT/APPLICATIONS

National Security

Marine animals do not observe any national boundaries. Our work has shown that a number of species freely migrate great distances over the ocean. The sooty shearwater alone migrates over 18 different national EEZs. These data are important to understand the potential of these organisms to carry disease vectors such as avian influenza.

Economic Development

This project has produced a number of completely new tags (CTD and GPA) and has vastly improved on the design of existing tags (Lotek Archival tags). These tags are now being used by a large number of investigators to study marine organisms. This grant has had a direct economic impact by the creation of these new tags that are now currently on the market.

Quality of Life

Our ability to identify oceanic hotspots used by marine predators will have significant implications for fisheries management and conservation. For example, areas that are deemed "sensitive" or critical to the proliferation of a given species could be protected or managed. However, because the oceans are so dynamic, it is important to identify key features or consistent phenomena (e.g. coastal upwelling or other physical forcing) that affect ocean productivity and the aggregation of predators and prey. Bio-logging science is an emerging field that bridges electronic tagging, biology, oceanography and computer science. Advances in microprocessor-driven electronic tags have advanced our ability to collect information on marine animals on an ecosystem scale. The TOPP team is equipping animals with satellite and archival tags that provide data on the location of the animals and the oceanic environment through which they travel. These newly developed electronic tagging techniques are providing the means to elucidate the structure and function of open-ocean ecosystems from the viewpoint of the top predators. In addition, the data collected by animals are contributing to our ability to observe the circulation and physical patterns in the ocean on a global scale. Animal-collected oceanographic data have a nearly untapped potential to contribute to the developing global ocean observing system. The TOPP team has taken the first steps to integrate across disciplines, combining the animal collected data with oceanographic data sources. Such information will provide needed data for poorly sampled regions that can then be used for developing and testing models. Sub-surface sampling has particular value, as these data are the scarcest yet are critical for models of ocean-atmospheric coupling and global heat balance.

Marine Ecosystem-based Resource Management

The major output of TOPP will be the ability examine or model the movements of pelagic predators leading to a greater understanding of the North Pacific ecosystems. Ultimately, TOPP will show us where marine mammals, tunas, seabirds, turtles and sharks go to feed and breed. Moreover, TOPP will examine how the physical dynamics of North Pacific influence the distribution and abundance of predators from a variety of trophic levels. The results should provide information on how animals at different levels of the food web are coupled to oceanic processes, where bottom-up effects enhance primary productivity. In essence, the results will provide information on the key physical and biological processes involved in open-ocean ecosystem dynamics. The data will be extremely important for fisheries management plans for the U.S. and Mexican EEZ.

The acquisition of data on 22 species over multiple years will provide the necessary foundation of information for developing predictive models of marine predator movements in relation to environmental conditions. This level of understanding is required for dynamic, adaptive fisheries management and provides the information needed to conserve and manage marine resources.

Science Education and Communication

The NOPP award has directly supported one post doctoral researcher, one engineer, and 2 graduate students. The results of this research are communicated to the public on the TOPP web page, www.toppcensus.org.

TOPP web efforts: Our web team regularly added content features to the public web site, providing coverage of squid, sea bird, tuna, whale and shark tagging efforts. Each feature included one or more written stories, supplemented by multimedia components. The most elaborate of these was in the salmon shark feature, which included animated figures from the Science paper, narrated by Kevin Weng, the paper's lead author.

Media relations: The TOPP program received national news coverage associated with the salmon shark paper in *Science*. AAAS decided to publish the paper side by side with another piece on white shark migrations by Ramón Bonfil et. al., and their own press release lumped both papers together. In 2006, our TOPP seabird researchers published a paper in the *Proceedings of the National Academy of Sciences USA* on sooty shearwater migrations. The paper was covered in over 70 websites and papers, as well as four radio interviews.

Informal education: The plan to share TOPP with aquarium visitors took a significant step forward when it was incorporated into the 2006 budget and planning process for the Monterey Bay Aquarium. TOPP will be incorporated into a remodeled “Vanishing Wildlife” exhibit, which surrounds the lower viewing window into the million-gallon Outer Bay exhibit. Planning for this will begin in November 2005, with the exhibit opening planned for Spring 2007.

Education and Outreach: In January 2006 the TOPP web site received a Level 2 Ultraweb Award. Of the nearly 5,600 applicants in 2005, fewer than 100 achieved this level. In January we also launched the “TOPP Communication Network,” a phone-based web site designed to facilitate information sharing within TOPP. Visit the award winning website at: <http://www.topp Census.org/>.

The white shark animation on TOPP’s home page was recognized this spring by the judges of the Web Marketing Association’s ninth annual Internet Advertising Competition (IAC) Awards as the best non-profit rich media online ad. The (IAC) Awards honor excellence in online advertising and recognize the individuals and organizations responsible for the best in Internet marketing. The IAC Awards are the first and only industry- based advertising award competition dedicated exclusively to online advertising. Other winners included Disney Online, Foote Cone & Belding, Walt Disney Parks and Resorts, America Online, Inc., and Ogilvy Worldwide.

E&O accomplishments also include a new print piece. The team completed a TOPP 8-1/2 by 11 inch single fold, full color, glossy brochure designed by the Monterey Bay Aquarium’s print graphics team.

TRANSITIONS

The GPS tag design developed by WildTrack Ltd (Leeds England) with support from this NOPP award is now commercially available from the Sea Mammal Research Unit, Wildlife Computers Inc (Redmond WA) and Sirtrack Ltd New Zealand.

RELATED PROJECTS

This project enabled the start of the Tagging of Pacific Pelagics program (TOPP) which is a pilot project of the Census of Marine Life (<http://www.topp.org>). TOPP is pioneering the application of biologging science to study pelagic habitat use by marine vertebrates and large squid in the North Pacific. Tags developed under this grant allowed the development of two major international research efforts. SEAOS and the IPY (International Polar Year) initiative MEOP (Marine Mammals as Explorers of the Ocean Pole to Pole). MEOP was developed from the success of SEAOS (Southern Elephant Seals as Ocean Sensors), which began in 2004 using elephant seals to collect a synoptic picture of the physical oceanography of the Southern Ocean. Participants have deployed the newly developed SMRU CTD tag on southern elephant seals from Australian, French, UK and USA research stations spanning the entire Southern Ocean.

Publications

2003

Block, B. A., Costa, D. P., Boehlert, G. W. and Kochevar, R. E. 2003. Revealing pelagic habitat use: the tagging of Pacific Pelagics program. *Oceanol. Acta*. 25: 255-266.

2004

Bograd, S. J., Foley, D. G., Schwing, F. B., Wilson, C., Laurs, R. M., Polovina, J. J., Howell, E. A. and Brainard, R. E. 2004. On the seasonal and interannual migrations of the transition zone chlorophyll front. *Geophys. Res. Lett.* 31, doi:10.1029/2004GL020637

Costa, D. P. and Sinervo, B. 2004. Field physiology: Physiological insights from animals in nature. *Ann. Rev. Physiol.* 66:23.1–23.30

Costa, D. P., Kuhn, C. E., Weise, M. J., Shaffer, S.A. Arnould, J. P. 2004. When does physiology limit the foraging behaviour of freely diving mammals? *Intl. Congr. Ser.* 1275:359– 366.

Shaffer, S. A. Costa, D. P., Suryan, R. M., Hyrenbach, K. D. 2004. North Pacific: breeding and non-breeding. In Global Procellariiform Tracking Workshop Report (ed J. P. Croxall). BirdLife International. Cambridge, pp. 47-49.

Shaffer, S. A. and Costa D. P. 2004. Tagging of Pacific Pelagics. In Global Procellariiform Tracking Workshop Report (ed J. P. Croxall). BirdLife International. Cambridge, pp. 83.

Teo, S. L. H., Boustany, A., Blackwell, S., Walli, A., Weng, K. C. & Block, B. A. 2004. Validation of geolocation estimates based on light level and sea surface temperature from electronic tags. *Mar. Ecol. Prog. Ser.* 283: 81-98.

Weng, K. and Block, B. A.. 2004. Tracking of big-eye thresher shark with satellite tags: *Fish Bull.* 102: 221-220.

2005

Block, B. A. 2005. Physiological Ecology in the 21st Century: Advancements in Biologging Science. *Integr. Comp. Biol.* 45: 305–320.

Markaida, U., Rosenthal, J. J. C., and Gilly, W. F. 2005. Tagging studies on the jumbo squid (*Dosidicus gigas*) in the Gulf of California, Mexico. *Fish. Bull.* 103: 219-226.

Palacios, D. M., and S. J. Bograd, 2005. A census of Tehuantepec and Papagayo eddies in the northeastern tropical Pacific. *Geophys. Res. Lett.* 32, doi:10.1029/2005GL024324.

Shaffer, S. A., Tremblay, Y., Awkerman, J. A., Henry, R. W., Teo, S. L. H., Anderson, D. J., Croll, D. A., Block, B. A., and Costa, D. P. 2005. Comparison of light- and SST-based geolocation with satellite telemetry in free-ranging albatrosses. *Mar. Biol.* 147: 833-843.

Wallace, B. P., Williams, C. L., Paladino, F. V., Morreale, S. J, Lindstrom, R. T. and Spotila, J. R. 2005. Bioenergetics and diving activity of internesting leatherback turtles *Dermochelys coriacea* at Parque Nacional Marino Las Baulas, Costa Rica. *J. Exp. Biol.* 208: 3873-3884.

Weng, K. C., Landiera, A., Castilho, P. C., Holts, D. B., Morrisette, J. M., Schallert, R. J., Goldman, K. J., & Block, B. A. 2005. Warm Sharks in Polar Seas: Satellite Tracking from the Dorsal Fins of Salmon Sharks. *Science* 310: 104-106.

2006

Arnould, J. P. Y. and Costa, D. P. 2006. Sea lions in drag, fur seals incognito: insights from the otariid deviants. Symposium Volume Sea Lions of the World, Univ of Alaska Press. Alaska Sea Grant College Program • AK-SG-06-01, 2006. Pages 111-125.

Costa, D. P., Weise, M. J. and Arnould, J. P. Y. 2006. Worldwide Pinniped Population Status and Trends Pages 342-357, in Whales, Whaling and Ocean Ecosystems ed Estes, J. A. Williams, T. M., Doak, D. and DeMaster, D. University of California Press Berkeley, CA

Costa, D. P. 2006. Elephant seals. World Book Encyclopedia.

Costa, D. P. 2006. The Physiology of Diving. Encyclopedia of Life Sciences. doi: 10.1002/9780470015902.a0004230. Pages 1-7

Crocker, D. E., Costa, D. P., Le Boeuf, B. J., Webb, P. M., and Houser, D. S. 2006. Impact of El Niño on the foraging behavior of female northern elephant seals. *Mar. Eco. Prog. Ser.* 309: 1-10.

Gilly, W. F., C. Elliger, C. Salinas, S. Camarillo-Coop, G. Bazzino and M. Beman. 2006. Spawning by jumbo squid *Dosidicus gigas* in the San Pedro Martir Basin, Gulf of California, Mexico. *Mar. Eco. Prog. Ser.* 313: 125-133.

Gilly, W. F., Markaida, U., Baxter, C. H., Block, B. A., Boustany, A., Zeidberg, L., Reisenbichler, K., Robison, B., Bazzino, G. and Salinas, C. 2006. Vertical and horizontal migrations by the jumbo squid *Dosidicus gigas* revealed by electronic tagging. *Mar. Eco. Prog. Ser.* 324: 1-17.

Kuhn, C. E., Auriolles-Gamboa, D. and Costa, D. P. 2006. Oxygen Stores of California sea lion pups: Implications for diving. Symposium Volume Sea Lions of the World, Univ of Alaska Press. Univ of Alaska Press. Alaska Sea Grant College Program • AK-SG-06-01, 2006. Pages 1-13.

Kuhn, C. E. and Costa, D. P. 2006. Identifying and quantifying prey consumption using stomach temperature change: a comparison between a seal and sea lion species. *J. Exp. Biol.* 209: 4524-4532.

Palacios, D. M., Bograd, S. J., Schwing, F. B. and Foley, D. G. 2006. Oceanographic characteristics of biological hot spots in the North Pacific: A remote sensing perspective. *Deep-Sea Res. II* 53: 250-269.

Shaffer, S. A. and Costa, D. P. 2006. A Database for the Study of Marine Mammal Behavior: Gap Analysis, Data Standardization, and Future Directions. *IEEE J Ocean Engineer.* Vol 31(1):82-86.

Shaffer, S. A. Tremblay, Y. Weimerskirch, H., Scott, D., Thompson, D. R. Sagar, P. M., Moller, H., Taylor, G. A., Foley, D. G., Block, B. A., and Costa, D. P. 2006. Migratory shearwaters integrate oceanic resources across the Pacific Ocean in an endless summer. *Proc. Natl. Acad. Sci. USA* 103: 12799-12802.

Tremblay, Y., Shaffer, S. A., Fowler, S. L., Kuhn, C. E., McDonald, B. I., Weise, M. J., Bost, C. -A., Weimerskirch, H., Crocker, D. E., Goebel, M. E., Costa, D. P. 2006. Interpolation of animal tracking data in a fluid environment. *J. Exp. Biol.* 209: 128-140.

Weise, M. J., D. P. Costa, and R. M. Kudela. 2006. Movement and diving behavior of male California sea lion (*Zalophus californianus*) during anomalous oceanographic conditions of 2005 compared to those of 2004. *Geophys. Res. Lett.* 33: L22S10.

Yen, P. P. W., Sydeman, W. J., Bograd, S. J. and Hyrenbach, K. D. 2006. Spring-time distributions of migratory marine birds in the southern California Current: Oceanic eddy associations and coastal habitat hot spots over 17 years. *Deep-Sea Res. II* 53: 399-418.

2007

Blank, J. M., Morrisette, J.M., Farwell, C.J., Price, M., Schallert, R. and B. A. Block. 2007. Temperature effects on metabolic rate of juvenile Pacific bluefin tuna (*T. Orientalis*) in the lab and wild. 2007. *J. Exp. Biol.* 210: 4254-4261.

Costa, D. P. 2007. Seals and Sea Lions. in Encyclopedia of Tidepools. University of California Press. Berkeley.

Davis, R. W., Jaquet, N., Gendron, D., Markaida, U., Bazzino, G. and Gilly, W. F. 2007. Diving behavior of sperm whales in relation to the behavior of a major prey-species, the jumbo squid, in the Gulf of California, Mexico. *Mar. Ecol. Prog. Ser.* 333:291-302.

Gilly, W.F. 2007. Horizontal and vertical migrations of *Dosidicus gigas* in the Gulf of California revealed by electronic tagging, in Olson, R.J. and Young, J.W. (Eds.), *The role of squid in open ocean ecosystems*, **GLOBEC Report** 24:3-6.

Gilly, W.F. and Markaida, U. 2007. Perspectives on *Dosidicus gigas* in a changing world, in Olson, R.J. and Young, J.W. (Eds.), *The role of squid in open ocean ecosystems*, **GLOBEC Report** 24: 81-90.

Hassrick, J. L., D. E. Crocker, R. L. Zeno, S. B. Blackwell, D. P. Costa, and B. J. Le Boeuf. 2007. Swimming speed and foraging strategies of northern elephant seals. *Deep-Sea Res. II* 54:369-383.

Kitagawa, T., Boustany, A., Farwell, C., Williams, T. D., Castleton, M., Block, B. A. 2007. Horizontal and vertical movement of bluefin tuna, *Thunnus thynnus orientalis*, in relationship to oceanography. *Fish. Oceanogr.* 16, 409-415.

Markaida, U., Rosas, R., Salinas, C. and Gilly, W. 2007. Trophic ecology of jumbo squid *Dosidicus gigas* in the Gulf of California and adjacent waters in Olson, R.J. and Young, J.W. (Eds.), *The role of squid in open ocean ecosystems*, **GLOBEC Report** 24: 3-6.

Peckham, S. H., D. Maldonado, A. Walli, G. Ruiz, W.J. Nichols and L. Crowder. 2007. Small-scale fisheries bycatch of Pacific loggerheads can rival that in large-scale oceanic fisheries. *PLoS Biology* **ONE** 2(10) 1:6.

Rasmussen, K., D.M. Palacios, J. Calambokidis, M. Saborio, L. Dalla-Rosa, E. Secchi, G. Steiger, J. Allen, and G. Stone. 2007. Southern Hemisphere humpback whales wintering off Central America: insights from water temperature into the longest mammalian migration. *Biol. Lett.* 3(3):302-305.

Robinson, P. W., Y. Tremblay, D. E. Crocker, M. A. Kappes, C. E. Kuhn, S. A. Shaffer, S. E. Simmons, and D. P. Costa. 2007. A comparison of indirect measures of feeding behaviour based on ARGOS tracking data. *Deep-Sea Res. II* 54:356-368.

Sato, K., Y. Watanuki, A. Takahashi, P. Miller, H. Tanaka, R. Kawabe, P. Ponganis, Y. Handrich, T. Akamatsu, Y. Watanabe, Y. Mitani, D. Costa, C. Bost, K. Aoki, M. Amano, P. Trathan, A. Shapiro,

- and Y. Naito. 2007. Stroke frequency, but not swimming speed, is related to body size in free-ranging seabirds, pinnipeds and cetaceans. *Proc. Roy. Soc. Lond. B* 274:471-477.
- Schaefer, K. M., Fuller, D. W and Block, B. A. 2007. Movements, behavior, and habitat utilization of yellowfin tuna (*Thunnus albacares*) in the northeastern Pacific Ocean, ascertained through archival tag data. *Mar. Biol.* 152 (3): 503-525.
- Simmons, SE, Crocker, DE, Kudela, RM, Costa, DP (2007) Linking Foraging Behaviour of the Northern Elephant seal with Oceanography and Bathymetry at Mesoscales. *Mar. Eco. Prog. Ser.* 346, 265-275.
- Sippel T. J., Davie P. S., Holdsworth J. C., Block B. A. (2007) Striped marlin (*Tetrapturus audax*) movements and habitat utilization during a summer and autumn in the Southwest Pacific Ocean. *Fish. Oceanogr.* 16: 459-472.
- Tremblay, Y., Roberts, A. J., and Costa, D. P. 2007. Fractal landscape method: an alternative approach to measuring area-restricted searching behavior. *J. Exp. Biol.* 210: 935-945.
- Weise, M. J. and D. P. Costa. 2007. Total body oxygen stores and physiological diving capacity of California sea lions as a function of sex and age. *J. Exp. Biol.* 210: 278-289.
- Weng, K.C., Boustany, A., Pyle, P., Anderson, S., Brown, A. and Block, B. A. 2007. Migration and habitat of white wharks (*Carcharodon carcharias*) in the Eastern Pacific Ocean. *Mar. Biol.* 152: 877-894.
- Weng, K. C., O'Sullivan, J., Lowe, C., Winkler, C., Dewar, H., Block, B. A. 2007. Movements, behavior and habitat preferences of juvenile white sharks in the eastern Pacific as revealed by electronic tags. *Mar. Eco. Prog. Ser.* 338: 211-224.

2008

- Bailey, H. R., Shillinger, G. L., Palacios, D. M., Bograd, S. J., Spotila, J. R., Wallace, B., Paladino, F. V., Eckert, S. A. and Block, B. A. 2008. Identifying and comparing phases of movement by leatherback turtles using state-space models, *J. Exp. Mar. Biol. Ecol.*, 356, 128-135.
- Burns, J. M., M. A. Hindell, C. J. A. Bradshaw, and D. P. Costa. 2008. Fine-scale habitat selection of crabeater seals as determined by diving behavior. *Deep Sea Research Part II: Topical Studies in Oceanography* 55:500-514.
- Costa, D. P., J. M. Klinck, E. E. Hofmann, M. S. Dinniman, and J. M. Burns. 2008. Upper ocean variability in West Antarctic Peninsula continental shelf waters as measured using instrumented seals. *Deep Sea Research Part II: Topical Studies in Oceanography* 55:323-337.
- McDonald, B. I., D. E. Crocker, J. M. Burns, and D. P. Costa. 2008. Body condition as an index of winter foraging success in crabeater seals (*Lobodon carcinophaga*). *Deep Sea Research Part II: Topical Studies in Oceanography* 55:515-522.
- Wilson, C., Villareal, T. A., Maximenko, N., Bograd, S. J., Montoya, J. P. and Schoenbaechler, C. A. 2008. Biological and physical forcings of late summer chlorophyll blooms at 30°N in the oligotrophic Pacific. *J. Mar. Syst.* 69, 164-176.
- Zeno, R. L., D. E. Crocker, D. L. Hassrick, S. G. Allen, and D. P. Costa. 2008. Development of foraging behavior in juvenile northern elephant seals. *Journal of Zoology (London)* 274:180-187.

Publications In Press

- Benoit-Bird, K. J., Gilly, W. F., Au, W. W. L. and Mate, B. 2008. Controlled and in situ target strengths of the jumbo squid *Dosidicus gigas* and identification of potential acoustic scattering sources. *J. Acoust. Soc. Am.*: In Press.
- Burger, A. E. and Shaffer S. A. 2008. The application of tracking and data-logging technology in research and conservation of seabirds. *Auk*, In Press.
- Charrassin, J.-B., Hindell, M. Rintoul, S.R., Roquet, F., Sokolov, S. Biuw, M., Costa, D., Boehme, L., Lovell, P., Coleman, R. Timmerman, R., Meijers, A., Meredith, M., Park, Y.-H., Bailleul, F., Tremblay, Y., Bost, C.-A., McMahon, C.R., Field, I.C., Fedak, M.A. and Guinet, C. in press. Southern Ocean frontal structure and sea ice formation rates revealed by elephant seals. Proceedings National Academy of Sciences.
- Hoskins, A.J., Dann, P. Ropert-Coudert, Y. Kato, A., Chiaradia, A. Costa, D.P. Arnould, J.P.Y. 2008. Foraging behaviour and habitat selection of the little penguin (*Eudyptula minor*) during early chick rearing. Marine Ecology Progress Series in press.
- Oleson, E. M., Calambokidis, J., Burgess, W. C., McDonald, M. A., LeDuc, C. A. and Hildebrand, J. A. 2008. Behavioral context of Northeast Pacific Blue Whale call production. *Mar. Eco. Prog. Ser.* In Press.
- Robinson, P.W., Villegas-Amtmann, S., Costa, D.P. in press. Field Validation of an Inexpensive Time-Depth Recorder. Marine Mammal Science.
- Saba, V. S., Shillinger, G. L., Swithenbank, A. M., Block, B. A., Spotila, J. R., Musick, .A., Paladino, F. V. An oceanographic context for the foraging ecology of eastern Pacific leatherback turtles: Consequences of ENSO and coastal gillnet fisheries. *Deep-Sea Res. I.* In Press.
- Shaffer, S. A. 2008. Albatross flight performance and energetics. In *Albatrosses: Their World, Their Ways* (De Roy, T., Jones, M., and Fitter, J. eds). Firefly Books, Ontario, Canada, In Press.
- Staaf, D.J., Camarillo-Coop, S., Haddock, S.H.D., Nyack, A.C., Payne, J., Salinas-Zavala, C.A., Seibel, B.A., Trueblood, L., Widmer, C. and Gilly, W. F. 2008. Natural egg mass deposition by the Humboldt squid (*Dosidicus gigas*) in the Gulf of California and characteristics of hatchlings and paralarvae. *U. Mar. Biol. Assn. U.K.*: In Press.
- Villegas-Amtmann, S, Costa, D. P, Tremblay, Y, Auriolles-Gamboa, D & Salazar, S. Multiple foraging strategies in a marine apex predator, the Galapagos sea lion. *Mar. Eco. Prog. Ser.* In Press.
- Wells, B. K., Field, J. C., Thayer, J. A., Grimes, C. B., Bograd, S. J., Sydeman, W. J., Schwing, F. B., and Hewitt, R. 2008. Untangling the relationship between climate, prey, and top predators in an ocean ecosystem, *Mar. Eco. Prog. Ser.* In Press.
- Shillinger, G. L., Palacios, D. M., Bailey, H. R., Bograd, S. J., Swithenbank, A., Gaspar, P., Wallace, B., Spotila, J. R., Paladino, F. V., Piedra, R., Eckert, S. A., and Block, B.A. Ocean currents shape the migration and dispersal of eastern Pacific leatherback turtles, *PLOS Biol.*

Dissertations

- Boustany, A. 2006. Migratory Movements, Population Structure and Environmental Preferences of Northern Bluefin Tuna Revealed through Electronic Tagging and Population Genetics. Ph.D. Dissertation Stanford University June 2006.

Kuhn, C.E. 2006. Measuring at Sea Feeding To Understand the Foraging Behavior of Pinnipeds. Ph.D. Dissertation University of California Santa Cruz June 2006.

Teo, S. L. H. 2006. The biology and oceanography of Atlantic bluefin tuna on their breeding grounds. Ph.D. Dissertation Stanford University June 2006.

Weise, M.J. 2006. Foraging ecology of California sea lion (*Zalophus californianus*): movement, diving and foraging behavior, and diving capacity. Ph.D. Dissertation University of California Santa Cruz June 2006.

Weng, K. 2006. Movements of Pacific pelagic sharks in relation their environment. Ph.D. Dissertation Stanford University 2006.

Simmons, S. E. Environmental and Individual Effects on The Foraging Success of an Apex Predator, The Northern Elephant Seal (*Mirounga angustirostris*). Ph.D. Dissertation University of California Santa Cruz June 2007.