



Binomial Distribution: Hypothesis Testing, Confidence Intervals (CI), and Reliability with Implementation in S-PLUS

by Joseph C. Collins

ARL-TR-5214

June 2010

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Aberdeen Proving Ground, MD 21005

ARL-TR-5214**June 2010**

Binomial Distribution: Hypothesis Testing, Confidence Intervals (CI), and Reliability with Implementation in S-PLUS

Joseph C. Collins

Survivability/Lethality Analysis Directorate, ARL

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) June 2010		2. REPORT TYPE Final		3. DATES COVERED (From - To) January to June 2009	
4. TITLE AND SUBTITLE Binomial Distribution: Hypothesis Testing, Confidence Intervals (CI), and Reliability with Implementation in S-PLUS				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Joseph C. Collins				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: RDRL-SLB-D Survivability/Lethality Analysis Directorate Aberdeen Proving Ground, MD 21005				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-5214	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT System requirements can be expressed in terms of an upper (lower) limit on the probability of failure (success). Statistical analysis of such binary (0/1, success/failure, go/no-go) data typically requires point and interval estimation and inference or hypothesis tests on the associated event probability. For identical independent trials, the proportion observed serves as an estimate of the event rate. Based on the method of Clopper-Pearson (CP) and the likelihood ratio (LR) technique properties of the binomial distribution are used to derive interval estimates, which are in turn used in inference. An application is the determination of sample size and maximum permissible number of failures (nf) required to establish a specific reliability (probability of success) with given probability (confidence).					
15. SUBJECT TERMS binomial reliability, probability, statistics					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 43	19a. NAME OF RESPONSIBLE PERSON Joseph C. Collins
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) (410) 278-6832

Contents

List of Figures	v
List of Tables	v
Summary	1
1. Binomial Distribution	3
1.1 Definitions and Properties	3
1.2 Implementation Issue	4
1.3 Beta Distribution	5
2. Hypothesis Testing	8
2.1 One Sided Upper	8
2.2 One Sided Lower	9
2.3 Two Sided.....	9
3. Confidence Intervals (CI)	10
3.1 Upper	10
3.2 Lower.....	10
3.3 Two Sided.....	11
3.4 Implementation.....	11
3.5 Conservative Coverage.....	11
4. The LR Approach	14
4.1 Hypothesis Tests.....	15
4.2 Confidence Intervals.....	16
4.3 Implementation.....	17
5. Reliability	18
5.1 Reliability Tables	18
5.2 Sample Size	20

6. References	22
Appendix A. S-PLUS Code	23
A.1 Library Functions	23
A.2 Upper-Tail Binomial Probabilities	23
A.3 Binomial Parameter Confidence Interval: CP	24
A.4 Binomial Reliability Table	24
A.5 Binomial Reliability Sample Size	24
A.6 Binomial LR	25
A.7 Binomial LR cdf Table	25
A.8 Binomial LR cdf Evaluation	26
A.9 Binomial LR Hypothesis Test	26
A.10 Binomial Parameter Confidence Interval: LR	26
Appendix B. Reliability Table	29
List of Symbols, Abbreviations, and Acronyms	34
Distribution List	35

List of Figures

Figure 1. Binomial cdf and quantile function (qf) examples with continuous interpolation.	7
Figure 2. Upper CI coverage.....	12
Figure 3. Lower CI coverage.	13
Figure 4. Two-sided CP CI coverage.....	14
Figure 5. Two-sided LR CI coverage.	17
Figure 6. Two-sided CI coverage comparison.	17

List of Tables

Table 1. LR cdf.	16
Table 2. CP cdf.	16

INTENTIONALLY LEFT BLANK.

Summary

The purpose of this report is to provide statisticians with the tools to solve certain reliability problems that arise in consultations with clients.

Statisticians often encounter clients seeking answers to questions such as, “I did an experiment 6 times with 3 successes and 3 failures, so the success rate is 50%, right?” A reasonable, useful, and informative response from the statistician would be, “yes, the estimate of success is 50%, but you can only state with 90% confidence that the true success rate lies between 15% and 85%. If you want to tighten that up, you need a larger sample.” The client asks, “how large?” and the conversation proceeds from there.

A client may pose a problem such as, “I need 90% confidence on 95% reliability. Can I do this with 10 runs?” The statistician can truthfully answer, no. You need 29 with no failures or 46 with 1 failure to assert 95% reliability with 90% confidence.” The client may respond, “well, what can I get with 1 failure in 20?” and so the consultation continues.

In fact, the solutions to both of these problems exploit properties of the binomial distribution. System requirements can be expressed in terms of an upper (lower) limit on the probability of failure (success). Statistical analysis of such binary (0/1, success/failure, go/no-go) data typically requires point and interval estimation and inference or hypothesis tests on the associated event probability. For identical independent trials, the proportion observed serves as an estimate of the event rate. Based on the method of Clopper-Pearson (CP) and the likelihood ratio (LR) technique properties of the binomial distribution are used to derive interval estimates, which are in turn used in inference. An application is the determination of sample size and maximum permissible number of failures (nf) required to establish a specific reliability (probability of success) with given probability (confidence).

The statistician needs to address such issues and answer these questions in real time. This report provides the necessary technology expressed in the S-PLUS / S / R statistical computing environment language family, implementations of which are available as both commercial and free open-source software.

INTENTIONALLY LEFT BLANK.

1. Binomial Distribution

1.1 Definitions and Properties

The random variable X is Bernoulli(p), or B_p , if $X \in \{0,1\}$ and $\Pr[X = 1] = p = 1 - \Pr[X = 0]$. The probability mass function, or discrete probability density function (pdf), is

$$f_p(x) = p^x (1 - p)^{1-x}. \quad (1)$$

The sum X of n independent and identically distributed (iid) B_p is Bernoulli(n, p), or $B_{n,p}$. Note that $B_p = B_{1,p}$. The binomial coefficient is $\binom{n}{k} = \frac{n!}{k!(n-k)!}$. Standard functions are defined for $k \in \{0,1,2, \dots, n\}$. The pdf is

$$f_{n,p}(x) = C(n, k) p^x (1 - p)^{1-x}. \quad (2)$$

The cumulative distribution function (cdf) is

$$F_{n,p}(k) = \Pr[X \leq k] = \sum_{j=0}^k f_{n,p}(j) \quad (3)$$

with $F_{n,p}(x) = 0$ if $x \leq 0$, and $F_{n,p}(x) = 1$ if $x \geq n$, and right-continuous extension to $\mathbb{R}$, as is customary for discrete distributions. The quantile function (qf) is the usual unique left-continuous pseudo-inverse of the cdf,

$$Q_{n,p}(u) = \inf \{ k: F_{n,p}(k) \geq u \} \quad (4)$$

It is also useful to have an upper-probability version of the cdf,

$$G_{n,p}(k) = \Pr[X \geq k] = \sum_{j=k}^n f_{n,p}(j). \quad (5)$$

So $G_{n,p}(k) = 1 - F_{n,p}(k)$ and $F_{n,p}(k) = 1 - G_{n,p}(k + 1)$. Note that $F_{n,p}(k)$ increases with k and decreases with n or p , and $G_{n,p}(k)$ decreases with k and increases with n or p .

Moments are

$$E[X^k] = \sum_{j=0}^n j^k f_{n,p}(j) \quad (6)$$

So $E[X] = np$, and $E[X^2] = np(1 - p) + n^2 p^2$, and $\text{Var}[X] = np(1 - p)$.

The log-likelihood function is

$$\begin{aligned}\mathcal{L} &= \sum_{j=0}^n (\log C(n, j) + X_j \log p + (n - X_j) \log(1 - p)) \\ &= \left(\sum_{j=0}^n \log C(n, j) \right) + k \log p + (n - k) \log(1 - p).\end{aligned}\tag{7}$$

The maximum likelihood estimator (MLE) of p is k/n .

1.2 Implementation Issue

Reasonable algorithms for $F_{n,p}(k)$ return the appropriate values for large n and small k but succumb to numeric underflow for large k .

For example, since $F_{n,p}(0) = (1 - p)^n$ and $F_{n,p}(n - 1) = 1 - p^n$, set $n = 100$ and $p = 0.6$ to get

$$F_{100,0.6}(0) = (1 - 0.6)^{100} = 0.4^{100} \cong 1.6 \times 10^{-40}\tag{8}$$

and

$$F_{100,0.6}(99) = 1 - 0.6^{100} \cong 1 - 6.5 \times 10^{-23}.\tag{9}$$

But with standard double precision resolution of about 16 decimal places, the result is $F_{100,0.6}(99) = 1$ exactly. The cure for lower tail probabilities too close to 1 is to use upper tail probabilities, which will be close to 0, and work with G instead of F , since $G_{n,p}(n) = p^n = 1 - F_{n,p}(n - 1)$, and so

$$G_{100,0.6}(100) = 0.6^{100} = 1 - F_{100,0.6}(99) = 1 - (1 - 0.6)^{100} \cong 6.5 \times 10^{-23}.\tag{10}$$

But naïve use of $G_{n,p}(k) = 1 - F_{n,p}(k - 1)$ directly will of course give $G_{100,0.6}(100) = 0$. To obtain a useful representation of $G(k)$ with k large in terms of an accurate algorithm for $F(k)$ with k small, let $X \sim B_{n,p}$ and $W = n - X$. Note that $F_W(k) = \Pr[W \leq k] = \Pr[n - X \leq k] = \Pr[X \geq n - k] = 1 - \Pr[X < n - k] = 1 - \Pr[X \leq n - k - 1] = 1 - F_X(n - k - 1)$. Then $G_X(k) = 1 - F_X(k - 1) = 1 - F_X(n - (n - k) - 1) = F_W(n - k)$, so

$$G_{n,p}(k) = \Pr[B_{n,p} \leq k] = \Pr[B_{n,1-p} \leq n - k] = F_{n,1-p}(n - k).\tag{11}$$

This gives $G_{100,0.6}(100) = F_{100,0.4}(0) = (1 - 0.4)^{100} = 0.6^{100} \cong 6.5 \times 10^{-23}$ as required.

1.3 Beta Distribution

The Gamma function, which has $\Gamma(n) = (n-1)!$ for $n = 1, 2, 3, \dots$, is given by

$$\Gamma(z) = \int_0^\infty u^{z-1} e^{-u} du. \quad (12)$$

The Beta function, which has $B(n, k) = (n-1)!(k-1)!/(n+k-1)!$ on positive integers, is

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} = \int_0^1 x^{a-1}(1-x)^{b-1} dx. \quad (13)$$

The Beta (a, b) distribution on $[0, 1]$, with $a > 0$ and $b > 0$, has pdf and cdf

$$\begin{aligned} f_{\text{Beta}(a,b)}(x) &= \frac{1}{B(a,b)} x^{a-1}(1-x)^{b-1} \\ F_{\text{Beta}(a,b)}(x) &= \int_0^x f_{\text{Beta}(a,b)}(u) du. \end{aligned} \quad (14)$$

The Binomial and Beta cdfs are related by $F_{n,p}(k-1) = F_{\text{Beta}(n-k+1,k)}(1-p)$, so

$$\begin{aligned} F_{n,p}(k) &= F_{\text{Beta}(n-k,k+1)}(1-p), \quad \text{or} \\ F_{n-k+1,1-p}(k-1) &= F_{\text{Beta}(n,k)}(p) \end{aligned} \quad (15)$$

and also

$$\begin{aligned} G_{n,p}(k) &= F_{\text{Beta}(k,n-k+1)}(p), \quad \text{or} \\ G_{n-k+1,p}(k) &= F_{\text{Beta}(n,k)}(p). \end{aligned} \quad (16)$$

To show $F_{n,p}(k-1) = F_{\text{Beta}(n-k+1,k)}(1-p)$, first note that $B(n-k+1, k) = (n-k)!(k-1)!/n!$, and also $F_{\text{Beta}(n,1)}(1-p) = n \int_0^1 x^{n-1} dx = (1-p)^n = f_{n,p}(0)$. Now, integrating by parts,

$$\begin{aligned}
F_{\text{Beta}(n-k+1,k)}(1-p) &= \frac{n!}{(n-k)!(k-n)!} \int_0^{1-p} x^{n-k}(1-x)^{k-1} dx \\
&= \frac{n!}{(n-k)!(k-n)!} \left[\frac{x^{n-k}(1-x)^{k-1}}{n-k+1} \Big|_0^{1-p} + \int_0^{1-p} \frac{(k-1)x^{n-k+1}(1-x)^{k-2}}{n-k+1} dx \right] \\
&= \frac{n! p^{k-1}(1-p)^{n-k+1}}{(n-k+1)!(k-1)!} + \frac{n!}{(k-2)!(n-k+1)!} \int_0^{1-p} x^{n-k+1}(1-x)^{k-2} dx \\
&= f_{n,p}(k-1) + \frac{1}{B(n-k+2, k-1)} \int_0^{1-p} x^{n-k+1}(1-x)^{k-2} dx \\
&= f_{n,p}(k-1) + F_{\text{Beta}(n-k+2, k-1)}(1-p) \\
&= f_{n,p}(k-1) + \dots + f_{n,p}(1) + F_{\text{Beta}(n,1)}(1-p) \\
&= f_{n,p}(k-1) + \dots + f_{n,p}(1) + f_{n,p}(0) \\
&= F_{n,p}(k-1).
\end{aligned} \tag{17}$$

See, for example, Stuart (1).

In fact, the function

$$F(x) = \begin{cases} 0, & x \leq -1 \\ F_{\text{Beta}(n-x+1,x)}(1-p), & -1 < x < n \\ 1, & n \leq x \end{cases} \tag{18}$$

is a continuous cdf on $[-1, n]$, and $F(k) = F_{n,p}(k)$ for $k = 0, 1, 2, \dots, n$. So F serves as a continuous version, or interpolation, of $F_{n,p}$, which is sometimes useful. The corresponding quantile function $Q(u) = F^{-1}(u)$ is not related to Q_{Beta} , but can be obtained by solving $F(x) = u$ numerically to get $x = Q(u)$. See figure 1.

Note that Q_{Beta} does provide the solution p of $u = F_{n,p}(k)$. Since $u = F_{n,p}(k) = F_{\text{Beta}(n-k,k+1)}(1-p)$, and so $Q_{\text{Beta}(n-k,k+1)}(u) = 1-p$, observe that

$$p = 1 - Q_{\text{Beta}(n-k,k+1)}(u). \tag{19}$$

Likewise for $u = G_{(n,p)}(k)$. Since $u = G_{n,p}(k) = F_{\text{Beta}(k,n-k+1)}(p)$ and so $Q_{\text{Beta}(k,n-k+1)}(u) = p$, it follows that

$$p = Q_{\text{Beta}(k, n-k+1)}(u). \quad (20)$$

Library implementations of Q_{Beta} provide for efficient and accurate evaluation of p in these situations. This will be useful later.




Figure 1. Binomial cdf and quantile function (qf) examples with continuous interpolation.

2. Hypothesis Testing

Tests of size γ and Type I error probability $\alpha = 1 - \gamma$ for

$$H_0: p = p_o \quad (21)$$

are based on data X . Here α is the null rejection probability, or probability of rejecting H_0 when it is true. See Mood (2) or Stuart (3) for an explanation.

2.1 One Sided Upper

The one-sided alternative

$$H_1: p < p_o \quad (22)$$

provides an upper limit for p upon rejection of H_0 . The critical value is k_U , and the decision rule is to reject H_0 if $X \leq k_U$ where $\Pr[X \leq k_U \mid H_0] = \alpha$. Since X is discrete, take

$$k_U = \sup \{ k \mid \Pr[X \leq k \mid H_0] \leq \alpha \} = \sup \{ k \mid F_{n,p_o}(k) \leq \alpha \}. \quad (23)$$

The rejection region is $I_1 = \{ 0, \dots, k_U \}$, and the non-rejection region is $I_0 = \{ k_U + 1, \dots, n \}$. Equivalently, the p-value for the test is $p_X = F_{n,p_o}(X)$, and the rule is to reject H_0 if $p_X \leq \alpha$. For example, let $p_o = 0.3$ and $\alpha = 0.1$ with $n = 100$. Since $F_{100,0.3}(23) = 0.0755$ and $F_{100,0.3}(24) = 0.114$, it follows that $k_U = 23$. These are also the p-values, $p_{23} = 0.0755$ and $p_{24} = 0.114$. The true value of α for this test is 0.0755.

For example, it is possible for the test to degenerate due to insufficient sample size. Consider $p_o = 0.03$ and $\alpha = 0.01$ with $n = 100$. Since $F_{100,0.03}(0) = 0.0476$, it follows that $F_{100,0.03}(k) > \alpha$ for all k , and so $I_1 = \emptyset$, and $I_0 = \{ 0, \dots, n \}$, and rejection never occurs. The true value of α for this test is 0.

But $F_{151,0.03}(0) = 0.0101$ and $F_{152,0.03}(0) = 0.00976$, so $n > 151$ is required for a non-degenerate test. With $n = 152$, since $F_{152,0.03}(1) = 0.0556$, it follows that $k_U = 0$. Then if $X = 0$ the decision is to reject H_0 and conclude that $p < 0.03$ with probability ≥ 0.99 . If $X > 0$ there is no indication that $p < 0.03$. The true value of α for this test is 0.00976.

Let $p_o = 1/2$ and $\alpha = 7/128$ with $n = 10$. Note that $F_{10,1/2}(2) = 7/128 = \alpha$. It follows that $k_U = 2$ and $p_X \leq \alpha$ when $X \leq 2$, in which case H_0 is rejected. The true value of α for this test is exactly 7/128.

2.2 One Sided Lower

The one-sided alternative

$$H_1: p > p_o \quad (24)$$

provides a lower limit for p upon rejection of H_0 . The critical value is k_L , and one rejects H_0 if $X \geq k_L$ where $\Pr[X \geq k_L | H_0] = \alpha$. Since X is discrete, take

$$k_L = \inf\{k | \Pr[X \geq k | H_0] \leq \alpha\} = \inf\{k | G_{n,p_o}(k) \leq \alpha\}. \quad (25)$$

The rejection region is $I_1 = \{k_L, \dots, n\}$, and the non-rejection region is $I_0 = \{0, \dots, k_L - 1\}$. Equivalently, the p-value for the test is $p_X = G_{n,p_o}(X)$, and one rejects H_0 if $p_X \leq \alpha$.

For example, let $p_o = 0.3$ and $\alpha = 0.1$ with $n = 100$. Since $G_{100,0.3}(36) = 0.116$ and $G_{100,0.3}(37) = 0.0799$, it follows that $k_L = 37$. These are also the p-values, $p_{36} = 0.116$ and $p_{37} = 0.0799$.

2.3 Two Sided

The two-sided alternative

$$H_1: p \neq p_o \quad (26)$$

provides upper and lower limits for p upon rejection of H_0 . The critical values are k_U and k_L , and one rejects H_0 if $X \leq k_U$ or $X \geq k_L$ where $\Pr[X \leq k_U \text{ or } X \geq k_L | H_0] = \alpha$. For symmetry, set $\Pr[X \leq k_U | H_0] = \Pr[X \geq k_L | H_0] = \alpha/2$. Since X is discrete, take

$$\begin{aligned} k_U &= \sup\{k | \Pr[X \leq k | H_0] \leq \alpha/2\} = \sup\{k | F_{n,p_o}(k) \leq \alpha/2\} \\ k_L &= \inf\{k | \Pr[X \geq k | H_0] \leq \alpha/2\} = \inf\{k | G_{n,p_o}(k) \leq \alpha/2\} \end{aligned} \quad (27)$$

The rejection region is $I_1 = \{0, \dots, k_U\} \cup \{k_L, \dots, n\}$, and the non-rejection region is $I_0 = \{k_U + 1, \dots, k_L - 1\}$. Equivalently, the p-value for the test is $p_X = 2 \min(F_{n,p_o}(X), G_{n,p_o}(X))$. One rejects H_0 if $p_X \leq \alpha$.

For example, let $p_o = 0.3$ and $\alpha = 0.1$ with $n = 100$. Since $F_{100,0.3}(22) = 0.0479$ and $F_{100,0.3}(23) = 0.0755$, it follows that $k_U = 22$. Since $G_{100,0.3}(38) = 0.0530$ and $G_{100,0.3}(39) = 0.0340$, it follows that $k_L = 39$. Note that $G_{100,0.3}(22) = 0.971$, $G_{100,0.3}(23) = 0.952$, $F_{100,0.3}(38) = 0.966$, and $F_{100,0.3}(39) = 0.979$. So the p-values are $p_{22} = 0.0957$, $p_{23} = 0.151$, $p_{38} = 0.106$, and $p_{39} = 0.0680$.

3. Confidence Intervals (CI)

The point estimate of p is of course X/n . CIs on p are obtained by pivoting or inverting hypothesis test critical regions as follows. See Clopper and Pearson (4).

3.1 Upper

Let I_1 be the set of p_o for which H_0 would be rejected in favor of $H_1: p < p_o$ with Type I error α , so $I_1 = \{ p \mid F_{n,p}(X) \leq \alpha \} = [p_U, 1]$ where

$$\alpha = F_{n,p_U}(X) = F_{\text{Beta}(n-X, X+1)}(1 - p_U), \quad (28)$$

so p_U can be expressed by equation 19 as a Beta distribution quantile

$$p_U = 1 - Q_{\text{Beta}(n-X, X+1)}(\alpha). \quad (29)$$

Null rejection occurs with probability α , so the non-rejection region

$$I_0 = [0, p_U) = \{ p \mid F_{n,p}(X) > \alpha \} \quad (30)$$

is a 100 γ % upper CI on p , and p_U is an upper confidence limit on p . To reject H_0 when $p \in I_1 = [p_U, 1]$ is precisely the p-value decision rule in section 0.

3.2 Lower

Let I_1 be the set of p_o for which H_0 would be rejected in favor of $H_1: p > p_o$ with Type I error α , so $I_1 = \{ p \mid G_{n,p}(X) \leq \alpha \} = [0, p_L]$ where

$$\alpha = G_{n,p_L}(X) = F_{\text{Beta}(X, n-X+1)}(p_L), \quad (31)$$

so p_L can be expressed by equation (20) as a Beta distribution quantile

$$p_L = Q_{\text{Beta}(X, n-X+1)}(\alpha).. \quad (32)$$

Null rejection occurs with probability α , so the non-rejection region

$$I_0 = (p_L, 1] = \{ p \mid G_{n,p}(X) > \alpha \} \quad (33)$$

is a 100 γ % lower CI on p , and p_L is a lower confidence limit on p . To reject H_0 when $p \in I_1 = [0, p_L]$ is precisely the p-value decision rule in section 2.2.

3.3 Two Sided

Let I_1 be the set of p_o for which H_0 would be rejected in favor of $H_1: p \neq p_o$ with Type I error α , so $I_1 = \{p \mid G_{n,p}(X) \leq \alpha/2\} \cup \{p \mid F_{n,p}(X) \leq \alpha/2\} = [0, p_L] \cup [p_U, 1]$ where

$$G_{n,p_L}(X) = F_{n,p_U}(X) = \alpha/2. \quad (34)$$

Then p_L and p_U can be expressed by equations 19 and 20 as Beta distributions quantiles

$$\begin{aligned} p_L &= Q_{\text{Beta}(X, n-X+1)}(\alpha/2) \\ p_U &= 1 - Q_{\text{Beta}(n-X, X+1)}(\alpha/2). \end{aligned} \quad (35)$$

Null rejection occurs with probability α , so the non-rejection region

$$I_0 = (p_L, p_U) = \{p \mid G_{n,p}(X) > \alpha/2\} \cap \{p \mid F_{n,p}(X) > \alpha/2\} \quad (36)$$

is a 100% two-sided CI on p . To reject H_0 when $p_o \in [0, p_L] \cup [p_U, 1]$ is precisely the p-value decision rule of section 2.3.

3.4 Implementation

Code listings are in appendix A.

The function `binoc.i(n, r, g, type)` uses equations 29, 32, or 35 to give the required CI. The arguments of `binoc.i` are the number of trials n , the number of successes r , confidence interval size g which should be between 0.5 and 1.0, and an integer `type` which can be 0, 1, or 2 to request a lower CI $(L, 1]$, upper CI $[0, U)$, or 2-sided CI (L, U) , respectively.

The return value from `binoc.i` is a named list holding the number of trials `$n`, number of successes `$r`, lower limit `$L`, parameter estimate `$p.hat`, upper limit `$U`, and confidence interval vector `$I`. For example,

```
binoc.i(n=100, r=30, g=0.9, type=2)
```

returns

```
$n:      100
$r:       30
$L:      0.22492
$p.hat:   0.3
$U:      0.38422
$I:      0.22492 0.38422
```

since $\Pr[B_{100,0.22492} \geq 30] = \Pr[B_{100,0.38422} \leq 30] = 0.05$.

3.5 Conservative Coverage

Coverage is the probability that a confidence interval captures the true parameter value.

Considering that $p_U(X)$ is random, and $[0, p_U(X))$ is a $100\gamma\%$ CI on p , one expects that $C_U(p) = \Pr[p < p_U(X)] = E[I_{[0, p_U(X))}(p)] \geq \gamma$. This coverage probability is

$$C_U(p) = \sum_{k=0}^n \Pr[X = k] \cdot I_{[0, p_U(X))}(p) = \sum_{p < p_U(k)} f_{n,p}(k). \quad (37)$$

Figure 2 illustrates C_U for $n = 10$ and $\gamma = 0.9$.




Figure 2. Upper CI coverage.

Likewise, $p_L(X)$ is random, and $(p_L(X), 1]$ is a $100\gamma\%$ CI on p . One expects that $C_L(p) = \Pr[p_L(X) < p] = E[I_{(p_L(X), 1]}(p)] \geq \gamma$. This coverage probability is

$$C_L(p) = \sum_{k=0}^n \Pr[X = k] \cdot I_{(p_L(X), 1]}(p) = \sum_{p_L(k) < p} f_{n,p}(k). \quad (38)$$

Figure 3 illustrates C_L for $n = 10$ and $\gamma = 0.9$.




Figure 3. Lower CI coverage.

For the 2-sided CI, $(p_L(X), p_U(X))$ is a $100\gamma\%$ CI on p , and one expects that $C_2(p) = \Pr[p_L(X) < p < p_U(X)] = E[I_{(p_L(X), p_U(X))}(p)] \geq \gamma$. This coverage probability is

$$C_2(p) = \sum_{k=0}^n \Pr[X = k] \cdot I_{(p_L(X), p_U(X))}(p) = \sum_{p_L(k) < p < p_U(k)} f_{n,p}(k). \quad (39)$$

Figure 4 illustrates C_2 for $n = 10$ and $\gamma = 0.9$.




Figure 4. Two-sided CP CI coverage.

4. The LR Approach

For the parameter θ in a space Θ with some subset $\Theta_o \subseteq \Theta$, a test of

$$\begin{aligned} H_0: \theta &\in \Theta_o \\ H_1: \theta &\notin \Theta_o \end{aligned} \tag{40}$$

can be conducted using the generalized LR

$$\Lambda = \frac{\sup \{ \mathcal{L} \mid \theta \in \Theta_o \}}{\sup \{ \mathcal{L} \mid \theta \in \Theta \}}. \tag{41}$$

for the binomial distribution, the likelihood is

$$\mathcal{L} = C(n, k) p^k (1 - p)^{n-k} \tag{42}$$

and, based on the maximum likelihood estimate $p = k/n$, the unconstrained supremum is

$$\sup \{ \mathcal{L} \mid \theta \in \Theta \} = C(n, k) \left(\frac{k}{n} \right)^k \left(1 - \frac{k}{n} \right)^{n-k}. \tag{43}$$

For the specific null

$$H_0 : p = p_o \quad (44)$$

the conditional supremum is

$$\sup \{ \mathcal{L} \mid \theta \in \theta_o \} = \mathcal{L} = C(n, k) p_o^k (1 - p_o)^{n-k}. \quad (45)$$

So the LR is

$$\Lambda = \left(\frac{p_o}{k/n} \right)^k \left(\frac{1 - p_o}{1 - k/n} \right)^{n-k} = \left(\frac{np_o}{k} \right)^k \left(\frac{n - np_o}{n - k} \right)^{n-k}. \quad (46)$$

For data $X \sim B_{n,p}$, the LR is

$$\Lambda = \left(\frac{np_o}{X} \right)^X \left(\frac{n - np_o}{n - X} \right)^{n-X}. \quad (47)$$

4.1 Hypothesis Tests

Consider the two-sided test

$$\begin{aligned} H_0 : p &= p_o \\ H_1 : p &\neq p_o. \end{aligned} \quad (48)$$

Since Λ has small values significant, one rejects H_0 if $\Lambda \leq \Lambda_o$ for the appropriate critical value. Note that the null probabilities are $\Pr[\Lambda = \Lambda(k)] = \Pr[X = k] = f_{n,p_o}(k)$ and the $\Lambda(k)$ can be sorted in ascending order and the probabilities summed to obtain the cdf F_Λ

$$F_\Lambda(t) = \sum_{\Lambda(k) \leq t} f_{n,p_o}(k) = \sum_{k=0}^n I_{[\Lambda(k), 1]}(t) \cdot f_{n,p_o}(k). \quad (49)$$

In particular with $n = 10$ and $p_o = 0.33$, for $\alpha = 0.1$, the critical region $\Lambda \leq \Lambda(6)$ corresponds to $k \in \{0, 6, 7, 8, 9, 10\}$ and the non-rejection region is $k \in \{1, 2, 3, 4, 5\}$. See table 1.

Table 1. LR cdf.

k	$f_{n,p_o}(k)$	$t = \Lambda(k)$	$F_{\Lambda}(t)$
10	0.01823	0.00002	0.00002
9	0.08978	0.00080	0.00033
8	0.19899	0.00941	0.00317
0	0.26136	0.01823	0.02140
7	0.22528	0.05765	0.03678
6	0.13315	0.21789	0.09143
1	0.05465	0.23174	0.18121
5	0.01538	0.54106	0.31436
2	0.00284	0.65894	0.51335
4	0.00031	0.89817	0.73864
3	0.00002	0.97952	1.00000

Now, based on the usual CP procedure $\alpha/2$ upper and lower tails of the B_{n,p_o} test statistic, the critical region is $\{0,7,8,9,10\}$ and the non-rejection region is $\{1,2,3,4,5,6\}$. See table 2. Note that $X = 6$ is critical for the LR test but not for the CP test.

Table 2. CP cdf.

k	$F_{n,p_o}(k)$	$G_{n,p_o}(k)$
0	0.01823	1.00000
1	0.10801	0.98177
2	0.30700	0.89199
3	0.56837	0.69300
4	0.79365	0.43163
5	0.92680	0.20635
6	0.98145	0.07320
7	0.99683	0.01855
8	0.99967	0.00317
9	0.99998	0.00033
10	1.00000	0.00002

4.2 Confidence Intervals

As usual, confidence intervals are obtained by inverting non-rejection regions. One-sided LR CIs coincide with CP CIs. Coverage for the two-sided LR CIs is illustrated in figure 5. Two-sided LR CIs are in general not as conservative as CP CIs. This is regarded as an advantage of the LR method, as the intervals are “tighter”.




Figure 5. Two-sided LR CI coverage.




Figure 6. Two-sided CI coverage comparison.

4.3 Implementation

Code listings are in appendix A.

The function `binom.LR` evaluates the LR of equation 42 given n , k , and p . The function `binom.LR.F.x` evaluates the cdf components as in table 1, given n , p , and a technical

parameter `up` which sets the orientation for breaking ties. The function `bino.LR.F.z` evaluates $F_{\Lambda}(\Lambda(k))$. The hypothesis test of equation 48 is computed with `bino.LR.ht`, which presents the critical and non-rejection regions for the given values of n , p , and α . The function `bino.LR.ci` inverts the cdf to find CIs.

5. Reliability

5.1 Reliability Tables

Cooke (5) presents a tabulation of lower confidence limits p_L for various numbers of trials n and successes r and confidence levels γ .

This standard language may be misleading. The binomial parameter p is a reliability, which is a probability of success. The endpoints p_L and p_U of CIs $[p_L, p_U]$ on p are reliability limits, or limits of reliability, at the given confidence level γ . These “confidence limits” are not “limits of confidence.”

Code listings are in appendix A. The functions `pbinom` and `gbinom` evaluate $F_{n,p}(k)$ and $G_{n,p}(k)$, respectively.

The function `bino.rel.tab(n, r, g, type)` uses `bino.ci` to evaluate p_L and arranges the results in the proper format to reproduce Cooke’s tables, adding a column p for the parameter point estimate. The arguments of `bino.rel.tab` are the number of trials n , the number of successes r , confidence interval size g which should be between 0.5 and 1.0, and an integer `type` which can be 0, 1, or 2 to request a lower CI $(L, 1]$, upper CI $[0, U)$, or 2-sided CI (L, U) , respectively. The arguments r and g can be vectors, in which case the function fills out the desired table.

For example, the first table, with $n = 4$, is given by

```
bino.rel.tab(4, 4:2)
```

n	r	p	80%	90%	95%	97.5%	99%	99.5%
4	4	1.00	0.66874	0.56234	0.472871	0.397635	0.316228	0.265915
4	3	0.75	0.41755	0.32046	0.248605	0.194120	0.140868	0.110885
4	2	0.50	0.21232	0.14256	0.097611	0.067586	0.041999	0.029445

For $n = 18$,

```
bino.rel.tab(18, 18:9)
```

n	r	p	80%	90%	95%	97.5%	99%	99.5%
18	18	1.00000	0.91447	0.87992	0.84668	0.81470	0.77426	0.74501
18	17	0.94444	0.84262	0.80053	0.76234	0.72706	0.68398	0.65365
18	16	0.88889	0.77700	0.73058	0.68974	0.65288	0.60881	0.57833
18	15	0.83333	0.71472	0.66559	0.62332	0.58582	0.54170	0.51159
18	14	0.77778	0.65469	0.60398	0.56112	0.52363	0.48011	0.45076

18	13	0.72222	0.59642	0.54498	0.50217	0.46520	0.42280	0.39452
18	12	0.66667	0.53962	0.48816	0.44595	0.40993	0.36909	0.34214
18	11	0.61111	0.48412	0.43328	0.39216	0.35745	0.31858	0.29318
18	10	0.55556	0.42982	0.38020	0.34060	0.30757	0.27101	0.24739
18	9	0.50000	0.37669	0.32885	0.29120	0.26019	0.22630	0.20465

Tables for $n = 4 - 50$ are in appendix B. Compare to Cooke.

For $n = 100$,

```
bino.rel.tab(100,c(100:90, 60:50))
```

n	r	p	80%	90%	95%	97.5%	99%	99.5%
100	100	1.00	0.98403	0.97724	0.97049	0.96378	0.95499	0.94840
100	99	0.99	0.97035	0.96166	0.95344	0.94554	0.93546	0.92804
100	98	0.98	0.95770	0.94765	0.93838	0.92962	0.91859	0.91057
100	97	0.97	0.94554	0.93441	0.92429	0.91482	0.90303	0.89452
100	96	0.96	0.93370	0.92165	0.91080	0.90074	0.88830	0.87937
100	95	0.95	0.92209	0.90923	0.89775	0.88717	0.87415	0.86486
100	94	0.94	0.91064	0.89706	0.88501	0.87397	0.86045	0.85083
100	93	0.93	0.89933	0.88510	0.87254	0.86108	0.84710	0.83720
100	92	0.92	0.88814	0.87330	0.86028	0.84844	0.83405	0.82389
100	91	0.91	0.87703	0.86165	0.84820	0.83602	0.82125	0.81085
100	90	0.90	0.86601	0.85012	0.83628	0.82378	0.80867	0.79805
...								
100	60	0.60	0.55309	0.53115	0.51298	0.49721	0.47890	0.46647
100	59	0.59	0.54302	0.52106	0.50289	0.48714	0.46888	0.45648
100	58	0.58	0.53297	0.51100	0.49284	0.47712	0.45890	0.44655
100	57	0.57	0.52293	0.50096	0.48282	0.46713	0.44897	0.43666
100	56	0.56	0.51291	0.49095	0.47284	0.45719	0.43908	0.42682
100	55	0.55	0.50291	0.48096	0.46289	0.44728	0.42924	0.41704
100	54	0.54	0.49292	0.47100	0.45297	0.43741	0.41944	0.40730
100	53	0.53	0.48295	0.46107	0.44309	0.42758	0.40969	0.39762
100	52	0.52	0.47300	0.45116	0.43323	0.41779	0.39999	0.38798
100	51	0.51	0.46306	0.44128	0.42341	0.40804	0.39033	0.37840
100	50	0.50	0.45314	0.43142	0.41362	0.39832	0.38072	0.36886

For $n = 500$,

```
bino.rel.tab(500,c(500:490, 260:250))
```

n	r	p	80%	90%	95%	97.5%	99%	99.5%
500	500	1.000	0.99679	0.99541	0.99403	0.99265	0.99083	0.98946
500	499	0.998	0.99402	0.99224	0.99055	0.98891	0.98680	0.98523
500	498	0.996	0.99146	0.98939	0.98746	0.98563	0.98330	0.98159
500	497	0.994	0.98900	0.98669	0.98457	0.98257	0.98005	0.97822
500	496	0.992	0.98659	0.98408	0.98179	0.97964	0.97697	0.97503
500	495	0.990	0.98423	0.98153	0.97909	0.97682	0.97399	0.97196
500	494	0.988	0.98191	0.97903	0.97645	0.97407	0.97111	0.96898
500	493	0.986	0.97960	0.97657	0.97387	0.97137	0.96829	0.96608
500	492	0.984	0.97732	0.97414	0.97132	0.96872	0.96552	0.96324
500	491	0.982	0.97505	0.97174	0.96880	0.96611	0.96280	0.96044
500	490	0.980	0.97280	0.96935	0.96631	0.96353	0.96012	0.95769
...								
500	260	0.520	0.50018	0.49035	0.48223	0.47520	0.46704	0.46148
500	259	0.518	0.49818	0.48835	0.48024	0.47321	0.46504	0.45949

500	258	0.516	0.49618	0.48635	0.47824	0.47121	0.46305	0.45751
500	257	0.514	0.49418	0.48435	0.47624	0.46922	0.46106	0.45552
500	256	0.512	0.49218	0.48236	0.47425	0.46723	0.45907	0.45353
500	255	0.510	0.49019	0.48036	0.47226	0.46523	0.45709	0.45155
500	254	0.508	0.48819	0.47836	0.47026	0.46324	0.45510	0.44956
500	253	0.506	0.48619	0.47637	0.46827	0.46125	0.45311	0.44758
500	252	0.504	0.48419	0.47437	0.46628	0.45926	0.45112	0.44559
500	251	0.502	0.48220	0.47238	0.46428	0.45727	0.44914	0.44361
500	250	0.500	0.48020	0.47038	0.46229	0.45529	0.44716	0.44163

5.2 Sample Size

One application of this is to find the minimal sample size N to obtain a lower reliability limit at least p_L with confidence γ for a certain $n_f = n - r$, so $r = n - n_f$.

For example, with $n_f = 0$ failures ($r = n$) and 90% reliability limit $p_L = 0.9$ with 95% confidence $\gamma = 0.95$, one can wade through Cooke's phone-book-sized report, or run `bino.rel.tab` to find suitable n , which bound the desired reliability limit,

`bino.rel.tab(n=28, r=28, g=0.95)` gives $p_L = 0.89853$ and
`bino.rel.tab(n=29, r=29, g=0.95)` gives $p_L = 0.90186$, so $N \geq 29$.

For $n_f = 1$ failure, $r = n - 1$

`bino.rel.tab(n=45, r=44, g=0.95)` gives $p_L = 0.89887$ and
`bino.rel.tab(n=46, r=45, g=0.95)` gives $p_L = 0.90098$, so $N \geq 46$.

By equation 31, this amounts to finding

$$N = \inf\{n \mid G_{n,p_L}(n - n_f) \leq 1 - \gamma\}. \quad (50)$$

For $n_f = 0$ failures, $p_L = 0.9$, and $\gamma = 0.95$,

$G_{28,0.9}(28) = \text{gbinom}(28, 28, 0.9) = 0.052335$ and
 $G_{29,0.9}(29) = \text{gbinom}(29, 29, 0.9) = 0.047101$, so $N \geq 29$.

For $n_f = 1$ failures, $p_L = 0.9$, and $\gamma = 0.95$,

$G_{45,0.9}(45) = \text{gbinom}(45, 45, 0.9) = 0.052368$ and
 $G_{46,0.9}(46) = \text{gbinom}(46, 46, 0.9) = 0.048004$, so $N \geq 46$.

By equation 11, this is equivalent to

$$N = \inf\{n \mid F_{n,1-p_L}(n_f) \leq 1 - \gamma\}. \quad (51)$$

For $n_f = 0$ failures, $p_L = 0.9$, and $\gamma = 0.95$,

$F_{28,0.9}(0) = \text{pbinom}(0, 28, 0.1) = 0.052335$ and
 $F_{29,0.9}(0) = \text{pbinom}(0, 29, 0.1) = 0.047101$, so $N \geq 29$.

For $n_f = 1$ failures, $p_L = 0.9$, and $\gamma = 0.95$,

$$F_{45,0.9}(1) = \text{pbinom}(1, 45, 0.1) = 0.052368 \text{ and} \\ F_{46,0.9}(1) = \text{pbinom}(1, 46, 0.1) = 0.048004, \text{ so } N \geq 46.$$

The function `binom.rel.size(nf, p, g, x)` takes arguments representing the `nf`, desired lower reliability limit `p`, confidence `g`, and an optional scale factor `x` for the numerical solution routine. It calculates the required sample size N and returns a named list holding

<code>\$par:</code>	parameters <code>nf</code> , <code>p</code> , and <code>g</code>
<code>\$n:</code>	required minimum sample size, N
<code>\$p:</code>	reliability limit for confidence <code>g</code> and sample size N
<code>\$g:</code>	confidence for reliability limit <code>p</code> and sample size N
<code>\$n1:</code>	the next lower sample size, $N - 1$
<code>\$p1:</code>	reliability limit for confidence <code>g</code> and sample size $N - 1$
<code>\$g1:</code>	confidence for reliability limit <code>p</code> and sample size $N - 1$

One expects that $\$p1 < p \leq \p and $\$g1 < g \leq \g .

For example,

```
binom.rel.size(nf = 1, p = 0.9, g = 0.95)
```

returns

```
$par$nf: 1
$par$p: 0.9
$par$g: 0.95
$n: 46
$p: 0.90098
$g: 0.952
$n1: 45
$p1: 0.89887
$g1: 0.94763
```

Consulting the tables for the sample sizes $\$n = 46$ and $\$n1 = 45$ and confidence $g = 0.95$ for the number of successes $r = \$n - n_f = 45$ and $r1 = \$n1 - n_f = 44$, one sees the bounding reliability limits $\$p = 0.90098$, and $\$p1 = 0.89887$.

6. References

1. Stuart, A.; Ord, J. K. *Kendall's Advanced Theory of Statistics: Distribution Theory*; Oxford University Press: New York, 1994.
2. Mood, A.; Graybill, F. A.; Boes, D. C. *Introduction to the Theory of Statistics. 3*; McGraw-Hill, New York, 1974.
3. Stuart, A.; Ord, J. K. *Kendall's Advanced Theory of Statistics: Classical Inference and Relationship*; Oxford University Press, New York, 1991.
4. Clopper, C. J.; Pearson, E. S. *The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial*. 1934, 4, *Biometrika*, Vol 26, pp 404–413.
5. Cooke, J. R.; Lee, M. T.; Vanderbeck, J. P. *Binomial Reliability Table (Lower Confidence Limits for the Binomial Distribution)*. China Lake, CA, Bureau of Naval Weapons, 1964.

Appendix A. S-PLUS Code

A.1 Library Functions

```
pbinom(q,  
      size = stop("no size arg"),  
      prob = stop("no prob arg"))
```

is the S-PLUS library implementation of the Binomial distribution cdf.

In the notation of this report,

$$\text{pbinom}(k, n, p) = F_{n,p}(k) = \Pr[B_{n,p} \leq k]$$

where, k is number of successes, n is the number of trials, and p is the probability of success.

```
qbeta(p,  
      shape1 = stop("no shape1 arg"),  
      shape2 = stop("no shape2 arg"))
```

is the S-PLUS library implementation of the Beta distribution qf.

In the notation of this report,

$$\text{qbeta}(u, a, b) = Q_{\text{Beta}(a,b)}(u) = \inf \{x \mid F_{\text{Beta}(a,b)}(x) \geq u\}$$

where, u is the desired probability, and a and b are the usual Beta distribution parameters.

A.2 Upper-Tail Binomial Probabilities

`gbinom` calculates upper tail binomial probabilities.

```
gbinom <- function(q,  
                  size = stop("no size arg"),  
                  prob = stop("no prob arg"))  
{  
  ##  
  ## Pr [ Binomial(size,prob) >= q ]  
  ##  
  ## underflow error  
  ## 1-pbinom(q-1,size,prob)  
  ##  
  ## no underflow  
  ##  
  pbinom ( size - q , size , 1 - prob )  
}
```

In the notation of this report,

$$\text{gbinom}(k, n, p) = G_{n,p}(k) = \text{pbinom}(n-k, n, 1-p)$$

where, k is number of successes, n is the number of trials, and p is the probability of success.

A.3 Binomial Parameter Confidence Interval: CP

`binom.ci` calculates a CP confidence interval on a binomial parameter estimate.

```
binom.ci <- function(n          ,          # sample size
                    r          ,          # number of successes
                    g = 0.8    ,          # CI size
                    type = 0    ,          # 0,1,2 for lower, upper, 2-sided
                    a = 1-g    )          # 1 - (CI size)
{
  ## 100g% Confidence Interval [ L , U ] on Binomial Parameter p
  ##
  ## type 0 : lower CI = ( L , 1 ] : Pr [ B(n,L) >= r ] = 1-g
  ## type 1 : upper CI = [ 0 , U ) : Pr [ B(n,U) <= r ] = 1-g
  ## type 2 : two-sided CI = (L,U) : Pr[B(n,L)>=r] = Pr[B(n,U)<=r]=(1-g)/2

  if(type==2) a <- a/2

  if ( type==1 || r==0 )
    L <- 0
  else
    L <- qbeta ( a , r , n-r+1 )

  if(type==0 || r==n)
    U <- 1
  else
    U <- 1 - qbeta ( a , n-r , r+1 )

  list(n=n, r=r, L=L, p.hat=r/n, U=U, I=c(L,U))
}
```

A.4 Binomial Reliability Table

`binom.rel.tab` tabulates binomial confidence limits.

```
binom.rel.tab <- function(n = 100 ,          # number of trials
                        r = n    ,          # number of successes
                        g = c(.8, .9, .95, .975, .99, .995), # confidence levels
                        type = 0    )          # 0,1 = lower, upper
{
  nr <- length(r)          # success vector length
  ng <- length(g)          # number of CI levels
  z <- matrix(NA, nc=ng, nr=nr) # reliability limits

  for( i in 1:nr ) for(j in 1:ng ) # compute reliabilities
    z[i,j] <- binom.ci(n,r[i],g[j],type)$I[type+1]

  z <- cbind(n, r, r/n, z)          # augmented table
  dimnames(z) <-                   # name the table columns
    list(NULL, c("n","r","p",paste(100*g,"%", sep="")))

  z                                # return table
}
```

A.5 Binomial Reliability Sample Size

`binom.rel.size` calculates sample size for reliability with a given confidence limit.

```

bino.rel.size <- function(nf = 0      , # number of failures
                          p = 0.9    , # reliability limit
                          g = 0.95   , # confidence level
                          x = 6      ) # log_2 ( search delta )
{
  ## N = inf { n : pbinom(nf, n, 1-p) } <= 1-g }

  a <- 1-g                                # confidence
  q <- 1-p                                # reliability

  f <- function(nf, n, q) list(n=n, a=pbinom(nf, n, q)) # ( n , a )

  rtn <- function(nf, p, g, z1, z)      # return list
    list(par=list(nf=nf, p=p, g=g),
          n=z$n, p=bino.ci(z$n, z$n-nf, g, 0)$L, g=1-z$a ,
          n1=z1$n, p1=bino.ci(z1$n, z1$n-nf, g, 0)$L, g1=1-z1$a)

  z <- f(nf, 2^x, q)                    # initial ( n , a ) guess
  while ( z$a < a ) z <- f(nf, z$n/2, q)
  while ( z$a > a ) z <- f(nf, z$n*2, q)

  z0 <- z                                # upper end at n
  z1 <- f(nf, z0$n/2, q)                 # lower end at n/2
  dn <- (z0$n-z1$n)/2                    # interval radius
  while ( dn >= 1 ) {
    z <- f(nf, z1$n+dn, q)               # midpoint
    if ( z$a < a )                      # one midpoint becomes new endpoint
      z0 <- z else z1 <- z
    dn <- dn/2                          # new radius
  }

  rtn(nf, p, g, z1, z0)
}

```

The algorithm finds sample sizes $n_1 = 2^{k-1}$ and $n = 2^k$ that bound the confidence for the desired reliability limit. Then it repeatedly bisects the interval $[n_1, n]$ and replaces one endpoint with the midpoint, maintaining the bound, until $n - n_1 = 1$. Reliability limits at the desired confidence level should now bound the desired reliability limit. Using powers of 2 ensures that the sample sizes are integers.

A.6 Binomial LR

```

bino.LR <- function(k , # successes
                    n , # trials
                    p ) # parameter
{
  ## binomial LR
  ( n*p/k ) ^ k * ( n*(1-p)/(n-k) ) ^ (n-k)
}

```

A.7 Binomial LR cdf Table

```

bino.LR.F.x <- function(n , # trials
                        p , # null parameter
                        up = T ) # tie-breaker direction
{

```

```

## binomial(n,p) LR cdf: complete table
k <- 0:n                                # successes
f <- dbinom(k,n,p)                      # probabilities
L <- bino.LR(k,n,p)                     # LR values

if(up) I <- order(L,k)                   # order LR values, tiebreaker
else I <- order(L,-k)
L <- L[I]                                # sort LR values
k <- k[I]                                # order successes by LR value
F.L <- cumsum(f[I])                      # LR cdf
cbind(k, I, f, L, F.L)
}

```

A.8 Binomial LR cdf Evaluation

```

bino.LR.F.z <- function(k      ,      # trials
                        n      ,      # successes
                        p      ,      # null parameter
                        x=F )      # extended return
{
  ## binomial(n,p) LR cdf (k)
  z <- bino.LR.F.x(n=n, p=p, k>n*p)
  u <- order(z[, "I"])
  if(x) z[u[k+1],]
  else z[u[k+1], "F.L"]
}

```

A.9 Binomial LR Hypothesis Test

```

bino.LR.ht <- function(n      ,      # trials
                       p      ,      # null parameter
                       a = 0.1 )      # significance
{
  ## binomial LR hypothesis test
  z <- bino.LR.F.x(n=n, p=p)

  I.rej <- z[, "F.L"] <= a
  k.rej <- sort(z[, "k"][I.rej])
  k.acc <- sort(z[, "k"][!I.rej])
  list(z=z, k.acc=k.acc, k.rej=k.rej)
}

```

A.10 Binomial Parameter Confidence Interval: LR

```

bino.LR.ci <- function(n = 100 ,      # sample size
                       r = 50  ,      # number of successes
                       g = 0.9 )      # CI size
{
  h <- r/n
  q <- 1-g

  ## [0,h] : L is increasing
  if(r == 0)
    p0 <- list(neg=0, pos=0, f0=q, f1=q)
  else
    p0 <- uniroot(function(x, n., r., q.) bino.LR.F.z(k=r., n=n., p=x)-q.,
                  lower=0, upper=h, q.=q, n.=n, r.=r)
}

```

```

## [h,1] : L is decreasing
if(r == n)
  p1 <- list(neg=1, pos=1, f0=q, f1=q)
else
  p1 <- uniroot(function(x, n., r., q.) bino.LR.F.z(k=r., n=n., p=x)-q.,
                lower=h, upper=1, q.=q, n.=n, r.=r)
list(n=n, r=r, L=p0$pos, p.hat=h, U=p1$neg, I=c(p0$pos, p1$neg))
}

```

INTENTIONALLY LEFT BLANK.

Appendix B. Reliability Table

n	r	p	80%	90%	95%	97.5%	99%	99.5%
1	1	1.00000	0.20000	0.10000	0.05000	0.02500	0.01000	0.00500
1	0	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000	0.00000
n	r	p	80%	90%	95%	97.5%	99%	99.5%
2	2	1.00000	0.44721	0.31623	0.22361	0.15811	0.10000	0.07071
2	1	0.50000	0.10557	0.05132	0.02532	0.01258	0.00501	0.00250
n	r	p	80%	90%	95%	97.5%	99%	99.5%
3	3	1.00000	0.58480	0.46416	0.36840	0.29240	0.21544	0.17100
3	2	0.66667	0.28714	0.19580	0.13535	0.09430	0.05890	0.04140
3	1	0.33333	0.07168	0.03451	0.01695	0.00840	0.00334	0.00167
n	r	p	80%	90%	95%	97.5%	99%	99.5%
4	4	1.00000	0.66874	0.56234	0.47287	0.39764	0.31623	0.26591
4	3	0.75000	0.41755	0.32046	0.24860	0.19412	0.14087	0.11088
4	2	0.50000	0.21232	0.14256	0.09761	0.06759	0.04200	0.02944
n	r	p	80%	90%	95%	97.5%	99%	99.5%
5	5	1.00000	0.72478	0.63096	0.54928	0.47818	0.39811	0.34657
5	4	0.80000	0.50981	0.41611	0.34259	0.28358	0.22207	0.18510
5	3	0.60000	0.32660	0.24664	0.18926	0.14663	0.10564	0.08283
5	2	0.40000	0.16861	0.11223	0.07644	0.05274	0.03268	0.02288
n	r	p	80%	90%	95%	97.5%	99%	99.5%
6	6	1.00000	0.76472	0.68129	0.60696	0.54074	0.46416	0.41352
6	5	0.83333	0.57755	0.48968	0.41820	0.35877	0.29431	0.25399
6	4	0.66667	0.41461	0.33319	0.27134	0.22278	0.17307	0.14360
6	3	0.50000	0.26865	0.20091	0.15316	0.11812	0.08473	0.06628
n	r	p	80%	90%	95%	97.5%	99%	99.5%
7	7	1.00000	0.79460	0.71969	0.65184	0.59038	0.51795	0.46912
7	6	0.85714	0.62914	0.54744	0.47930	0.42128	0.35664	0.31509
7	5	0.71429	0.48324	0.40382	0.34126	0.29042	0.23632	0.20297
7	4	0.57143	0.35009	0.27860	0.22532	0.18405	0.14227	0.11770
7	3	0.42857	0.22833	0.16964	0.12876	0.09899	0.07080	0.05530
n	r	p	80%	90%	95%	97.5%	99%	99.5%
8	8	1.00000	0.81777	0.74989	0.68766	0.63058	0.56234	0.51567
8	7	0.87500	0.66963	0.59375	0.52932	0.47349	0.41006	0.36848
8	6	0.75000	0.53790	0.46178	0.40031	0.34914	0.29323	0.25783
8	5	0.62500	0.41634	0.34462	0.28924	0.24486	0.19820	0.16970
8	4	0.50000	0.30323	0.23966	0.19290	0.15701	0.12095	0.09987
n	r	p	80%	90%	95%	97.5%	99%	99.5%
9	9	1.00000	0.83625	0.77426	0.71687	0.66373	0.59948	0.55505
9	8	0.88889	0.70223	0.63164	0.57086	0.51750	0.45597	0.41503
9	7	0.77778	0.58232	0.50992	0.45036	0.39991	0.34369	0.30739
9	6	0.66667	0.47086	0.40058	0.34494	0.29930	0.25003	0.21914
9	5	0.55556	0.36609	0.30097	0.25137	0.21201	0.17097	0.14606
9	4	0.44444	0.26755	0.21040	0.16875	0.13700	0.10526	0.08679
n	r	p	80%	90%	95%	97.5%	99%	99.5%
10	10	1.00000	0.85134	0.79433	0.74113	0.69150	0.63096	0.58870
10	9	0.90000	0.72901	0.66315	0.60584	0.55498	0.49565	0.45571
10	8	0.80000	0.61906	0.55040	0.49310	0.44390	0.38826	0.35180
10	7	0.70000	0.51634	0.44827	0.39338	0.34755	0.29712	0.26489
10	6	0.60000	0.41913	0.35422	0.30354	0.26238	0.21834	0.19092
10	5	0.50000	0.32683	0.26732	0.22244	0.18709	0.15044	0.12831
n	r	p	80%	90%	95%	97.5%	99%	99.5%
11	11	1.00000	0.86389	0.81113	0.76160	0.71509	0.65793	0.61775
11	10	0.90909	0.75140	0.68976	0.63564	0.58722	0.53018	0.49144
11	9	0.81818	0.64993	0.58484	0.52991	0.48224	0.42768	0.39151
11	8	0.72727	0.55478	0.48924	0.43563	0.39026	0.33958	0.30672
11	7	0.63636	0.46431	0.40053	0.34981	0.30790	0.26220	0.23320
11	6	0.54545	0.37787	0.31772	0.27125	0.23379	0.19398	0.16931
11	5	0.45455	0.29526	0.24053	0.19958	0.16749	0.13439	0.11447
n	r	p	80%	90%	95%	97.5%	99%	99.5%
12	12	1.00000	0.87449	0.82540	0.77908	0.73535	0.68129	0.64305
12	11	0.91667	0.77038	0.71250	0.66132	0.61520	0.56046	0.52297
12	10	0.83333	0.67622	0.61448	0.56189	0.51586	0.46266	0.42705
12	9	0.75000	0.58765	0.52473	0.47267	0.42814	0.37781	0.34478
12	8	0.66667	0.50315	0.44100	0.39086	0.34888	0.30240	0.27248
n	r	p	80%	90%	95%	97.5%	99%	99.5%
13	13	1.00000	0.88355	0.83768	0.79418	0.75295	0.70170	0.66527
13	12	0.92308	0.78668	0.73216	0.68366	0.63970	0.58717	0.55098
13	11	0.84615	0.69886	0.64022	0.58990	0.54553	0.49383	0.45896
13	10	0.76923	0.61606	0.55574	0.50535	0.46187	0.41224	0.37936
13	9	0.69231	0.53686	0.47657	0.42738	0.38574	0.33910	0.30872
13	8	0.61538	0.46061	0.40176	0.35480	0.31578	0.27289	0.24543
13	7	0.53846	0.38700	0.33086	0.28705	0.25135	0.21288	0.18870
13	6	0.46154	0.31596	0.26373	0.22396	0.19223	0.15882	0.13827
n	r	p	80%	90%	95%	97.5%	99%	99.5%
14	14	1.00000	0.89140	0.84834	0.80736	0.76836	0.71969	0.68492
14	13	0.92857	0.80083	0.74933	0.70327	0.66132	0.61090	0.57597
14	12	0.85714	0.71856	0.66279	0.61461	0.57187	0.52174	0.48769
14	11	0.78571	0.64085	0.58302	0.53434	0.49202	0.44333	0.41082
14	10	0.71429	0.56636	0.50803	0.45999	0.41896	0.37257	0.34206
14	9	0.64286	0.49446	0.43689	0.39041	0.35138	0.30797	0.27986
14	8	0.57143	0.42486	0.36913	0.32503	0.28861	0.24880	0.22343
14	7	0.50000	0.35741	0.30455	0.26358	0.23036	0.19472	0.17240
n	r	p	80%	90%	95%	97.5%	99%	99.5%
15	15	1.00000	0.89826	0.85770	0.81896	0.78198	0.73564	0.70242
15	14	0.93333	0.81322	0.76443	0.72060	0.68052	0.63211	0.59841
15	13	0.86667	0.73585	0.68271	0.63656	0.59540	0.54683	0.51367
15	12	0.80000	0.66265	0.60721	0.56022	0.51911	0.47149	0.43947
15	11	0.73333	0.59237	0.53603	0.48925	0.44900	0.40311	0.37269
15	10	0.66667	0.52441	0.46829	0.42256	0.38380	0.34029	0.31184
15	9	0.60000	0.45846	0.40353	0.35957	0.32287	0.28229	0.25613
15	8	0.53333	0.39436	0.34152	0.29999	0.26586	0.22873	0.20514
15	7	0.46667	0.33207	0.28218	0.24373	0.21267	0.17946	0.15873
n	r	p	80%	90%	95%	97.5%	99%	99.5%
16	16	1.00000	0.90430	0.86596	0.82925	0.79409	0.74989	0.71810
16	15	0.93750	0.82417	0.77783	0.73604	0.69768	0.65116	0.61864
16	14	0.87500	0.75115	0.70044	0.65617	0.61652	0.56951	0.53724
16	13	0.81250	0.68198	0.62878	0.58343	0.54354	0.49706	0.46564
16	12	0.75000	0.61548	0.56108	0.51560	0.47623	0.43103	0.40087
16	11	0.68750	0.55107	0.49649	0.45165	0.41338	0.37005	0.34151
16	10	0.62500	0.48844	0.43456	0.39101	0.35435	0.31341	0.28677
16	9	0.56250	0.42745	0.37504	0.33337	0.29878	0.26069	0.23623
16	8	0.50000	0.36801	0.31783	0.27860	0.24651	0.21172	0.18969
n	r	p	80%	90%	95%	97.5%	99%	99.5%
17	17	1.00000	0.90967	0.87333	0.83843	0.80494	0.76270	0.73223
17	16	0.94118	0.83390	0.78979	0.74988	0.71311	0.66837	0.63697
17	15	0.88235	0.76478	0.71630	0.67381	0.63559	0.59008	0.55871
17	14	0.82353	0.69923	0.64813	0.60436	0.56568	0.52038	0.48960
17	13	0.76471	0.63613	0.58361	0.53945	0.50101	0.45661	0.42682
17	12	0.70588	0.57493	0.52193	0.47808	0.44042	0.39749	0.36901
17	11	0.64706	0.51535	0.46265	0.41971	0.38328	0.34229	0.31541
17	10	0.58824	0.45722	0.40551	0.36401	0.32925	0.29062	0.26558
17	9	0.52941	0.40044	0.35039	0.31083	0.27812	0.24225	0.21928
17	8	0.47059	0.34500	0.29726	0.26011	0.22983	0.19711	0.17644
n	r	p	80%	90%	95%	97.5%	99%	99.5%
18	18	1.00000	0.91447	0.87992	0.84668	0.81470	0.77426	0.74501
18	17	0.94444	0.84262	0.80053	0.76234	0.72706	0.68398	0.65365
18	16	0.88889	0.77700	0.73058	0.68974	0.65288	0.60881	0.57833
18	15	0.83333	0.71472	0.66559	0.62332	0.58582	0.54170	0.51159
18	14	0.77778	0.65469	0.60398	0.56112	0.52363	0.48011	0.45076
18	13	0.72222	0.59642	0.54498	0.50217	0.46520	0.42280	0.39452
18	12	0.66667	0.53962	0.48816	0.44595	0.40993	0.36909	0.34214
18	11	0.61111	0.48412	0.43328	0.39216	0.35745	0.31858	0.29318
18	10	0.55556	0.42982	0.38020	0.34060	0.30757	0.27101	0.24739
18	9	0.50000	0.37669	0.32885	0.29120	0.26019	0.22630	0.20465
n	r	p	80%	90%	95%	97.5%	99%	99.5%
19	19	1.00000	0.91878	0.88587	0.85413	0.82353	0.78476	0.75665
19	18	0.94737	0.85046	0.81023	0.77363	0.73972	0.69820	0.66889
19	17	0.89474	0.78802	0.74349	0.70420	0.66862	0.62595	0.59632
19	16	0.84211	0.72869	0.68141				

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
19	11	0.57895	0.45654	0.40754	0.36812	0.33500	0.29805	0.27399	
19	10	0.52632	0.40558	0.35793	0.32009	0.28864	0.25395	0.23160	
19	9	0.47368	0.35564	0.30983	0.27395	0.24447	0.21235	0.19189	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
20	20	1.00000	0.92268	0.89125	0.86089	0.83157	0.79433	0.76727	
20	19	0.95000	0.85757	0.81904	0.78389	0.75127	0.71121	0.68286	
20	18	0.90000	0.79800	0.75523	0.71738	0.68302	0.64166	0.61287	
20	17	0.85000	0.74136	0.69581	0.65634	0.62107	0.57927	0.55053	
20	16	0.80000	0.68670	0.63934	0.59897	0.56339	0.52172	0.49339	
20	15	0.75000	0.63354	0.58511	0.54442	0.50895	0.46789	0.44024	
20	14	0.70000	0.58162	0.53273	0.49218	0.45721	0.41714	0.39039	
20	13	0.65000	0.53078	0.48197	0.44197	0.40781	0.36906	0.34343	
20	12	0.60000	0.48093	0.43267	0.39358	0.36054	0.32342	0.29909	
20	11	0.55000	0.43200	0.38475	0.34693	0.31528	0.28008	0.25723	
20	10	0.50000	0.38397	0.33817	0.30195	0.27196	0.23896	0.21775	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
21	21	1.00000	0.92622	0.89615	0.86705	0.83890	0.80309	0.77701	
21	20	0.95238	0.86403	0.82706	0.79327	0.76184	0.72316	0.69571	
21	19	0.90476	0.80709	0.76595	0.72945	0.69623	0.65614	0.62815	
21	18	0.85714	0.75291	0.70898	0.67079	0.63658	0.59589	0.56784	
21	17	0.80952	0.70059	0.65478	0.61559	0.58093	0.54021	0.51242	
21	16	0.76190	0.64967	0.60267	0.56302	0.52834	0.48802	0.46076	
21	15	0.71429	0.59991	0.55229	0.51261	0.47825	0.43869	0.41217	
21	14	0.66667	0.55114	0.50339	0.46406	0.43032	0.39185	0.36627	
21	13	0.61905	0.50327	0.45584	0.41720	0.38435	0.34724	0.32277	
21	12	0.57143	0.45623	0.40954	0.37190	0.34021	0.30472	0.28153	
21	11	0.52381	0.41000	0.36443	0.32811	0.29781	0.26421	0.24245	
21	10	0.47619	0.36457	0.32051	0.28580	0.25713	0.22567	0.20549	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
22	22	1.00000	0.92946	0.90063	0.87269	0.84563	0.81113	0.78597	
22	21	0.95455	0.86993	0.83441	0.80188	0.77156	0.73416	0.70757	
22	20	0.90909	0.81539	0.77578	0.74053	0.70839	0.66950	0.64229	
22	19	0.86364	0.76348	0.72106	0.68409	0.65088	0.61127	0.58389	
22	18	0.81818	0.71331	0.66896	0.63091	0.59715	0.55737	0.53013	
22	17	0.77273	0.66445	0.61883	0.58020	0.54630	0.50674	0.47990	
22	16	0.72727	0.61667	0.57030	0.53151	0.49778	0.45879	0.43256	
22	15	0.68182	0.56982	0.52316	0.48454	0.45128	0.41316	0.38771	
22	14	0.63636	0.52379	0.47725	0.43913	0.40658	0.36960	0.34510	
22	13	0.59091	0.47853	0.43248	0.39516	0.36355	0.32795	0.30456	
22	12	0.54545	0.43399	0.38881	0.35254	0.32210	0.28813	0.26598	
22	11	0.50000	0.39017	0.34619	0.31126	0.28221	0.25008	0.22932	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
23	23	1.00000	0.93242	0.90474	0.87788	0.85181	0.81855	0.79425	
23	22	0.95652	0.87534	0.84116	0.80980	0.78051	0.74433	0.71856	
23	21	0.91304	0.82302	0.78481	0.75075	0.71962	0.68188	0.65541	
23	20	0.86957	0.77318	0.73219	0.69636	0.66411	0.62555	0.59882	
23	19	0.82609	0.72499	0.68203	0.64507	0.61219	0.57332	0.54663	
23	18	0.78261	0.67804	0.63374	0.59610	0.56297	0.52419	0.49779	
23	17	0.73913	0.63210	0.58695	0.54902	0.51595	0.47758	0.45166	
23	16	0.69565	0.58703	0.54144	0.50356	0.47081	0.43313	0.40787	
23	15	0.65217	0.54271	0.49709	0.45954	0.42734	0.39060	0.36615	
23	14	0.60870	0.49910	0.45378	0.41685	0.38542	0.34985	0.32635	
23	13	0.56522	0.45616	0.41147	0.37539	0.34495	0.31076	0.28835	
23	12	0.52174	0.41386	0.37012	0.33515	0.30588	0.27329	0.25210	
23	11	0.47826	0.37219	0.32971	0.29609	0.26820	0.23742	0.21756	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
24	24	1.00000	0.93514	0.90852	0.88265	0.85753	0.82540	0.80191	
24	23	0.95833	0.88032	0.84738	0.81711	0.78880	0.75375	0.72875	
24	22	0.91667	0.83003	0.79315	0.76020	0.73003	0.69337	0.66761	
24	21	0.87500	0.78212	0.74246	0.70773	0.67639	0.63883	0.61274	
24	20	0.83333	0.73576	0.69412	0.65819	0.62616	0.58819	0.56205	
24	19	0.79167	0.69058	0.64754	0.61086	0.57849	0.54048	0.51454	
24	18	0.75000	0.64635	0.60237	0.56531	0.53289	0.49515	0.46959	
24	17	0.70833	0.60292	0.55840	0.52127	0.48905	0.45185	0.42683	
24	16	0.66667	0.56021	0.51551	0.47858	0.44678	0.41035	0.38601	
24	15	0.62500	0.51814	0.47359	0.43711	0.40594	0.37049	0.34698	
24	14	0.58333	0.47669	0.43258	0.39679	0.36643	0.33218	0.30961	
24	13	0.54167	0.43582	0.39245	0.35756	0.32821	0.29534	0.27383	
24	12	0.50000	0.39553	0.35317	0.31942	0.29124	0.25994	0.23962	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
25	25	1.00000	0.93765	0.91201	0.88707	0.86281	0.83176	0.80902	
25	24	0.96000	0.88491	0.85313	0.82388	0.79648	0.76251	0.73824	
25	23	0.92000	0.83652	0.80086	0.76896	0.73969	0.70407	0.67899	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
30	30	1.00000	0.94777	0.92612	0.90497	0.88430	0.85770	0.83811	
30	29	0.96667	0.90345	0.87643	0.85140	0.82783	0.79841	0.77725	
30	28	0.93333	0.86271	0.83219	0.80467	0.77926	0.74810	0.72600	
30	27	0.90000	0.82380	0.79070	0.76140	0.73471	0.70238	0.67969	
30	26	0.86667	0.78609	0.75101	0.72038	0.69278	0.65967	0.63662	
30	25	0.83333	0.74925	0.71264	0.68103	0.65279	0.61920	0.59597	
30	24	0.80000	0.71311	0.67531	0.64299	0.61433	0.58049	0.55724	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
25	22	0.88000	0.79038	0.75198	0.71828	0.68781	0.65122	0.62574	
25	21	0.84000	0.74573	0.70533	0.67039	0.63917	0.60208	0.57649	
25	20	0.80000	0.70219	0.66034	0.62459	0.59296	0.55573	0.53024	
25	19	0.76000	0.65954	0.61669	0.58048	0.54871	0.51163	0.48643	
25	18	0.72000	0.61765	0.57418	0.53779	0.50612	0.46944	0.44468	
25	17	0.68000	0.57643	0.53266	0.49636	0.46500	0.42894	0.40476	
25	16	0.64000	0.53581	0.49205	0.45607	0.42521	0.38997	0.36651	
25	15	0.60000	0.49576	0.45228	0.41684	0.38665	0.35244	0.32979	
25	14	0.56000	0.45625	0.41332	0.37862	0.34928	0.31626	0.29455	
25	13	0.52000	0.41725	0.37514	0.34139	0.31306	0.28141	0.26074	
25	12	0.48000	0.37878	0.33774	0.30513	0.27797	0.24786	0.22835	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
26	26	1.00000	0.93998	0.91525	0.89117	0.86773	0.83768	0.81564	
26	25	0.96154	0.88917	0.85847	0.83017	0.80363	0.77068	0.74709	
26	24	0.92308	0.84253	0.80803	0.77711	0.74870	0.71406	0.68962	
26	23	0.88462	0.79804	0.76083	0.72810	0.69846	0.66279	0.63791	
26	22	0.84615	0.75497	0.71575	0.68176	0.65132	0.61508	0.59002	
26	21	0.80769	0.71296	0.67226	0.63741	0.60649	0.57002	0.54500	
26	20	0.76923	0.67179	0.63003	0.59465	0.56352	0.52710	0.50228	
26	19	0.73077	0.63134	0.58888	0.55323	0.52213	0.48598	0.46152	
26	18	0.69231	0.59151	0.54866	0.51300	0.48210	0.44646	0.42248	
26	17	0.65385	0.55225	0.50929	0.47384	0.44333	0.40837	0.38500	
26	16	0.61538	0.51351	0.47071	0.43566	0.40571	0.37161	0.34896	
26	15	0.57692	0.47527	0.43286	0.39842	0.36918	0.33612	0.31429	
26	14	0.53846	0.43751	0.39574	0.36209	0.33371	0.30184	0.28094	
26	13	0.50000	0.40022	0.35932	0.32664	0.29927	0.26876	0.24888	

	n	r	p	80%	90%	95%	97.5%	99%	99.5%
27	27	1.00000	0.94213	0.91825	0.89498	0.87230	0.84319	0.82182	
27	26	0.96296	0.89312	0.86343	0.83603	0.81029	0.77830	0.75536	
27	25	0.92593	0.84811	0.81470	0.78470	0.75710	0.72339	0.69958	
27	24	0.88889	0.80516	0.76906	0.73726	0.70841	0.67363	0.64933	
27	23	0.85185	0.76357	0.72546	0.69237	0.66269	0.62727	0.60273	
27	22	0.81481	0.72298	0.68337	0.64938	0.61917	0.58344	0.55888	
27	21	0.77778	0.68319	0.64248	0.60790	0.57742	0.54164	0.51722	
27	20	0.74074	0.64408	0.					

30	23	0.76667	0.67756	0.63886	0.60605	0.57716	0.54327	0.52012
30	22	0.73333	0.64250	0.60316	0.57007	0.54111	0.50734	0.48440
30	21	0.70000	0.60790	0.56813	0.53493	0.50604	0.47256	0.44992
30	20	0.66667	0.57370	0.53372	0.50056	0.47188	0.43882	0.41658
30	19	0.63333	0.53988	0.49987	0.46691	0.43856	0.40605	0.38430
30	18	0.60000	0.50642	0.46657	0.43395	0.40603	0.37421	0.35302
30	17	0.56667	0.47330	0.43378	0.40163	0.37427	0.34325	0.32270
30	16	0.53333	0.44052	0.40149	0.36995	0.34326	0.31315	0.29331
30	15	0.50000	0.40807	0.36970	0.33889	0.31297	0.28390	0.26485

n	r	p	80%	90%	95%	97.5%	99%	99.5%
31	31	1.00000	0.94941	0.92841	0.90789	0.88781	0.86195	0.84289
31	30	0.96774	0.90647	0.88023	0.85591	0.83298	0.80433	0.78370
31	29	0.93548	0.86698	0.83731	0.81054	0.78578	0.75539	0.73381
31	28	0.90323	0.82925	0.79704	0.76850	0.74246	0.71088	0.68868
31	27	0.87097	0.79267	0.75850	0.72863	0.70166	0.66927	0.64669
31	26	0.83871	0.75694	0.72123	0.69036	0.66273	0.62981	0.60702
31	25	0.80645	0.72188	0.68496	0.65335	0.62527	0.59205	0.56919
31	24	0.77419	0.68737	0.64953	0.61739	0.58904	0.55571	0.53290
31	23	0.74194	0.65334	0.61481	0.58234	0.55387	0.52060	0.49795
31	22	0.70968	0.61973	0.58074	0.54810	0.51964	0.48658	0.46418
31	21	0.67742	0.58651	0.54724	0.51458	0.48627	0.45355	0.43149
31	20	0.64516	0.55365	0.51428	0.48175	0.45370	0.42144	0.39980
31	19	0.61290	0.52113	0.48183	0.44956	0.42187	0.39021	0.36906
31	18	0.58065	0.48892	0.44986	0.41797	0.39076	0.35980	0.33922
31	17	0.54839	0.45703	0.41835	0.38698	0.36034	0.33019	0.31025
31	16	0.51613	0.42545	0.38731	0.35657	0.33061	0.30138	0.28214
31	15	0.48387	0.39417	0.35673	0.32674	0.30155	0.27334	0.25488

n	r	p	80%	90%	95%	97.5%	99%	99.5%
32	32	1.00000	0.95095	0.93057	0.91063	0.89112	0.86596	0.84741
32	31	0.96875	0.90930	0.88380	0.86015	0.83783	0.80991	0.78980
32	30	0.93750	0.87098	0.84213	0.81606	0.79193	0.76227	0.74119
32	29	0.90625	0.83437	0.80301	0.77518	0.74977	0.71891	0.69719
32	28	0.87500	0.79887	0.76556	0.73640	0.71005	0.67835	0.65622
32	27	0.84375	0.76417	0.72933	0.69916	0.67212	0.63986	0.61749
32	26	0.81250	0.73012	0.69406	0.66313	0.63561	0.60300	0.58052
32	25	0.78125	0.69660	0.65959	0.62810	0.60027	0.56750	0.54504
32	24	0.75000	0.66354	0.62581	0.59394	0.56595	0.53318	0.51083
32	23	0.71875	0.63088	0.59263	0.56055	0.53253	0.49999	0.47775
32	22	0.68750	0.59858	0.56001	0.52786	0.49992	0.46756	0.44570
32	21	0.65625	0.56663	0.52790	0.49581	0.46807	0.43610	0.41460
32	20	0.62500	0.53499	0.49626	0.46436	0.43692	0.40546	0.38439
32	19	0.59375	0.50366	0.46508	0.43349	0.40645	0.37559	0.35503
32	18	0.56250	0.47262	0.43433	0.40317	0.37663	0.34649	0.32649
32	17	0.53125	0.44186	0.40401	0.37339	0.34744	0.31812	0.29875
32	16	0.50000	0.41139	0.37412	0.34415	0.31887	0.29047	0.27180

n	r	p	80%	90%	95%	97.5%	99%	99.5%
33	33	1.00000	0.95240	0.93260	0.91322	0.89424	0.86975	0.85167
33	32	0.96970	0.91196	0.88717	0.86415	0.84241	0.81519	0.79556
33	31	0.93939	0.87476	0.84667	0.82127	0.79774	0.76878	0.74817
33	30	0.90909	0.83919	0.80864	0.78150	0.75668	0.72651	0.70526
33	29	0.87879	0.80470	0.77222	0.74375	0.71798	0.68695	0.66526
33	28	0.84848	0.77099	0.73697	0.70748	0.68101	0.64938	0.62742
33	27	0.81818	0.73789	0.70265	0.67237	0.64540	0.61339	0.59129
33	26	0.78788	0.70530	0.66910	0.63824	0.61092	0.57870	0.55657
33	25	0.75758	0.67315	0.63620	0.60493	0.57741	0.54513	0.52308
33	24	0.72727	0.64139	0.60388	0.57235	0.54476	0.51257	0.49068
33	23	0.69697	0.60998	0.57209	0.54044	0.51289	0.48090	0.45925
33	22	0.66667	0.57889	0.54078	0.50914	0.48173	0.45007	0.42872
33	21	0.63636	0.54809	0.50993	0.47842	0.45125	0.42001	0.39904
33	20	0.60606	0.51759	0.47950	0.44823	0.42139	0.39069	0.37017
33	19	0.57576	0.48735	0.44948	0.41856	0.39215	0.36208	0.34206
33	18	0.54545	0.45739	0.41986	0.38940	0.36351	0.33416	0.31471
33	17	0.51515	0.42768	0.39064	0.36074	0.33544	0.30691	0.28809
33	16	0.48485	0.39824	0.36181	0.33258	0.30796	0.28034	0.26221

n	r	p	80%	90%	95%	97.5%	99%	99.5%
34	34	1.00000	0.95377	0.93452	0.91566	0.89718	0.87333	0.85570
34	33	0.97059	0.91448	0.89035	0.86793	0.84673	0.82018	0.80101
34	32	0.94118	0.87832	0.85095	0.82619	0.80323	0.77494	0.75479
34	31	0.91176	0.84374	0.81396	0.78747	0.76322	0.73372	0.71290
34	30	0.88235	0.81021	0.77851	0.75070	0.72550	0.69511	0.67384

n	r	p	80%	90%	95%	97.5%	99%	99.5%
38	36	0.94737	0.89074	0.86595	0.84344	0.82251	0.79663	0.77814
38	35	0.92105	0.85964	0.83257	0.80841	0.78623	0.75912	0.73993
38	34	0.89474	0.82944	0.80056	0.77510	0.75195	0.72391	0.70421
38	33	0.86842	0.79991	0.76954	0.74304	0.71914	0.69040	0.67033
38	32	0.84211	0.77089	0.73929	0.71196	0.68747	0.65821	0.63789
38	31	0.81579	0.74230	0.70968	0.68168	0.65674	0.62711	0.60663
38	30	0.78947	0.71406	0.68061	0.65209	0.62681	0.59694	0.57639
38	29	0.76316	0.68614	0.65202	0.62309	0.59759	0.56759	0.54703
38	28	0.73684	0.65850	0.62385	0.59463	0.56899	0.53897	0.51847
38	27	0.71053	0.63112	0.59606	0.56666	0.54097	0.51101	0.49063
38	26	0.68421	0.60397	0.56863	0.53914	0.51347	0.48367	0.46347
38	25	0.65789	0.57704	0.54153	0.51204	0.48647	0.45690	0.43693
38	24	0.63158	0.55032	0.51474	0.48534	0.45994	0.43068	0.41099
38	23	0.60526	0.52381	0.48826	0.45902	0.43386	0.40499	0.38562
38	22	0.57895	0.49748	0.46207	0.43307	0.40821	0.37979	0.36080

34	22	0.64706	0.56048	0.52289	0.49177	0.46489	0.43390	0.41306
34	21	0.61765	0.53076	0.49318	0.46225	0.43564	0.40512	0.38467
34	20	0.58824	0.50130	0.46386	0.43321	0.40697	0.37700	0.35700
34	19	0.55882	0.47209	0.43492	0.40466	0.37886	0.34953	0.33004
34	18	0.52941	0.44312	0.40635	0.37657	0.35129	0.32269	0.30377
34	17	0.50000	0.41440	0.37814	0.34894	0.32427	0.29649	0.27819

n	r	p	80%	90%	95%	97.5%	99%	99.5%
35	35	1.00000	0.95506	0.93633	0.91797	0.89997	0.87671	0.85952
35	34	0.97143	0.91685	0.89335	0.87150	0.85083	0.82491	0.80618
35	33	0.94286	0.88168	0.85501	0.83085	0.80843	0.78079	0.76108
35	32	0.91429	0.84805	0.81899	0.79312	0.76942	0.74055	0.72017
35	31	0.88571	0.81541	0.78447	0.75728	0.73262	0.70285	0.68200
35	30	0.85714	0.78350	0.75104	0.72282	0.69743	0.66701	0.64584
35	29	0.82857	0.75217	0.71847	0.68944	0.66350	0.63263	0.61127
35	28	0.80000	0.72130	0.68661	0.65695	0.63062	0.59947	0.57802
35	27	0.77143	0.69084	0.65536	0.62523	0.59864	0.56734	0.54589
35	26	0.74286	0.66073	0.62464	0.59418	0.56744	0.53613	0.51477
35	25	0.71429	0.63094	0.59439	0.56374	0.53696	0.50574	0.48453
35	24	0.68571	0.60144	0.56459	0.53385	0.50712	0.47611	0.45513
35	23	0.65714	0.57222	0.53519	0.50448	0.47789	0.44718	0.42648
35	22	0.62857	0.54324	0.50618	0.47560	0.44923	0.41892	0.39856
35	21	0.60000	0.51452	0.47753	0.44718	0.42112	0.39129	0.37133
35	20	0.57143	0.48602	0.44924	0.41920	0.39353	0.36427	0.34477
35	19	0.54286	0.45776	0.42130	0.39167	0.36646	0.33784	0.31886
35	18	0.51429	0.42973	0.39369	0.36457	0.33989	0.31201	0.29359
35	17	0.48571	0.40192	0.36643	0.33789	0.31383	0.28676	0.26895

n	r	p	80%	90%	95%	97.5%	99%	99.5%
36	36	1.00000	0.95628	0.93804	0.92015	0.90261	0.87992	0.86314
36	35	0.97222	0.91910	0.89620	0.87488	0.85471	0.82940	0.81109
36	34	0.94444	0.88486	0.85885	0.83527	0.81336	0.78633	0.76705
36	33	0.91667	0.85212	0.82375	0.79848	0.77531	0.74705	0.72708
36	32	0.88889	0.82034	0.79011	0.76352	0.73939	0.71022	0.68976
36	31	0.86111	0.78926	0.75753	0.72990	0.70503	0.67519	0.65439
36	30	0.83333	0.75874	0.72577	0.69732	0.67188	0.64157	0.62056
36	29	0.80556	0.72867	0.69470	0.66561	0.63975	0.60912	0.58800
36	28	0						

38 21 0.55263 0.47135 0.43616 0.40748 0.38299 0.35510 0.33652
38 20 0.52632 0.44540 0.41054 0.38224 0.35818 0.33088 0.31277
38 19 0.50000 0.41964 0.38519 0.35736 0.33379 0.30715 0.28954

n	r	p	80%	90%	95%	97.5%	99%	99.5%
39	39	1.00000	0.95957	0.94267	0.92606	0.90975	0.88862	0.87297
39	38	0.97436	0.92516	0.90389	0.88405	0.86524	0.84158	0.82444
39	37	0.94872	0.89345	0.86924	0.84723	0.82676	0.80142	0.78330
39	36	0.92308	0.86312	0.83666	0.81302	0.79130	0.76473	0.74591
39	35	0.89744	0.83366	0.80540	0.78048	0.75779	0.73029	0.71095
39	34	0.87179	0.80484	0.77511	0.74915	0.72570	0.69749	0.67777
39	33	0.84615	0.77652	0.74557	0.71876	0.69472	0.66597	0.64598
39	32	0.82051	0.74861	0.71664	0.68916	0.66465	0.63551	0.61534
39	31	0.79487	0.72105	0.68823	0.66021	0.63536	0.60594	0.58568
39	30	0.76923	0.69379	0.66029	0.63185	0.60674	0.57717	0.55688
39	29	0.74359	0.66680	0.63274	0.60400	0.57873	0.54909	0.52884
39	28	0.71795	0.64006	0.60557	0.57661	0.55126	0.52166	0.50150
39	27	0.69231	0.61354	0.57874	0.54966	0.52431	0.49482	0.47481
39	26	0.66667	0.58723	0.55223	0.52311	0.49783	0.46853	0.44871
39	25	0.64103	0.56113	0.52602	0.49695	0.47180	0.44276	0.42319
39	24	0.61538	0.53521	0.50009	0.47114	0.44619	0.41749	0.39821
39	23	0.58974	0.50948	0.47445	0.44569	0.42100	0.39270	0.37375
39	22	0.56410	0.48393	0.44907	0.42058	0.39621	0.36838	0.34981
39	21	0.53846	0.45855	0.42395	0.39581	0.37181	0.34452	0.32636
39	20	0.51282	0.43335	0.39910	0.37136	0.34780	0.32111	0.30341
39	19	0.48718	0.40832	0.37451	0.34725	0.32418	0.29815	0.28095

n	r	p	80%	90%	95%	97.5%	99%	99.5%
40	40	1.00000	0.96056	0.94406	0.92784	0.91190	0.89125	0.87594
40	39	0.97500	0.92699	0.90620	0.88681	0.86841	0.84527	0.82848
40	38	0.95000	0.89604	0.87237	0.85085	0.83080	0.80599	0.78823
40	37	0.92500	0.86643	0.84055	0.81741	0.79614	0.77009	0.75163
40	36	0.90000	0.83767	0.81002	0.78560	0.76336	0.73638	0.71739
40	35	0.87500	0.80954	0.78042	0.75497	0.73197	0.70426	0.68488
40	34	0.85000	0.78188	0.75155	0.72526	0.70165	0.67339	0.65372
40	33	0.82500	0.75463	0.72327	0.69629	0.67221	0.64354	0.62368
40	32	0.80000	0.72770	0.69550	0.66797	0.64352	0.61456	0.59458
40	31	0.77500	0.70108	0.66817	0.64021	0.61549	0.58634	0.56631
40	30	0.75000	0.67471	0.64124	0.61294	0.58804	0.55800	0.53879
40	29	0.72500	0.64858	0.61465	0.58612	0.56112	0.53187	0.51193
40	28	0.70000	0.62266	0.58840	0.55972	0.53468	0.50552	0.48569
40	27	0.67500	0.59695	0.56245	0.53370	0.50871	0.47969	0.46003
40	26	0.65000	0.57143	0.53679	0.50805	0.48316	0.45436	0.43492
40	25	0.62500	0.54609	0.51140	0.48275	0.45801	0.42951	0.41032
40	24	0.60000	0.52093	0.48628	0.45778	0.43327	0.40512	0.38623
40	23	0.57500	0.49593	0.46141	0.43314	0.40890	0.38117	0.36262
40	22	0.55000	0.47110	0.43679	0.40881	0.38491	0.35765	0.33949
40	21	0.52500	0.44644	0.41242	0.38480	0.36128	0.33457	0.31682
40	20	0.50000	0.42194	0.38830	0.36109	0.33802	0.31190	0.29461

n	r	p	80%	90%	95%	97.5%	99%	99.5%
41	41	1.00000	0.96151	0.94539	0.92954	0.91396	0.89376	0.87877
41	40	0.97561	0.92872	0.90841	0.88945	0.87145	0.84878	0.83234
41	39	0.95122	0.89851	0.87536	0.85429	0.83467	0.81035	0.79294
41	38	0.92683	0.86959	0.84426	0.82160	0.80075	0.77522	0.75709
41	37	0.90244	0.84150	0.81442	0.79049	0.76869	0.74220	0.72355
41	36	0.87805	0.81401	0.78549	0.76053	0.73796	0.71074	0.69169
41	35	0.85366	0.78699	0.75726	0.73146	0.70827	0.68049	0.66114
41	34	0.82927	0.76036	0.72960	0.70311	0.67944	0.65123	0.63167
41	33	0.80488	0.73405	0.70244	0.67539	0.65133	0.62281	0.60312
41	32	0.78049	0.70802	0.67570	0.64820	0.62386	0.59513	0.57537
41	31	0.75610	0.68225	0.64935	0.62149	0.59695	0.56810	0.54833
41	30	0.73171	0.65670	0.62333	0.59522	0.57056	0.54167	0.52194
41	29	0.70732	0.63136	0.59763	0.56935	0.54463	0.51578	0.49615
41	28	0.68293	0.60622	0.57222	0.54385	0.51913	0.49041	0.47092
41	27	0.65854	0.58126	0.54709	0.51869	0.49405	0.46551	0.44620
41	26	0.63415	0.55647	0.52222	0.49388	0.46936	0.44107	0.42199

n	r	p	80%	90%	95%	97.5%	99%	99.5%
45	45	1.00000	0.96487	0.95012	0.93560	0.92129	0.90273	0.88893
45	44	0.97778	0.93492	0.91629	0.89887	0.88230	0.86139	0.84619
45	43	0.95556	0.90730	0.88602	0.86662	0.84851	0.82601	0.80986
45	42	0.93333	0.88085	0.85753	0.83661	0.81732	0.79362	0.77676
45	41	0.91111	0.85515	0.83016	0.80802	0.78779	0.76315	0.74574
45	40	0.88889	0.82999	0.80361	0.78046	0.75946	0.73407	0.71623
45	39	0.86667	0.80525	0.77769	0.75370	0.73208	0.70697	0.68790
45	38	0.84444	0.78085	0.75228	0.72759	0.70545	0.67897	0.66054
45	37	0.82222	0.75674	0.72731	0.70203	0.67947	0.65261	0.63399
45	36	0.80000	0.73287	0.70271	0.67694	0.65404	0.62690	0.60815
45	35	0.77778	0.70923	0.67845	0.65227	0.62911	0.60176	0.58294
45	34	0.75556	0.68579	0.65448	0.62798	0.60463	0.57715	0.55830
45	33	0.73333	0.66252	0.63079	0.60404	0.58055	0.55301	0.53418
45	32	0.71111	0.63943	0.60735	0.58042	0.55685	0.52931	0.51054
45	31	0.68889	0.61649	0.58414	0.55710	0.53351	0.50603	0.48734
45	30	0.66667	0.59370	0.56116	0.53406	0.51050	0.48313	0.46458
45	29	0.64444	0.57105	0.53839	0.51129	0.48780	0.46060	0.44221
45	28	0.62222	0.54853	0.51582	0.48878	0.46541	0.43843	0.42024
45	27	0.60000	0.52615	0.49345	0.46652	0.44331	0.41660	0.39864
45	26	0.57778	0.50389	0.47128	0.44451	0.42150	0.39511	0.37741
45	25	0.55556	0.48176	0.44929	0.42273	0.39997	0.37394	0.35653
45	24	0.53333	0.45975	0.42748	0.40118	0.37872	0.35310	0.33601
45	23	0.51111	0.43786	0.40586	0.37987	0.35774	0.33258	0.31584

42	36	0.85714	0.79187	0.76271	0.73738	0.71461	0.68729	0.66824
42	35	0.83333	0.76583	0.73565	0.70963	0.68636	0.65860	0.63933
42	34	0.80952	0.74011	0.70907	0.68248	0.65882	0.63072	0.61131
42	33	0.78571	0.71466	0.68290	0.65585	0.63188	0.60356	0.58406
42	32	0.76190	0.68945	0.65710	0.62968	0.60550	0.57703	0.55750
42	31	0.73810	0.66446	0.63163	0.60393	0.57960	0.55107	0.53157
42	30	0.71429	0.63968	0.60646	0.57857	0.55416	0.52564	0.50621
42	29	0.69048	0.61508	0.58157	0.55356	0.52914	0.50070	0.48139
42	28	0.66667	0.59066	0.55695	0.52889	0.50451	0.47623	0.45707
42	27	0.64286	0.56640	0.53258	0.50454	0.48026	0.45218	0.43323
42	26	0.61905	0.54230	0.50845	0.48050	0.45637	0.42856	0.40984
42	25	0.59524	0.51836	0.48455	0.45674	0.43282	0.40534	0.38690
42	24	0.57143	0.49457	0.46088	0.43328	0.40961	0.38252	0.36439
42	23	0.54762	0.47092	0.43743	0.41009	0.38673	0.36008	0.34229
42	22	0.52381	0.44742	0.41419	0.38719	0.36418	0.33801	0.32061
42	21	0.50000	0.42406	0.39118	0.36455	0.34195	0.31633	0.29935

n	r	p	80%	90%	95%	97.5%	99%	99.5%
43	43	1.00000	0.96326	0.94786	0.93270	0.91779	0.89844	0.88407
43	42	0.97674	0.93196	0.91253	0.89437	0.87711	0.85536	0.83957
43	41	0.95349	0.90310	0.88093	0.86073	0.84189	0.81852	0.80176
43	40	0.93023	0.87547	0.85119	0.82944	0.80939	0.78481	0.76734
43	39	0.90698	0.84863	0.82264	0.79964	0.77865	0.75312	0.73511
43	38	0.88372	0.82236	0.79495	0.77093	0.74917	0.72289	0.70446
43	37	0.86047	0.79653	0.76792	0.74306	0.72068	0.69381	0.67506
43	36	0.83721	0.77106	0.74144	0.71587	0.69299	0.66566	0.64668
43	35	0.81395	0.74590	0.71542	0.68927	0.66599	0.63831	0.61917
43	34	0.79070	0.72100	0.68979	0.66318	0.63958	0.61165	0.59241
43	33	0.76744	0.69634	0.66452	0.63753	0.61369	0.58560	0.56631
43	32	0.74419	0.67188	0.63957	0.61228	0.58828	0.56011	0.54082
43	31	0.72093	0.64763	0.61491	0.58741	0.56331	0.53512	0.51589
43	30	0.69767	0.62355	0.59053	0.56288	0.53875	0.51061	0.49147
43	29	0.67442	0.59964	0.56639	0.53868	0.51456	0.48654	0.46753
43	28	0.65116	0.57589	0.54250	0.51478	0.49073	0.46289	0.44406
43	27	0.62791	0.55230	0.51884	0.49117	0.46725	0.43964	

45 22 0.48889 0.41609 0.38442 0.35879 0.33703 0.31237 0.29601

n	r	p	80%	90%	95%	97.5%	99%	99.5%
46	46	1.00000	0.96562	0.95118	0.93695	0.92294	0.90474	0.89120
46	45	0.97826	0.93630	0.91805	0.90098	0.88473	0.86422	0.84931
46	44	0.95652	0.90926	0.88841	0.86939	0.85161	0.82953	0.81367
46	43	0.93478	0.88337	0.86050	0.83998	0.82104	0.79776	0.78119
46	42	0.91304	0.85820	0.83369	0.81196	0.79208	0.76786	0.75075
46	41	0.89130	0.83356	0.80768	0.78494	0.76430	0.73933	0.72178
46	40	0.86957	0.80933	0.78228	0.75870	0.73743	0.71185	0.69395
46	39	0.84783	0.78544	0.75737	0.73309	0.71131	0.68523	0.66707
46	38	0.82609	0.76182	0.73290	0.70802	0.68581	0.65934	0.64099
46	37	0.80435	0.73844	0.70878	0.68341	0.66085	0.63409	0.61559
46	36	0.78261	0.71528	0.68499	0.65921	0.63638	0.60939	0.59080
46	35	0.76087	0.69231	0.66148	0.63537	0.61233	0.58520	0.56656
46	34	0.73913	0.66951	0.63825	0.61187	0.58868	0.56146	0.54283
46	33	0.71739	0.64688	0.61525	0.58868	0.56540	0.53815	0.51956
46	32	0.69565	0.62440	0.59249	0.56578	0.54245	0.51524	0.49673
46	31	0.67391	0.60206	0.56993	0.54315	0.51983	0.49271	0.47430
46	30	0.65217	0.57985	0.54759	0.52078	0.49751	0.47053	0.45227
46	29	0.63043	0.55778	0.52544	0.49866	0.47548	0.44869	0.43060
46	28	0.60870	0.53583	0.50347	0.47678	0.45374	0.42719	0.40930
46	27	0.58696	0.51400	0.48169	0.45513	0.43227	0.40600	0.38835
46	26	0.56522	0.49229	0.46009	0.43371	0.41107	0.38513	0.36774
46	25	0.54348	0.47070	0.43867	0.41251	0.39013	0.36456	0.34747
46	24	0.52174	0.44922	0.41742	0.39154	0.36946	0.34430	0.32753
46	23	0.50000	0.42786	0.39634	0.37078	0.34904	0.32434	0.30792

n	r	p	80%	90%	95%	97.5%	99%	99.5%
47	47	1.00000	0.96634	0.95219	0.93825	0.92451	0.90666	0.89339
47	46	0.97872	0.93763	0.91974	0.90300	0.88706	0.86694	0.85230
47	45	0.95745	0.91115	0.89070	0.87204	0.85459	0.83291	0.81733
47	44	0.93617	0.88578	0.86335	0.84321	0.82461	0.80174	0.78545
47	43	0.91489	0.86113	0.83708	0.81574	0.79621	0.77240	0.75556
47	42	0.89362	0.83699	0.81158	0.78924	0.76895	0.74438	0.72711
47	41	0.87234	0.81325	0.78667	0.76350	0.74259	0.71740	0.69978
47	40	0.85106	0.78983	0.76226	0.73838	0.71694	0.69126	0.67336
47	39	0.82979	0.76669	0.73826	0.71378	0.69191	0.66583	0.64772
47	38	0.80851	0.74378	0.71461	0.68963	0.66740	0.64101	0.62275
47	37	0.78723	0.72108	0.69127	0.66587	0.64336	0.61673	0.59837
47	36	0.76596	0.69856	0.66821	0.64247	0.61974	0.59294	0.57453
47	35	0.74468	0.67622	0.64541	0.61940	0.59650	0.56960	0.55117
47	34	0.72340	0.65403	0.62285	0.59662	0.57362	0.54668	0.52827
47	33	0.70213	0.63198	0.60051	0.57413	0.55106	0.52413	0.50578
47	32	0.68085	0.61008	0.57837	0.55190	0.52882	0.50195	0.48369
47	31	0.65957	0.58830	0.55643	0.52992	0.50687	0.48011	0.46198
47	30	0.63830	0.56665	0.53468	0.50817	0.48520	0.45860	0.44062
47	29	0.61702	0.54512	0.51311	0.48666	0.46380	0.43741	0.41962
47	28	0.59574	0.52371	0.49172	0.46537	0.44266	0.41653	0.39895
47	27	0.57447	0.50241	0.47049	0.44430	0.42178	0.39594	0.37860
47	26	0.55319	0.48122	0.44944	0.42344	0.40116	0.37565	0.35858
47	25	0.53191	0.46014	0.42855	0.40279	0.38078	0.35565	0.33887
47	24	0.51064	0.43917	0.40783	0.38236	0.36064	0.33594	0.31948
47	23	0.48936	0.41831	0.38727	0.36212	0.34076	0.31651	0.30041

n	r	p	80%	90%	95%	97.5%	99%	99.5%
48	48	1.00000	0.96703	0.95316	0.93950	0.92603	0.90852	0.89549
48	47	0.97917	0.93890	0.92136	0.90494	0.88930	0.86955	0.85517
48	46	0.95833	0.91295	0.89290	0.87459	0.85746	0.83616	0.82085

n	r	p	80%	90%	95%	97.5%	99%	99.5%
49	49	1.00000	0.96769	0.95410	0.94069	0.92748	0.91030	0.89751
49	48	0.97959	0.94012	0.92292	0.90681	0.89146	0.87206	0.85794
49	47	0.95918	0.91469	0.89501	0.87703	0.86021	0.83929	0.82424
49	46	0.93878	0.89032	0.86872	0.84929	0.83134	0.80925	0.79349
49	45	0.91837	0.86664	0.84345	0.82285	0.80399	0.78096	0.76465
49	44	0.89796	0.84344	0.81892	0.79734	0.77772	0.75393	0.73718
49	43	0.87755	0.82062	0.79496	0.77256	0.75231	0.72789	0.71079
49	42	0.85714	0.79811	0.77147	0.74836	0.72758	0.70265	0.68526
49	41	0.83673	0.77586	0.74836	0.72465	0.70343	0.67809	0.66047
49	40	0.81633	0.75383	0.72559	0.70137	0.67978	0.65410	0.63631
49	39	0.79592	0.73200	0.70311	0.67846	0.65657	0.63063	0.61272
49	38	0.77551	0.71034	0.68090	0.65589	0.63376	0.60762	0.58963
49	37	0.75510	0.68885	0.65893	0.63362	0.61130	0.58503	0.56700
49	36	0.73469	0.66750	0.63719	0.61164	0.58918	0.56283	0.54479
49	35	0.71429	0.64628	0.61565	0.58991	0.56737	0.54099	0.52298
49	34	0.69388	0.62520	0.59430	0.56844	0.54585	0.51949	0.50154
49	33	0.67347	0.60423	0.57313	0.54720	0.52460	0.49831	0.48044
49	32	0.65306	0.58339	0.55215	0.52618	0.50361	0.47743	0.45969
49	31	0.63265	0.56265	0.53132	0.50537	0.48288	0.45685	0.43925
49	30	0.61224	0.54202	0.51067	0.48477	0.46239	0.43655	0.41913
49	29	0.59184	0.52150	0.49016	0.46437	0.44213	0.41653	0.39930
49	28	0.57143	0.50107	0.46982	0.44416	0.42210	0.39678	0.37978
49	27	0.55102	0.48075	0.44962	0.42415	0.40231	0.37729	0.36054
49	26	0.53061	0.46053	0.42958	0.40432	0.38273	0.35807	0.34159
49	25	0.51020	0.44040	0.40968	0.38469	0.36338	0.33910	0.32292
49	24	0.48980	0.42038	0.38993	0.36524	0.34425	0.32040	0.30454

n	r	p	80%	90%	95%	97.5%	99%	99.5%
50	50	1.00000	0.96832	0.95499	0.94184	0.92888	0.91201	0.89945
50	49	0.98000	0.94130	0.92442	0.90860	0.89353	0.87448	0.86060
50	48	0.96000	0.91635	0.89704	0.87939	0.86286	0.84230	0.82750
50	47	0.94000	0.89246	0.87124	0.85216	0.83452	0.81279	0.79729
50	46	0.92000	0.86923	0.84645	0.82621	0.80766	0.78500	0.76895
50	45	0.90000	0.84648	0.82238	0.80117	0.78186	0.75845	0.74195
50	44	0.88000	0.82410	0.79887	0.77683	0.75690	0.73285	0.71600
50	43	0.86000	0.80201	0.77581	0.75306	0.73260	0.70804	0.69089
50	42	0.84000	0.78018	0.75313	0.72978	0.70887	0.68389	0.66651
50	41	0.82000	0.75857	0.73077	0.70691	0.68563	0.66030	0.64274
50	40	0.80000	0.73715	0.70870	0.68440	0.66282	0.63721	0.61952
50	39	0.78000	0.71590	0.68689	0.66223	0.64039	0.61457	0.59679
50	38	0.76000	0.69480	0.66532	0.64034	0.61831	0.59235	0.57451
50	37	0.74000	0.67385	0.64396	0.61874	0.59655	0.57049	0.55264
50	36	0.72000	0.65303	0.62280	0.59738	0.57509	0.54899	0.53115
50	35	0.70000	0.63233	0.60182	0.57627	0.55392	0.52781	0.51002
50	34	0.68000	0.61175	0.58103	0.55538	0.53301	0.50695	0.48923
50	33	0.66000	0.59128	0.56040	0.53470	0.51235	0.48638	0.46876
50	32	0.64000	0.57092	0.53994	0.51423	0.49193	0.46609	0.44860
50	31	0.62000	0.55066	0.51963	0.49396	0.47175	0.44608	0.42874
50	30	0.60000	0.53051	0.49947	0.47388	0.45179	0.42633	0.40918
50	29	0.58000	0.51045	0.47946	0.45399	0.43206	0.40684	0.38989
50	28	0.56000	0.49049	0.45960	0.43428	0.41254	0.38761	0.37089
50	27	0.54000	0.47062	0.43988	0.41476	0.39324	0.36863	0.35216
50	26	0.52000	0.45084	0.42030	0.39541	0.37415	0.34989	0.33370
50	25	0.50000	0.43116	0.40086	0.37625	0.35527	0.33140	0.31551

List of Symbols, Abbreviations, and Acronyms

cdf	cumulative distribution function
CI	confidence interval
CP	Clopper-Pearson
iid	independent and identically distributed
LR	likelihood ratio
MLE	maximum likelihood estimator
nf	number of failures
pdf	probability density function
qf	quantile function

NO. OF
COPIES ORGANIZATION

1 DEFENSE TECHNICAL
(PDF INFORMATION CTR
only) DTIC OCA
8725 JOHN J KINGMAN RD
STE 0944
FORT BELVOIR VA 22060-6218

1 DIRECTOR
US ARMY RESEARCH LAB
IMNE ALC HRR
2800 POWDER MILL RD
ADELPHI MD 20783-1197

1 DIRECTOR
US ARMY RESEARCH LAB
RDRL CIM L
2800 POWDER MILL RD
ADELPHI MD 20783-1197

1 DIRECTOR
US ARMY RESEARCH LAB
RDRL CIM P
2800 POWDER MILL RD
ADELPHI MD 20783-1197

1 ASST SECY ARMY
(CD ACQSTN LOGISTICS & TECH
only) SAAL ZP RM 2E661
103 ARMY PENTAGON
WASHINGTON DC 20310-0103

1 ASST SECY ARMY
(CD ACQSTN LOGISTICS & TECH
only) SAAL ZS RM 3E448
103 ARMY PENTAGON
WASHINGTON DC 20310-0103

1 DIRECTOR FORCE DEV
(CD DAPR FDZ
only) RM 3A522
460 ARMY PENTAGON
WASHINGTON DC 20310-0460

1 US ARMY TRADOC ANL CTR
ATRC W
A KEINTZ
WSMR NM 88002-5502

1 USARL
RDRL SLE
R FLORES
WSMR NM 88002-5513

ABERDEEN PROVING GROUND

1 US ARMY DEV TEST COM
CSTE DTC TT T
314 LONGS CORNER RD
APG MD 21005-5055

1 US ARMY EVALUATION CTR
CSTE AEC SVE
R LAUGHMAN
4120 SUSQUEHANNA AVE
APG MD 21005-3013

22 DIR USARL
RDRL SL
J BEILFUSS
J FEENEY
J FRANZ
M STARKS
P TANENBAUM
RDRL SLB
G MANNIX
RDRL SLB A
D FARENWALD
RDRL SLB D
R GROTE
RDRL SLB E
M PERRY
RDRL SLB G
P MERGLER
RDRL SLB S
S SNEAD
RDRL SLB W
L ROACH
RDRL CIM G (BLDG 4600)
RDRL SLB D
J COLLINS (5 CPS)
J EDWARDS
T HOLDREN
L MOSS
E SNYDER