

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAR 2010		2. REPORT TYPE		3. DATES COVERED 00-01-2010 to 00-03-2010	
4. TITLE AND SUBTITLE First Record of the Gulf Coast Tick, Amblyomma maculatum, from the Lark Bunting, Calamospiza melanocorys			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Medical Center,DPMIAC/AFPMB,Washington,DC,20307-5001			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

**First Record of the Gulf Coast Tick,
Amblyomma maculatum, from the
Lark Bunting, *Calamospiza
melanocorys***

On 16 Dec 2006, we removed a single engorged ixodid tick from the head of an adult female Lark Bunting, *Calamospiza melanocorys*, that had been captured accidentally in a Museum Special snap trap baited with peanut butter and rolled oats to collect small mammals. The capture site was located on a tract of land north of U.S. Highway 90, approximately 2 km east of Alpine, Brewster County, TX. This land is grazed by cattle, and the desert scrub vegetation is characterized by grasses (Poaceae), mesquite (*Prosopis glandulosa*), cactus (*Opuntia* spp.), yucca (*Yucca* spp.), fendlerbush (*Fendlera* sp.), and agarito (*Mahonia trifoliolata*). The specimen was shipped to Richard G. Robbins, who identified it as a nymph of *Amblyomma maculatum*, the so-called Gulf Coast tick, which in North America chiefly occurs along the Gulf Coast and in the southern Atlantic coastal states. Adults parasitize a variety of large wild and domestic mammals (Keirans and Litwak 1989); immature stages (nymphs, larvae) feed on small- to medium-sized mammals, but also on ground-frequenting birds, sometimes accompanying the latter on migrations as far north as southern Canada (Keirans and Durden 1998). A thorough search of the literature has revealed that our specimen of *A. maculatum* is the first to have been recorded from Lark Bunting.

The Lark Bunting feeds and nests on the ground, breeds in plains, prairies, meadows, and sagebrush, and winters in cultivated or brushy areas and deserts (Shane 2000). It should, therefore, be parasitized by any of numerous North American ticks specific to small mammals and lagomorphs, its ecological associates (Cooley and Kohls 1945, Cooley 1946, Rainwater et al. 2007). However, we have found only one brief reference to tick parasitism of this bird: a single larva of the Rocky Mountain wood tick, *Dermacentor andersoni*, that was removed from one of four Lark Buntings

examined for ticks on the grounds of the Idaho National Laboratory (formerly the National Reactor Testing Station) in southeastern Idaho (Allred 1968).

Amblyomma maculatum commonly parasitizes a great variety of ground-foraging birds, including members of the families Corvidae, Emberizidae, Icteridae, Laniidae, Mimidae, Odontophoridae, and Troglodytidae (Hixson 1940, Durden et al. 2001). Surprisingly, Lark Bunting is only the third North American emberizid from which *A. maculatum* has been recorded. We hope that our novel collection will prompt additional investigations of the ectoparasite fauna associated with the Lark Bunting, a bird that appears to be declining in concert with its open-country habitats.

In summary, the Gulf Coast tick, *Amblyomma maculatum*, is reported from the Lark Bunting for the first time. Only one other tick species, the Rocky Mountain wood tick, *Dermacentor andersoni*, is known from this bunting, a ground-nesting bird that should theoretically support a diverse tick fauna.

We thank Lewis Medlock for assisting us with trapping. Our Lark Bunting specimen has been assigned accession number CUSC 4016 and deposited in the Clemson University Museum of Natural Sciences. Our tick specimen has been assigned accession number YPM-ENT 300690 and deposited in the Division of Entomology, Peabody Museum of Natural History, Yale University, New Haven, CT.

LITERATURE CITED

- Allred, D.M. 1968. Ticks of the National Reactor Testing Station. *Brigham Young University Science Bulletin* 10:1-29.
- Cooley, R.A. 1946. The genera *Boophilus*, *Rhipicephalus*, and *Haemaphysalis* (Ixodidae) of the new world. *National Institute Health Bulletin* 187:1-54.
- Cooley, R.A. and G.M. Kohls. 1945. The genus *Ixodes* in North America. *National Institute Health Bulletin* 184:1-246.

- Durden, L.A., J.H. Oliver, Jr., and A.A. Kinsey. 2001. Ticks (Acari: Ixodidae) and spirochetes (Spirochaetaceae: Spirochaetales) recovered from birds on a Georgia barrier island. *Journal of Medical Entomology* 38:231-236.
- Hixson, H. 1940. Field biology and environmental relationships of the Gulf Coast tick in southern Georgia. *Journal of Economic Entomology* 33:179-189.
- Keirans, J.E. and L.A. Durden. 1998. Illustrated key to nymphs of the tick genus *Amblyomma* (Acari: Ixodidae) found in the United States. *Journal of Medical Entomology* 35:489-495.
- Keirans, J.E. and T.R. Litwak. 1989. Pictorial key to the adults of hard ticks, family Ixodidae (Ixodida: Ixodoidea), east of the Mississippi River. *Journal of Medical Entomology* 26:435-448.
- Rainwater, T.R., R.G. Robbins, J.L. Ingold, S.G. Platt and T.L. Walker. 2007. First record of the Gulf Coast tick, *Amblyomma maculatum*, from Lincoln's Sparrow, *Melospiza lincolnii*, with a review of tick parasitism of this sparrow. *North American Bird Bander* 32:118-119.
- Shane, T.G. 2000. Lark Bunting (*Calamospiza melanocorys*). In *The birds of North America*, No. 542 (A. Poole and F. Gill, eds). The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.

Richard G. Robbins
DPMIAC/AFPMB
Walter Reed Army Medical Center
Washington, DC 20307-5001
 richard.robbsins@osd.mil

Steven G. Platt
Department of Biology
Sul Ross State University
Alpine, TX 79832

Stanlee M. Miller
Department Biological Sciences
Clemson University
Clemson, SC 29634