

Acoustic Tomography With Navy Sonars

John L. Spiesberger
Department of Meteorology and the Applied Research Laboratory
The Pennsylvania State University, University Park, PA 16802
phone: (814) 863-8601 fax: (814) 863-9527 email: jspies@ems.psu.edu
Award #: N00014-97-1-0484

LONG-TERM GOAL

Utilize Navy sonars to map and understand temperature variability in ocean basins using acoustical tomography and hydrographic data. The data can be used to check and update ocean models for nowcasts and forecasts, and to check complimentary information from satellites.

Most of the data come from U.S. submarines which tow arrays and measure temperature at 1-s intervals. Additional data come from the U.S. Navy's Sound Surveillance System (SOSUS).

OBJECTIVES

A principal objective of this grant is to show it is possible to use U.S. submarines to collect temperature and acoustical tomography data over basin-scales. A secondary objective is to show it possible to read existing Navy data tapes at SOSUS stations to process tomography signals from ATOC sources and sources deployed by Naval Air.

APPROACH

Data are to be collected from submarines and SOSUS stations, and combined with hydrographic data to map ocean temperatures using a Kalman filter.

WORK COMPLETED

Acoustical data have been collected from a submarine. These data show high signal-to-noise ratios at 3000 km from electronically generated signals from the Kauaii ATOC source. ATOC source signals have been successfully processed from the Navy's data tapes at SOSUS stations. Acoustic signals generated from Naval Air exercises have been successfully interpreted from the SOSUS data tapes.

RESULTS

We have found that submarines provide excellent platforms for receiving acoustic tomography data over basin-scales. We have learned that existing hydrographic data sets show westward-propagating Rossby waves which emanate from the west coast of the U.S. following El Nino's and La Nina's. These Rossby waves are evident in these data sets since the 1970's, leading to temperature variations of a few degrees Celsius. The speeds of these waves are consistently faster than predicted using standard linear theory, as the work by Chelton and Schax have observed using the TOPEX/POSEIDON altimeter. The Naval Research Laboratory's Layered Ocean Model (Hurlburt et al., 1996) shows similar

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Acoustic Tomography With Navy Sonars				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Pennsylvania State University, Applied Research Laboratory, PO Box 30, State College, PA, 16804				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

westward-propagating Rossby waves following ENSO events. However, the speed of the modeled Rossby waves are significantly slower than observed, despite the fact that the ocean model incorporates physical mechanisms that have been proposed to explain why the Rossby waves travel faster than given by standard linear theory (Jacobson and Spiesberger, 1998).

We have found that it is usually the case that ray theory is adequate for interpreting multipath transmissions over basin scales in the northeast Pacific (Norris et al, 1998). We have found that Rossby waves linked to ENSO are a dominant parameter affecting climatic temperature variability in the northeast Pacific (Spiesberger et al. 1998). These waves lead to temperature variations of a few degrees Celsius at 300 m depth (Spiesberger et al., 1998; Jacobson and Spiesberger, 1998). We have found that acoustic thermometry data can measure changes in the sea surface height, due to thermal expansion, to an accuracy of about 0.3 cm. This is about ten times more accurate than is possible from the one billion dollar TOPEX/POSEIDON satellite altimeter (Spiesberger et al, 1998).

IMPACT/APPLICATIONS

Utilizing the U.S. submarine fleet to collect oceanographic data will greatly increase the quality of our images of the temperature field in the ocean's interior. These images will make it possible to better test theories, models, and other sensor's estimates of the ocean, such as satellites.

TRANSITIONS

This work is at a start, and it is too early for a transition to have occurred. Discussions with the Navy are underway.

RELATED PROJECTS

The ATOC project is utilizing SOSUS stations to map the ocean's temperature field from basin-scale transmissions.

REFERENCE

Hurlburt, H.E., Wallcraft, A.J., Schmitz, W.J., Jr., Hogan, P.J., Metzger, E.J., Teague, W.J., Dynamics of the Kuroshio/Oyashio current system using eddy-resolving models of the North Pacific Ocean, *J. Geophys. Res.*, 101, 941-976, (1996).

PUBLICATIONS

Jacobson, A. R., and Spiesberger, J. L., Observations of ENSO-induced Rossby Waves in the Northeast Pacific Using in situ Data, *J. Geophys. Res.*, in press (1998).

Spiesberger, J.L., Hurlburt, H.E., Johnson, M., Keller, M., Meyers, S., and O'Brien, J.J., Acoustic thermometry data compared with two ocean models: The importance of Rossby waves and ENSO in modifying the ocean interior, *Dynamics of Atmospheres and Oceans*, 26(4), 209-240, (1998).

Fabrikant, A., Spiesberger, J.L., and Silivra, A., and Hurlburt, H.E., Estimating Climatic Temperature

Change in the Ocean with Synthetic Acoustic Apertures, J. IEEE Ocean Eng. 23(1), 20-25, (1998).

Norris, D.E., Spiesberger, J.L., and Merdes, D.W., Comparison of basin-scale acoustic transmissions with rays and further evidence for a structured thermal field in the northeast Pacific, J. Acoust. Soc. Am., 103(1), 182-194, (1998).

Spiesberger, J.L., Fabrikant, A., Silivra, A., and Hurlburt, H.E., Mapping Climatic Temperature Changes in the Ocean with Acoustic Tomography: Navigational Requirements, J. IEEE Ocean. Eng., 22, No. 1, 128-142, (1997).

Silivra, A., Spiesberger, J.L., and Fabrikant, A., and Hurlburt, H.E., Acoustic tomography at basin-scales and clock errors, J. IEEE Ocean Eng., 22, No. 1, 143-150 (1997).