

Dynamical Evaluation of Gulf Stream Simulations in Models with High Vertical Resolution

Harley E. Hurlburt¹, Eric P. Chassignet², E. Joseph Metzger¹, James G. Richman¹, William J. Schmitz, Jr.³, Jay F. Shriver¹, Alan J. Wallcraft¹, and Xiaobiao Xu⁴

1 Naval Research Laboratory, Stennis Space Center, MS

2 FSU/COAPS, Tallahassee, FL

3 WHOI, emeritus, Corpus Christi, TX

4 USM/DMS, Stennis Space Center, MS

Layered Ocean Model Workshop

RSMAS, University of Miami

Miami, FL


1-3 June 2009

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE JUN 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Dynamical Evaluation of Gulf Stream Simulations in Models with High Vertical Resolution				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,Stennis Space Center,MS,39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 17	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Focus of the evaluation


- Mean Gulf Stream pathway
- Inertial character of the surface currents
- Sea surface height variability
- Mean abyssal currents

Summer average pycnocline depth contours over 1980-1983 from a $1/12^\circ$ MICOM Atlantic simulation (CI=50 m)




From Haza et al. (2007, Ocean Modelling)


Velocity vectors of the 1980-1983 deep mean flow from a 1/12° MICOM simulation superimposed on lower layer isopachs, or contours of potential thickness ($h_2 \times f_0 / (f_0 + \beta_y)$)



From Haza et al. (2007, Ocean Modelling)

Observed Gulf Stream pathway using the 12°C isotherm at 400 m (Watts et al., 1995; JGR-O) (—) vs simulations 14a (top) and 14c (bottom) over 1998-2000 from Bryan et al. (2007, Ocean Modelling) (—). Simulations by 1/10° Atlantic POP model







RMS SSH variability over 1998-2000 for (a) simulation 14a, (b) simulation 14c, and (c) AVISO processed T/P and ERS altimeter data (Bryan et al., 2007, Ocean Model.).

From Atlantic simulations using the Los Alamos POP model, a z-level model on a B-grid.

Gulf Stream region mean abyssal transport in the 3°-4°C class from a 1/10° Atlantic POP simulation



From Hecht et al. (2008, AGU Monograph 177)
using Smith and Gent (2004, JPO) simulation A'
Arrows are plotted for values of transport per unit
width above an unspecified threshold



RAFOS float trajectories at
~1000 m (a-d) and ~1500 m (e-f)
depth overlaid on topography
contoured at 1 km intervals


Circles on trajectories at 1 mo.
intervals

Star where float is entrained into
the Gulf Stream

(—) Contemporaneous Gulf
Stream IR northwall pathway

(- -) Mean pathway

From Bower and Hunt (2000b,
JPO)



RAFOS float trajectories at 3500 m that do not retroreflect into the interior overlaid on topography contoured at 1 km intervals


Circles on trajectories at 1 mo. intervals

Star where float crosses under the Gulf Stream

(—) Contemporaneous Gulf Stream IR northwall pathway

(- -) Mean pathway

From Bower and Hunt (2000b, JPO)



RAFOS float trajectories at 3500 m that retrofect into the interior overlaid on topography contoured at 1 km intervals

Circles on trajectories at 1 mo. intervals


Star where float crosses under the Gulf Stream

(—) Contemporaneous Gulf Stream IR northwall pathway


(- -) Mean pathway

From Bower and Hunt (2000b, JPO)


Mean SSH over years 12-15 from 1/12° global HYCOM simulation 9.4 with the mean Gulf Stream IR northwall pathway by Cornillon and Sirkes overlaid





contour interval = 5 cm



1/12° global HYCOM years 12-15, Exp. 9.4




1/25° global HYCOM year 3, Exp. 1.0



SSH variability observed by altimetry vs 1/12° and 1/25° global HYCOM

Variability from altimetry is calculated along altimeter tracks from satellites in 4 different orbits


Layer 6: speed and currents over topography



1/12° global HYCOM mean currents from Exp. 9.4 yrs 12-15

(top) near surface (layer 6) and (bottom) abyssal (layer 27) with topographic contours and the mean Gulf Stream IR northwall pathway by Cornillon and Sirkes overlaid.


Layer 27: speed and currents over topography



Mean max. current speed (m/s)

	Near 75°W	Mean 75°-70°W
1/12° HYCOM	1.34	.89
1/16° NLOM	1.70	1.39
Ratio N/H	1.27	1.56
N/H = NLOM/HYCOM		

Mass Transport Streamfunction from a Barotropic Linear Simulation Forced by a QuikSCAT-Corrected ECMWF ERA40 Wind Stress Climatology




Sverdrup (1947) interior flow with Munk (1950) western boundary layers

— Observed mean IR north wall pathway (1982-1996) $\pm$ 1 std. dev. by Cornillon and Sirkes


CI = 2 Sv

Atlantic Meridional Overturning Streamfunction from 1/12° Global HYCOM



Transport Streamfunction (Sv)
Exp 9.4 Years 12-15

1/12° vs 1/25° global HYCOM mean current speed and vectors overlaid on topography in upper ocean (layer 6, 20-25 m) and abyssal ocean (layer 27, 3100-3650 m)



Dynamical Evaluation in the Gulf Stream Region: Summary and Conclusions

- Given an understanding of Gulf Stream dynamics
 - supported by a wide range of observational evidence in the upper and abyssal ocean
- We can evaluate Gulf Stream simulations dynamically
 - Allows us to identify relationships between flaws in the simulation and flaws in the dynamics
 - Helps us identify specific improvements that are needed
 - Much more powerful approach than the traditional focus on improving parameterizations of subgrid-scale processes and on parameter tuning