Modeling For Both Deep Ocean And Coastal Wave-Wave Interactions: Implications For WAM

Ray Q. Lin

Hydromechanics Directorate (Code 5500)

Carderock Division

Naval Surface Warfare Center (David Taylor Model Basin)

9500 MacArthur Boulevard

West Bethesda, MD 20817-5700

Phone: (301) 227-3945 Fax: (301) 227-5442

e-mail address: rlin@wave2.dt.navy.mil or linr@oasys.dt.navy.mil

Award # N0001498WX30038

LONG-TERM GOAL

My long term goal is to construct an accurate and efficient new WAM model, suitable for both deep ocean and coastal region.

OBJECTIVES

The central emphasis of the project is the construction of an efficient accurate source function for nonlinear wave-wave interactions (S_{nl}) . We also will construct the proper numerical scheme, kinematics, and the various source terms (wind input function, and dissipation), based on rigorous physical principles and statistical data analysis. Ultimately, we will achieve a completely new WAM wave model for deep ocean and coastal regions, with potential capability to include the current-wave and atmosphere-wave interactions in a truly interactive mode.

Surface waves in coastal waters play a far more important role than in deep water. The motions of the waves can penetrate the whole water column to influence mixing, sediment

Report Documentation Page				Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.						
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVE 00-00-1998	ered 3 to 00-00-1998	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Modeling for Both Deep Ocean and Coastal Wave-Wave Interactions: Implications for WAM				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Surface Warfare Center (David Taylor Model Basin),Carderock Division,9500 MacArthur Blvd,West Bethesda,MD,20817-5700				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited						
13. SUPPLEMENTARY NO See also ADM0022	DTES 52.					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER	19a. NAME OF			
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	ABSTRACT Same as Report (SAR)	OF PAGES 5	RESPONSIBLE PERSON	

Standard Form 298 (Rev. 8-9	8
Prescribed by ANSI Std Z39-	18

transportation, turbidity, and circulation of the water mass. Most past efforts in wave modeling apply to deep water. However, there are crucial differences between deep water and shallow water wave dynamics. Our scientific objectives also include clarification of these differences and establishment of the proper representations of their dynamics in both deep ocean and finite-depth coastal waters.

APPROACH

We have investigated both deep ocean and coastal kinematics and dynamics, in preparation for construction of a new WAM model.

In order to obtain an accurate and efficient S_{nl} , we are developing a "Reduced Integration Approximation" (RIA), which is based on the highly nonlinear dependence the interaction coefficient, within S_{nl} , has on the wave number. From an analysis of the nonlinearity, we reduce the 3-dimensional integrations by Resio and Tracy (1982), hereafter denoted R-T, to a quasi-line integration. Resio and Tracy (1982) integrated the nonlinear Boltzmann integral for the wave-wave interactions along the resonant locus of interactions in wave-number space and reduced the Hasselmann and Hasselmann (1981)'s classical 5-dimensional integrations, hereafter denoted H-H, to 3-dimensional integrations. In reducing the dimensions of integration, this method significantly reduces the CPU time. Thus RIA provides substantial savings.

WORK COMPLETED

We had completed studies on the numerics and kinematics in FY97 (Lin and Huang, 1996a and b). In FY98, we completed the studies on the nonlinear four-wave interactions and five-wave nonlinear. We also have produced seven manuscripts, associated with the finished work (Lin and Perrie, 1997a, b, and c; 1998a, b, and c; Lin and Su, 1998).

We have developed a preliminary "RIA" source code for calculating nonlinear source functions. The documentation for RIA is in preparation. We have compared our model results with some published model results for a wide range spectra (wide, narrow, and split spreading angles). We are currently developing an independent pseudospectral method to calculate the nonlinear source function. Our object is to verify the accuracy of our nonlinear source function S_{nl} and to reduce the CPU time.

RESULTS

(1) Theory:

Resonant wave-wave interaction processes were studied, using the nonlinear dispersion relationship for finite depth water. The formulation was derived from a Hamiltonian representation. Our results show that four waves are needed for resonant interactions at all depths. Furthermore, when we consider the nonlinear dispersion relationship for waves in intermediate water, two interaction modes can result, depending on the water depth and the nonlinearity. The first mode is the classic Phillips (1960) interaction, involving four waves of comparable wave lengths, which dominates in deep water. The second mode, which prevails in shallow water, still involves four waves, but with one component of vanishingly small wave number. Triadic shoaling interactions are an approximate asymptotic limit of the latter case.

Three-dimensional (five-waves) interactions may dominate when the spreading angle of the spectrum is very narrow and nonlinearity is greater than 0.3 (Lin and Su, 1998).

(2) Numerical Method:

The RIA method is based on Hamiltonian principles. To analyze the major solutions which we locate, we reduce 3 integrations of R-T to quasi-line integrations. Our method is both accurate and efficient, and also suitable for both deep ocean and shallow water(1997a and b). We compare our model results with published model results for a wide range spectra (both wide, narrow, and split spreading angles), and with 4 well-known methods, H-H, R-T, DIA (standard WAM model), and DIA2 (proposed 'new' WAM model). Our results showed:

- a. We got very good agreement with H-H, R-T, Masuda (1980), and Masuda and Komatsu (1998) when the dispersion is linear. The most exciting thing about our results is that when the spectrum is split, all exact solutions (H-H, R-T, and RIA) generated similar horse shoe patterns. For a narrow spreading angle spectrum, our model results agree with those of Herterich and Hasselmann (1980). This is true in both deep ocean water and shallow water.
- b. Our model results differ significantly from the results of DIA and DIA2.
- c. Our model results agree well with the winter storm data from DUCK94 experiment (Lin and Perrie, 1998a).

IMPACT/APPLICATION

The means by which we have studied the coastal kinematics, dynamics, wave-wave interactions and our development of the RIA method is novel and appropriate for future investigations. Nonlinear wave-wave interactions need 4-wave interactions for all water depths. When the water depth becomes very small, one component must become of vanishingly small wave number. Previous work on triadic shoaling interactions represent an approximate asymptotic limit. We first developed this theory in 1997, which is supported by Zakharov (1998). The accuracy of this theory is a guiding light for future source code development.

RIA method is based on theory: the nonlinear interaction coefficient depends on the wave number in a highly nonlinear manner. Therefore, we are able to reduce the 3-dimensional integrations of R-T to a quasi-line integration. This is different from the (previous) approach using piece-wide approximations, which motivate DIA and DIA2 (Snyder et al., 1981). Recent tests comparing RIA to DIA and DIA2 at Vicksburg show (Jenssen et al, 1998) that the former is correct, but the latter is incorrect.

TRANSITIONS

The theory which we suggested above is accepted by the S_{nl} BE group. We hope that we can use it to significantly improve S_{nl} in WAM.

RELATED PROJECT

I have recently begun a collaboration with Dr. Scott Chubb who is in the remote sensing group in NRL (Washington DC) to study effects of seamounts to trap waves. I also have collaborated with Dr. Perrie at BIO (Dartmouth, Nova Scotia, Canada) to model and predict wave-current interactions in the St. Lawrence River and Gulf.

REFERENCES

- Hasselmann, S. and K. Hasselmann, 1981: A Symmetrical Method of Computing the Nonlinear Transfer in a Gravity Wave Spectrum. *Hamburger Geophys.* Einzelschr. A 52, 163 pp.
- 2 Hasselmann et al: The SWAMP Group. (1985) <u>Ocean Wave Modeling</u>, *Plenum*, New York. 256 pp.
- 3 Jenssen, R. E., D. Resio, R.-Q. Lin, van Vledder, T. H. C. Herbers, and B. Tracy, 1998: A Report of the Computational Test of the Nonlinear Source Functions. A Workshop in Vicksburg.
- 4 Lin, R.-Q., and N. E. Huang, 1996a: The Goddard Coastal Wave Model. Part I. Numerics. J. Physical Oceanography, Vol. 26, 833-847.
- 5 Lin, R.-Q., and N. E. Huang, 1996b: The Goddard Coastal Wave Model. Part II. Kinematics. J. Physical Oceanography, Vol. 26, 848-862.
- 6 Resio, D and B. Tracy, 1998: Wave-wave interactions in finite water depth. In preparation.
- 7 Tracy, B. A. and D. T. Resio, 1982: Theory and calculation of the nonlinear energy transfer between sea waves in deep water. WIS REP. 11, US Army Engineering Waterways Experiment Station, Vicksburg, MS. 50pp.
- 8 Snyder, R., F. Dobson, J. Elliott and R. Long, 1981: Array Measurements of Atmospheric Pressure Fluctuations above Surface Gravity Waves. J. Fluid Mech., Vol. 102, p. 1-59.
- 9 Zakharov, V. E., 1998: Weakly Nonlinear Waves on the Surface of an Ideal Finite Depth Fluid. Amer. Math. Soc. Transl. Vol 182, 167-197.

PUBLICATIONS:

- Lin, R.-Q., and W. Perrie, 1997a: A New Coastal Wave Model. Part III. Nonlinear Wave-Wave interaction. J. Physical Oceanography Vol. 27, 1813-1826.
- 2 Lin, R.-Q., and W. Perrie, 1997b: On the Mathematics and Approximation of the Nonlinear Wave-wave Interactions. In press in <u>Nonlinear Ocean Waves</u>. Ed. W. Perrie. Computational Fluid Mechanics, Southampton. 27pp.
- 3 Lin, R.-Q., and W. Perrie, 1997c: A New Coastal Wave Model. Part V. Five Wave Interactions. J. Physical Oceanography. Vol. 27, 2169-2186.
- 4 Lin, R.-Q., and W. Perrie, 1998a: Case studies of Hurricanes and Winter Storms using a New coastal Wave model. *Proceeding of 22nd Naval Symposium*.
- 5 Lin, R.-Q., and W. Perrie, 1998b: An Equilibrium Energy Spectral Submitted to J. of Geophys. Res..
- 6 Lin, R.-Q., and W. Perrie, 1998c: A New Coastal Wave Model. Part VI. Edge Waves. Submitted to J. Geophys. Res..
- 7 Lin, R.-Q., and M. Su, 1998: Five-Wave Interactions in Deep Water. Submitted to IUTAM Symposium, France, Nice.