

Peer Reviewed

Title:

Os Odontoideum: Rare Cervical Lesion

Journal Issue:

Western Journal of Emergency Medicine, 12(4)

Author:

Robson, Kristie A LCDR, MC, USN, Naval Medical Center Portsmouth, Department of Emergency Medicine, Portsmouth Virginia

Publication Date:

2011

Publication Info:

Western Journal of Emergency Medicine, Department of Emergency Medicine, UC Irvine

Permalink:

http://www.escholarship.org/uc/item/76k070f7

Acknowledgements:

Corresponding address: The author can be contacted by email at Kristie.robson@med.navy.mil Disclaimer (Must be printed alongside manuscript): The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government. Copyright statement: We are military service members. This work was prepared as part of our official duties. Title 17, USC, §105 provides that 'Copyright protection under this title is not available for any work of the United States Government.' Title 17, USC, §101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person's official duties.

Keywords:

Os Odontoideum, atlantoaxillary instability

Preferred Citation:

Robson KA. Os Odontoideum: Rare Cervical Lesion. West J Emerg Med. 2011;12(4):520–522.

Abstract:

We report the case of a 22-year-old Marine who presented to the emergency department, after a martial arts exercise, with transient weakness and numbness in all extremities. Computed tomography cervical spine radiographs revealed os odontoideum. Lateral flexion—extension radiographs identified atlanto-axillary instability. This abnormality is rare and can be career ending for military members who do not undergo surgical fusion. [West J Emerg Med. 2011;12(4):520—522.]



eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comments arters Services, Directorate for Info	regarding this burden estimate rmation Operations and Reports	or any other aspect of the 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE NOV 2011		2. REPORT TYPE		3. DATES COVERED 00-11-2011 to 00-11-2011	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Os Odontoideum: Rare Cervical Lesion				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Medical Center Portsmouth, Department of Emergency Medicine, Portsmouth, VA, 23701				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NO Western Journal o	otes f Emergency Medic	ine, Vol 12, no. 4, N	ovember 2011		
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC	17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF		
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	6	RESPONSIBLE PERSON

Report Documentation Page

Form Approved OMB No. 0704-0188

License Statement:

This is an Open Access article distributed under the terms of the Creative Commons Non-Commercial Attribution License, which permits its use in any digital medium, provided the original work is properly cited and not altered. For details, please refer to http://creativecommons.org/licenses/by-nc/3.0/. Authors grant Western Journal of Emergency Medicine as well as the National Library of Medicine a nonexclusive license to publish the manuscript. Western Journal of Emergency Medicine is produced by the eScholarship Repository and bepress.



Os Odontoideum: Rare Cervical Lesion

Kristie A. Robson, MD, LCDR, MC, USN

Naval Medical Center Portsmouth, Department of Emergency Medicine, Portsmouth, Virginia

Supervising Section Editor: Sean Henderson, MD

Submission history: Submitted July 7, 2010; Revision received January 5, 2011; Accepted April 4, 2011

Reprints available through open access at http://escholarship.org/uc/uciem westjem

DOI: 10.5811/westjem.2011.4.2029

We report the case of a 22-year-old Marine who presented to the emergency department, after a martial arts exercise, with transient weakness and numbness in all extremities. Computed tomography cervical spine radiographs revealed os odontoideum. Lateral flexion—extension radiographs identified atlanto-axillary instability. This abnormality is rare and can be career ending for military members who do not undergo surgical fusion. [West J Emerg Med. 2011;12(4):520–522.]

INTRODUCTION

Os odontoideum is a congenital or posttraumatic abnormality of the second cervical vertebrae in which the odontoid process is separated from the body of the axis by a transverse gap. This lesion can look like a fracture at the base of the odontoid. The lesion is frequently asymptomatic. Discovery of this cervical abnormality in an active-duty Marine Corps Private First Class occurred after trauma from martial arts training. This case report will review clinical presentations, diagnostic studies, and treatment options for os odontoideum.

CASE REPORT

A 22-year-old man presented to the emergency department (ED) for evaluation of diffuse numbness and weakness in all extremities after head trauma during a martial arts exercise. He was thrown by his right arm and landed on the top of his head. The patient stated that he was unable to move for approximately 8 minutes secondary to the "wind being knocked out of him." He regained all sensation and strength by the time he reached the ED 20 minutes later by ambulance. His only complaint on arrival was pain in the midline, occipito-cervical region of his posterior neck. The patient was an active duty Marine Corps Private First Class with no significant medical history or history of head or neck trauma.

Physical examination revealed a fit young man on a backboard with a rigid cervical spinal collar in place. Vital signs were all within normal limits. His head was atraumatic. Patient had palpable midline posterior cervical spinal tenderness along C1 to C2 with no step-offs noted. Cranial nerves were intact, and he had no sensory or motor deficits on

his neurologic examination. The remainder of his examination was without abnormalities.

A noncontrast cervical spine computed tomography (CT) scan was obtained for further evaluation of his midline spinal tenderness and transient neurologic complaints. The CT revealed os odontoideum, which is demonstrated on the existing Figure 1. The radiologist could not exclude an acute subluxation and associated cord injury on CT alone and recommended a magnetic resonance image (MRI) of the cervical spine. MRI obtained in the ED showed a focal 7-mm craniocaudal axis region of myelomalacia at the C1 to C2 level (Figure 2). Findings were not thought to represent acute cord edema, but rather a chronic condition associated with the os odontoideum.

After neurosurgery was consulted, the patient was discharged home in a rigid cervical spinal collar and evaluated by neurosurgery the next day. Flexion/extension radiographs performed the next day showed instability, exacerbated on flexion, between the articulation of C1 and the C2 os odontoideum (Figure 3). The remainder of his cervical vertebral bodies had normal alignment with no additional abnormal translation. The patient was offered surgical fixation for his atlantoaxial instability.

DISCUSSION

Os odontoideum is an independent ossicle of variable size with smooth circumferential margins separated from a foreshortened odontoid peg. The ossicle stands apart from the hypoplastic dens and can adopt 2 anatomic types: orthotopic and dystopic. An ossicle located in the position of the normal odontoid is referred to as orthotopic. The ossicle is considered dystopic if it appears near the occiput in the area of the foramen

Robson Os Odontoideum



Figure 1. Noncontrast computed tomography of cervical spine. This shows an unfused dens (os odontoideum).




Figure 2. Magnetic resonance imaging of cervical spine. This shows 7 mm of focal myelomalacia at the C1 to C2 level associated with thinning of the cord, compatible with myelomalacia and not definitive acute cord edema. The cord above and below this level appears normal.



Figure 3. Flexion view of plain cervical spine. This image shows abnormal translation of the articulation between C1 and the C2 os odontoideum in relation to the remainder of the C2 vertebral body.

magnum.² The incidence of os odontoideum is unknown because the lesion is usually asymptomatic.²

The presentation of os odontoideum can vary from an incidental radiographic finding to severe findings of myelopathy, vertebral artery compression, and intracranial manifestations. ¹ Neurologic symptoms can vary from transient episodes of paresis after trauma to complete myelopathy from cord compression. ² Neck pain, torticollis, or headache can be caused from local irritation of the atlantoaxial joint. ² Compression of the vertebral artery can cause cervical and brainstem ischemia presenting as seizure, syncope, visual disturbances, and vertigo. Our patient experienced a transient episode of paresis associated with mild trauma.

Radiographic findings for os odontoideum consist of a free ossicle with smooth, sclerotic borders, usually half the size of a normal dens. An acute fracture would have a thinner and irregular space instead of the wide and smooth space of os odontoideum. CT scans with reconstruction views and MRIs are helpful in making the diagnosis. But initial routine cervical spine radiographs with an open-mouth odontoid view will also make the diagnosis. Lateral flexion and extension radiographs can detect instability. Instability in an adult is defined as greater than 3 mm of anterior and posterior displacement of the atlas on the axis.² This number increases to greater than 4 to 5 mm in children. Our young Marine had 5.1 mm of anterior translation of the articulation between C1 and the os odontoideum on flexion imaging. The remainder of his cervical vertebral bodies had normal alignment with no additional abnormal translation.

Indications for operative stabilization of the os odontoideum are symptomatic neurologic involvement, including transient symptoms; instability of >5 mm posteriorly or anteriorly; progressive instability, and

Os Odontoideum Robson

persistent neck complaints.² The primary concern in this odontoid abnormality is the ability of the abnormal atlantoaxial joint to sublux or even dislocate with minor trauma, resulting in death or permanent neurologic damage.² Service members who undergo surgical fixation will be able to continue on active duty on a case-by-case basis. Aviators or service members who require full atlantoaxial range of motion will undergo a military medical board review to determine fitness for duty. This Marine will be able to continue on service after surgical fixation. If he had, however, decided not to undergo surgery, he would not be recommended for continued military service by neurosurgery.

In summary, os odontoideum is an uncommon, but important cervical spine abnormality that can be associated with significant spinal cord and vertebral artery injuries and should be considered when a young person has symptoms consistent with upper cervical spine compression. Surgical fixation for patients with instability can save a military career and prevent catastrophic neurologic insult after minor trauma in the future.

Address for Correspondence: Kristie A. Robson, MD, LCDR, MC, USN, Naval Medical Center Portsmouth, Department of Emergency Medicine, 620 John Paul Jones Cir, Portsmouth, VA 23708-2197. E-mail: Kristie.robson@med.navy.mil.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources, and financial or management relationships that could be perceived as potential sources of bias. Disclaimer: The views expressed in this article are those of the author and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, or the US Government.

REFERENCES

- Arvin B, Fournier-Gosselin MP, Fehlings M. Os odontoideum: etiology and surgical management. *Neurosurgery*. 2010;66:A22–A31.
- Warner W. Pediatric cervical spine. In: Canale ST, Beaty JH, eds. Campbell's Operative Orthopaedics. 11th ed. Philadelphia, PA: Mosby; 2007:1879–1898.