

Award Number: W81XWH-09-2-0112

TITLE: Regional Center of Excellence for PTSD:Phoebe Putney Memorial Hospital

PRINCIPAL INVESTIGATOR: Steven Ziemba, Ph.D., FACHE

CONTRACTING ORGANIZATION: Phoebe Putney Memorial Hospital
Albany, GA 31701

REPORT DATE: September 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE September 2011		2. REPORT TYPE Annual		3. DATES COVERED 13 August 2010 – 12 August 2011	
4. TITLE AND SUBTITLE Regional Center of Excellence for PTSD:Phoebe Putney Memorial Hospital				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-09-2-0112	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Lori-Ann Landry, MSW, LCSW Steven Ziemba, PhD, FACHE E-Mail: sziemba@ppmh.org				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Phoebe Putney Memorial Hospital Albany, GA 31701				8. PERFORMING ORGANIZATION REPORT NUMBER	
10. SPONSOR/MONITOR'S ACRONYM(S)				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Post Traumatic Stress Disorder (PTSD) has been recognized by the Armed Forces as a significant mental health condition among returning US military personnel from Iraq and Afghanistan. Multiple challenges exist in the successful diagnosis and treatment of soldiers prior to symptoms becoming unmanageable and leading to dysfunction within families, homes, work, and in severe cases increased suicide rates. The current study will investigate the use of telepsychiatry services for military personnel who have served in Iraq and Afghanistan, demonstrate early signs of PTSD, and now reside in geographically isolated communities of Southwest Georgia. The study will follow a non-inferiority design, with 174 subjects enrolled to receive CBT either face-to-face or via telepsychiatry. Our sample size of 174 subjects consists of Active Duty, Veterans, Reservists, and National Guard personnel (within the Marine Corps, Navy and Army) between the ages of 18-45 of varying ethnicities who have served in Iraq and/or Afghanistan. The study is currently under regulatory review, with initial patient enrollment predicted for Fall 2011.					
15. SUBJECT TERMS Cognitive Behavioral Therapy, Telepsychiatry, PTSD, Videoconferencing, Access to Care, Veterans					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)
			UU	69	

Table of Contents

Introduction	4
Body.....	5
Key Research Accomplishments.....	9
Reportable Outcomes.....	10
Conclusion.....	11
References.....	12
Appendices.....	14

Introduction

United States Veterans living in rural areas of the United States often have difficulty accessing needed care. Professional resources, distance to providers, and limits on transportation options are identified as some of the barriers to care (Rosenheck & Stolar 1998; McCarthy & Blow, 2004; Tessler, Rosenheck, & Gamache, 2005). Access to care and barriers to receiving this care for returning soldiers from military operations in Iraq and Afghanistan with suspected or projected symptoms of Post Traumatic Stress Disorder (PTSD) has been highly publicized (Maurer, K, and Watson, J., 2010, August 26). Numerous agencies, both military and civilian are collaboratively mobilizing to increase options for medical and mental health care for returning soldiers (Maurer K., and Watson, J., 2010, August, 26). One particular example is seen in the treatment of soldiers with PTSD using telepsychiatry. The purpose of the present study is to demonstrate that Cognitive Behavioral Therapy (CBT) delivered via telepsychiatry can be as effective for the treatment of PTSD as traditional face-to-face therapy sessions for military personnel who have served in Iraq and Afghanistan, demonstrate early signs of PTSD, and now reside in geographically isolated communities of Southwest Georgia. Previous studies have typically utilized only VA institutions. The present study will provide telepsychiatry for up to 202 subjects through a civilian healthcare network using a non-inferiority design. Eligible populations include active duty, Reserve, National Guard, and Veterans of Iraq and Afghanistan who are between the ages of 18-45. The population will be drawn from the Southwest region of Georgia, which is predominantly rural. The study will be conducted at six Phoebe Putney Memorial Hospital (PPMH) clinics located throughout the region with subjects attending the clinic most conveniently located to them. The study's significance lies in its utilization of civilian hospital services to assess and compare feasibility and efficacy of CBT to treat PTSD using telepsychiatry in military populations.

Body

PTSD had been identified as a condition of concern for soldiers returning from Operation Iraqi Freedom (OIF) and Operation Enduring Freedom (OEF). Georgia is home to 12 active military installations with more than 100,000 Active Duty, Reservist and National Guard personnel and nearly one million Veterans and their dependents. Symptoms of PTSD can be debilitating and may include insomnia, intense anxiety, and difficulty coping with work, social, and family relationships. Left untreated, PTSD can lead to substance abuse, severe depression, and in some cases suicide. Symptoms may appear within months of the traumatic event or be delayed for years. While there is no cure for PTSD, experts believe early identification and treatment of PTSD symptoms may lessen their severity and improve the overall quality of life for individuals with this disorder (Department of Veteran Affairs, Department of Defense, 2004).

Studies have shown that Cognitive Behavioral Therapy (CBT) is effective in the treatment of PTSD for symptom reduction and increasing an individual's overall quality of life. Numerous other studies have demonstrated a reduction in PTSD symptoms as measured on various scales with CBT treatment in comparison to no treatment (Basoglu, Salcioglu, Livanou, Kalender & Acar, 2005; McDonagh et al. 2005; Foa et al., 2005; Hinton, Chhean, Pich, Safren, Hofmann, Pollack, 2005; Monson et al., 2006; Maercker A, Zollner T, Menning H., Rabe, S., and Karl, A., 2006).


Despite this effective treatment, many veterans may keep symptoms secret for a number of reasons, not the least of which is the stigma associated with psychiatric disorders (Hoge et al., 2004). Others may believe their symptoms are simply a private matter and will refrain from seeking treatment. Still others may fear that their careers may be adversely affected if they disclose signs or symptoms of any psychiatric disorder, including PTSD. A significant factor contributing to the lack of patients receiving care is limited accessibility of mental health services in many locations, especially rural locations. This challenge is especially present in Southwest Georgia. One method that can address this challenge is the use of videoconferencing as a means to provide psychiatric services or, as it is more commonly known, telepsychiatry. Specific forms of treatments and assessments, such as CBT delivered via telepsychiatry, have been reported for diagnoses such as Panic Disorder, Agoraphobia, Major Depressive Disorder, Oppositional Defiance Disorder, and Attention Deficit Disorder. (Cowain, 2001; Bouchard et al., 2000; Alessi, 2002; Hilty, Nesbitt, Canning & Hales, 2000). Telepsychiatry shows great promise as a model of therapeutic delivery and may be as effective as conventional face-to-face psychotherapy treatment delivery. Several studies have investigated the effectiveness of CBT delivered via telemedicine. However no studies have been conducted that use a community based treatment facility partnering with the Department of Defense. Further research in this area is warranted considering the US Army and Marine Corp's immediate need to provide psychiatric services to personnel in geographically remote regions in provider shortage areas (Maurer, K. and Watson, J., 2010, August 26).

Phoebe Putney Memorial Hospital (PPMH) is a 450-bed tertiary care hospital with a psychiatric and behavioral health component of 38 inpatient beds and a complete continuum of outpatient services. The hospital is located in Dougherty County in Southwest Georgia. This location provides a potential referral base extending from the immediate region to throughout the state, with several Department of Defense Facilities within Georgia, including the Marine Corps Logistics Base, Fort Benning, Fort Gordon, Fort McPherson, Fort Stewart, Moody AFB, Robins AFB and other DoD operated health clinics.

Phoebe's Behavioral Health Center has been serving the area for many years and includes inpatient and outpatient psychiatric care services, including a Structured Outpatient Program (SOP) in addition to traditional clinic visits. Counselors, social workers, marriage and family therapists, addiction specialists, nurse practitioners and psychiatrists are employed or available as contractors to provide care in the

programs. Phoebe Putney Memorial Hospital is well positioned to provide healthcare to rural areas with over six clinics spread throughout the Southwest Georgia region.

Phoebe Clinic Locations



The present study incorporates CBT delivered via telepsychiatry as compared with traditional face-to-face administration among military personnel diagnosed with PTSD. Subjects will be randomized into one of these two groups. The study will follow a non-inferiority design with pre and post assessments to test the hypothesis that CBT administered via telepsychiatry is as effective as CBT administered face-to-face in treating PTSD. A one sided design was selected, as opposed to an equivalency study, as the intent is to demonstrate effectiveness and not superior efficacy. This design approach has resulted in a minimum target population of 174. However, an attrition rate of twenty percent is expected, based on similar studies, resulting in a maximum target of up to 202. Subjects will be recruited from public and private area clinics, Veterans Administration and CBOC clinics, the local Marine Corps Logistics Base (MCLB) as well as U.S. Army installations, including Reservist and National Guard. The subjects will undergo 10 weekly sessions of CBT provided by trained and independently licensed study therapists within a 15 week period. Initial measures will include assessment using the Primary Care PTSD Screen (PC-PTSD)(Appendix A), the PTSD Checklist-Military (PCL-M)(Appendix B), and the DSM-IV diagnostic criteria (Appendix C)(Prins, A., Ouimette, P., Kimerling, R., Cameron, R.P., Hugelshofer, D.S., Shaw-Hegwer, J., Thrailkill, A., Gunsman, F.D., & Sheikh, J.I., 2003; Weathers, F., Litz, B., Herman, D., Huska, J., & Keane, T., 1993; American Psychological Association, 2000). Any effect upon PTSD symptoms of subjects will be measured by the Clinician-Administered PTSD Scale (CAPS)(Appendix D), Hamilton Anxiety Rating Scale (HAM-A)(Appendix E), Montgomery Asberg Depression Rating Scale (MADRS)(Appendix F), and the SF-36 Quality of Life Indicator (Appendix G) before initiation of treatment and following completion of all therapy sessions (Dudley D. B., Weathers, F.W., Nagy, L.M., Kaloupek, D.G., Charney, D.S., Keane, T.M., 1998; Hamilton, M, 1959; Montgomery, S.A. & Asberg, M., 1979; Quality Metric, 2009). The use of civilian facilities to provide telepsychiatry care to military personnel is unique in its approach.

An essential early objective of the study was the creation of an interactive, telepsychiatric network at several Primary Care clinics throughout Southwest Georgia. The telepsychiatry units were purchased from NuPhysicia, located in Houston, Texas, and consist of equipment mounted on mobile lockable carts as well as desktop units. Sites identified for equipment installation were checked for network access, and Ethernet ports were installed as required. Following installation, the network was tested for accessibility and reliability. Each unit can be contacted and controlled remotely from a unit operated by study staff. In this manner, the network has been continuously tested to ensure each unit is operational. Security is maintained as the network is encrypted and operates as an internal system.

Study personnel are currently in place and undergoing training in anticipation of the initiation of enrollment. Study personnel include a research rater to independently conduct the pre and post test therapy assessments; a research therapist to provide guided and standardized CBT therapy to enrolled subjects, both face-to-face and via telepsychiatry; and an independently contracted LCSW to measure fidelity in the use of CBT. In addition to the research personnel in place, the Research Coordinator, one Research Rater, and one Research Therapist position remains vacant with the management team actively recruiting to fill these positions.

The study is delayed in initiating enrollment as it awaits final regulatory approval from HRPO. It was indicated that approval by the local organization's (Phoebe Putney Memorial Hospital) IRB was required, followed by subsequent reviews by the Department of the Navy (DON) IRB and finally the U.S. Army Office of Research Protections, Human Research Protection Office (ORP, HRPO). DON IRB approval is required due to recruitment of potential subjects from the local Marine Corps Logistics Base (MCLB). Phoebe Putney Memorial Hospital IRB approval has been obtained and DON IRB approval was granted in December 2010. However, upon submission to ORP, HRPO, the necessity of the DON IRB oversight was questioned. Following a lengthy discussion with both ORP, HRPO and the DON IRB staff, the DON IRB

withdrew their approval recommending ORP HRPO oversight of the project with a letter of support from the MCLB Commander. Currently Project staff are awaiting this letter of support for resubmission to ORP HRPO. It is anticipated that all approvals will be received and subject enrollment will commence in the coming quarter.

Initiation of enrollment was anticipated for August 2010. However, multiple IRB submissions as well as coordination between DON IRB and HRPO has posed significant delays for this study. Following AIBS Scientific review and subsequent contracting in August 2009, the previous two years have been spent preparing for multiple submission and reviews for PPMH IRB, DON IRB submission and finally, ORP HRPO submission. No other significant issues causing delay in the study initiation have occurred.

KEY RESEARCH ACCOMPLISHMENTS

Accomplishments to date have been predominantly administrative and logistic in nature. Regardless, significant milestones have been met. These include the following:

- Design and construction of the study telepsychiatry network at PPMH Clinic sites in Albany, GA, Camilla, GA, Ashburn, GA, Sylvester, GA, and Lee County, GA
 - Receipt of equipment, delivery to PPMH clinics, design of network, and daily testing of equipment
- Hiring and ongoing training of study personnel to include initial research coordinator, two research raters, two research therapists (the research coordinator, one research therapist and one research rater have vacated their positions)
- Obtaining local (PPMH) IRB approval; DON IRB approval; and initial submission to ORP, HRPO
- Contracts for the Research Instruments have been purchased and licensed use permitted to the study for the SF-36 Quality of Life Indicator with Quality Metric and the Patient Satisfaction Questionnaire with Press Ganey. These contracts have been extended until August 2012 in anticipation of a one year no-cost extension
- Professional presentations:
 - May 2010: American Telemedicine Association Annual Conference May Poster Presentation: *Challenges of Implementing a PTSD Telepsychiatry Project: A Hospital Perspective*(Appendix H)
 - April 2010: TATRC Product Line Review
- Database developed, tested, and ready for use
- Fidelity Consultant and Project Manager completion of Beck Institute's Cognitive Therapy training with Drs. Judith and Aaron Beck

REPORTABLE OUTCOMES

The study is currently in its startup phase. The enrollment of study subjects is pending final ORP HRPO approval. As such, at present there are no outcomes to report.

CONCLUSION

The present study contributes to the ongoing validation of telepsychiatry in the treatment of PTSD among returning military personnel. Operationally, it is currently undergoing regulatory review. The required network, staff and data collection instruments have been put into place and are undergoing their respective testing and training. It is anticipated that enrollment can begin in the fall of 2011. However, if regulatory approval is granted earlier, our resources will be in place to activate the study. It is expected that up to a one-year time period is needed to meet the enrollment target of 174, although the rate of enrollment will be closely monitored. As such, project staff have formally requested a one year no-cost extension. It is expected that the next annual report will include final results. The expected implication of these results includes demonstrating a model of increased access to care for Active Duty, National Guard, Reservists, and Veterans via telepsychiatry. The results will further validate existing literature on the treatment of this population via telepsychiatry. Overall, increasing access to care through non-military providers in assisting in the management of symptoms, and restoration to functioning levels assists in the fight against PTSD and its disabling effects. To date literature searches have not yielded any peer reviewed results assessing the comparability of combat-related PTSD being treated with CBT face-to-face or via telepsychiatry using a community based institution in a rural area.

REFERENCES

- Alessi N.E. (2002). Telepsychiatric care for a depressed adolescent. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41(8), 894–895.
- American Psychological Association. (2000). *Diagnostic & Statistical Manual of Mental Disorders Text Revision (4th Edition)*. Washington, DC: APA.
- Basoglu M, Salcioglu E, Livanou M, Kalender, D., & Acar, G. (2005). Single-session behavioral treatment of earthquake-related posttraumatic stress disorder: A randomized waiting list controlled trial. *Journal of Trauma Stress*, 18(1), 1–11.
- Bouchard S., Payeur R., Rivard V., Allard, M., Paquin, B., Renaud, P., and Goyer, L. (2000). Cognitive behavior therapy for panic disorder with agoraphobia in videoconference: Preliminary results. *Cyberpsychology and Behavior*, 3(6), 999–1007.
- Cowain T. (2001). Cognitive-behavioural therapy via videoconferencing to a rural area. *The Australian and New Zealand Journal of Psychiatry*, 35(1), 62–64.
- Department of Veteran Affairs, Department of Defense. (2004). VA/DoD Clinical Practice Guideline for the Management of Post-Traumatic Stress. *Department of Veteran Affairs, Department of Defense*.
- Dudley D. B., Weathers, F.W., Nagy, L.M., Kaloupek, D.G., Charney, D.S., Keane, T.M. (1998). National Center for Post Traumatic Stress Disorder. Behavioral Science Division -- Boston VA Medical Center, Neurosciences Division -- West Haven VA Medical Center *Clinician-Administered PTSD Scale for DSM-IV*.
- Foa EB, Hembree EA, Cahill SP, Rauch, SA, Riggs, DS, Feeny, NC, and Yadin, E. (2005). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: outcomes at academic and community clinics. *Journal of Consulting and Clinical Psychology*, 73(5), 953–964.
- Hamilton, M. (1959). The assessment of anxiety states by rating. *British Journal of Medical Psychology*, 32, 50-55.
- Hilty DM, Nesbitt TS, Canning RD, and Hales, R. (2000). Telepsychiatry for the management of a liver transplantation candidate in the primary care setting. *General Hospital Psychiatry*, 22(2), 122–123.
- Hinton DE, Chhean D, Pich V, Safren, SA, Hofmann, SG, Pollack, MH. (2005). A randomized controlled trial of cognitive-behavior therapy for Cambodian refugees with treatment-resistant PTSD and panic attacks: A cross-over design. *Journal of Trauma Stress*, 18(6), 617–629.
- Hoge, C.W., Castro, C.A., Messer, S.C., McGurk, D., Cotting, D. I. & Koffman, R.L. (2004). Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *New England Journal of Medicine* 351(1), 13-22.

- Maercker A, Zollner T, Menning H., Rabe, S., and Karl, A. (2006). Dresden PTSD treatment study: Randomized controlled trial of motor vehicle accident survivors. *BMC Psychiatry*, 6(29), 1-8.
- Maurer, K. and Watson, J. (2010, August 26). Marines Pour Resources into Mental Health Care. *The Albany Herald*, pp.10A.
- McCarthy, J.F. & Blow, F.C. (2004). Older patients with serious mental illness: Sensitivity to distance barriers for outpatient care. *Medical Care*, 42(11), 1073-1080.
- McDonagh A, Friedman M, McHugo G, Ford, J., Sen Gupta, A., Meuser, K., Demment, C., Fournier, D., and Schnurr, P. (2005). Randomized trial of cognitive-behavioral therapy for chronic posttraumatic stress disorder in adult female survivors of childhood sexual abuse. *Journal of Consulting and Clinical Psychology*, 73(3), 515–524.
- Monson CM, Schnurr PP, Resick PA, Friedman, M., Yinong, Y., and Stevens, S. (2006). Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 74(5), 898–907.
- Montgomery, S.A. & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *British Journal of Psychiatry*, 134, 382-389.
- Prins, A., Ouimette, P., Kimerling, R., Cameron, R.P., Hugelshofer, D.S., Shaw-Hegwer, J., Thrailkill, A., Gunsman, F.D., & Sheikh, J.I. (2003) *The primary care PTSD screen*. Retrieved March 16, 2009, from United States Department of Veteran Affairs: National Center for PTSD Assessments Web site: http://www.ncptsd.va.gov/ncmain/ncdocs/assmnts/the_primary_care_ptsd_screen_pcptsd.html
- Quality Metric. (2009). *SF-36v2 Health Survey*. Retrieved March 19, 2009, from: Quality Metric Web site: <http://www.qualitymetric.com/WhatWeDo/GenericHealthSurveys/SF36v2HealthSurvey/tabid/185/Default.aspx>.
- Rosenheck R, Stolar M.(1998). Access to public mental health services: Determinants of population coverage. *Medical Care*, 36(4), 503–512.
- Tessler R, Rosenheck R, Gamache G. (2005). Declining access to alcohol and drug abuse services among veterans in the general population. *Military Medicine*, 170(3), 234–238.
- Weathers, F., Litz, B., Herman, D., Huska, J., & Keane, T. (1993). *PTSD Checklist(PCL)* Retrieved January 23, 2009 from United States Department of Veteran Affairs: National Center for PTSD Web site: <ftp://PTSD:Assessments@Files8.cyberlynk.net>.

APPENDICES

A.	Primary Care PTSD (PC-PTSD).....	15
B.	PTSD Checklist-Military (PCL-M).....	16
C.	DSM-IV Criteria for PTSD.....	17
D.	Clinician Administered PTSD Scale (CAPS).....	18
E.	Hamilton Anxiety Rating Scale (HAM-A).....	38
F.	Montgomery-Asberg Depression Rating Scale (MADRS).....	47
G.	SF-36 Quality of Life Indicator.....	53
H.	ATA Poster Presentation.....	59
I.	Product Line review (PLR) Behavioral Health (2010).....	60

The Primary Care PTSD Screen (PC-PTSD)

Prins, Ouimette, Kimerling et al., 2003

Description

The PC-PTSD is a 4-item screen that was designed for use in primary care and other medical settings and is currently used to screen for PTSD in veterans at the VA. The screen includes an introductory sentence to cue respondents to traumatic events. The authors suggest that in most circumstances the results of the PC-PTSD should be considered "positive" if a patient answers "yes" to any 3 items. Those screening positive should then be assessed with a structured interview for PTSD. The screen does not include a list of potentially traumatic events.

Scale

Instructions

In your life, have you ever had any experience that was so frightening, horrible, or upsetting that, in the past month, you:

1. Have had nightmares about it or thought about it when you did not want to?

YES / NO

2. Tried hard not to think about it or went out of your way to avoid situations that reminded you of it?

YES / NO

3. Were constantly on guard, watchful, or easily startled?

YES / NO

4. Felt numb or detached from others, activities, or your surroundings?

YES / NO

Current research suggests that the results of the PC-PTSD should be considered "positive" if a patient answers "yes" to any three items.

References

Prins, A., Ouimette, P., Kimerling, R., Cameron, R. P., Hugelshofer, D. S., Shaw-Hegwer, J., Thrailkill, A., Gusman, F.D., Sheikh, J. I. (2004). The primary care PTSD screen (PC-PTSD): development and operating characteristics. *Primary Care Psychiatry*, 9, 9-14

Appendix B

PCL-M

INSTRUCTIONS: Below is a list of problems and complaints that veterans sometimes have in response to stressful military experiences. Please read each one carefully, then circle one of the numbers to the right to indicate how much you have been bothered by that problem in the past month.

	Not at all	A little bit	Moderately	Quite a bit	Extremely
1. Repeated, disturbing <i>memories, thoughts, or images</i> of a stressful military experience?	1	2	3	4	5
2. Repeated, disturbing <i>dreams</i> of a stressful military experience?	1	2	3	4	5
3. Suddenly <i>acting or feeling</i> as if a stressful military experience <i>were happening again</i> (as if you were reliving it)?	1	2	3	4	5
4. Feeling <i>very upset</i> when <i>something reminded you</i> of a stressful military experience?	1	2	3	4	5
5. Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, sweating) when <i>something reminded you</i> of a stressful military experience?	1	2	3	4	5
6. Avoiding <i>thinking about</i> or <i>talking about</i> a stressful military experience or avoiding <i>having feelings</i> related to it?	1	2	3	4	5
7. Avoiding <i>activities</i> or <i>situations</i> because <i>they reminded you</i> of a stressful military experience?	1	2	3	4	5
8. Trouble <i>remembering important parts</i> of a stressful military experience?	1	2	3	4	5
9. <i>Loss of interest</i> in activities that you used to enjoy?	1	2	3	4	5
10. Feeling <i>distant</i> or <i>cut off</i> from other people?	1	2	3	4	5
11. Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?	1	2	3	4	5
12. Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?	1	2	3	4	5
13. Trouble <i>falling</i> or <i>staying asleep</i> ?	1	2	3	4	5
14. Feeling <i>irritable</i> or having <i>angry outbursts</i> ?	1	2	3	4	5
15. Having <i>difficulty concentrating</i> ?	1	2	3	4	5
16. Being " <i>super-alert</i> " or watchful or on guard?	1	2	3	4	5
17. Feeling <i>jumpy</i> or easily startled?	1	2	3	4	5

PCL-M for DSM-IV (11/1/94) Weathers, Litz, Huska, & Keane National Center for PTSD - Behavioral Science Division

Appendix C

PTSD Diagnostic Criteria DSM-IV-TR

Subject ID _____ Date: _____

Diagnostic criteria for PTSD include a history of exposure to a traumatic event (meeting two criteria) and symptoms from each of three symptom clusters: intrusive recollections, avoidant/numbing symptoms, and hyper-arousal symptoms. A fifth criterion concerns duration of symptoms and a sixth assesses functioning.

Criterion A: stressor

The person has been exposed to a traumatic event in which **both** of the following have been present:

- The person has experienced, witnessed, or been confronted with an event or events that involve actual or threatened death or serious injury, or a threat to the physical integrity of oneself or others;
- The person's response involved intense fear, helplessness, or horror.

Criterion B: intrusive recollection

The traumatic event is persistently re-experienced in at least **one (or more)** of the following ways:

- Recurrent and intrusive distressing recollections of the event, including images, thoughts, or perceptions. Note: in young children, repetitive play may occur in which themes or aspects of the trauma are expressed;
- Recurrent distressing dreams of the event. Note: in children, there may be frightening dreams without recognizable content;
- Acting or feeling as if the traumatic event were recurring (includes a sense of reliving the experience, illusions, hallucinations, and dissociative flashback episodes, including those that occur upon awakening or when intoxicated). Note: in children, trauma-specific reenactment may occur;
- Intense psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event;
- Physiologic reactivity upon exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event.

Criterion C: avoidant/numbing

Persistent avoidance of stimuli associated with the trauma and numbing of general responsiveness (not present before the trauma), as indicated by at least **three** of the following:

- Efforts to avoid thoughts, feelings, or conversations associated with the trauma;
- Efforts to avoid activities, places, or people that arouse recollections of the trauma;
- Inability to recall an important aspect of the trauma;
- Markedly diminished interest or participation in significant activities;
- Feeling of detachment or estrangement from others;
- Restricted range of affect (e.g., unable to have loving feelings);
- Sense of foreshortened future (e.g., does not expect to have a career, marriage, children, or a normal life span).

Criterion D: hyper-arousal

Persistent symptoms of increasing arousal (not present before the trauma), indicated by at least **two** of the following:

- Difficulty falling or staying asleep;
- Irritability or outbursts of anger;
- Difficulty concentrating;
- Hyper-vigilance;
- Exaggerated startle response.

Criterion E: duration

- Duration of the disturbance (symptoms in B, C, and D) is more than one month.

Criterion F: functional significance

- The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Specify if:

- Acute:** if duration of symptoms is less than three months (length of sxs: _____)
- Chronic:** if duration of symptoms is three months or more (length of sxs: _____)

Specify if:

- With or
- Without delay onset: Onset of symptoms at least six months after the stressor

Appendix D

National Center for PTSD

CLINICIAN-ADMINISTERED PTSD SCALE FOR DSM-IV

Name: _____ ID #: _____
Interviewer: _____ Date: _____
Study: _____

Dudley D. Blake, Frank W. Weathers, Linda M. Nagy, Danny
G. Kaloupek, Dennis S. Charney, & Terence M. Keane

National Center for Posttraumatic Stress Disorder

Behavioral Science Division -- Boston VA Medical Center
Neurosciences Division -- West Haven VA Medical Center

Revised July 1998

Criterion A. The person has been exposed to a traumatic event in which both of the following were present:
 (1) the person experienced, witnessed, or was confronted with an event or events that involved actual or threatened death or serious injury, or a threat to the physical integrity of self or others
 (2) the person’s response involved intense fear, helplessness, or horror. Note: In children, this may be expressed instead by disorganized or agitated behavior

I’m going to be asking you about some difficult or stressful things that sometimes happen to people. Some examples of this are being in some type of serious accident; being in a fire, a hurricane, or an earthquake; being mugged or beaten up or attacked with a weapon; or being forced to have sex when you didn’t want to. I’ll start by asking you to look over a list of experiences like this and check any that apply to you. Then, if any of them do apply to you, I’ll ask you to briefly describe what happened and how you felt at the time.

Some of these experiences may be hard to remember or may bring back uncomfortable memories or feelings. People often find that talking about them can be helpful, but it’s up to you to decide how much you want to tell me. As we go along, if you find yourself becoming upset, let me know and we can slow down and talk about it. Also, if you have any questions or you don’t understand something, please let me know. Do you have any questions before we start?

ADMINISTER CHECKLIST, THEN REVIEW AND INQUIRE UP TO THREE EVENTS. IF MORE THAN THREE EVENTS ENDORSED, DETERMINE WHICH THREE EVENTS TO INQUIRE (E.G., FIRST, WORST, AND MOST RECENT EVENTS; THREE WORST EVENTS; TRAUMA OF INTEREST PLUS TWO OTHER WORST EVENTS, ETC.)

IF NO EVENTS ENDORSED ON CHECKLIST: *(Has there ever been a time when your life was in danger or you were seriously injured or harmed?)*

IF NO: *(What about a time when you were threatened with death or serious injury, even if you weren’t actually injured or harmed?)*

IF NO: *(What about witnessing something like this happen to someone else or finding out that it happened to someone close to you?)*

IF NO: *(What would you say are some of the most stressful experiences you have had over your life?)*

EVENT #1

<p>What happened? <i>(How old were you? Who else was involved? How many times did this happen? Life threat? Serious injury?)</i></p> <p>How did you respond emotionally? <i>(Were you very anxious or frightened? Horrified? Helpless? How so? Were you stunned or in shock so that you didn’t feel anything at all? What was that like? What did other people notice about your emotional response? What about after the event - how did you respond emotionally?)</i></p>	<p>Describe (e.g., event type, victim, perpetrator, age, frequency):</p> <p>A. (1)</p> <p>Life threat? NO YES [self other]</p> <p>Serious injury? NO YES [self other]</p> <p>Threat to physical integrity? NO YES [self other]</p> <p>A. (2)</p> <p>Intense fear/help/horror? NO YES [during after]</p> <p>Criterion A met? NO PROBABLE YES</p>
--	---

EVENT #2

<p>What happened? <i>(How old were you? Who else was involved? How many times did this happen? Life threat? Serious injury?)</i></p> <p>How did you respond emotionally? <i>(Were you very anxious or frightened? Horrified? Helpless? How so? Were you stunned or in shock so that you didn't feel anything at all? What was that like? What did other people notice about your emotional response? What about after the event - how did you respond emotionally?)</i></p>	<p>Describe (e.g., event type, victim, perpetrator, age, frequency):</p> <p>A. (1) Life threat? NO YES [self other]</p> <p>Serious injury? NO YES [self other]</p> <p>Threat to physical integrity? NO YES [self other]</p> <p>A. (2) Intense fear/help/horror? NO YES [during after]</p> <p>Criterion A met? NO PROBABLE YES</p>
---	---

EVENT #3

<p>What happened? <i>(How old were you? Who else was involved? How many times did this happen? Life threat? Serious injury?)</i></p> <p>How did you respond emotionally? <i>(Were you very anxious or frightened? Horrified? Helpless? How so? Were you stunned or in shock so that you didn't feel anything at all? What was that like? What did other people notice about your emotional response? What about after the event - how did you respond emotionally?)</i></p>	<p>Describe (e.g., event type, victim, perpetrator, age, frequency):</p> <p>A. (1) Life threat? NO YES [self other]</p> <p>Serious injury? NO YES [self other]</p> <p>Threat to physical integrity? NO YES [self other ___]</p> <p>A. (2) Intense fear/help/horror? NO YES [during after ___]</p> <p>Criterion A met? NO PROBABLE YES</p>
---	---

For the rest of the interview, I want you to keep (EVENTS) in mind as I ask you some questions about how they may have affected you.

I'm going to ask you about twenty-five questions altogether. Most of them have two parts. First, I'll ask if you've ever had a particular problem, and if so, about how often in the past month (*week*). Then I'll ask you how much distress or discomfort that problem may have caused you.

Criterion B. The traumatic event is persistently reexperienced in one (or more) of the following ways:

1. (B-1) recurrent and intrusive distressing recollections of the event, including images, thoughts, or perceptions.
Note: In young children, repetitive play may occur in which themes or aspects of the trauma are expressed.

<p>Frequency Have you ever had unwanted memories of (EVENT)? What were they like? (<i>What did you remember?</i>) [IF NOT CLEAR:] (<i>Did they ever occur while you were awake, or only in dreams?</i>) [EXCLUDE IF MEMORIES OCCURRED ONLY DURING DREAMS] How often have you had these memories in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much distress or discomfort did these memories cause you? Were you able to put them out of your mind and think about something else? (<i>How hard did you have to try?</i>) How much did they interfere with your life?</p> <p>0 None 1 Mild, minimal distress or disruption of activities 2 Moderate, distress clearly present but still manageable, some disruption of activities 3 Severe, considerable distress, difficulty dismissing memories, marked disruption of activities 4 Extreme, incapacitating distress, cannot dismiss memories, unable to continue activities</p> <p>QV (specify) _____</p>	<p>Past week</p> <p>F I</p>
		<p>Past month</p> <p>F I Sx: Y N</p> <p>Lifetime</p> <p>F I Sx: Y N</p>

2. (B-2) recurrent distressing dreams of the event. **Note:** In children, there may be frightening dreams without recognizable content.

<p>Frequency Have you ever had unpleasant dreams about (EVENT)? Describe a typical dream. (<i>What happens in them?</i>) How often have you had these dreams in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much distress or discomfort did these dreams cause you? Did they ever wake you up? [IF YES:] (<i>What happened when you woke up? How long did it take you to get back to sleep?</i>) [LISTEN FOR REPORT OF ANXIOUS AROUSAL, YELLING, ACTING OUT THE NIGHTMARE] (<i>Did your dreams ever affect anyone else? How so?</i>)</p> <p>0 None 1 Mild, minimal distress, may not have awoken 2 Moderate, awoke in distress but readily returned to sleep 3 Severe, considerable distress, difficulty returning to sleep 4 Extreme, incapacitating distress, did not return to sleep</p> <p>QV (specify) _____</p>	<p>Past week</p> <p>F I</p>
		<p>Past month</p> <p>F I Sx: Y N</p> <p>Lifetime</p> <p>F I Sx: Y N</p>

3. (B-3) acting or feeling as if the traumatic event were recurring (includes a sense of reliving the experience, illusions, hallucinations, and dissociative flashback episodes, including those that occur on awakening or when intoxicated). **Note:** In young children, trauma-specific reenactment may occur.

<p>Frequency Have you ever suddenly acted or felt as if (EVENT) were happening again? (<i>Have you ever had flashbacks about [EVENT]?</i>) [IF NOT CLEAR:] (<i>Did this ever occur while you were awake, or only in dreams?</i>) [EXCLUDE IF OCCURRED ONLY DURING DREAMS] Tell me more about that. How often has that happened in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much did it seem as if (EVENT) were happening again? (<i>Were you confused about where you actually were or what you were doing at the time?</i>) How long did it last? What did you do while this was happening? (<i>Did other people notice your behavior? What did they say?</i>)</p> <p>0 No reliving 1 Mild, somewhat more realistic than just thinking about event 2 Moderate, definite but transient dissociative quality, still very aware of surroundings, daydreaming quality 3 Severe, strongly dissociative (reports images, sounds, or smells) but retained some awareness of surroundings 4 Extreme, complete dissociation (flashback), no awareness of surroundings, may be unresponsive, possible amnesia for the episode (blackout)</p> <p>QV (specify) _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
--	---	--

4. (B-4) intense psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event

<p>Frequency Have you ever gotten emotionally upset when something reminded you of (EVENT)? (<i>Has anything ever triggered bad feelings related to [EVENT]?</i>) What kinds of reminders made you upset? How often in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much distress or discomfort did (REMINDERS) cause you? How long did it last? How much did it interfere with your life?</p> <p>0 None 1 Mild, minimal distress or disruption of activities 2 Moderate, distress clearly present but still manageable, some disruption of activities 3 Severe, considerable distress, marked disruption of activities 4 Extreme, incapacitating distress, unable to continue activities</p> <p>QV (specify) _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	---	--

5. (B-5) physiological reactivity on exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event

<p>Frequency Have you ever had any physical reactions when something reminded you of (EVENT)? <i>(Did your body ever react in some way when something reminded you of [EVENT]?)</i> Can you give me some examples? <i>(Did your heart race or did your breathing change? What about sweating or feeling really tense or shaky?)</i> What kinds of reminders triggered these reactions? How often in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong were (PHYSICAL REACTIONS)? How long did they last? <i>(Did they last even after you were out of the situation?)</i></p> <p>0 No physical reactivity 1 Mild, minimal reactivity 2 Moderate, physical reactivity clearly present, may be sustained if exposure continues 3 Severe, marked physical reactivity, sustained throughout exposure 4 Extreme, dramatic physical reactivity, sustained arousal even after exposure has ended</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i> F I</p> <p><i>Past month</i> F I Sx: Y N</p> <p><i>Lifetime</i> F I Sx: Y N</p>
---	---	---

Criterion C. Persistent avoidance of stimuli associated with the trauma and numbing of general responsiveness (not present before the trauma), as indicated by three (or more) of the following:

6. (C-1) efforts to avoid thoughts, feelings, or conversations associated with the trauma

<p>Frequency Have you ever tried to avoid thoughts or feelings about (EVENT)? <i>(What kinds of thoughts or feelings did you try to avoid?)</i> What about trying to avoid talking with other people about it? <i>(Why is that?)</i> How often in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much effort did you make to avoid (THOUGHTS/FEELINGS/CONVERSATIONS)? <i>(What kinds of things did you do? What about drinking or using medication or street drugs?)</i> [CONSIDER ALL ATTEMPTS AT AVOIDANCE, INCLUDING DISTRACTION, SUPPRESSION, AND USE OF ALCOHOL/DRUGS] How much did that interfere with your life?</p> <p>0 None 1 Mild, minimal effort, little or no disruption of activities 2 Moderate, some effort, avoidance definitely present, some disruption of activities 3 Severe, considerable effort, marked avoidance, marked disruption of activities, or involvement in certain activities as avoidant strategy 4 Extreme, drastic attempts at avoidance, unable to continue activities, or excessive involvement in certain activities as avoidant strategy</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i> F I</p> <p><i>Past month</i> F I Sx: Y N</p> <p><i>Lifetime</i> F I Sx: Y N</p>
---	--	---

7. (C-2) efforts to avoid activities, places, or people that arouse recollections of the trauma

<p>Frequency Have you ever tried to avoid certain activities, places, or people that reminded you of (EVENT)? <i>(What kinds of things did you avoid? Why is that?)</i> How often in the past month (week)?</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How much effort did you make to avoid (ACTIVITIES/PLACES/PEOPLE)? <i>(What did you do instead?)</i> How much did that interfere with your life?</p> <p>0 None 1 Mild, minimal effort, little or no disruption of activities 2 Moderate, some effort, avoidance definitely present, some disruption of activities 3 Severe, considerable effort, marked avoidance, marked disruption of activities or involvement in certain activities as avoidant strategy 4 Extreme, drastic attempts at avoidance, unable to continue activities, or excessive involvement in certain activities as avoidant strategy</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	--	--

8. (C-3) inability to recall an important aspect of the trauma

<p>Frequency Have you had difficulty remembering some important parts of (EVENT)? Tell me more about that. <i>(Do you feel you should be able to remember these things? Why do you think you can't?)</i> In the past month (week), how much of the important parts of (EVENT) have you had difficulty remembering? <i>(What parts do you still remember?)</i></p> <p>0 None, clear memory 1 Few aspects not remembered (less than 10%) 2 Some aspects not remembered (approx 20-30%) 3 Many aspects not remembered (approx 50-60%) 4 Most or all aspects not remembered (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How much difficulty did you have recalling important parts of (EVENT)? <i>(Were you able to recall more if you tried?)</i></p> <p>0 None 1 Mild, minimal difficulty 2 Moderate, some difficulty, could recall with effort 3 Severe, considerable difficulty, even with effort 4 Extreme, completely unable to recall important aspects of event</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	--	--

9. (C-4) markedly diminished interest or participation in significant activities

<p>Frequency Have you been less interested in activities that you used to enjoy? (What kinds of things have you lost interest in? Are there some things you don't do at all anymore? Why is that?) [EXCLUDE IF NO OPPORTUNITY, IF PHYSICALLY UNABLE, OR IF DEVELOPMENTALLY APPROPRIATE CHANGE IN PREFERRED ACTIVITIES] In the past month (week), how many activities have you been less interested in? (What kinds of things do you still enjoy doing?) When did you first start to feel that way? (After the [EVENT]?)</p> <p>0 None 1 Few activities (less than 10%) 2 Some activities (approx 20-30%) 3 Many activities (approx 50-60%) 4 Most or all activities (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How strong was your loss of interest? (Would you enjoy [ACTIVITIES] once you got started?)</p> <p>0 No loss of interest 1 Mild, slight loss of interest, probably would enjoy after starting activities 2 Moderate, definite loss of interest, but still has some enjoyment of activities 3 Severe, marked loss of interest in activities 4 Extreme, complete loss of interest, no longer participates in any activities</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p>Past week F I</p> <p>Past month F I Sx: Y N</p> <p>Lifetime F I Sx: Y N</p>
--	---	---

10. (C-5) feeling of detachment or estrangement from others

<p>Frequency Have you felt distant or cut off from other people? What was that like? How much of the time in the past month (week) have you felt that way? When did you first start to feel that way? (After the [EVENT]?)</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How strong were your feelings of being distant or cut off from others? (Who do you feel closest to? How many people do you feel comfortable talking with about personal things?)</p> <p>0 No feelings of detachment or estrangement 1 Mild, may feel "out of synch" with others 2 Moderate, feelings of detachment clearly present, but still feels some interpersonal connection 3 Severe, marked feelings of detachment or estrangement from most people, may feel close to only one or two people 4 Extreme, feels completely detached or estranged from others, not close with anyone</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p>Past week F I</p> <p>Past month F I Sx: Y N</p> <p>Lifetime F I Sx: Y N</p>
--	--	---

11. (C-6) restricted range of affect (e.g., unable to have loving feelings)

<p>Frequency Have there been times when you felt emotionally numb or had trouble experiencing feelings like love or happiness? What was that like? (What feelings did you have trouble experiencing?) How much of the time in the past month (week) have you felt that way? When did you first start having trouble experiencing (EMOTIONS)? (After the [EVENT]?)</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How much trouble did you have experiencing (EMOTIONS)? (What kinds of feelings were you still able to experience?) [INCLUDE OBSERVATIONS OF RANGE OF AFFECT DURING INTERVIEW]</p> <p>0 No reduction of emotional experience 1 Mild, slight reduction of emotional experience 2 Moderate, definite reduction of emotional experience, but still able to experience most emotions 3 Severe, marked reduction of experience of at least two primary emotions (e.g., love, happiness) 4 Extreme, completely lacking emotional experience</p> <p>QV (specify)</p> <hr/> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
--	---	--

12. (C-7) sense of a foreshortened future (e.g., does not expect to have a career, marriage, children, or a normal life span)

<p>Frequency Have there been times when you felt there is no need to plan for the future, that somehow your future will be cut short? Why is that? [RULE OUT REALISTIC RISKS SUCH AS LIFE-THREATENING MEDICAL CONDITIONS] How much of the time in the past month (week) have you felt that way? When did you first start to feel that way? (After the [EVENT]?)</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How strong was this feeling that your future will be cut short? (How long do you think you will live? How convinced are you that you will die prematurely?)</p> <p>0 No sense of a foreshortened future 1 Mild, slight sense of a foreshortened future 2 Moderate, sense of a foreshortened future definitely present, but no specific prediction about longevity 3 Severe, marked sense of a foreshortened future, may make specific prediction about longevity 4 Extreme, overwhelming sense of a foreshortened future, completely convinced of premature death</p> <p>QV (specify)</p> <hr/> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
--	--	--

Criterion D. Persistent symptoms of increased arousal (not present before the trauma), as indicated by two (or more) of the following:

13. (D-1) difficulty falling or staying asleep

<p>Frequency Have you had any problems falling or staying asleep? How often in the past month (week)? When did you first start having problems sleeping? (After the [EVENT]?)</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Sleep onset problems? Y N Mid-sleep awakening? Y N Early a.m. awakening? Y N Total # hrs sleep/night _____ Desired # hrs sleep/night _____</p>	<p>Intensity How much of a problem did you have with your sleep? (How long did it take you to fall asleep? How often did you wake up in the night? Did you often wake up earlier than you wanted to? How many total hours did you sleep each night?)</p> <p>0 No sleep problems 1 Mild, slightly longer latency, or minimal difficulty staying asleep (up to 30 minutes loss of sleep) 2 Moderate, definite sleep disturbance, clearly longer latency, or clear difficulty staying asleep (30-90 minutes loss of sleep) 3 Severe, much longer latency, or marked difficulty staying asleep (90 min to 3 hrs loss of sleep) 4 Extreme, very long latency, or profound difficulty staying asleep (> 3 hrs loss of sleep)</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p>Past week</p> <p>F I</p>
		<p>Past month</p> <p>F I</p> <p>Sx: Y N</p>
		<p>Lifetime</p> <p>F I</p> <p>Sx: Y N</p>

14. (D-2) irritability or outbursts of anger

<p>Frequency Have there been times when you felt especially irritable or showed strong feelings of anger? Can you give me some examples? How often in the past month (week)? When did you first start feeling that way? (After the [EVENT]?)</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong was your anger? (How did you show it?) [IF REPORTS SUPPRESSION:] (How hard was it for you to keep from showing your anger?) How long did it take you to calm down? Did your anger cause you any problems?</p> <p>0 No irritability or anger 1 Mild, minimal irritability, may raise voice when angry 2 Moderate, definite irritability or attempts to suppress anger, but can recover quickly 3 Severe, marked irritability or marked attempts to suppress anger, may become verbally or physically aggressive when angry 4 Extreme, pervasive anger or drastic attempts to suppress anger, may have episodes of physical violence</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p>Past week</p> <p>F I</p>
		<p>Past month</p> <p>F I</p> <p>Sx: Y N</p>
		<p>Lifetime</p> <p>F I</p> <p>Sx: Y N</p>

15. (D-3) difficulty concentrating

<p>Frequency Have you found it difficult to concentrate on what you were doing or on things going on around you? What was that like? How much of the time in the past month (week)? When did you first start having trouble concentrating? (After the [EVENT]?)</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How difficult was it for you to concentrate? [INCLUDE OBSERVATIONS OF CONCENTRATION AND ATTENTION IN INTERVIEW] How much did that interfere with your life?</p> <p>0 No difficulty with concentration 1 Mild, only slight effort needed to concentrate, little or no disruption of activities 2 Moderate, definite loss of concentration but could concentrate with effort, some disruption of activities 3 Severe, marked loss of concentration even with effort, marked disruption of activities 4 Extreme, complete inability to concentrate, unable to engage in activities</p> <p>QV (specify)</p> <p>_____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	--	--

16. (D-4) hypervigilance

<p>Frequency Have you been especially alert or watchful, even when there was no real need to be? (Have you felt as if you were constantly on guard?) Why is that? How much of the time in the past month (week)? When did you first start acting that way? (After the [EVENT]?)</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How hard did you try to be watchful of things going on around you? [INCLUDE OBSERVATIONS OF HYPERVIGILANCE IN INTERVIEW] Did your (HYPERVIGILANCE) cause you any problems?</p> <p>0 No hypervigilance 1 Mild, minimal hypervigilance, slight heightening of awareness 2 Moderate, hypervigilance clearly present, watchful in public (e.g., chooses safe place to sit in a restaurant or movie theater) 3 Severe, marked hypervigilance, very alert, scans environment for danger, exaggerated concern for safety of self/family/home 4 Extreme, excessive hypervigilance, efforts to ensure safety consume significant time and energy and may involve extensive safety/checking behaviors, marked watchfulness during interview</p> <p>QV (specify)</p> <p>_____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	--	--

17. (D-5) exaggerated startle response

<p>Frequency Have you had any strong startle reactions? When did that happen? (What kinds of things made you startle?) How often in the past month (week)? When did you first have these reactions? (After the [EVENT]?)</p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong were these startle reactions? (How strong were they compared to how most people would respond?) How long did they last?</p> <p>0 No startle reaction 1 Mild, minimal reaction 2 Moderate, definite startle reaction, feels "jumpy" 3 Severe, marked startle reaction, sustained arousal following initial reaction 4 Extreme, excessive startle reaction, overt coping behavior (e.g., combat veteran who "hits the dirt")</p> <p>QV (specify)</p> <p>_____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p>
		<p><i>Past month</i></p> <p>F I</p> <p>Sx: Y N</p>
		<p><i>Lifetime</i></p> <p>F I</p> <p>Sx: Y N</p>

Criterion E. Duration of the disturbance (symptoms in Criteria B, C, and D) is more than 1 month.

18. onset of symptoms

<p>[IF NOT ALREADY CLEAR:] When did you first start having (PTSD SYMPTOMS) you've told me about? (How long after the trauma did they start? More than six months?)</p>	<p>_____ <i>total # months delay in onset</i></p> <p>With delayed onset (> 6 months)? NO</p> <p>YES</p>
---	---

19. duration of symptoms

<p>[CURRENT] How long have these (PTSD SYMPTOMS) lasted altogether?</p>	<p><i>Duration more than 1 month?</i></p> <p><i>Total # months duration</i></p> <p><i>Acute (< 3 months) or chronic (> 3 months)?</i></p>	<p>Current</p> <p>NO YES</p> <p>_____</p>	<p>Lifetime</p> <p>NO YES</p> <p>_____</p>
<p>[LIFETIME] How long did these (PTSD SYMPTOMS) last altogether?</p>		<p>acute chronic</p>	<p>acute chronic</p>

Criterion F. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

20. subjective distress

<p>[CURRENT] Overall, how much have you been bothered by these (PTSD SYMPTOMS) you've told me about? [CONSIDER DISTRESS REPORTED ON EARLIER ITEMS]</p> <p>[LIFETIME] Overall, how much were you bothered by these (PTSD SYMPTOMS) you've told me about? [CONSIDER DISTRESS REPORTED ON EARLIER ITEMS]</p>	<p>0 None 1 Mild, minimal distress 2 Moderate, distress clearly present but still manageable 3 Severe, considerable distress 4 Extreme, incapacitating distress</p>	<p><i>Past week</i></p> <p>_____</p> <p><i>Past month</i></p> <p>_____</p> <p><i>Lifetime</i></p>

21. impairment in social functioning

<p>[CURRENT] Have these (PTSD SYMPTOMS) affected your relationships with other people? How so? [CONSIDER IMPAIRMENT IN SOCIAL FUNCTIONING REPORTED ON EARLIER ITEMS]</p> <p>[LIFETIME] Did these (PTSD SYMPTOMS) affect your social life? How so? [CONSIDER IMPAIRMENT IN SOCIAL FUNCTIONING REPORTED ON EARLIER ITEMS]</p>	0	No adverse impact	<i>Past week</i>
	1	Mild impact, minimal impairment in social functioning	—
	2	Moderate impact, definite impairment, but many aspects of social functioning still intact	<i>Past month</i>
	3	Severe impact, marked impairment, few aspects of social functioning still intact	
	4	Extreme impact, little or no social functioning	<i>Lifetime</i>
			—

22. impairment in occupational or other important area of functioning

<p>[CURRENT -- IF NOT ALREADY CLEAR] Are you working now?</p> <p>IF YES: Have these (PTSD SYMPTOMS) affected your work or your ability to work? How so? [CONSIDER REPORTED WORK HISTORY, INCLUDING NUMBER AND DURATION OF JOBS, AS WELL AS THE QUALITY OF WORK RELATIONSHIPS. IF PREMORBID FUNCTIONING IS UNCLEAR, INQUIRE ABOUT WORK EXPERIENCES BEFORE THE TRAUMA. FOR CHILD/ADOLESCENT TRAUMAS, ASSESS PRE-TRAUMA SCHOOL PERFORMANCE AND POSSIBLE PRESENCE OF BEHAVIOR PROBLEMS]</p> <p>IF NO: Have these (PTSD SYMPTOMS) affected any other important part of your life? [AS APPROPRIATE, SUGGEST EXAMPLES SUCH AS PARENTING, HOUSEWORK, SCHOOLWORK, VOLUNTEER WORK, ETC.] How so?</p> <p>[LIFETIME -- IF NOT ALREADY CLEAR] Were you working then?</p> <p>IF YES: Did these (PTSD SYMPTOMS) affect your work or your ability to work? How so? [CONSIDER REPORTED WORK HISTORY, INCLUDING NUMBER AND DURATION OF JOBS, AS WELL AS THE QUALITY OF WORK RELATIONSHIPS. IF PREMORBID FUNCTIONING IS UNCLEAR, INQUIRE ABOUT WORK EXPERIENCES BEFORE THE TRAUMA. FOR CHILD/ADOLESCENT TRAUMAS, ASSESS PRE-TRAUMA SCHOOL PERFORMANCE AND POSSIBLE PRESENCE OF BEHAVIOR PROBLEMS]</p> <p>IF NO: Did these (PTSD SYMPTOMS) affect any other important part of your life? [AS APPROPRIATE, SUGGEST EXAMPLES SUCH AS PARENTING, HOUSEWORK, SCHOOLWORK, VOLUNTEER WORK, ETC.] How so?</p>	0	No adverse impact	<i>Past week</i>
	1	Mild impact, minimal impairment in occupational/other important functioning	—
	2	Moderate impact, definite impairment, but many aspects of occupational/other important functioning still intact	<i>Past month</i>
	3	Severe impact, marked impairment, few aspects of occupational/other important functioning still intact	
	4	Extreme impact, little or no occupational/other important functioning	<i>Lifetime</i>
			—

Global Ratings

23. global validity

ESTIMATE THE OVERALL VALIDITY OF RESPONSES. CONSIDER FACTORS SUCH AS COMPLIANCE WITH THE INTERVIEW, MENTAL STATUS (E.G., PROBLEMS WITH CONCENTRATION, COMPREHENSION OF ITEMS, DISSOCIATION), AND EVIDENCE OF EFFORTS TO EXAGGERATE OR MINIMIZE SYMPTOMS.	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20px; text-align: center;">0</td> <td>Excellent, no reason to suspect invalid responses</td> </tr> <tr> <td style="text-align: center;">1</td> <td>Good, factors present that may adversely affect validity</td> </tr> <tr> <td style="text-align: center;">2</td> <td>Fair, factors present that definitely reduce validity</td> </tr> <tr> <td style="text-align: center;">3</td> <td>Poor, substantially reduced validity</td> </tr> <tr> <td style="text-align: center;">4</td> <td>Invalid responses, severely impaired mental status or possible deliberate "faking bad" or "faking good"</td> </tr> </table>	0	Excellent, no reason to suspect invalid responses	1	Good, factors present that may adversely affect validity	2	Fair, factors present that definitely reduce validity	3	Poor, substantially reduced validity	4	Invalid responses, severely impaired mental status or possible deliberate "faking bad" or "faking good"
0	Excellent, no reason to suspect invalid responses										
1	Good, factors present that may adversely affect validity										
2	Fair, factors present that definitely reduce validity										
3	Poor, substantially reduced validity										
4	Invalid responses, severely impaired mental status or possible deliberate "faking bad" or "faking good"										

24. global severity

ESTIMATE THE OVERALL SEVERITY OF PTSD SYMPTOMS. CONSIDER DEGREE OF SUBJECTIVE DISTRESS, DEGREE OF FUNCTIONAL IMPAIRMENT, OBSERVATIONS OF BEHAVIORS IN INTERVIEW, AND JUDGMENT REGARDING REPORTING STYLE.	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20px; text-align: center;">0</td> <td>No clinically significant symptoms, no distress and no functional impairment</td> <td style="text-align: center;"><i>Past week</i></td> </tr> <tr> <td style="text-align: center;">1</td> <td>Mild, minimal distress or functional impairment</td> <td style="text-align: center;">—</td> </tr> <tr> <td style="text-align: center;">2</td> <td>Moderate, definite distress or functional impairment but functions satisfactorily with effort</td> <td style="text-align: center;"><i>Past month</i></td> </tr> <tr> <td style="text-align: center;">3</td> <td>Severe, considerable distress or functional impairment, limited functioning even with effort</td> <td></td> </tr> <tr> <td style="text-align: center;">4</td> <td>Extreme, marked distress or marked impairment in two or more major areas of functioning</td> <td style="text-align: center;"><i>Lifetime</i></td> </tr> </table>	0	No clinically significant symptoms, no distress and no functional impairment	<i>Past week</i>	1	Mild, minimal distress or functional impairment	—	2	Moderate, definite distress or functional impairment but functions satisfactorily with effort	<i>Past month</i>	3	Severe, considerable distress or functional impairment, limited functioning even with effort		4	Extreme, marked distress or marked impairment in two or more major areas of functioning	<i>Lifetime</i>
0	No clinically significant symptoms, no distress and no functional impairment	<i>Past week</i>														
1	Mild, minimal distress or functional impairment	—														
2	Moderate, definite distress or functional impairment but functions satisfactorily with effort	<i>Past month</i>														
3	Severe, considerable distress or functional impairment, limited functioning even with effort															
4	Extreme, marked distress or marked impairment in two or more major areas of functioning	<i>Lifetime</i>														

25. global improvement

RATE TOTAL OVERALL IMPROVEMENT PRESENT SINCE THE INITIAL RATING. IF NO EARLIER RATING, ASK HOW THE SYMPTOMS ENDORSED HAVE CHANGED OVER THE PAST 6 MONTHS. RATE THE DEGREE OF CHANGE, WHETHER OR NOT, IN YOUR JUDGMENT, IT IS DUE TO TREATMENT.	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20px; text-align: center;">0</td> <td>Asymptomatic</td> </tr> <tr> <td style="text-align: center;">1</td> <td>Considerable improvement</td> </tr> <tr> <td style="text-align: center;">2</td> <td>Moderate improvement</td> </tr> <tr> <td style="text-align: center;">3</td> <td>Slight improvement</td> </tr> <tr> <td style="text-align: center;">4</td> <td>No improvement</td> </tr> <tr> <td style="text-align: center;">5</td> <td>Insufficient information</td> </tr> </table>	0	Asymptomatic	1	Considerable improvement	2	Moderate improvement	3	Slight improvement	4	No improvement	5	Insufficient information
0	Asymptomatic												
1	Considerable improvement												
2	Moderate improvement												
3	Slight improvement												
4	No improvement												
5	Insufficient information												

Current PTSD Symptoms

<i>Criterion A met (traumatic event)?</i>	NO	YES
# <i>Criterion B sx (> 1)?</i>	NO	YES
# <i>Criterion C sx (> 3)?</i>	NO	YES
# <i>Criterion D sx (> 2)?</i>	NO	YES
<i>Criterion E met (duration > 1 month)?</i>	NO	YES
<i>Criterion F met (distress/impairment)?</i>	NO	YES

CURRENT PTSD (Criteria A-F met)? NO YES

IF CURRENT PTSD CRITERIA ARE MET, SKIP TO ASSOCIATED FEATURES.

IF CURRENT CRITERIA ARE NOT MET, ASSESS FOR LIFETIME PTSD. IDENTIFY A PERIOD OF AT LEAST A MONTH SINCE THE TRAUMATIC EVENT IN WHICH SYMPTOMS WERE WORSE.

Since the (EVENT), has there been a time when these (PTSD SYMPTOMS) were a lot worse than they have been in the past month? When was that? How long did it last? (At least a month?)

IF MULTIPLE PERIODS IN THE PAST: **When were you bothered the most by these (PTSD SYMPTOMS)?**

IF AT LEAST ONE PERIOD, INQUIRE ITEMS 1-17, CHANGING FREQUENCY PROMPTS TO REFER TO WORST PERIOD: **During that time, did you (EXPERIENCE SYMPTOM)? How often?**

Lifetime PTSD Symptoms

<i>Criterion A met (traumatic event)?</i>	NO	YES
# <i>Criterion B sx (> 1)?</i>	NO	YES
# <i>Criterion C sx (> 3)?</i>	NO	YES
# <i>Criterion D sx (> 2)?</i>	NO	YES
<i>Criterion E met (duration > 1 month)?</i>	NO	YES
<i>Criterion F met (distress/impairment)?</i>	NO	YES

LIFETIME PTSD (Criteria A-F met)? NO YES

Associated Features

26. guilt over acts of commission or omission

<p>Frequency Have you felt guilty about anything you did or didn't do during (EVENT)? Tell me more about that. (What do you feel guilty about?) How much of the time have you felt that way in the past month (week)?</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%)</p> <p>Description/Examples</p>	<p>Intensity How strong were these feelings of guilt? How much distress or discomfort did they cause?</p> <p>0 No feelings of guilt 1 Mild, slight feelings of guilt 2 Moderate, guilt feelings definitely present, some distress but still manageable 3 Severe, marked feelings of guilt, considerable distress 4 Extreme, pervasive feelings of guilt, self-condemnation regarding behavior, incapacitating distress</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i></p> <p>F</p> <p>I</p>
		<p><i>Past month</i></p> <p>F</p> <p>I</p> <p>Sx: Y N</p>
		<p><i>Lifetime</i></p> <p>F</p> <p>I</p> <p>Sx: Y N</p>

27. survivor guilt [APPLICABLE ONLY IF MULTIPLE VICTIMS]

<p>Frequency Have you felt guilty about surviving (EVENT) when others did not? Tell me more about that. (What do you feel guilty about?) How much of the time have you felt that way in the past month (week)?</p> <p>0 None of the time 1 Very little of the time (less than 10%) 2 Some of the time (approx 20-30%) 3 Much of the time (approx 50-60%) 4 Most or all of the time (more than 80%) 8 N/A</p> <p>Description/Examples</p>	<p>Intensity How strong were these feelings of guilt? How much distress or discomfort did they cause?</p> <p>0 No feelings of guilt 1 Mild, slight feelings of guilt 2 Moderate, guilt feelings definitely present, some distress but still manageable 3 Severe, marked feelings of guilt, considerable distress 4 Extreme, pervasive feelings of guilt, self-condemnation regarding survival, incapacitating distress</p> <p>QV (specify)</p> <hr/>	<p><i>Past week</i></p> <p>F</p> <p>I</p>
		<p><i>Past month</i></p> <p>F</p> <p>I</p> <p>Sx: Y N</p>
		<p><i>Lifetime</i></p> <p>F</p> <p>I</p> <p>Sx: Y N</p>

28. a reduction in awareness of his or her surroundings (e.g., “being in a daze”)

<p>Frequency Have there been times when you felt out of touch with things going on around you, like you were in a daze? What was that like? [DISTINGUISH FROM FLASHBACK EPISODES] How often has that happened in the past month (week)? [IF NOT CLEAR:] <i>(Was it due to an illness or the effects of drugs or alcohol?)</i> When did you first start feeling that way? <i>(After the [EVENT]?)</i></p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong was this feeling of being out of touch or in a daze? <i>(Were you confused about where you actually were or what you were doing at the time?)</i> How long did it last? What did you do while this was happening? <i>(Did other people notice your behavior? What did they say?)</i></p> <p>0 No reduction in awareness 1 Mild, slight reduction in awareness 2 Moderate, definite but transient reduction in awareness, may report feeling “spacy” 3 Severe, marked reduction in awareness, may persist for several hours 4 Extreme, complete loss of awareness of surroundings, may be unresponsive, possible amnesia for the episode (blackout)</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
---	---	--

29. derealization

<p>Frequency Have there been times when things going on around you seemed unreal or very strange and unfamiliar? [IF NO:] <i>(What about times when people you knew suddenly seemed unfamiliar?)</i> What was that like? How often has that happened in the past month (week)? [IF NOT CLEAR:] <i>(Was it due to an illness or the effects of drugs or alcohol?)</i> When did you first start feeling that way? <i>(After the [EVENT]?)</i></p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong was (DEREALIZATION)? How long did it last? What did you do while this was happening? <i>(Did other people notice your behavior? What did they say?)</i></p> <p>0 No derealization 1 Mild, slight derealization 2 Moderate, definite but transient derealization 3 Severe, considerable derealization, marked confusion about what is real, may persist for several hours 4 Extreme, profound derealization, dramatic loss of sense of reality or familiarity</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p><i>Past week</i></p> <p>F I</p> <p><i>Past month</i></p> <p>F I Sx: Y N</p> <p><i>Lifetime</i></p> <p>F I Sx: Y N</p>
--	--	--

30. depersonalization

<p>Frequency Have there been times when you felt as if you were outside of your body, watching yourself as if you were another person? [IF NO:] <i>(What about times when your body felt strange or unfamiliar to you, as if it had changed in some way?)</i> What was that like? How often has that happened in the past month (week)? [IF NOT CLEAR:] <i>(Was it due to an illness or the effects of drugs or alcohol?)</i> When did you first start feeling that way? <i>(After the [EVENT]?)</i></p> <p>0 Never 1 Once or twice 2 Once or twice a week 3 Several times a week 4 Daily or almost every day</p> <p>Description/Examples</p>	<p>Intensity How strong was (DEPERSONALIZATION)? How long did it last? What did you do while this was happening? <i>(Did other people notice your behavior? What did they say?)</i></p> <p>0 No depersonalization 1 Mild, slight depersonalization 2 Moderate, definite but transient depersonalization 3 Severe, considerable depersonalization, marked sense of detachment from self, may persist for several hours 4 Extreme, profound depersonalization, dramatic sense of detachment from self</p> <p>QV (specify) _____</p> <p>Trauma-related? 1 definite 2 probable 3 unlikely Current _____ Lifetime _____</p>	<p>Past week F I</p> <p>Past month F I Sx: Y N</p> <p>Lifetime F I Sx: Y N</p>
---	---	---

CAPS SUMMARY SHEET

Name: _____ ID#: _____ Interviewer: _____ Study: _____ Date: _____

A. Traumatic event:

B. Reexperiencing symptoms	PAST WEEK			PAST MONTH			LIFETIME		
	Freq	Int	F+I	Freq	Int	F+I	Freq	Int	F+I
(1) intrusive recollections									
(2) distressing dreams									
(3) acting or feeling as if event were recurring									
(4) psychological distress at exposure to cues									
(5) physiological reactivity on exposure to cues									
<i>B subtotals</i>									
<i>Number of Criterion B symptoms (need 1)</i>									

C. Avoidance and numbing symptoms	PAST WEEK			PAST MONTH			LIFETIME		
	Freq	Int	F+I	Freq	Int	F+I	Freq	Int	F+I
(6) avoidance of thoughts or feelings									
(7) avoidance of activities, places, or people									
(8) inability to recall important aspect of trauma									
(9) diminished interest in activities									
(10) detachment or estrangement									
(11) restricted range of affect									
(12) sense of a foreshortened future									
<i>C subtotals</i>									
<i>Number of Criterion C symptoms (need 3)</i>									

D. Hyperarousal symptoms	PAST WEEK			PAST MONTH			LIFETIME		
	Freq	Int	F+I	Freq	Int	F+I	Freq	Int	F+I
(13) difficulty falling or staying asleep									
(14) irritability or outbursts of anger									
(15) difficulty concentrating									
(16) hypervigilance									
(17) exaggerated startle response									
<i>D subtotals</i>									
<i>Number of Criterion D symptoms (need 2)</i>									

Total Freq, Int, and Severity (F+I)	PAST WEEK			PAST MONTH			LIFETIME		
	Freq	Int	F+I	Freq	Int	F+I	Freq	Int	F+I
<i>Sum of subtotals (B+C+D)</i>									

E. Duration of disturbance	CURRENT		LIFETIME	
(19) duration of disturbance at least one month	NO	YES	NO	YES

F. Significant distress or impairment in functioning	PAST WEEK		PAST MONTH		LIFETIME	
(20) subjective distress						
(21) impairment in social functioning						
(22) impairment in occupational functioning						
<i>AT LEAST ONE > 2?</i>	NO	YES	NO	YES	NO	YES

PTSD diagnosis	CURRENT		LIFETIME	
<i>PTSD PRESENT -- ALL CRITERIA (A-F) MET?</i>	NO	YES	NO	YES
<i>Specify:</i>				
(18) with delayed onset (> 6 months delay)	NO	YES	NO	YES
(19) acute (< 3 months) or chronic (> 3 months)	acute	chronic	acute	chronic

<i>Global ratings</i>	<i>PAST WEEK</i>	<i>PAST MONTH</i>	<i>LIFETIME</i>
(23) global validity			
(24) global severity			
(25) global improvement			

<i>Associated features</i>	<i>PAST WEEK</i>			<i>PAST MONTH</i>			<i>LIFETIME</i>		
	<i>Freq</i>	<i>Int</i>	<i>F+I</i>	<i>Freq</i>	<i>Int</i>	<i>F+I</i>	<i>Freq</i>	<i>Int</i>	<i>F+I</i>
(26) guilt over acts of commission or omission									
(27) survivor guilt									
(28) reduction in awareness of surroundings									
(29) derealization									
(30) depersonalization									

Appendix E

HAMILTON ANXIETY RATING SCALE (HAM-A)

HAMILTON ANXIETY RATING SCALE (HAM- A)

Patient Information										
Patient	Date			Time			Hour		Min	
				Day	Mth.	Year				
Personal notes										

1. Anxious mood	
This item covers the emotional condition of uncertainty about the future, ranging from worry, insecurity, irritability and apprehension to overpowering dread.	
0 – The patient is neither more or less insecure or irritable than usual.	<input type="checkbox"/>
1 – Doubtful whether the patient is more insecure or irritable than usual.	<input type="checkbox"/>
2 – The patient expresses more clearly to be in a state of anxiety, apprehension or irritability, which he may find difficult to control. However, the worrying still is about minor matters and thus without influence on the patient's daily life.	<input type="checkbox"/>
3 – At times the anxiety or insecurity is more difficult to control because the worrying is about major injuries or harms which might occur in the future. Has occasionally interfered with the patient's daily life.	<input type="checkbox"/>
4 – The feeling of dread is present so often that it markedly interferes with the patient's daily life.	<input type="checkbox"/>

2. Tension

This item includes inability to relax, nervousness, bodily tensions, trembling and restless fatigue.

0 – The patient is neither more nor less tense than usual	<input type="checkbox"/>
1 – The patient seems somewhat more nervous and tense than usual.	<input type="checkbox"/>
2 – Patient expresses clearly unable to relax and full of inner unrest, which he finds difficult to control, but it is still without influence on the patient's daily life.	<input type="checkbox"/>
3 – The inner unrest and nervousness is so intense or frequent that it occasionally interferes with the patient's daily work.	<input type="checkbox"/>
4 – Tensions and unrest interfere with the patient's life and work at all times.	<input type="checkbox"/>

3. Fears

This item includes fear of being in a crowd, of animals, of being in public places, of being alone, of traffic, of strangers, of dark etc. It is important to note whether there has been more phobic anxiety during the present episode than usual.

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present.	<input type="checkbox"/>
2 – The patient experiences phobic anxiety but is able to fight it.	<input type="checkbox"/>
3 – It is difficult to fight or overcome the phobic anxiety, which thus to some extent interferes with the patient's daily life and work.	<input type="checkbox"/>
4 – The phobic anxiety clearly interferes with the patient's daily life and work.	<input type="checkbox"/>

4. Insomnia

This item covers the patient's subjective experience of sleep duration and sleep depth during the three preceding nights. Note: Administration of hypnotics or sedatives is disregarded

0 – Usual sleep duration and sleep depth	<input type="checkbox"/>
1 – Sleep duration is doubtfully or slightly reduced (e.g. due to difficulties falling asleep), but no change in sleep depth.	<input type="checkbox"/>
2 – Sleep depth is also reduced, sleep being more superficial. Sleep as a whole is somewhat disturbed.	<input type="checkbox"/>
3 – Sleep duration and sleep depth is markedly changed. Sleep periods total only a few hours per 24 hours.	<input type="checkbox"/>
4 – Sleep depth is so shallow that the patient speaks of short periods of slumber or dozing, but no real sleep.	<input type="checkbox"/>

5. Difficulties in concentration and memory

This item covers difficulties in concentration, making decision about everyday matters, and memory

0 – The patient has neither more nor less difficulty in concentration and/or memory that usual.	<input type="checkbox"/>
1 – Doubtful whether the patient has difficulty in concentration and/or memory.	<input type="checkbox"/>
2 – Even with a major effort it is difficult for the patient to concentrate on his daily routine work.	<input type="checkbox"/>
3 – The patient has pronounced difficulties with concentration, memory, or decision making, e.g. in reading a newspaper article or watching a television programme to the end.	<input type="checkbox"/>
4 – During the interview the patient shows difficulty in concentration, memory or decision making.	<input type="checkbox"/>

6. Depressed mood

This item covers both the verbal and the non-verbal communication of sadness, depression, despondency, helplessness and hopelessness

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether the patient is more despondent or sad than usual, or is only vaguely so.	<input type="checkbox"/>
2 – The patient is more clearly concerned with unpleasant experiences, although he still lacks helplessness or hopelessness.	<input type="checkbox"/>
3 – The patient shows clear non-verbal signs of depression and/or hopelessness.	<input type="checkbox"/>
4 – The patient remarks on despondency and helplessness or the non-verbal signs dominate the interview and the patient cannot be distracted.	<input type="checkbox"/>

7. General somatic symptoms: Muscular

Weakness, stiffness, soreness or real pain, more or less diffusely localized in the muscles, such as jaw ache or neck ache.

0 – The patient is neither more nor less sore or stiff in the muscles than usual.	<input type="checkbox"/>
1 – The patient seems somewhat more stiff or sore in the muscles than usual.	<input type="checkbox"/>
2 – The symptoms have the character of pain.	<input type="checkbox"/>
3 – Muscle pain interferes to some extent with the patient's daily work and life.	<input type="checkbox"/>
4 – Muscle pain is present most of the time and clearly interferes with the patient's daily work and life.	<input type="checkbox"/>

8. General somatic symptoms: Sensory

This item includes increased fatigability and weakness or real functional disturbances of the senses, including tinnitus, blurring of vision, hot and cold flashes and prickling sensations

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether the patient's indications of symptoms are more pronounced than usual	<input type="checkbox"/>
2 – The sensations of pressure reach the character of buzzing in the ears, visual disturbances and prickling or itching sensations in the skin.	<input type="checkbox"/>
3 – The generalized sensory symptoms interfere to some extent with the patient's daily life and work.	<input type="checkbox"/>
4 – The generalized sensory symptoms are present most of the time and clearly interfere with the patient's daily life and work.	<input type="checkbox"/>

9. Cardiovascular symptoms

This item includes tachycardia, palpitations, oppression, chest pain, throbbing in the blood vessels, and feelings of faintness.

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present.	<input type="checkbox"/>
2 – Cardiovascular symptoms are present, but the patient can still control them.	<input type="checkbox"/>
3 – The patient has occasional difficulty controlling the cardiovascular symptoms, which thus to some extent interfere with his daily life and work.	<input type="checkbox"/>
4 – Cardiovascular symptoms are present most of the time and clearly interfere with the patient's daily life and work.	<input type="checkbox"/>

10. Respiratory symptoms

Feelings of constriction or contraction in throat or chest, dyspnoea or choking sensations and sighing respiration

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present.	<input type="checkbox"/>
2 – Respiratory symptoms are present, but the patient can still control them.	<input type="checkbox"/>
3 – The patient has occasional difficulty controlling the respiratory symptoms, which thus to some extent interfere with his daily life and work.	<input type="checkbox"/>
4 – Respiratory symptoms are present most of the time and clearly interfere with the patient's daily life and work.	<input type="checkbox"/>

11. Gastro-intestinal symptoms

This item covers difficulties in swallowing, "sinking" sensation in stomach, dyspepsia (heartburn or burning sensation in the stomach, abdominal pains related to meals, fullness, nausea and vomiting), abdominal rumbling and diarrhoea.

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present (or doubtful whether different from usual).	<input type="checkbox"/>
2 – One or more gastro-intestinal symptoms are present, but the patient can still control them.	<input type="checkbox"/>
3 – The patient has occasional difficulty controlling the gastro-intestinal symptoms, which to some extent interfere with his daily life and work.	<input type="checkbox"/>
4 – The gastro-intestinal symptoms are present most of the time and interfere clearly with the patient's daily life and work.	<input type="checkbox"/>

12. Genito-urinary symptoms

This item includes non-organic or psychic symptoms such as frequent or more pressing passing of urine, menstrual irregularities, anorgasmia, dyspareunia, premature ejaculation, loss of erection.

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present (or doubtful whether different from usual).	<input type="checkbox"/>
2 – One or more genito-urinary symptoms are present, but do not interfere with the patient's daily life and work.	<input type="checkbox"/>
3 – Occasionally, one or more genito-urinary symptoms are present to such a degree that they interfere to some extent with the patient's daily life and work.	<input type="checkbox"/>
4 – The genito-urinary symptoms are present most of the time and interfere clearly with the patient's daily life and work.	<input type="checkbox"/>

13. Other autonomic symptoms

This item includes dryness of the mouth, blushing or pallor, sweating and dizziness

0 – Not present.	<input type="checkbox"/>
1 – Doubtful whether present.	<input type="checkbox"/>
2 – One or more autonomic symptoms are present, but they do not interfere with the patient's daily life and work.	<input type="checkbox"/>
3 – Occasionally, one or more autonomic symptoms are present to such a degree that they interfere to some extent with the patient's daily life and work.	<input type="checkbox"/>
4 – Autonomic symptoms are present most of the time and clearly interfere with the patient's daily life and work.	<input type="checkbox"/>

14. Behaviour during interview

The patient may appear tense, nervous, agitated, restless, tremulous, pale, hyperventilating or sweating during the interview. Based on such observations a global estimate is made.

0 – The patient does not appear anxious.	<input type="checkbox"/>
1 – It is doubtful whether the patient is anxious.	<input type="checkbox"/>
2 – The patient is moderately anxious.	<input type="checkbox"/>
3 – The patient is markedly anxious.	<input type="checkbox"/>
4 – Patient is overwhelmed by anxiety, for example with shaking and trembling all over.	<input type="checkbox"/>

Total score

HAM-A score level of anxiety

<17: mild

18 – 24: mild to moderate

25 – 30: moderate to severe

Appendix F

Montgomery-Åsberg Depression Rating Scale (MADRS)

Montgomery-Åsberg Depression Rating Scale (MADRS)

1. Apparent sadness

Representing despondency, gloom and despair (more than just ordinary transient low spirits), reflected in speech, facial expression, and posture. Rate by depth and inability to brighten up.

0 = No sadness.	<input type="checkbox"/>
2 = Looks dispirited but does brighten up without difficulty.	<input type="checkbox"/>
4 = Appears sad and unhappy most of the time.	<input type="checkbox"/>
6 = Looks miserable all the time. Extremely despondent	<input type="checkbox"/>

2. Reported sadness

Representing reports of depressed mood, regardless of whether it is reflected in appearance or not. Includes low spirits, despondency or the feeling of being beyond help and without hope.

0 = Occasional sadness in keeping with the circumstances.	<input type="checkbox"/>
2 = Sad or low but brightens up without difficulty.	<input type="checkbox"/>
4 = Pervasive feelings of sadness or gloominess. The mood is still influenced by external circumstances.	<input type="checkbox"/>
6 = Continuous or unvarying sadness, misery or despondency.	<input type="checkbox"/>

3. Inner tension

Representing feelings of ill-defined discomfort, edginess, inner turmoil, mental tension mounting to either panic, dread or anguish. Rate according to intensity, frequency, duration and the extent of reassurance called for.

0 = Placid. Only fleeting inner tension.	<input type="checkbox"/>
2 = Occasional feelings of edginess and ill-defined discomfort.	<input type="checkbox"/>
4 = Continuous feelings of inner tension or intermittent panic which the patient can only master with some difficulty.	<input type="checkbox"/>
6 = Unrelenting dread or anguish. Overwhelming panic.	<input type="checkbox"/>

4. Reduced sleep

Representing the experience of reduced duration or depth of sleep compared to the subject's own normal pattern when well.

0 = Sleeps as normal.	<input type="checkbox"/>
2 = Slight difficulty dropping off to sleep or slightly reduced, light or fitful sleep.	<input type="checkbox"/>
4 = Moderate stiffness and resistance	<input type="checkbox"/>
6 = Sleep reduced or broken by at least 2 hours.	<input type="checkbox"/>

5. Reduced appetite

Representing the feeling of a loss of appetite compared with when-well. Rate by loss of desire for food or the need to force oneself to eat.

0 = Normal or increased appetite.	<input type="checkbox"/>
2 = Slightly reduced appetite.	<input type="checkbox"/>
4 = No appetite. Food is tasteless.	<input type="checkbox"/>
6 = Needs persuasion to eat at all.	<input type="checkbox"/>

6. Concentration difficulties

Representing difficulties in collecting one's thoughts amounting to an incapacitating lack of concentration. Rate according to intensity, frequency, and degree of incapacity produced.

0 = No difficulties in concentrating.	<input type="checkbox"/>
2 = Occasional difficulties in collecting one's thoughts.	<input type="checkbox"/>
4 = Difficulties in concentrating and sustaining thought which reduced ability to read or hold a conversation.	<input type="checkbox"/>
6 = Unable to read or converse without great difficulty.	<input type="checkbox"/>

<p>7. Lassitude Representing difficulty in getting started or slowness in initiating and performing everyday activities.</p>	
<p>0 = Hardly any difficulty in getting started. No sluggishness.</p>	<input type="checkbox"/>
<p>2 = Difficulties in starting activities.</p>	<input type="checkbox"/>
<p>4 = Difficulties in starting simple routine activities which are carried out with effort.</p>	<input type="checkbox"/>
<p>6 = Complete lassitude. Unable to do anything without help.</p>	<input type="checkbox"/>

<p>8. Inability to feel Representing the subjective experience of reduced interest in the surroundings, or activities that normally give pleasure. The ability to react with adequate emotion to circumstances or people is reduced.</p>	
<p>0 = Normal interest in the surroundings and in other people.</p>	<input type="checkbox"/>
<p>2 = Reduced ability to enjoy usual interests.</p>	<input type="checkbox"/>
<p>4 = Loss of interest in the surroundings. Loss of feelings for friends and acquaintances.</p>	<input type="checkbox"/>
<p>6 = The experience of being emotionally paralyzed, inability to feel anger, grief or pleasure and a complete or even painful failure to feel for close relatives and friends.</p>	<input type="checkbox"/>

9. Pessimistic thoughts Representing thoughts of guilt, inferiority, self-reproach, sinfulness, remorse and ruin.	
0 = No pessimistic thoughts.	<input type="checkbox"/>
2 = Fluctuating ideas of failure, self-reproach or self- depreciation.	<input type="checkbox"/>
4 = Persistent self-accusations, or definite but still rational ideas of guilt or sin. Increasingly pessimistic about the future.	<input type="checkbox"/>
6 = Delusions of ruin, remorse or irredeemable sin. Self- accusations which are absurd and unshakable.	<input type="checkbox"/>

10. Suicidal thoughts Representing the feeling that life is not worth living, that a natural death would be welcome, suicidal thoughts, and preparations for suicide. Suicide attempts should not in themselves influence the rating.	
0 = Enjoys life or takes it as it comes.	<input type="checkbox"/>
2 = Weary of life. Only fleeting suicidal thoughts.	<input type="checkbox"/>
4 = Probably better off dead. Suicidal thoughts are common, and suicide is considered as a possible solution, but without specific plans or intention.	<input type="checkbox"/>
6 = Explicit plans for suicide when there is an opportunity. Active preparations for suicide.	<input type="checkbox"/>

Your Health and Well-Being

This survey asks for your views about your health. This information will help keep track of how you feel and how well you are able to do your usual activities. *Thank you for completing this survey!*

For each of the following questions, please mark an in the one box that best describes your answer.

1. In general, would you say your health is:

Excellent	Very good	Good	Fair	Poor
▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

2. Compared to one year ago, how would you rate your health in general now?

Much better now than one year ago	Somewhat better now than one year ago	About the same as one year ago	Somewhat worse now than one year ago	Much worse now than one year ago
▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

The following questions are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?

Yes, limited a lot	Yes, limited a little	No, not limited at all
▼	▼	▼

- a Vigorous activities, such as running, lifting heavy objects, participating in strenuous sports ₁ ₂ ₃
- b Moderate activities, such as moving a table, pushing a vacuum cleaner, bowling, or playing golf..... ₁ ₂ ₃
- c Lifting or carrying groceries..... ₁ ₂ ₃
- d Climbing several flights of stairs ₁ ₂ ₃
- e Climbing one flight of stairs ₁ ₂ ₃
- f Bending, kneeling, or stooping ₁ ₂ ₃
- g Walking more than a mile..... ₁ ₂ ₃
- h Walking several hundred yards..... ₁ ₂ ₃
- i Walking one hundred yards..... ₁ ₂ ₃
- j Bathing or dressing yourself..... ₁ ₂ ₃

SF-36v2™ Health Survey © 1996, 2000 by QualityMetric Incorporated and Medical Outcomes Trust. All Rights Reserved.
 SF-36® is a registered trademark of Medical Outcomes Trust.
 (SF-36v2 Standard, US Version 2.0)

3. During the past 4 weeks, how much of the time have you had any of the following problems with your work or other regular daily activities as a result of your physical health?

All of the time	Most of the time	Some of the time	A little of the time	None of the time
▼	▼	▼	▼	▼

- a Cut down on the amount of time you spent on work or other activities..... ₁ ₂ ₃ ₄ ₅
- b Accomplished less than you would like..... ₁ ₂ ₃ ₄ ₅
- c Were limited in the kind of work or other activities ₁ ₂ ₃ ₄ ₅
- d Had difficulty performing the work or other activities (for example, it took extra effort) ₁ ₂ ₃ ₄ ₅

4. During the past 4 weeks, how much of the time have you had any of the following problems with your work or other regular daily activities as a result of any emotional problems (such as feeling depressed or anxious)?

All of the time	Most of the time	Some of the time	A little of the time	None of the time
▼	▼	▼	▼	▼

- a Cut down on the amount of time you spent on work or other activities..... ₁ ₂ ₃ ₄ ₅
- b Accomplished less than you would like..... ₁ ₂ ₃ ₄ ₅
- c Did work or other activities less carefully than usual..... ₁ ₂ ₃ ₄ ₅

SF-36v2™ Health Survey © 1996, 2000 by QualityMetric Incorporated and Medical Outcomes Trust. All Rights Reserved.
 SF-36® is a registered trademark of Medical Outcomes Trust.
 (SF-36v2 Standard, US Version 2.0)

6. During the past 4 weeks, to what extent has your physical health or emotional problems interfered with your normal social activities with family, friends, neighbors, or groups?

Not at all	Slightly	Moderately	Quite a bit	Extremely
▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

7. How much bodily pain have you had during the past 4 weeks?

None	Very mild	Mild	Moderate	Severe	Very Severe
▼	▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅	<input type="checkbox"/> ₆

8. During the past 4 weeks, how much did pain interfere with your normal work (including both work outside the home and housework)?

Not at all	A little bit	Moderately	Quite a bit	Extremely
▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

9. These questions are about how you feel and how things have been with you during the past 4 weeks. For each question, please give the one answer that comes closest to the way you have been feeling. How much of the time during the past 4 weeks...

All of the time	Most of the time	Some of the time	A little of the time	None of the time
-----------------	------------------	------------------	----------------------	------------------



- a Did you feel full of life? ₁ ₂ ₃ ₄ ₅
- b Have you been very nervous? ₁ ₂ ₃ ₄ ₅
- c Have you felt so down in the dumps that nothing could cheer you up? ₁ ₂ ₃ ₄ ₅
- d Have you felt calm and peaceful? ₁ ₂ ₃ ₄ ₅
- e Did you have a lot of energy? ₁ ₂ ₃ ₄ ₅
- f Have you felt downhearted and depressed? ₁ ₂ ₃ ₄ ₅
- g Did you feel worn out? ₁ ₂ ₃ ₄ ₅
- h Have you been happy? ₁ ₂ ₃ ₄ ₅
- i Did you feel tired? ₁ ₂ ₃ ₄ ₅

10. During the past 4 weeks, how much of the time has your physical health or emotional problems interfered with your social activities (like visiting friends, relatives, etc.)?

All of the time	Most of the time	Some of the time	A little of the time	None of the time
▼	▼	▼	▼	▼
<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

SF-36v2™ Health Survey © 1996, 2000 by QualityMetric Incorporated and Medical Outcomes Trust. All Rights Reserved.
 SF-36® is a registered trademark of Medical Outcomes Trust.
 (SF-36v2 Standard, US Version 2.0)

11. How TRUE or FALSE is each of the following statements for you?

	Definitely true	Mostly true	Don't know	Mostly false	Definitely false
	▼	▼	▼	▼	▼
a I seem to get sick a little easier than other people.....	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
b I am as healthy as anybody I know	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
c I expect my health to get worse	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
d My health is excellent.....	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

THANK YOU FOR COMPLETING THESE QUESTIONS!

Challenges of Implementing a PTSD Telepsychiatry Project: A HOSPITAL PERSPECTIVE

PHOEBE PUTNEY MEMORIAL HOSPITAL


Study Description

PURPOSE:
The purpose of this project is to develop a telemental health services network for military personnel who have served in Iraq and Afghanistan, demonstrate early signs of PTSD, and now reside in geographically isolated communities of Southwest Georgia.

SIGNIFICANCE:
The significance of this study is its use of telemental health for military personnel in the community health-care provider setting.

HYPOTHESIS:
Cognitive Behavioral Therapy (CBT) administered face-to-face versus CBT administered via telemental health will demonstrate equivalent efficacy in reducing symptoms of PTSD among military personnel returning from operations in Iraq and Afghanistan.

STUDY DESIGN:
This study will follow a pre-test / post-test equivalency model. Subjects are randomized to receive CBT via standard face-to-face administration or via telemental health, with assessments before and after completion of therapy. Target sample size is 202, allowing for a 20% attrition rate.



Background

DESCRIPTION OF PTSD

- Post Traumatic Stress Disorder (PTSD) has been cited as a significant condition of many returning service personnel from Iraq and Afghanistan.
- The National Institutes of Mental Health describes PTSD as an anxiety disorder that can develop following actual or threatened physical harm. Treatment includes psychotherapy and/or medication.
- Treatment of returning military personnel will be challenging.
 - Expected significant increase in patient volume
 - Limited resources available for treatment due to increase in volume without an increase in providers or treatment options in this rural area
 - Many veterans reside or are located in rural or remote localities

REGIONAL DEMOGRAPHICS

- Southwest Georgia demonstrates a mostly rural demographic as defined by HPSA scores

Ashburn Clinic	17
Leshburg Clinics	16
Camilla Clinic	16
Sylvester Clinic	16
PPMH Main Campus	16
Behavioral Health Center	16
Phoebe Northwest	16
Phoebe East	16

Challenges for region in treatment of PTSD

- Limited access to education
- Limited access to mental health care in general


SOLUTION:
Mental health care for PTSD provided by increasing access to services via telemental health



Equipment and Network

- NUPHYSICA T-CART**
- Physically secure and mobile: ideal for use in busy exam rooms used for other purposes
 - Mobile camera that can be controlled remotely
 - Will be used by subjects and therapists for treatment sessions
- NUPHYSICA PHYSICIAN SMALL STUDIO**
- Desktop unit - stationary camera
 - Will be used in study by psychiatrists / raters for subject interaction

- CONNECTIVITY**
- Network backbone
 - WAN circuits
- SECURITY**
- Encryption
 - VPN
- QUALITY**
- Quality of Service
 - Bandwidth monitoring



Challenges

MET TO DATE



- Initiated assembly of telemental health research team
- Acquired and installed telemental health equipment
- Received approval from hosting organization IRB
- Network initiated and tested



YET TO OVERCOME

- Receive research review approval from Navy and Army
- Complete assembly of research team
- Market study to Southwest Georgia region
- Recruit and enroll first patient

Portions of this work are sponsored by the U.S. Dept. of the Army, under (Contract number W11XWH-09-2-0112). The information does not necessarily reflect the position of the government, and no official endorsement should be inferred. Please direct questions or comments to Steven Ziemba, Principal Investigator, Phoebe Putney Memorial Hospital, 427 West Third Avenue, Suite 100, Albany GA, 31701. (229) 312-0284, sziembai@ppmh.org

Appendix I

  Product Line Review (PLR) Meeting Behavioral Health 27 April 2010 1	<h3>Regional Center of Excellence for PTSD Phoebe Putney Memorial Hospital</h3> <p>Steven Ziemba Director of Clinical Research</p> <p>Period of Performance August 2009 - August 2011</p>
--	---

  Product Line Review (PLR) Meeting Behavioral Health 27 April 2010 2	<h3>Military Relevant Issue to be Solved</h3> <ul style="list-style-type: none">■ Studies have estimated that as many as 19% and 11% of soldiers returning from Iraq and Afghanistan, respectively, experience PTSD. Additional research (September 2009) conducted by the Naval Postgraduate School and Stanford University collectively project that rates of PTSD among veterans of Iraq and Afghanistan could potentially be as high as 35%.■ Specialty mental health services may be available only in urban referral centers and access to even routine mental health services may be limited to Veterans Affairs facilities. As recent as November 2009, the Washington Post reported approximately 408 psychiatrists for 545,000 troops.■ Travel distances for rural veterans who are remote from referral centers or parent facilities may restrict veterans' access to mental health services.
--	--



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

3

Solution

- The use of interactive videoconferencing to provide psychiatric services to geographically remote regions, often referred to as telemental health, is growing with both Medicare and TRICARE currently reimbursing providers for mental health care delivered via this method.
- TRICARE not only launched a Telemental health program in August 2009, but has also launched TRIAP involving providing beneficiaries access via the Internet to a licensed counselor (Association of the United States Army News, April 2010).
- Specific forms of mental health treatments and assessments, such as Cognitive Behavioral Therapy (CBT) delivered via telemental health, have been reported since 2000. Frueh and colleagues (November, 2007) conducted research showing that CBT delivered using telemental health equipment were effective in terms of therapist's adherence and competence.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

4

Solution

- Telemental health continues to show great promise and may be as effective as conventional face to face psychotherapy. ATA's Practice Guidelines for Video-Conferencing-Based Telemental Health (October 2009) cited 15 references showing that telemental health can impact quality, timeliness, and availability of services in a wide variety of mental health care settings.
- Further research in this area, considering the US Army's immediate need to provide psychiatric and mental health services to personnel in geographically remote regions, can assist in not only providing OIF/OEF veterans and active duty personnel with access to care, but also provide additional literature substantiating evidence-based telemental health practices.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

5

Project Description

- The overarching goal of this project is to demonstrate that Cognitive Behavioral Therapy (CBT) delivered via telemental health is equally effective as when delivered face-to-face, with the long-term goal of increasing access to mental health services for OIF/OEF veterans and active duty service members services who show early signs of PTSD, and now reside in geographically isolated communities.
- These services will be unique in that they will be offered by a civilian hospital at seven office based-rural clinics encompassing a 40 mile radius.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010


6

Project Description

NuPhysicia T-Cart



Physician Small Studio






Product Line Review (PLR) Meeting

Behavioral Health

27 April 2010

7

Project Description



Product Line Review (PLR) Meeting

Behavioral Health

27 April 2010

8

Validation Strategy

- **Hypothesis:** CBT (face-to-face administration) and CBT (telemental health administration) will be equally effective in reducing symptoms of PTSD among military personnel returning from operations in Iraq and Afghanistan.
- **Technical Objectives:**
 - Create an interactive, telemental health network at several office-based Phoebe Putney Memorial Hospital Primary Care Clinics.
 - Quantify the effectiveness of CBT (face-to-face administration) versus CBT (telemental health administration) in reducing symptoms of PTSD among military personnel returning from Iraq and Afghanistan.
 - The study follows pretest / post-test design to assess for equivalency between the two treatment regimens.
 - 150 subjects will be randomized to either the CBT (face-to-face administration) or CBT (telemental health administration) groups.




Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

9

Validation Strategy



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

10

Validation Strategy

- CBT will consist of one week of psycho-education, seven-eight weeks of cognitive restructuring, and one-two weeks of relapse prevention using Beck's CBT method containing collaborative structured agendas during each session and structured homework assignments for cognitive restructuring and relapse prevention.
- Subjects randomly assigned to the CBT Group (face-to-face administration) will receive CBT therapy from a licensed therapist trained in CBT 1X/week for a period up to 15 weeks.
- Subjects randomly assigned to the CBT Group (telemental health administration) will receive CBT from a licensed therapist trained in CBT 1X/week for a period up to 15 weeks.
- The 15-week time period was chosen to allow for cancellations, emergencies, and both therapist and patient vacation time. There is a 10-session maximum during this time period.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

11

Validation Strategy

- **Rater Measures:**

Frequency and intensity of individual symptoms will be assessed pre/post treatment:

- Clinician-Administered PTSD Scale (CAPS)
- Montgomery-Asberg Depression Rating Scale (MADRS)
- Hamilton Rating Scale for Anxiety (HAM-A)

- **Self-Report Measures:**

- PCL-M. This is a standardized evaluation tool used by the military to assess PTSD symptoms
- SF-36 Quality of Life Assessment
- Patient Satisfaction Questionnaire (post-treatment only) developed by Press Ganey and Associates

- **Fidelity:**

- Cognitive Therapy Scale-Revised (CTS-R) chosen on ability to measure fidelity of both adherence to CBT model and competence of therapist with model
- Consultant hired as fidelity rater to complete CTS-R on 15-minute audio segments of every session for each subject.
- Fidelity consultant trained at Beck Institute




Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

12

Research/Development Timeline





Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

13

Successes to Date

- PI response to AIBS review submitted and accepted
- Contract with ARMY awarded
- PPMH IRB approved
- Telemental health equipment purchased, installed at all sites, and configured
- Preliminary training of PPMH staff in telemental health equipment use



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

14

Challenges

- Obtaining DON HRPP Approval
- Obtaining Army OHRP approval.
- Hiring qualified, licensed mental health professionals in a rural area willing to travel for distances up to 80 miles round-trip for the face-to-face control group of subjects.
- Configuring study psychiatrists research schedules with their out-patient and in-patient schedules.
- Recruiting and providing therapy to 150 PTSD subjects.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

15

What's Next

- Submit research protocol back to Phoebe IRB for expedited review regarding study protocol revisions
- Submit research protocol to DON HRPP
- Submit research protocol to Army OHRP
- Planning and hosting marketing and Kick-off event for community
- Recruitment of study mental health staff and study subjects



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

16

Intellectual Property / Publications Deriving from this Project

List any Confidentiality Agreements - N/A

Patents Filed – N/A

List Invention Disclosures Submitted –N/A

List all Publications –N/A



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

17

Transition/ Business/ Marketing Plan

- Georgia is home to 12 active military installations with more than 100,000 Active Duty, Reservists and National Guard personnel and almost a million veterans and their dependents.
- 3100 Soldiers from The National Guard's 48th Infantry Brigade are returning back to GA from a one year deployment in Afghanistan.
- A marketing campaign will be developed that will target potentially rich referral sources such as primary care physicians, spouses and family members, clergy, local fraternal organizations, etc.
- A Community Kick-off event will be hosted to publicly announce to the community the implementation phase of the study and demonstrate use of the telemental health equipment.



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

18

Project Funding



Product Line
Review
(PLR)
Meeting

Behavioral
Health

27 April 2010

19

Additional Project Information

Lab/Company/Group: Phoebe Putney Memorial Hospital

Principal Investigator: Steven Ziemba, PhD(c)

Government COR: Jeanette Rasche, MS

Government Project Officer: Edward Kensinger

Contract Instrument: Cooperative Agreement

Period of Performance: AUG 2009--SEP-2011

Contract Specialist: Chris Helman

EDMS# : 3893

Contract #: W81XWH-09-2-0112