AD_____

AWARD NUMBER: W81XWH-08-1-0563

TITLE: Autocrine and Paracrine Hh Signaling Regulate Prostate Development

PRINCIPAL INVESTIGATOR: Min Yu, Ph.D.

CONTRACTING ORGANIZATION: University of Wisconsin, Madison Madison, WI 53715

REPORT DATE: September 201€

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

P			Form Approved				
Dublic reporting burden for this		wing instructions, econ	OMB No. 0704-0188				
Prubic reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202- 4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.							
1. REPORT DATE	2	2. REPORT TYPE		3. E	DATES COVERED		
September 201€	1	Annual Summary		11	August 200J – 10 August 201€		
4. TITLE AND SUBTIT	LE			5a.	CONTRACT NUMBER		
Autocrine and Para	acrine Hh Signaling	Regulate Prostate	Development	5b. W8	5b. GRANT NUMBER W81XWH-08-1-0563		
				5c.	PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d.	PROJECT NUMBER		
Min Yu, Ph.D.				5e.	TASK NUMBER		
Email: ^˘O∙˘¦*^¦^È â	8Èåĭ			5f. '	WORK UNIT NUMBER		
7 PERFORMING ORG	ANIZATION NAME(S)			8 6	FREORMING ORGANIZATION REPORT		
	,, , , , , , , , , , , , , , , , , , ,			N	IUMBER		
University of Wisco Madison, WI 5371	onsin, Madison 5						
9. SPONSORING / MC U.S. Army Medica Fort Detrick, Marvl	NITORING AGENCY N Research and Ma and 21702-5012	AME(S) AND ADDRES: teriel Command	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)		
				11.	SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT OEJ] ¦[ç^åÁų ¦ÁÚčà ã&ÁÜ^ ^æ•^LÁÖã:dãačqã} ÁW} ãĮãe^å Á							
13. SUPPLEMENTAR	YNOTES						
14. ABSTRACT Hedgehog (Hh) signaling has been reported to play an important role in prostate development and prostate cancer. Studies from several laboratories have shown that Hh ligands are expressed in the epithelium of the developing prostate; activate Hh target genes expressed in the surrounding mesenchyme and influence prostate ductal growth. However, publications from these different laboratories have reported conflicting effects on growth. One major goal of this research was to elucidate the reason for those conflicting data. We addressed this issue by: 1) Defining Hh mesenchymal responsive genes in fetal prostate by microarray; 2) Characterizing epithelial growth effects of Hh signaling pre- and postnatal in both in vitro explants culture and in vivo transgenic mice model; 3) Proposing the potential molecular mechanisms underlying this growth regulation by Hh signaling. Herein, we identify a total of 89 genes that are regulated by Hh signaling in UGSM-2 cell line, the immortalized mesenchymal cells from the E16 UGS mesenchyme. We observe stage-specific growth effects of Hh signaling. Specifically, Hh pathway activation stimulates epithelial proliferation prenatally, while inhibits epithelial proliferation postanatally. We propose this temporal growth effects is mediated by the discordant regulation of a subset of target genes by Hh signaling in the prenatal and postnatal prostate. In addition, we provide the evidence of autocrine Hh signaling in the developing prostate and identify its potential role in maintaining a progenitor cell population. The findings of this research resolve the long-term discrepancies on growth effects of Hh signaling in the developing prostate, and provide a novel interpretation of both autocrine and paracrine Hh signaling mechanism in regulating prostate development, which will offer critically important insight into the actions of Hh signaling in advanced prostate cancer							
15. SUBJECT TERMS							
Prostate, neogenog signaling, proliferation, morphogenesis, regeneration							
16. SECURITY CLASS	SIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC		
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	UU	2F	19b. TELEPHONE NUMBER (include area code)		

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. Z39.18

Table of Contents

<u>Page</u>

Introduction	4
Body	4-5
Key Research Accomplishments	5
Reportable Outcomes	5
Appendices	6-21

INTRODUCTION

Hedgehog (Hh) signaling has been reported to play an important role in prostate development and prostate cancer. Studies from several laboratories have shown that Hh ligands are expressed in the epithelium of the developing prostate; activate Hh target genes expressed in the surrounding mesenchyme and influence prostate ductal growth. However, publications from these different laboratories have reported conflicting effects on growth. One major goal of this research was to elucidate the reason for those conflicting data. We addressed this issue by: 1) Defining Hh mesenchymal responsive genes in fetal prostate by microarray; 2) Characterizing epithelial growth effects of Hh signaling pre- and postnatal in both in vitro explants culture and in vivo transgenic mice model; 3) Proposing the potential molecular mechanisms underlying this growth regulation by Hh signaling. Herein, we identify a total of 89 genes that are regulated by Hh signaling in UGSM-2 cell line, the immortalized mesenchymal cells from the E16 UGS mesenchyme. We observe stage-specific growth effects of Hh signaling. Specifically, Hh pathway activation stimulates epithelial proliferation prenatally, while inhibits epithelial proliferation postanatally. We propose this temporal growth effects is mediated by the discordant regulation of a subset of target genes by Hh signaling in the prenatal and postnatal prostate. In addition, we provide the evidence of autocrine Hh signaling in the developing prostate and identify its potential role in maintaining a progenitor cell population.

BODY

To examine the influence of dynamic interaction with the epithelium on the mesenchymal target gene response to Hh signaling, we investigated the target gene response in recombinant tissues composed of epithelium and mesenchyme from different stages of development and cultured in the presence or absence of AZ75. In order to selectively measure mesenchymal gene expression, we utilized the mesenchyme from the mouse and epithelium from the rat. The developmental stages used were mouse E16 UGS and the developmental stage equivalent rat E18 UGS and both mouse and rat P1 prostate. The expectation that recombinants of mouse mesenchyme and rat epithelium would recapitulate the molecular interactions of the developing mouse prostate is based upon the finding that such recombinants grafted under the renal capsule generate normal appearing prostate tissue with rat-derived prostate ductal epithelium and mouse-derived stromal element. Freshly separated mesenchyme and epithelium were recombined and cultured for seven days in testosterone supplemented media in the presence or absence of AZ75. The four tissue recombinants were: E16 mouse mesenchyme with either E18 or P1 rat epithelium and P1 mouse mesenchyme with either E18 or P1 rat epithelium. Hh target gene expression was assayed by RT-PCR using mouse target gene primers. Expression of the canonical target genes Gli1 and Ptc1 was strongly inhibited by AZ75 in all cultured tissue. This finding indicates both that robust Hh signaling was re-established in the tissue recombinants and was effectively inhibited by addition of AZ75 to the media. All of the five non-canonical Hh target genes assayed (Igfbp6, Igfbp3, Fbn2, Fgf5 and Agpt4) showed significantly regulated expression by Hh signaling in at least one of the recombinants. However, the canonical patterns of Hh regulated target gene expression in

the cultured E16 UGS and P1 prostate was not recapitulated in the cultured (M)E16m/(R)E18e and (M)P1m/(R)P1e recombinants, respectively. Only Agpt4 was significantly regulated by Hh signaling in the cultured (M)E16m/(R)E18e recombinant. Both Igfbp3 and Agpt4 were positively regulated by Hh signaling in the (M)P1m/(R)P1e recombinant but this contrasts with their negative regulation in the mouse P1 prostate. Hh regulated expression of several target genes was observed in the temporally heterotypic recombinants (M) 3 E16m/(R) P1e and (M) P1m/(R) E18e but this was inconsistent. These findings indicate that even though paracrine Hh signaling is re-established in the tissue recombinants, canonical patterns of positive and negative target gene regulation are only sporadically recapitulated in the temporally homotypic M)E16m/(R)E18e and (M)P1m/(R)P1e recombinants. While we cannot discount the influence of species differences, these data suggest that the physical disruption associated with mesenchymal-epithelial separation and recombination perturbs stage-specific patterns of Hh target gene regulation determined by the architecture of the developing tissue and the mesenchymal cell microenvironment.

KEY RESEARCH ACCOMPLISHMENTS

1. We found that the paracrine Hh signaling affect the prostate morphogenesis in a stagespecific

way.

2. Our gene analysis studies on primary culture mesenchymal cells suggested that the mesenchymal character is responsible but not sufficient for the discordant response of those genes.

REPORTALBLE OUTCOMES

• We published one paper on JBC journal. Yu M, Gipp J, Yoon JW, Iannaccone P, Walterhouse D, Bushman W. Sonic Hedgehog-responsive Genes in the Fetal Prostate. J Biol Chem. 2009 Feb 27; 284(9): 5620-9.

• Abstract presentations:

o Min Yu, Wade Bushman. Molecular and Environmental Toxicology Center, University of Wisconsin-Madison.Hedgehog signaling regulates prostate development: interplay of paracrine and autocrine mechanism. (Platform presentation presented at 2009- 48th annual meeting of SOT in Baltimore by Min Yu)

o Min Yu, Bushman Wade. Molecular and Environmental Toxicology Center, University of Wisconsin-Madison. The study of hedgehog signaling in prostate development: a role of both paracrine and autocrine mechanism. (Poster presented at 2009- Annual meeting of AUA in Chicago by Min Yu) Supplemental Material can be found at: http://www.jbc.org/content/suppl/2008/12/22/M809172200.DC1.html

> THE JOURNAL OF BIOLOGICAL CHEMISTRY VOL. 284, NO. 9, pp. 5620–5629, February 27, 2009 © 2009 by The American Society for Biochemistry and Molecular Biology, Inc. Printed in the U.S.A.

Sonic Hedgehog-responsive Genes in the Fetal Prostate*S

Received for publication, December 5, 2008, and in revised form, December 15, 2008 Published, JBC Papers in Press, December 18, 2008, DOI 10.1074/jbc.M809172200

Min Yu^{+§}, Jerry Gipp[§], Joon Won Yoon[¶], Phillip Iannaccone[¶], David Walterhouse[¶], and Wade Bushman^{§1}

From the [‡]University of Wisconsin Medical School, Molecular and Environmental Toxicology Center, Madison, Wisconsin 53705, the [§]Department of Urology, Clinical Sciences Center, University of Wisconsin Medical School, Madison, Wisconsin 53792, and the [¶]Northwestern University Feinberg School of Medicine and the Children's Memorial Research Center, Chicago, Illinois 60614

The Hedgehog (Hh) signaling pathway plays an important role in prostate development and appears to play an equally important role in promoting growth of advanced prostate cancer. During prostate development, epithelial cells in the urogenital sinus (UGS) express Sonic Hedgehog (Shh) and secrete Shh peptide. The secreted Hh peptide acts on adjacent mesenchymal cells to activate the Hh signal transduction pathway and elicit paracrine effects on epithelial proliferation and differentiation. To identify mesenchymal targets of Shh signaling, we performed microarray analysis on a Shh-responsive, immortalized urogential sinus mesenchymal cell line. We found 68 genes that were up-regulated by Shh and 21 genes that were down-regulated. Eighteen of those were selected for further study with Ptc1 and Gli1 serving as reference controls. We found 10 of 18 were also Hh-regulated in primary UGS mesenchymal cells and 13 of 18 in the cultured UGS. Seven of 18 exhibited Shh-regulated expression in both assays (Igfbp-6, Igfbp-3, Fbn2, Ntrk3, Agpt4, Dmp1, and Mmp13). Three of the 18 genes contained putative Gli binding motifs that bound Gli1 peptide in electrophoretic mobility shift assays. With the exception of Tiam1, target gene expression generally showed no differences in the concentration dependence of ligand-induced expression, but we observed strikingly different responses to direct pathway activation by transfection with activated Smo, Gli1, and Gli2.

Sonic Hedgehog $(Shh)^2$ is a secreted signaling protein that regulates epithelial-mesenchymal interactions during embryonic development in a variety of organs (1). During fetal prostate development, Shh is expressed and secreted by the epithelium of the urogenital sinus (UGS) and acts on the surrounding mesenchyme (2–5). Binding of Hh ligand to its receptor Patched 1 (Ptc1), a 12-span transmembrane protein, activates an intracellular signal transduction mechanism involving a second transmembrane protein Smoothened (Smo), and results in changes in transcriptional regulation of specific target genes through the coordinated activities of three highly related Gli transcription factors (Gli1, Gli2, and Gli3) (6). The transcriptional changes not only affect proliferation and differentiation of UGS mesenchyme but elicit paracrine signals that regulate proliferation and differentiation of the adjacent UGS epithelium (3, 7–9). Emerging evidence suggests that proper spatial and temporal patterns of Hh signaling are required for normal prostate ductal development (1, 10).

Although the diversity of Shh effects is clear, molecules that function downstream of Shh in different contexts remains incompletely characterized. Gli1, Ptc1, and Hedgehog interacting protein are the canonical targets of Hh signaling expressed in nearly all cell types examined to date (11). However, other target genes regulated by the Shh signaling pathway likely vary between tissues and cell types, influenced by the state of differentiation and the presence or absence of co-regulators (1, 12). A number of target genes have been described. The actin-binding protein Missing in Metastasis (MIM) has been identified as a Shh-responsive gene that potentiates Gli-dependent transcriptional activation in skin development and tumorigenesis (13). The forkhead transcription factor Foxe1 was established as a downstream target of the Shh pathway in hair follicle morphogenesis (14). Other Hh target genes include Foxa2 (Hnf-3 β) and Coup-Tfii during floor plate development (15, 16), Foxd2 (Mf-2) during somitogenesis (17), Foxm1 in basal cell carcinoma (18), and Foxf1 (Freac-1 or Hfh-8) during lung and foregut organogenesis (19). Other identified target genes include SFRP1, SFRP2, Igfbp-6, and Sil (1, 20-23). For the developing prostate, however, the only target gene identified thus far is Igfbp-6 (24).

We performed a microarray analysis to identify Shh-induced transcriptional changes in a previously characterized, immortalized UGS mesenchymal cell line, UGSM-2 (9). We identified a combination of previously described and novel Shh targets, some of which appear to be prostate selective. The novel Shh-responsive genes provide insight into the molecular mechanism of Shh-regulated prostate development, suggesting roles in cell growth regulation as well as in providing cross-talk with the Wnt and Notch signaling pathways.

EXPERIMENTAL PROCEDURES

Microarray Cell Growth and RNA Isolation—UGSM-2 cells, an immortalized mouse E16 prostate mesenchymal cell lines (10), were set at confluence and allowed to attach overnight in Dulbecco's modified Eagle's medium/F-12 containing 10% fetal bovine serum. The next day cells were treated with Shh (R&D Systems 1845-SH/CF) or vehicle in Dulbecco's modified Eagle's

ASBMB

^{*} This work was supported, in whole or in part, by National Institutes of Health Grants P50 DK065303 and RO1 DK056238. This work was also supported by the Robert and Delores Schnoes Chair in Urologic Research. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

S The on-line version of this article (available at http://www.jbc.org) contains supplemental Table S1 and Figs. S1–S3.

¹ To whom correspondence should be addressed: K6/562 CSC, 600 Highland Ave., Madison, WI 53792. Fax: 608-265-8133; E-mail: bushman@ surgery.wisc.edu.

² The abbreviations used are: Shh, Sonic Hedgehog; UGS, urogential sinus; MES, 4-morpholineethanesulfonic acid; Ptc1, Patched 1; MEF, mouse embryonic fibroblast; Smo, smoothened; PKA, protein kinase A; IGFBP, insulin-like growth factor binding protein.

medium/F-12 containing 0.1% fetal bovine serum. Following 24 h treatment total RNA was isolated using the Qiagen RNeasy RNA isolation kit with on column DNase treatment and eluted in nuclease-free water.

Microarray cDNA Synthesis—First strand synthesis was performed using the SuperScript Choice system (Invitrogen). The manufacturer's protocol was modified by using a high performance liquid chromatography purified T7-(dT)₂₄ primer (Genset Corp.) and incubating at 42 °C. The 20- μ l first-strand reactions consisted of 100 pmol of T7-(dT)₂₄ primer, 10 μ g of total RNA, 1× first-strand buffer, 10 mM dithiothreitol, 500 μ M each dNTP, and 400 units of Superscript II reverse transcriptase. The 150- μ l second-strand reactions included 1× second-strand reaction buffer, 200 μ M each dNTP, 10 units of DNA ligase, 40 units of DNA polymerase I, and 2 units of RNase H. A phenol/chloroform extraction using phase lock gels (Eppendorf-5 Prime, Inc.) and ethanol precipitation were used to purify the double-stranded cDNA.

Microarray Synthesis of Biotin-labeled cRNA—An in vitro transcription reaction was performed starting with 5 μ l of purified cDNA using the Enzo BioArray high yield RNA transcript labeling kit (Affymetrix). The *in vitro* transcription product was purified using RNeasy spin columns (Qiagen). 20 μ g of cRNA was fragmented (0.5 μ g/ μ l) according to Affymetrix instructions.

Microarray Hybridization-Quantification of the cRNA was adjusted to reflect carryover of unlabeled total RNA with an equation provided by Affymetrix: adjusted cRNA yield = cRNA measured after *in vitro* transcription (μg) – (starting amount of total RNA) \times (fraction of cDNA reaction used in *in vitro* transcription). 20 μ g of adjusted fragmented cRNA was added to 300 μ l of hybridization mixture that included 0.1 mg/ml herring sperm DNA (Promega/Fisher), 0.5 mg/ml acetylated bovine serum albumin (Invitrogen), and 2× MES hybridization buffer (Sigma). The mixture also contained the following hybridization controls: 50 pM oligonucleotide B2 (Genset Corp.) and 1.5, 5, 25, and 100 рм cRNA BioB, BioC, BioD, and Cre, respectively (ATCC). 200 µl of this mixture was hybridized to the chips (mouse genome 430 2.0 Array) with 24×24 - μ m probe cells for 16 h according to Affymetrix procedures. The 50×50 -µm test chips were hybridized with 80 µl for 16 h.

Microarray Washing, Staining, and Scanning-The probe arrays were washed with stringent (100 mM MES, 0.1 M [Na⁺], 0.01% Tween 20) and non-stringent ($6 \times$ SSPE, 0.01% Tween 20, 0.005% antifoam) buffers in the Affymetrix GeneChip Fluidics station using pre-programmed Affymetrix protocols. The probe arrays were then stained with streptavidin phycoerythrin, and the signal amplified using an antibody solution. The streptavidin stain contained $2 \times$ stain buffer (final $1 \times$ concentration 100 mм MES, 1 м [Na⁺], 0.05% Tween 20, 0.005% antifoam, 2 μ g/ μ l acetylated bovine serum albumin, and 10 μ g/ml streptavidin (Molecular Probes)). The antibody amplification solution contained $2\times$ stain buffer, 2 μ g/ μ l acetylated bovine serum albumin (Invitrogen), 0.1 mg/ml normal goat IgG (Sigma), and 3 µg/ml biotinylated antibody (Vector Laboratories). Staining was done in the GeneChip fluidics station using pre-programmed Affymetrix protocols. The probe arrays were scanned in the Affymetrix GeneChip scanner.

Microarray Data Analysis-Microarray analysis using Affymetrix mouse 430 2.0 chip was performed in triplicate using RNA from Shh-treated and control cells. Six data files (triplicate of control and Shh-treated samples) were uploaded into Affymetrix Suite 5.0 (MAS5) software. Unknown expressed sequence tags were identified by searching human genome databases. Genes whose signal intensity were lower than 40 were eliminated from further analyses because the signal was not significantly different from base line (no signal). Data with more than 2 present calls in either the control or Shh-treated group were selected for further analysis. Mean signals of the controls and Shh-treated samples were calculated and compared. Student's t tests were performed and genes with p < 0.05 were selected for further analysis. Fold change was calculated by comparing signals from the Shh-treated group with signals from the control group. Genes showing fold changes greater than ± 2 were selected.

Cell Treatment for Array Gene Validation-To validate the array data and to examine the Shh response of the array-identified genes several cells lines were used. All cells were treated at confluence for 24 h ±50 nM Shh (R&D Systems catalog number 1845-SH/CF) and $\pm 5 \mu M$ cyclopamine (Toronto Research Chemicals) as described above for the microarray. Following treatment, RNA was isolated with Qiagen RNeasy kit with on-column DNase treatment as per the manufacturers protocol. UGSM-2 cells were grown and treated as described above for the array. Mouse embryonic fibroblast (MEF) were isolated and grown as previously described (25, 26). Balb/c 3T3 fibroblast were obtained from ATCC and grown according to ATCC protocols. Primary UGM cells were made from the mouse E16 mesenchyme, as described (9). Primary cells were in culture for 1 week to expand them before exposure to Shh and cyclopamine. All experiments were performed in triplicate and reported differences were statistically significant (p < 0.05; Student's t test).

Prostate Organ Culture—E16 UGS tissues were isolated from CD-1 mice (Charles River) placed on Millicell-CM filters (Millipore) and grown in serum-free media with supplements as described (26). Tissues were incubated for 3 days $\pm 10 \ \mu$ M cyclopamine; RNA was harvested with Qiagen RNeasy columns as per the manufacturers recommendations.

Effect of Gli1, Gli2, Smo Overexpression, and Forskolin Treatment—UGSM-2 cells were infected with retrovirus that expressed green fluorescent protein alone or co-expressed green fluorescent protein and Gli1, Gli2, or Smo. Cells overexpressing green fluorescent protein were identified by flow cytometry and analyzed for expression of the gene of interest. These cells were grown to confluence, maintained at confluence for 24 h, and RNA harvested as described above. To examine the effect of forskolin, UGSM-2 cells were treated with or without 50 μ M forskolin (LC Laboratories, Woburn, MA) during the 24-h period of confluence.

Gene Expression Analysis by Reverse Transcriptase-PCR— Following RNA isolation reverse transcriptase-PCR was performed with gene specific primers (supplemental data Table S1) with normalization to glyceraldehyde-3-phosphate dehydrogenase for each sample. Total RNA was reverse transcribed with Moloney murine leukemia virus-reverse transcriptase fol-

ASBMB



lowing standard protocols followed by real time PCR. PCR products were detected with Power SYBR Green (Applied Biosystems) using a Bio-Rad iCycler thermocycler.

Immunofluorescence Staining-UGS and prostate tissue sections were fixed with 10% formalin, embedded in paraffin, and processed for immunofluorescence staining (27). The following antibodies were used: rabbit anti-Tiam1 polyclonal antibody (1:250, Calbiochem) and mouse anti-smooth muscle actin (1:200, Sigma). Sections were 4',6-diamidino-2-phenylindole counterstained, coverslipped, and imaged.

Electrophoretic Mobility Shift Assays-Electrophoretic mobility shift assays were performed as previously described (21), using purified Gli1 protein (amino acids 211-1106) or PinPoint protein (Promega, Madison, WI) fused to Gli1 amino acids 879–1106, lacking the zinc finger DNA binding domain. Probes were designed using MacVector software (MacVector, Inc., Cary, NC). Probe sequences are shown, listing the sense sequence $(5' \rightarrow 3')$ followed by the antisense sequence $(5' \rightarrow 3')$ used to produce the double-stranded probe: Artn, TTGGGT-CCTGGAACCCCCAACTCCCCACA, TGTGGGGAGTTG-GGGGTTCCAGGACCCAA; Dner, AGACCAGGCTGACC-TCCAACACTGCCTCT, AGAGGCAGTGTTGGAGGTCA-GCCTGGTCT; Fbn2, CCATCTCTCTGACCACCAAGTTT-CCTCAC, GTGAGGAAACTTGGTGGTCAGAGAGAGATGG; Hsd11b1, TATACAAACTCACCTCCCAGAAAAGAACT, AGT-TCTTTTCTGGGAGGTGAGTTGTATA; Igfbp-3, TGCG-CCGGCCCACCCCCACCCTCGCCGT, ACGGCGAGGG-TGGGGGGTGGGCCGGCGCA; Mmp13, ACCATGAGAT-GACCACCAAGAGCATCAGC, GCTGATGCTCTTGGTG-GTCATCTCATGGT; Plxna2, CAATACAAGGGACCTCCC-ACTTGGTAAAG, CTTTACCAAGTGGGAGGTCCCTTG-TATTG; Rgs4, CCTCCCACTCCACCACCAAGGACAGC-TCT, AGAGCTGTCCTTGGTGGTGGAGTGGGAGG; Tiam1, CCGATCCACAGACCACCCAGCACCAGAGC, GCTCTG-GTGCTGGGTGGTCTGTGGATCGG; Tnmd-1, GTGTTG-TGGTGACCACCAAACATAAAAT, AATTTTATGTTTG-GTGGTCACCACAACAC.

Mutant probe sequences are shown, listing the sense sequence $(5' \rightarrow 3')$ followed by the antisense sequence $(5' \rightarrow$ 3'): Artn, TTGGGTCCTGGAAGGGGGAACTCCCCACA, TGTGGGGAGTTCCCCCTTCCAGGACCCAA; Dner, AGA-CCAGGCTGACGAGGTACACTGCCTCT, AGAGGCAGT-GTACCTCGTCAGCCTGGTCT; Fbn2, CCATCTCTCTGA-CGTGGTAGTTTCCTCAC, GTGAGGAAACTACCACGT-CAGAGAGATGG; Hsd11b1, TATACAAACTCACCTCCC-AGAAAAGAACT, AGTTCTTTTCTGGGAGGTGAGTTT-GTATA; Igfbp-3, TGCGCCGGCCCACGGGGGACC-CTCGCCGT, ACGGCGAGGG<u>TCCCCCGTG</u>GGCCGGC-GCA; Mmp13, ACCATGAGATGACGTGGTAGAGCATC-AGC, GCTGATGCTCTACCACGTCATCTCATGGT; Plxna2, CAATACAAGGGACGAGGGACTTGGTAAAG, CTTTAC-CAAGTCCCTCGTCCCTTGTATTG; Rgs4, CTCCCGAAG-AAGTGGGACAATGATGGTTC, GAACCATCATTGTCC-CACTTCTTCGGGAG; Tiam1, CCGATCCACAGACGTGG-GAGCACCAGAGC, GCTCTGGTGCTCCCACGTCTGTG-GATCGG; Tnmd, GTGTTGTGGTGGTGGTGGTAACATAA-AATT, AATTTTATGT<u>TACCACGTC</u>ACCACAACAC.

Whole Mount in Situ Hybridization—The following cDNAs were used as probes: Igfbp-3 (accession number BC058261), Timp3 (accession number BC014713), Plxna2 (accession number BC051045), Cxcl14 (accession number BC0799661), and Dner (accession number BC034634). Those plasmids were linearized, and digoxigenin-labeled probes were prepared by transcription using T7 or T3 RNA polymerase in the presence of DIG RNA labeling mixture. In situ hybridizations were performed as previously described (3). The stained tissues were incubated with 30% sucrose at 4 °C overnight, embedded in Tissue-Tek OCT compound and cryosectioned at 10 μ M.

RESULTS

Identification of Shh Target Genes in UGS Mesenchyme-To identify Shh target genes in the mesenchyme of the developing prostate, we performed microarray analysis in triplicate using RNA from UGSM-2 cells cultured for 24 h in the presence or absence of Shh. Genes showing significantly different signal intensities (p < 0.05) and fold changes greater than ± 2 following treatment with Shh were selected. A total of 89 genes were identified by the array: 68 were up-regulated by Shh and 21 were down-regulated (Table 1). Eighteen of the genes whose transcription was significantly altered by Shh in UGSM-2 cells, including 14 up-regulated genes and 4 down-regulated genes, were selected for further study (Table 2).

Shh-regulated Expression in Primary Mesenchymal Cells and the UGS—To evaluate how well the target gene profile of the UGSM-2 cell line represents the general population of mesenchymal cells in the UGS, we isolated primary mesenchymal cells from the separated E16 UGS mesenchyme and cultured them in the presence or absence of Shh for 24 h. We then analyzed expression of Gli1, Ptc1, and the 18 validated target genes by reverse transcriptase-PCR. Of the 18 genes, 10 demonstrated Shh- regulated expression in the primary UGS mesenchyme cells (Table 2). We then analyzed expression of the 18 genes in intact E16 UGS cultured in serum-free media ± cyclopamine (28), an inhibitor of Hh signaling, to identify those genes whose expression is regulated by Hh signaling during prostate development. Thirteen exhibited significantly altered expression when Hh signaling was inhibited with cyclopamine. In each case, genes induced by Shh in the microarray (n = 10) showed repression when the UGS was exposed to cyclopamine and genes repressed by Shh (n = 3) were induced by exposure of the UGS to cyclopamine (Table 2).

Hedgehog Target Gene Expression in the Newborn Prostate— We performed *in situ* hybridization to examine the expression of selected Hedgehog target genes in the newborn (P1) prostate. This time point was selected because of the robust and distinctive pattern of Shh and Gli1 expression in the nascent buds and adjacent mesenchyme, respectively (3). Expression of all five target genes examined (Timp3, Plxna2, Dner, Cxcl14, and Igfbp-3) exhibited mesenchyme-specific expression localized around the nascent buds, mimicking the expression of Ptc1 and Gli1 (Fig. 1).

Prostate Selective Expression—To determine whether any of the 18 identified target genes exhibit prostate selective Hh-regulated expression, we examined their Shh-dependent expression in primary E13 WT MEFs and NIH3T3 cells. The cells

ASBMB

TABLE 1

ASBMB

The Journal of Biological Chemistry

jbc

Genes regulated by Shh in UGSM-2 cells after 24 h treatment 89 genes changed at least 2-fold between the 24-h control sample and 24-h Shh treatment samples listed according to fold change 89 ge

Gene Name	Gene Symbol	Fold change(S hh/cont)	Gene Name	Gene Symbol	Fold change(S hh/cont)
Upregulated genes			Insulin-like growth factor binding protein 6	Igfbp6	2.28
Chemokine (C-X-C motif) ligand 14	BRAK	27.60	Solute carrier family 7 (cationic amino acid	Slc7a12	2.27
Delta/notch-like EGF-related receptor	Dner	24.95	transporter, y+ system), member 12		
GLI-Kruppel family member GLI1	Gli1	17.09	Fatty acid 2-hydroxylase	Fa2h	2.22
Fibroblast growth factor 5	Fgf5	9.06	Major histocompatibility complex, class I-	Mr1	2.21
Patched homolog 1	Ptc1	8.61	related		
RAS-like, family 11, member B	Rasl11b	4.10	Plexin A2	Plxna2	2.20
Transcribed locus		4.08	Procollagen lysine, 2-oxoglutarate 5-	Plod2	2.20
Mitogen activated protein kinase kinase	Map3k12	3.97	dioxygenase 2		
kinase 12	maponia	0107	Hairy and enhancer of split 1 (Drosophila)	Hes1	2.20
CD24a antigen	Cd24a	3.85	PREDICTED: Rattus norvegicus RNA	Rbm24 pr	2.19
Tissue inhibitor of metalloproteinase 3	Timp3	3.81	binding motif protein 24 (predicted)	edicted,	
Claudin 2	Cldn2	3.70	(Rbm24 predicted)	LOC38084	1
Coiled-coil domain containing 3	Ccdc3	3.67		3	
Angiopoietin 4	Angpt4	3.51	RIKEN cDNA 1110007A13 gene	1110007A	2.18
Hydroxysteroid 11-beta dehydrogenase 1	Hsd11b1	3.48	U U	13Rik	
Calcium/calmodulin-dependent protein	Camk1d	3.39	Weakly similar to POL1 HUMAN		2.18
kinase ID			RETROVIRUS-RELATED POL		
Tenomodulin	Tnmd	3.34	POLYPROTEIN (H.sapiens)		
Ribonucleotide reductase M2 B (TP53	Rrm2b	3.27	Glutamic pyruvate transaminase (alanine	Gpt2	2.16
inducible)			aminotransferase) 2	*	
SPARC related modular calcium binding 2	Smoc2	3.26	Flavin containing monooxygenase 1	Fmo1	2.13
Very low density lipoprotein receptor	Vldlr	3.19	Sperm associated antigen 9	Spag9	2.09
Leucine-rich repeat LGI family. member 2	Lgi2	3.08	Ribonucleotide reductase M2 B (TP53	Rrm2b	2.06
RIKEN cDNA 5133400D11 gene	5133400D	3.05	inducible)		
	11Rik		Lipoma HMGIC fusion partner	Lhfp	2.04
Forkhead box D1	Foxd1	2.97	Son of sevenless homolog 1 (Drosophila)	Sos1	2.04
CDC14 cell division cycle 14 homolog A (S.	Cdc14a	2.96	Syntaxin binding protein 6 (amisyn)	(amisyn)	2.03
cerevisiae)			Fatty acid desaturase 2	Fads2	2.03
NYGGF4 homolog [Homo sapiens]	AL024069	2.90	Scavenger receptor class B, member 1	Scarb1	2.00
Solute carrier family 6 (neurotransmitter	Slc6a6	2.76			
transporter, taurine), member 6					
Syntaxin binding protein 6 (amisyn)	Stxbp6	2.75	Demma mileted and a		
Artemin	Artn	2.67	Downregulated genes		
Kinase suppressor of ras	Ksr	2.67			
STEAP family member 4	Steap4	2.65	Destinguest in section 1	D 1	())
Small nuclear ribonucleoprotein N, full insert	i i	2.63	Dentin matrix protein 1	Dmp1	0.23
sequence			Matrix metalloproteinase 13	Mmp13	4.71
Mus musculus transmembrane protein 69	Tmem69	2.62	WD repeat domain 61	W dr61	3.48
RNA binding protein gene with multiple	Rbpms	2.60	I ranscribed locus		2.73
splicing	.1		Suppressor of fused homolog (Drosophila)	Suru Eldari	2.71
Serine (or cysteine) proteinase inhibitor	Serpinb9b	2.59	FK506 binding protein 1a	rkbpla	2.40
clade B, member 9b	r0 5 0		Autophagy-related 12-like (yeast)	Atg121	2.31
Rattus norvegicus similar to melanoma	17000800	2.57	KIKEN CDINA 4950523CU/ gene	4930323C	2.24
antigen family A, 5	16Rik		Permission of Connection and the	U/KIK Das 4	2.34
Fibrillin 2	Fbn2	2.55	Chamalaina (C. V. C. resting line and 4	Kg84 Cwal4	2.27
Inhibitor of DNA binding 4	Id4	2.51	Chemokine (U-X-U motif) ligand 4 Similar to $Di B = 2$	UXCI4	2.23
Insulin-like growth factor binding protein 3	Igfbp3	2.51	Similar to DI-Kas2	LUC54493	, , , ,
Transcribed locus		2.47	A distance and mat-11-matrix-set d	Z Adam 10	2.22
Phosphodiesterase 9A	Pde9a	2.46	A disintegrin and metalloproteinase domain	Adam12	2.22
Tribbles homolog 2 (Drosophila)	Trib2	2.45	12 (mettrin alpha)		
PR domain containing 8	Prdm8	2.45	rypoineucal Arginine-rich region containing		2.17
G protein-coupled receptor 73	Gpr73	2.44	protein	T-6 012	2.12
Ephrin B2	Efnb2	2.40	Transcription factor CP2-like 3	retep213	2.12
Homo sapiens neurotrophic tyrosine kinase	Ntrk3	2.40	Chloride channel calcium activated 1 ///	Cleal ///	0.10
receptor. type 3			chioride channel calcium activated 2	Clca2	2.10
Scavenger receptor class A member 5	Scara5	2.37	Expressed sequence AU022121	AU022121	2.10
(putative) (Scara5)	4933425F	,	Chemokine (C-X3-C motif) ligand 1	Cx3el1	2.09
(parate) (pourae)	03Rik		Tumor necrosis factor alpha induced protein	Tnfaip6	
	0.5 IVIK	2.27	6		2.06
Filamin hinding I IM protein I	Fhlim1	/ 1/		NT. 11. 4	
Filamin binding LIM protein 1 SLIT and NTRK-like family member 6	Fblim1 Slitrk6	2.37	Nuclear receptor subfamily 1, group H,	Nr1n4	2.06
Filamin binding LIM protein 1 SLIT and NTRK-like family, member 6 Ubiquitin carboxyl-terminal esterase 1.3	Fblim1 Slitrk6 Uch13	2.37 2.36 2.32	Nuclear receptor subfamily 1, group H, member 4	Nr1n4	2.06
Filamin binding LIM protein 1 SLIT and NTRK-like family, member 6 Ubiquitin carboxyl-terminal esterase L3 (ubiquitin thiolesterase)	Fblim1 Slitrk6 Uchl3	2.37 2.36 2.32	Nuclear receptor subfamily 1, group H, member 4 Kringle containing transmembrane protein 1	Nr1n4 Kremen1	2.06 2.02

9

Vaseme)

TABLE 2

ASBMB

The Journal of Biological Chemistry

|--|

	Gene expression ^a					
Gene	Res	ponse to Shh	Response to cyclopamine			
	Array	Primary UGM	UGS organ culture			
Gli1	Ŷ	\uparrow	Ļ			
Ptc1	↑	↑	Ļ			
Igfbp-6	↑	↑	Ļ			
Igfbp-3	ŕ	Ϋ́ τ	Ļ			
Fbn2	↑	\uparrow	ļ			
Ntrk3	↑	\uparrow	ļ			
Agpt4	ŕ	Ϋ́ τ	Ļ			
Dner	↑	\uparrow	ò			
Hsd11b1	↑	↑	0			
Artn	↑	Ϋ́ τ	0			
Tnmd	↑	Ó	Ļ			
Timp3	↑	0	Ļ			
Tiam1	ŕ	0	Ļ			
Plxna2	Ϋ́ Υ	0	Ļ			
BRAK	Ϋ́ Υ	0	\downarrow			
Fgf5	Ϋ́ Υ	0	0			
Dmp1	\downarrow	\downarrow	1			
Mmp13	\downarrow	\downarrow				
Rgs4	\downarrow	0	Ö			
Sufu	\downarrow	0	\uparrow			

^{*a*} Note: gene expression induced by Shh (\uparrow) is expected to be inhibited by cyclopamine (\downarrow). \uparrow , increased expression; \downarrow , decreased expression; 0, no significant change. \uparrow or \downarrow signifies at least 2-fold statistically significant (p < 0.05) change in expression in replicate experiments.

were cultured with and without Shh, and expression of Gli1, Ptc1, and the 18 target genes was examined. Both Ptc1 and Gli1 exhibited robust Shh-induced expression in MEFs and NIH3T3 cells (Fig. 2). Fifteen of the 18 genes demonstrated Shh-regulated expression in MEFs and/or NIH3T3 cells (data not shown). One gene (Dner) that exhibited Shh-induced expression in both UGSM-2 cells and primary UGS mesenchymal cells was not responsive to Shh in either MEFs or NIH3T3 cells. Two additional genes (Tnmd and Timp3) that showed Shh-regulated expression in UGSM-2 cells but not in primary UGS mesenchymal cells also did not show significant Shh-induced change in expression in MEFs or NIH3T3 cells (Fig. 2).

Differential Regulation of Shh-responsive Genes—Direct regulation of Shh target genes is believed to result from the combined activities of Gli1, Gli2, and Gli3. Smo regulates the activities of all three Gli genes by mechanisms that remain incompletely understood. To understand the convoluted actions of Hh signaling in simultaneously regulating proliferation, differentiation, and morphogenesis during development, there is considerable interest in determining the mechanisms by which differential regulation of Shh target genes is achieved in different cells. We first examined the concentration dependence of Shh-induced changes in expression of several of the newly identified Hh target genes and found that most followed the same pattern as Gli1 and Ptc1 with a dose-dependent expression that reached a plateau and remained elevated even with additional ligand (supplemental data Figs. S1 and S2). Tiam1 exhibited a uniquely different concentration dependence of expression. Expression increased to a maximum at 10-50 nM Shh and then progressively decreased to uninduced levels at higher concentrations (supplemental data Fig. S1). Localization of Tiam1 in the P10 and adult prostate by immunofluorescence staining shows expression co-localizing with smooth muscle actin in stromal cells surrounding the prostatic ducts (Fig. 3).



FIGURE 1. Localization of target gene expressions in the newborn prostate. Whole-mount *in situ* hybridization reveals focused domains of Timp3 (A), Cxcl14 (C), Plxna2 (E), Dner (G), and Igfbp3 (I) expression in the mesenchyme surrounding the nascent prostate buds. Sectioning confirms the expression of Timp3 (B), Cxcl14 (D), Plxna2 (F), Dner (H), and Igfbp3 (J) localized to the mesenchyme (m) surrounding the epithelial buds (e). Scale bar = $500 \ \mu m$ for A, C, E, G, and I. Scale bar = $40 \ \mu m$ for B, D, F, H, and J. Data are representative of 3 specimens per gene analyzed.

We next assayed the transcriptional response by transfecting UGSM-2 cells with an activated form of Smo, full-length Gli1, or an activated form of Gli2. All three manipulations would generally be considered methods of achieving Hh target gene activation and, indeed, the canonical target genes Gli1 and Ptc1 exhibited significantly increased expression in response to transfection with all three constructs. Overall, induced increases in expression were lowest following Gli1 transfection and greatest after Smo transfection. When we assayed for changes in expression of the 18 target genes using the same cDNA made from those cells, most of the genes exhibited consistent induction or repression by the three manipulations. This included Igfbp-6, Igfbp-3, Ntrk3, Agpt4, Artn, Timp3, Tiam1, Plxna2, Fgf5, Dmp1, Mmp13, Rgs4, and Sufu (data not shown). However, we uncovered some intriguing differences. Some of the target genes (Dner, Fbn2, Brak (Cxcl14), and Hsd11b1) were not induced by either Gli1 or Gli2 transfection, but were induced by Smo transfection. In striking contrast,





Tnmd showed the highest level of induced expression with Gli1 transfection and the lowest level of induced expression with Smo transfection (Fig. 4).

Gli Binding Elements in the Promoters of Shh Target Genes— The three Gli proteins share a 9-bp consensus binding sequence (GACCACCCA) (29). We performed *in silico* analysis of the remaining validated target genes and identified putative Gli-binding sites upstream of the promoter in 10 additional genes (Tnmd, Tiam1, Fbn2, Mmp13, Plxna2, Dner, Artn, Hsd11b1, Rgs4, and Igfbp-3). Each of the putative Gli binding sites contains at least 7-bp homology with the 9-bp consenture the E16 UGS mesenchyme (3), but the yield of RNA from this is extremely low and would have required an amplification step before array analysis. We considered using freshly isolated E16 UGS mesenchymal cells cultured in monolayer, but our studies of freshly isolated cells suggest that there is significant variability in the growth and behavior of these mixed cell populations that could make validation of the array analysis problematic. Ultimately, we elected to use the previously characterized UGSM-2 cell line that we have shown recapitulates the androgen and Hh responsiveness of the normal UGS mesenchyme and can participate in glandular morphogenesis of the

Shh-responsive Genes

sus (GACCACCCA) (supplemental data Fig. S3). Electrophoretic mobility shift assays were performed to determine the ability of these 10 putative binding sites to interact with the DNA binding motif of Gli protein. The putative Gli-binding elements from the promoter region of target genes Igfbp-3, Fbn2, and Rgs4 bound Gli protein specifically (Fig. 5).

Unique Target Gene Response to PKA Activation-Fbn2 contains a bona fide Gli binding element but neither Gli1 nor an activated form of Gli2 is able to activate Fbn2 expression (Fig. 6). Because gene activation can be achieved by alleviating repression by Gli3 and Gli3 processing is regulated by PKA (30, 31), we examined the effect of the PKA activator forskolin on Shh-induced Fbn2 expression. Forskolin reduced basal expression and blocked Shh-induced expression of Fbn2, consistent with its regulation by Gli3 (Fig. 6). Hsd11b1 is another identified target gene that exhibits activation by Smo but not by either Gli1 or activated Gli2. Forskolin increased basal expression of Hsdllb1 but no further increase in expression was induced by Shh (Fig. 6).

DISCUSSION

We used RNA microarray analysis of an immortalized mesenchymal cell line to identify Shh-responsive genes in the fetal prostate. We could not perform a direct analysis of the cultured E16 UGS because the endogenous Hh ligand would obscure the effect of exogenous Shh peptide and the use of a chemical inhibitor such as cyclopamine could be complicated by nonspecific effects. We are able to isolate and cul-

ASBMB



FIGURE 3. **Tiam1 expression in the developing prostate.** Immunofluorescence staining of Tiam1 and smooth muscle actin (*SMA*) were performed on sections of P10 and adult prostate. Expression of Tiam1 (*green*) co-localized with smooth muscle actin (*red*) expression. *Scale bar* = 100 μ m.



FIGURE 4. **Target gene expression in UGSM-2 cells transfected with Hh pathway components.** UGSM-2 cells were transfected with Gli1, activated Gli2, activated Smo, or green fluorescent protein alone. Expression of the selected 18 genes was analyzed by real time PCR on the extracted RNA from transfected cells (*, p < 0.05, n = 3). *GFP*, green fluorescent protein; *GAPDH*, glyceraldehyde-3-phosphate dehydrogeanse.



FIGURE 5. Identification of Gli1-binding motifs in the 5' region of the Shh target genes. Electrophoretic mobility shift assays demonstrated two shifted bands (*arrows*) in Igfbp-3, Fbn2, and Rgs-4 genes. Specific shifted bands were abrogated by non-radiolabeled oligonucleotide specific to each gene. Control protein or a mutant oligonucleotide DNA did not affect the mobility shift, indicating the specificity of the Gli binding interaction.

developing prostate (9). The time point selected for analysis of the transcriptional response to Hh ligand was based on our previously published studies characterizing the kinetics of induction of Ptc1, Gli1, and Hip in Shh-treated cells and on preliminary studies examining the expression of known target genes such as Igfbp-6 (25). These studies suggested that 24 h after Shh stimulation was the earliest time point that we could expect a robust transcriptional response of novel Hh target genes, a time point that might include both primary and secondary response genes.

To the extent that the UGS mesenchyme is a heterogeneous tissue layer, we expected that the transcriptional response of the UGSM-2 cells might not accurately represent the response of the intact UGS mesenchyme. We were surprised to find that 16 of 18 genes identified in the microarray as putative Hh target genes exhibited Hh-regulated expression in either primary UGS mesenchyme cells, the intact UGS or both. Even more remarkable, all five genes (Timp3, Plxna2, Dner, Igfbp-3, and Cxcl14) selected for analysis of expression in vivo exhibited mesenchyme-specific expression adjacent to sites of robust Shh expression reminiscent of the distribution of Gli1 and Ptc expression in the newborn prostate (3, 29). One explanation is that UGSM-2 cells faithfully represent the dominant Hh-responsive cell type in the E16 UGS mesenchyme. However, another possibility is that the target genes identified in the array are generic Hh target genes and not specific to the mesenchyme of the developing prostate. Several previous studies used microarrays to investigate the target genes of Shh signaling: Ingram et al. (32) identified 11 induced genes and 4 repressed genes in Shh-treated C3H/10T1/2 cells; Yoon et al. (21) identified Gli1-regulated genes in RK3E cells; Oliver et al. (33) identified 134 genes regulated by Shh in granule cell precursors; and Hochman et al. (34) treated C166 endothelial cells with Shh and identified 54 genes exhibiting at least 1.5-fold induction. Indeed, several genes shown to be targets of Hh regulation here were previously identified as Hh target genes in other cells or tissues. These include Camk1d, Foxd1, Igfbp-3, Igfbp-6, Trib2,

bc

ASBMB

Supplemental Material can be found at: http://www.jbc.org/content/suppl/2008/12/22/M809172200.DC1.html



FIGURE 6. **Regulation of target gene expression by forskolin treatment.** UGSM-2 cells were treated with Shh in the presence or absence of forskolin. RNA were extracted after 24 h, followed by performing real time PCR to analyze selected gene expression (*, p < 0.05, n = 3). *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase.

Mr1, and Rgs4 (34). However, our studies identified several genes that had not been previously identified as Hh targets.

To identify genes for which regulation by Hh signaling is prostate selective, we examined expression of the validated genes in MEFs and NIH3T3 cells. Three genes (Dner, Tnmd, and Timp3) that exhibited a diminished response to Shh treatment in both cell lines may be considered as potentially prostate-selective Hh target genes, but this remains to be established by further studies.

Dner, Tnmd, and Timp3 have been studied in other contexts and have all been shown to have roles in growth regulation. Tnmd is a regulator of tenocyte proliferation (35); Timp3 has been shown to regulate cell proliferation, inflammation, and innate immunity; both Tnmd and Timp3 function as angiogenesis inhibitors (35–38). Dner has been previously characterized as a neuron-specific Notch ligand and is essential for precise cerebellar development by stimulating morphological differentiation of Bergmann glia (39). Our study provides the first evidence for its expression in the developing prostate.

Among the other putative target genes newly identified on the array but not necessarily prostate selective are genes related to Notch and Wnt signaling. Hes1 and Efnb2 are both known targets of Notch signaling, with the expression of Hes1 also having been shown to be dependent on Shh in tissues such as retina and cerebellum (40, 41). Another gene, Kremen1, is a Dickkopf receptor that negatively regulates the Wnt signaling pathway (42). These findings suggest previously unrecognized cross-talk or points of conversion between Hh signaling and the Wnt and Notch signaling pathways during prostate development. Another notable feature in the genes identified here are three involved in regulating angiogenesis, including Efnb2, Rgs4, and Angiopoietin-4 (Agpt4) (43–45).

ence on specific, albeit as yet unidentified transcriptional cofactors working in combination to influence the transcriptional machinery and produce a unique pattern of target gene activation. However, other mechanisms for differential gene regulation exist. In Drosophila, selective gene activation results from a differential responsiveness of target genes to a concentration gradient of Hh ligand. Our studies identify Tiam1 (T lymphoma invasion and metastasis 1), a guanine nucleotide exchange factor, as a gene whose Hh-regulated expression may be gradient dependent and it is notable that Tiam1 expression is found in a restricted zone of stroma around the ducts of the developing prostate. Expression of Hh target genes in the periductal stroma is not unexpected because Shh is expressed by the epithelium of the developing ducts, but co-localization with the band of smooth muscle actin expression around the ducts suggests the intriguing possibility that a restricted band of Tiam1 expression induced by the gradient of secreted Shh ligand plays a role in radial patterning of the duct and determining the location of smooth muscle induction.

Differential target gene regulation in mammals can also occur from varying affinities of the three different Gli factors for Gli binding sites in different promoter regions. Indeed, recent studies have shown a differential regulation of Ptc1 and Bcl2 in human epidermal cells that is attributed to relative affinities of Gli1 and Gli2 for the Gli binding sites in the promoters of the two genes (46, 47). Here we tested whether Gli1, Ptc1, and the non-canonical Hh target genes identified in our array are similarly responsive to retroviral transfection of UGSM-2 cells with activated Smo, Gli1, or activated Gli2. The data show that this is clearly not the case and, furthermore, reveal three distinctive response patterns among the genes tested. 1) Genes that are induced (or repressed) by all three constructs. Genes in this category include Gli1 and Ptc1 as well as Igfbp-6, Igfbp-3,

Shh-responsive Genes

Hh regulation of early target gene transcription is thought to be mediated by the activities of the three members of the Gli family whose combined activities are regulated by Smo and perhaps other members of the mammalian Hh signal transduction machinery by a combination of phosphorylation events, protein processing, and nuclear translocation. In previously published studies, we examined the transcriptional regulation of the canonical Hh target genes Gli1, Ptc1, and Hip in wild type, single Gli allele mutant, and double Gli allele mutant MEFs (25). Those studies demonstrated similar patterns of Hh ligand-induced transcription of Gli1 and Ptc1 in the different mutant cells, arguing for a common mechanism of regulation. But it is clear that other target genes are differentially regulated. How is this achieved? Presumably, a major mechanism of regulation is depend-

ASBMB



Ntrk3, Agpt4, Artn, Timp3, Tiam1, Plxna2, Fgf5, Dmp1, Mmp13, Rgs4, and Sufu. 2) Genes that are induced by Smo but not by either Gli1 or Gli2. These genes include Dner, Fbn2, Brak, and Hsd11b1. 3) Genes such as Tnmd that are induced by Gli1 and Gli2 transfection but not by Smo.

There are likely a variety of reasons for the different patterns of response, but the data presented here provide a starting place for functional studies to examine this issue. Indeed, we explored the role of Gli3-mediated regulation in expression of two genes, Fbn2 and Hsd11b1, which exhibited Smo-induced expression but were not activated by either Gli1 or activated Gli2. The PKA activator forskolin enhances the proteolysis of Gli3 to generate a repressor form. As would be expected for a gene regulated by Gli3, Fbn2 induction by Shh was blocked by forskolin treatment. The induction of Hsd11b1 by Shh was also blocked by forskolin, whereas constitutive expression was increased, suggesting multiple roles for PKA in the regulation of Hsd11b1.

In previously published studies Igfbp-6 and Foxd1 have been shown to have conserved Gli1 binding motifs that were confirmed by gel shift binding assays (21). To determine whether any of our identified target genes might be regulated directly by Gli proteins, we examined the promoter regions of those genes. Putative Gli binding motifs were found in 10 of them. Functional Gli binding sites were found in 3 of them after performing gel shift assay.

Identification of mesenchymal target genes of Hh signaling in the developing prostate was undertaken to advance our studies of how Hh signaling regulates prostate budding and ductal morphogenesis. In addition to identifying specific Hh target genes, these studies provide the first evidence to our knowledge for cross-talk and/or convergence between Hh signaling and the Wnt and Notch pathways in the fetal prostate and offer novel opportunities to begin elucidating the importance of these connections in regulating growth. These findings are particularly relevant to ongoing studies of Hh signaling in prostate cancer. We have previously shown that paracrine Hh signaling occurs in human prostate cancer and can promote tumor growth in a xenograft model (8). In more recent studies, we have shown that expression of multiple target genes identified here correlate significantly with Hh signaling in specimens of human prostate cancer, but not benign prostate tissue, suggesting a re-activation of fetal signaling mechanisms in cancer.³ It is hoped that identification of these Hh target genes and the connections of Hh signaling with the Wnt and Notch pathways, both implicated in prostate cancer, will advance the effort to devise new targeted therapies for advanced disease not amenable to ablative therapy.

REFERENCES

- 1. Ingham, P. W., and McMahon, A. P. (2001) Genes Dev. 15, 3059-3087
- 2. Podlasek, C. A., Barnett, D. H., Clemens, J. Q., Bak, P. M., and Bushman, W. (1999) Dev. Biol. 209, 28-39
- 3. Lamm, M. L., Catbagan, W. S., Laciak, R. J., Barnett, D. H., Hebner, C. M., Gaffield, W., Walterhouse, D., Iannaccone, P., and Bushman, W. (2002) Dev. Biol. 249, 349-366
- 4. Freestone, S. H., Marker, P., Grace, O. C., Tomlinson, D. C., Cunha, G. R.,
- ³ A. Shaw, J. Gipp, and W. Bushman, unpublished observations.

Harnden, P., and Thomson, A. A. (2003) Dev. Biol. 264, 352-362

- Berman, D. M., Desai, N., Wang, X., Karhadkar, S. S., Reynon, M., Abate-5. Shen, C., Beachy, P. A., and Shen, M. M. (2004) Dev. Biol. 267, 387-398
- 6. Hooper, J. E., and Scott, M. P. (2005) Nat. Rev. Mol. Cell. Biol. 6, 306-317
- 7. Wang, B. E., Shou, J., Ross, S., Koeppen, H., De Sauvage, F. J., and Gao, W. Q. (2003) J. Biol. Chem. 278, 18506-18513
- 8. Fan, L., Pepicelli, C. V., Dibble, C. C., Catbagan, W., Zarycki, J. L., Laciak, R., Gipp, J., Shaw, A., Lamm, M. L., Munoz, A., Lipinski, R., Thrasher, J. B., and Bushman, W. (2004) Endocrinology 145, 3961-3970
- 9. Shaw, A., Papadopoulos, J., Johnson, C., and Bushman, W. (2006) Prostate 66, 1347-1358
- 10. Shaw, A., and Bushman, W. (2007) J. Urol. 177, 832-838
- 11. Chuang, P. T., and McMahon, A. P. (1999) Nature 397, 617-621
- 12. Ingham, P. W., and Placzek, M. (2006) Nat. Rev. Genet. 7, 841-850
- Callahan, C. A., Ofstad, T., Horng, L., Wang, J. K., Zhen, H. H., Coulombe, 13. P. A., and Oro, A. E. (2004) Genes Dev. 18, 2724-2729
- 14. Eichberger, T., Regl, G., Ikram, M. S., Neill, G. W., Philpott, M. P., Aberger, F., and Frischauf, A. M. (2004) J. Investig. Dermatol. 122, 1180-1187
- 15. Roelink, H., Porter, J. A., Chiang, C., Tanabe, Y., Chang, D. T., Beachy, P. A., and Jessell, T. M. (1995) Cell 81, 445-455
- 16. Krishnan, V., Pereira, F. A., Qiu, Y., Chen, C. H., Beachy, P. A., Tsai, S. Y., and Tsai, M. J. (1997) Science 278, 1947-1950
- 17. Wu, S. C., Grindley, J., Winnier, G. E., Hargett, L., and Hogan, B. L. (1998) Mech. Dev. 70, 3-13
- 18. Teh, M. T., Wong, S. T., Neill, G. W., Ghali, L. R., Philpott, M. P., and Quinn, A. G. (2002) Cancer Res. 62, 4773-4780
- 19. Mahlapuu, M., Enerback, S., and Carlsson, P. (2001) Development 128, 2397-2406
- 20. Izraeli, S., Lowe, L. A., Bertness, V. L., Campaner, S., Hahn, H., Kirsch, I. R., and Kuehn, M. R. (2001) Genesis 31, 72-77
- 21. Yoon, J. W., Kita, Y., Frank, D. J., Majewski, R. R., Konicek, B. A., Nobrega, M. A., Jacob, H., Walterhouse, D., and Iannaccone, P. (2002) J. Biol. Chem. 277, 5548-5555
- 22. Gaur, T., Rich, L., Lengner, C. J., Hussain, S., Trevant, B., Ayers, D., Stein, J. L., Bodine, P. V., Komm, B. S., Stein, G. S., and Lian, J. B. (2006) J. Cell. Physiol. 208, 87-96
- 23. He, J., Sheng, T., Stelter, A. A., Li, C., Zhang, X., Sinha, M., Luxon, B. A., and Xie, J. (2006) J. Biol. Chem. 281, 35598-35602
- 24. Lipinski, R. J., Cook, C. H., Barnett, D. H., Gipp, J. J., Peterson, R. E., and Bushman, W. (2005) Dev. Dyn. 233, 829-836
- 25. Lipinski, R. J., Gipp, J. J., Zhang, J., Doles, J. D., and Bushman, W. (2006) Exp. Cell Res. 312, 1925–1938
- 26. Doles, J. D., Vezina, C. M., Lipinski, R. J., Peterson, R. E., and Bushman, W. (2005) Prostate 65, 390-399
- 27. Garrett, W. M., and Guthrie, H. D. (1998) Biochemica 1, 17-20
- 28. Cooper, M. K., Porter, J. A., Young, K. E., and Beachy, P. A. (1998) Science 280, 1603-1607
- 29. Kinzler, K. W., and Vogelstein, B. (1990) Mol. Cell. Biol. 10, 634-642
- 30. Wang, B., Fallon, J. F., and Beachy, P. A. (2000) Cell 100, 423-434
- 31. Dai, P., Akimaru, H., Tanaka, Y., Maekawa, T., Nakafuku, M., and Ishii, S. (1999) J. Biol. Chem. 274, 8143-8152
- 32. Ingram, W. J., Wicking, C. A., Grimmond, S. M., Forrest, A. R., and Wainwright, B. J. (2002) Oncogene 21, 8196-8205
- 33. Oliver, T. G., Grasfeder, L. L., Carroll, A. L., Kaiser, C., Gillingham, C. L., Lin, S. M., Wickramasinghe, R., Scott, M. P., and Wechsler-Reya, R. J. (2003) Proc. Natl. Acad. Sci. U. S. A. 100, 7331-7336
- 34. Hochman, E., Castiel, A., Jacob-Hirsch, J., Amariglio, N., and Izraeli, S. (2006) J. Biol. Chem. 281, 33860-33870
- 35. Docheva, D., Hunziker, E. B., Fassler, R., and Brandau, O. (2005) Mol. Cell. Biol. 25, 699-705
- 36. Gill, S. E., Pape, M. C., and Leco, K. J. (2006) Dev. Biol. 298, 540-554
- Spurbeck, W. W., Ng, C. Y., Strom, T. S., Vanin, E. F., and Davidoff, A. M. (2002) Blood 100, 3361-3368
- 38. Shukunami, C., Oshima, Y., and Hiraki, Y. (2005) Biochem. Biophys. Res. Commun. 333, 299-307
- 39. Eiraku, M., Tohgo, A., Ono, K., Kaneko, M., Fujishima, K., Hirano, T., and Kengaku, M. (2005) Nat. Neurosci. 8, 873-880
- 40. Wang, Y., Dakubo, G. D., Thurig, S., Mazerolle, C. J., and Wallace, V. A.

SBMB

The Journal of Biological Chemistry

5628 JOURNAL OF BIOLOGICAL CHEMISTRY

Supplemental Material can be found at: http://www.jbc.org/content/suppl/2008/12/22/M809172200.DC1.html

Shh-responsive Genes

Downloaded from www.jbc.org at University of Wisconsin-Madison, on January 15, 2010

(2005) Development 132, 5103-5113

- Dakubo, G. D., Mazerolle, C. J., and Wallace, V. A. (2006) J. Neurooncol. 79, 221–227
- Mao, B., Wu, W., Davidson, G., Marhold, J., Li, M., Mechler, B. M., Delius, H., Hoppe, D., Stannek, P., Walter, C., Glinka, A., and Niehrs, C. (2002) *Nature* 417, 664–667
- 43. Gerety, S. S., and Anderson, D. J. (2002) Development 129, 1397-1410
- 44. Albig, A. R., and Schiemann, W. P. (2005) Mol. Biol. Cell 16, 609-625
- Olsen, M. W., Ley, C. D., Junker, N., Hansen, A. J., Lund, E. L., and Kristjansen, P. E. (2006) *Neoplasia* 8, 364–372
- Regl, G., Kasper, M., Schnidar, H., Eichberger, T., Neill, G. W., Philpott, M. P., Esterbauer, H., Hauser-Kronberger, C., Frischauf, A. M., and Aberger, F. (2004) *Cancer Res.* 64, 7724–7731
- Bigelow, R. L., Chari, N. S., Unden, A. B., Spurgers, K. B., Lee, S., Roop, D. R., Toftgard, R., and McDonnell, T. J. (2004) *J. Biol. Chem.* 279, 1197–1205



Supporting data:

(Fig. 1)





UGS tissues from e16 and P1 stage were treated with AZ75 (Hh inhibitor) for 7 days in organ culture. (Left) Quantitive results showed a decreased ductal tips in e16 UGS tissues but an increased dutal tips in P1 UGS tissues.(Top) Real-time PCR results showing that though the conserved downstream genes Gli1 and Ptc1 were all inhibited in both e16 and P1 UGS tissues by AZ75 treament, a differential regulation pattern were found in other downstream genes(Hes1, Fgf5, Hsd11b1 and Agpt4).



(Top) X-gal staining on UGS tissues from Ptc1.Lacz and Gli1.Lacz mice showed positive staining in epithelium, indicating the existence of autocrine Hh signaling. (Bottom left) Typical phenotype of null autocrine signaling mutant mice (P1); ventral prostate was missing in the mutant mice.(Bottom right) The UGS from mutant mice showed impaired growth ability in organ culture system (without Testosterone).









HE stains on the grafted UGS tissues from control and Smo conditional null mutant mice. (A and A'): 8 weeks graft UGS tissues ; (B and B') 8 weeks graft followed by 2 weeks castration; (C and C') 8 weeks graft with 2 weeks castration and 2 weeks Testosterone resupply.

Conclusions:

Emerging evidences have suggested an important role of Hh signaling in promoting human prostate cancer growth. However, lack of proper *in vivo* animal model and *in vitro* cell lines hampered its research progress. My interesting in study Hh signaling pathway in normal prostate development is originated from the connection between embryogenesis and oncogenesis; most signaling pathway that is crucial for developmental process is mostly the one that frequently activated in the cancer process. Noting the decade's discrepancies in the field, I initiated this work to re-visit the role of Hh signaling in the developing prostate. Major findings of this research are: 1) Paracrine Hh signaling bidirectionally tunes prostate ductal growth 2) Autocrine Hh signaling has been neglected in the whole story. (3) Some non-canonical target genes are temporally regulated. In the following part, I listed the conclusions by answering each research question. Limitations of research are also discussed in each section.

What are Hh responsive genes in developing prostate?

This would be the first question to answer when starting to investigate paracrine Hh signaling in prostate tissue. By microarray analysis, we acquired lists of Hh responsive genes in UGSM-2 cells, and selected genes were further proved in cultured E16 UGS tissues and primary E16 UGM cells by RT-PCR gene analysis. The identification of Hh downstream genes in fetal mesenchymal cells provided a platform to perform functional study of each responsive gene, and further for understanding how Hh signaling exert its specific role by up or down-regulation those genes. However, it need to keep in mind that the responsive genes in colonial UGSM-2 cells may not well represent those in intact heterogeneous UGM. Notably, just because those genes are identified in mesenchymal cells doesn't mean they don't express in the epithelium. Finally, realizing the composition of mesenchymal cells changes along development, it is possible that the target genes or the responsiveness of each target gene or both may varied during the development time course.

What is the role of paracrine Hh signaling on prostate ductal growth? Is it stimulatory? Is it inhibitory?

We demonstrated that Hh signaling exerts a stage-specific effect on prostatic duct growth. Utilizing *in vitro* explants culture and *in vivo* gain of function animal models, we observed consistent results, which showed promotion effect of Hh signaling on prostatic dutal growth prenatally, and inhibition effect on prostatic ductal growth postanatally. A possible interpretation for this stage-specific effect could be that during embryonic development stage, Hh signaling acts as a mitogen to expand epithelial population and stimulate budding and branching morphogenesis; while in later postnatal development stage, its role switches to promote ductal differentiation; as its function in promote epithelial differentiation was suggested earlier. This temporal controlled tissue response is by no means unique to prostate tissue. It was suggested in pancreas, Hh signaling functions early in control morphogenesis by regulation of epithelial and beta-cell proliferation and modulate insulin secretion in adult mice. However, without temporal controlled cre mice, we are not able to decide what specific timing is the turning point, which will be of crucial important both for development process and prostate cancer process.

What is the mechanism?

Our studies have provided a list of target genes that showed discordant response to Hh signaling in prenatal and postnatal stage UGS tissues. Since transcriptional genes are thought to be responsible for the function of cell signaling, it is possible that the stage-specific effect is mediated by differentially regulated those genes. The genes (Igfbp3, Agpt4, Fgf5 and Fbn2) we identified are involved in control cell proliferation, extracellular matrix remodeling and angiogenesis process. Their association with reactive stroma in prostate cancer is also being suggested. Future functional studies of each gene need to be done to correlate it with the net stage-specific epithelial response to Hh signaling, which will be of equal interest to their function in cancer process. Our gene analysis studies on primary culture mesenchymal cells suggested that the mesenchymal character is responsible but not sufficient for the discordant response of those genes. While failure of recapitulate the discordant gene regulation pattern in our tissue recombination experiment study suggested that the tissue recombination model is not suitable for our purpose.

Is it all paracrine?

Our localization studies using Gli1-lacZ and Ptc1-lacZ mice clearly evidenced the positive epithelial Hh signaling, suggested the possibility of autocrine Hh signaling. Functional studies implicated a role in control adult progenitor cell population and a possible role in control the androgen-sensitivity of epithelium. In mice with ablation of epithelial Hh signaling, a severe epithelial involution response was observed. The degenerate prostatic ducts cannot recover to its normal morphology with androgen supplementation. To strengthen our conclusions, specifically labeling those progenitor cells and directly comparison of their population will be of great helpful. So what?

We picture the role of Hh signaling in prostate development as a story of both paracrine and autocrine mechanism; with paracrine Hh signaling, it promotes prostatic ductal growth prenatally and inhibits postnatally; with autocrine Hh signaling, it controls the adult progenitor cell population which is important for tissue regeneration.

In prostate development field, results from this work resolved the long-term discrepancies on the role of Hh signaling in prostate development, by concreting a novel understanding of this concept, and thus advanced the field to another stage.

In the prostate cancer field, the stage-specific results of Hh signaling in prostate development may suggest that the embryonic role of Hh signaling is reactivated, and the potential mechanism underlying this shift may also offer promising molecular targeting for prostate cancer treatment. Our autocrine Hh signaling theory in development posed a possible role of autocrine Hh signaling in prostate cancer which may be involved in control the cancer stem cell population.