

# 1/12° Global HYCOM Evaluation and Validation

Joe Metzger<sup>1</sup>, Harley Hurlburt<sup>1</sup>, Alan Wallcraft<sup>1</sup>, Ole Martin Smedstad<sup>2</sup>, Birol Kara<sup>1</sup>, Jay Shriver<sup>1</sup>, Lucy Smedstad<sup>1</sup>, Debbie Franklin<sup>2</sup>, Bill Schmitz, Jr.<sup>3</sup>, and Prasad Thoppil<sup>4</sup>

<sup>1</sup>Naval Research Laboratory, <sup>2</sup>Planning Systems, Inc.,  
<sup>3</sup>Texas A&M-CC, <sup>4</sup>University of Southern Mississippi

11<sup>th</sup> HYCOM Consortium Meeting  
24-26 April 2007  
Stennis Space Center, MS

# Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*


Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>APR 2007</b>	2. REPORT TYPE	3. DATES COVERED <b>00-00-2007 to 00-00-2007</b>	
4. TITLE AND SUBTITLE <b>1/12degree Global HYCOM Evaluation and Validation</b>		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory,Stennis Space Center,MS,39529</b>		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>			
13. SUPPLEMENTARY NOTES			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	
			18. NUMBER OF PAGES <b>23</b>
			19a. NAME OF RESPONSIBLE PERSON

# 1/12° HYCOM/NCODA/PIPS

- **Capability:** Provide accurate 3D temperature, salinity and current structure; depict the location of mesoscale features such as oceanic eddies and fronts
- **Progress:** 1/12° global HYCOM/NCODA running in real-time in the NAVOCEANO operational queues; Validation testing has begun
- **Issues:**
  - Complete coupling of HYCOM/PIPS via ESMF (NRL)
  - Get NCODA working in curvilinear part of grid (NRL)
  - Need OcnQC running operationally (NAVOCEANO)

# HYCOM/NCODA Runstream




- 1) Perform first NCODA analysis centered on  $\tau = -120$
- 2) Run HYCOM for 24 hours using incremental updating (■) over the first 6 hrs
- 3) Repeat steps 1) and 2) until the nowcast time
- 4) Run HYCOM in forecast mode out to  $\tau = 96$ , eventually to  $\tau = 120$

Approximate run times\* (using 379 IBM Power 5+ processors):

- 1) Six NCODA analyses: 0.9 hrs/analysis = 5.4 hrs
- 2) Five HYCOM hindcast days @ 150 sec  $\Delta t$ : 1.1 hrs/day = 5.5 hrs
- 3) Four HYCOM forecast days @ 150 sec  $\Delta t$ : 1.1 hrs/day = 4.4 hrs
- 4) Total: 15.3 hrs

\* Timings do not include PIPS coupling; assimilation in the Mercator part of grid only

# HYCOM/NCODA Runstream



- 1) Perform first NCODA analysis centered on tau = -126, i.e. 18Z
- 2) Run HYCOM for 24 hours using incremental updating (■) over the first 6 hrs starting at 18Z
- 3) Repeat steps 1) and 2) until the nowcast time
- 4) Run HYCOM in forecast mode out to tau = 96, eventually to tau = 120


Under this scheme the incremental updating ends at the nowcast time (00Z) whereas in the previous scheme incremental updating ended at 06Z and the 00Z nowcast actually represents an 18-hour forecast from the previous day. **Most results shown in this presentation are from 18-hour forecasts.**

# FY07 Validation Tasks

1. Mixed layer depth / sonic layer depth / deep sound channel
  - Compare simulated vs. observed for non-assimilated buoys
2. Vertical profiles of T&S
  - Quantitative comparison of simulated vs. observed for non-assimilated buoys
3. Large scale circulation features
  - Determine correct placement of large scale features
4. Eddy kinetic energy / sea surface height variability
  - Determine if the system has a realistic level and distribution of energy at depths
5. Sea surface temperature
  - Evaluate whether the models are producing acceptable nowcasts and forecasts of sea surface temperature
6. Coastal sea level
  - Assess the model's ability to represent observed sea surface heights

# Mean Sea Surface Evaluation


1992-2002 Mean dynamic ocean topography (0.5°)



Mean ocean dynamic topography data has been obtained from Nikolai Maximenko (IPRC) and Peter Niiler (SIO)

# Mean Sea Surface Evaluation

2004 Mean sea level from 1/12° global HYCOM/NCODA




From the 1/12° global HYCOM/NCODA hindcast simulation  
Mean shifted by 8.7 cm; standard deviation of difference = 9.6 cm




# SSH Variability Evaluation

Sea surface height variability





Oct 92 – Nov 98 SSH variability based on T/P, ERS-1 and ERS-2 altimeters (from Collecte, Localisation, Satellites (CLS))



SSH variability over 2004 from the 1/12° global HYCOM/NCODA hindcast simulation

# Western Boundary Current Comparison

Sea surface height – 9 March 2007






IR frontal analysis overlaid:  
White  $\leq 4$  days  
Black  $> 4$  days

# Eddy Kinetic Energy Comparison

## Surface EKE in the Gulf Stream




Observations from  
Fratantoni (2001) –  
Based on 1990-99  
surface drifters



# Eddy Kinetic Energy Comparison


EKE at ~700 m in the Gulf Stream

Observations from  
Schmitz (1996)



# Mixed Layer Depth Comparison


2004 MLD difference: HYCOM minus unassimilated MEDS profiles



MLD = negative temperature difference of  $0.5^{\circ}\text{C}$  between the surface and depth;  
data averaged in  $0.5^{\circ}$  bins

# Mixed Layer Depth Comparison


2004 MLD difference: HYCOM minus unassimilated MEDS profiles



MLD = negative temperature difference of  $0.5^{\circ}\text{C}$  between the surface and depth;  
data averaged in  $0.5^{\circ}$  bins

# Temperature Structure Comparison

Locations of TAO and PIRATA buoys used in this evaluation



Buoys are divided into two sets based on the vertical sampling and continuity of the time series over calendar year 2004



Set 1 (denoted by o's): 1, 20, 40, 60, 80, 100, 120, 140, 180, 300, 500 m.

Set 2 (denoted by x's): 1, 25, 50, 75, 100, 125, 150, 200, 250, 300, 500 m.




# Temperature Structure Comparison

2004 subsurface temp at 140°W, 2°N  
Buoy / HYCOM / nonassim HYCOM

Temperature difference  
Buoy - HYCOM / Buoy - nonassim HYCOM



Black = assimilation white = no assimilation





Significant impact of temperature profile assimilation via NCODA





# Temperature Structure Comparison

HYCOM vs. non-assim HYCOM – Mean error – 47 TAO/PIRATA buoys 2004




# Temperature Structure Comparison

HYCOM vs. non-assim HYCOM – Skill score – 47 TAO/PIRATA buoys 2004



# Sea Surface Temperature Comparison


HYCOM vs. MODAS – Mean error – white area =  $\pm .25^{\circ}\text{C}$



Over 2004 from the  $1/12^{\circ}$  global HYCOM/NCODA hindcast simulation

# Sea Surface Temperature Comparison


HYCOM vs. MODAS – Skill score



Over 2004 from the 1/12° global HYCOM/NCODA hindcast simulation

# Sea Surface Temperature Comparison

Unassimilated MEDS SST vs. HYCOM vs. NCOM




	HYCOM	NCOM
ME	-.1°C	.2°C
RMSE	.9°C	2.2°C
R	.99	.93
SS	.98	.86

Over 2004 from the 1/12° global HYCOM/NCODA hindcast simulation and operational 1/8° global NCOM; MEDS = Marine Environmental Data Services


# Coastal/Island Sea Level Comparison

Simulated vs. observed sea level at 84 coastal / island stations during 2004



# Coastal/Island Sea Level Comparison

RMSE improvement  
(HYCOM – observed) – (NCOM – observed)



Simulated vs. observed sea level at 84 coastal / island stations during 2004

# FY08 Validation Tasks

1. Below layer depth gradient
  - Compare simulated vs. observed for non-assimilated buoys
2. Comparison with drifting buoys
  - Evaluate the model's ability to produce ocean currents that yield drifter and ARGO float trajectories similar to observations
3. Current cross sections
  - Evaluate model velocity cross-sections through qualitative and quantitative comparisons
4. Provide boundary conditions to nested models
  - Nest East Asian Seas NCOM and Relocatable NCOM within HYCOM and compare inner model with the solution when forced NCOM
5. Eddy tracking
  - Evaluate the model's ability to track mesoscale eddies
6. Ice drift, thickness and concentration
  - Assess the model's ability to represent sea ice