

Effects of histamine receptor blockade on cardiovascular changes induced by 35 GHz radio frequency radiation heating

J. R. Jauchem¹, K. L. Ryan² & M. R. Tehrany²

¹Air Force Research Laboratory, Directed Energy Bio-Effects Division, Radio Frequency Radiation Branch, Brooks City-Base, TX 78235-5147 and ²Department of Biology, Trinity University, San Antonio, TX 78212, USA

Correspondence:
J.R. Jauchem

Summary

1 The role of histamine in heat-induced cardiovascular changes is uncertain. The purpose of this study was to examine effects of histamine H-1- and H-2-antagonism on heart rate, mean arterial blood pressure (MAP), localized body temperature changes, survival times, and lethal body temperatures that occur during the exposure of anaesthetized rats to 35 GHz radio frequency radiation (RFR).

2 Forty-eight ketamine-anaesthetized Sprague–Dawley rats were exposed, in several different treatment groups ($n = 8$ in each), to 35 GHz RFR at a level that resulted in significant body heating and subsequent death. During irradiation, a continuous increase in heart rate and a biphasic response in blood pressure (initial increase followed by a decrease) were observed in all groups of animals.

3 An H-1-antagonist, diphenhydramine (1 mg kg^{-1} body wt) and an H-2-antagonist, cimetidine (5 mg kg^{-1}), administered after sustained RFR exposure, failed to reverse the RFR-induced hypotension. High doses of the drugs (5 and 10 mg kg^{-1} , respectively) also did not alter the response. Post-RFR survival time was significantly decreased in the high-dose drug-treated group, compared with vehicle-treated (0.9% NaCl, 50% ethanol and 50% D5W) controls.

4 In experiments in which the two drugs were administered prior to RFR exposure, MAP in animals receiving high-dose antihistamines was significantly depressed compared with that of vehicle-treated animals during the first 35 min of RFR exposure. Antihistamine pretreatment, however, did not alter the total RFR exposure time required for death to occur.

5 In summary, pharmacological blockade of H-1 and H-2 receptors is not beneficial in anaesthetized rats made hypotensive by RFR exposure. This indicates that activation of H-1 and H-2 receptors by histamine does not occur to any significant extent and does not mediate the hypotensive response developed in this model of hyperthermia.

Keywords: histamine, H₂ receptors, H₁ receptors, radio frequency radiation, blood pressure, heart rate

Introduction

The increased use of radio frequency radiation (RFR) exposure sources capable of generating millimetre wavelengths for military and civilian purposes (e.g. communications, radar imaging/mapping, space power beaming, radiometry) will continue to generate interest in possible biological effects of exposure to these sources. Among the potential effects, an increase in body temperature is the primary consequence of whole body exposure

to high levels of RFR. The use of RFR to study physiological changes during heat stress has been discussed previously (e.g. Jauchem & Frei, 1994).

Cardiovascular responses due to RFR induced heating are, in general, similar to responses to environmental heating as described by Kregel & Gisolfi (1990). Exposure to short wavelength RFR (including millimetre waves), however, can result in energy deposition at the body surface without deposition at deeper sites. Thus, there will be differences in temperature gradients within the

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 JAN 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Effects of histamine receptor blockade on cardiovascular changes induced by 35-GHz radio-frequency radiation heating.				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jauchem J. R., Tehrany M. R., Ryan K. L.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 13	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

body resulting from short wavelength RFR *vs.* environmental heating. These differences may, in turn, lead to quantitative variations in the magnitude of cardiovascular changes (Jauchem & Frei, 1992). In addition, extended RFR exposure of this type will result in increased core body temperature due to circulatory transfer of heat. Ryan, D'Andrea, Jauchem & Mason (2000) have reviewed the topic of millimetre wave heating.

Frei, Ryan, Berger & Jauchem (1995) previously demonstrated that sustained whole body exposure to RFR energy of approximately millimetre wave length (35 GHz) produces shock like hypotension in anaesthetized Sprague Dawley rats. In these studies, circulatory collapse occurred when skin temperature was rapidly elevated to levels much greater than those measured at deeper sites within the body. Heart rate (HR) and mean arterial blood pressure (MAP) changes were similar to those that occur during environmental heat stress. It should be noted, however, that circulatory collapse occurred at much lower core temperatures than observed in heatstroke models of environmental heating (Kregel & Gisolfi, 1990; Frei *et al.*, 1995).

Histamine is a potent vasodilator that has been implicated as a mediator in several different forms of shock. Previous studies (the majority listed here using the Sprague Dawley rat) have elucidated the role of histamine in cardiovascular changes occurring during such insults as endotoxaemia (Brackett, Schaefer & Wilson, 1985), hepatic inflow occlusion (Kogure & Suzuki, 1992), haemorrhagic shock (Johnson, Charya, Atkins, Wiesmann & Pearce, 1997), mesenteric ischaemia (Kaszaki, Boros, Szabó & Nagy, 1994), anaphylactic shock (Mink *et al.*, 1999), and polyamine induced shock (Figueroa, Stone, Cohly, Lehan & Markov, 2001). The role of histamine in heat induced hypotension, however, is not clear.

Histamine is produced in and released from mast cells, which are found in large numbers in the skin of Sprague Dawley rats (Shakhanbeh & Abo Galyon, 1996). A variety of noxious stimuli can cause skin mast cells to release histamine. Substance P, a neuropeptide present in C fibres of the skin, can mediate such release (Dunnick, Gibran & Heimbach, 1996). In addition, histamine immunoreactive nerves have been identified in the skin of Sprague Dawley rats (Johansson, Virtanen & Hilliges, 1995). Thermal injury can cause release of histamine from the skin (Ward & Till, 1990). Serum histamine is significantly higher in rats with superficial dermal burns than in those with deep burns (Shimizu *et al.*, 2002).

Exposure to RFR at relatively lower wavelengths (2450 MHz) results in disruption of mast cells in the peritoneal fluid of rats (Valtonen, 1966). Because of the high skin temperatures reached during 35 GHz RFR exposure, it is plausible such exposure could disrupt skin mast cells. Subsequent

release of histamine may play a role in producing the circulatory failure that results.

In support of this hypothesis, Sharma & Cervos Navarro (1991) reported a beneficial effect of H₂ receptor antagonism by cimetidine during heat stress (caused by environmental heating) in rats. In addition, cimetidine can prevent harmful sequelae caused by scald injury to skin in rodents (Boykin, Eriksson, Sholley & Pittman, 1980). Pretreatment with a combination of an H₁ and H₂ antagonist attenuates harmful events (such as microvascular leakage) caused by injury in rodent skin (He & Walls, 1997).

The purpose of the current experiments was to examine the effects of combined H₁ and H₂ receptor blockade, with diphenhydramine and cimetidine, respectively, on responses to heating of anaesthetized Sprague Dawley rats with 35 GHz RFR. HR, MAP, localized body temperature changes, survival times, and lethal body temperatures were measured.

Methods

Animal care

All experiments and animal care procedures were approved by the Institutional Animal Care and Use Committee of the Air Force Research Laboratory, Brooks City Base, TX, and were conducted according to the 'Guide for the Care and Use of Laboratory Animals' prepared by the Institute of Laboratory Animal Resources National Research Council.

Male Sprague Dawley rats (total $n = 48$) were obtained from the colonies of Charles River (Wilmington, MA). They were individually housed in standard plastic cages (26 × 23 × 20.5) with water and food available *ad libitum*. Rats weighed between 350 and 400 g at the time of the experiments. A 12:12 h light dark cycle (light on at 06.00) was used, and the room temperature was maintained at 22–24 °C.

Instrumentation and preparation

Ketamine HCl (Ketalar; Parke Davis, Morris Plains, NJ, USA; 150 mg kg⁻¹, intramuscularly) was administered, with supplemental doses provided as necessary during experimentation. Ketamine administration at this dose level has been shown to produce prolonged surgical anaesthesia in Sprague Dawley rats (Smith, Pekoe, Martin & Coalgate, 1980; Frei & Jauchem, 1989; Jauchem & Frei, 1994), and results in a stable animal preparation compatible with physiological monitoring (Jauchem & Frei, 1991). Ketamine exerts minimal effects on temperature regulation in rats (Refinetti & Carlisle, 1989), and is known for its lack of significant autonomic, cardiovascular, or

respiratory effects in this animal model. Although some anaesthetics, such as enflurane and pento barbital, can affect brain histaminergic systems (which are involved in blood pressure and body temperature regulation), ketamine is ineffective in this respect in rodents (Baba, Nishibori, Oishi, Saeki & Kosaka, 1987).

Colonic temperature (T_c) was maintained at 37.0 ± 0.5 °C during all surgical procedures. A Teflon catheter (PE 50) was placed into the right jugular vein for drug administration. Additionally, the left carotid artery was instrumented with a Teflon catheter (PE 50) for measurement of arterial blood pressure and collection of arterial blood. The catheter was attached to a precalibrated blood pressure transducer (Century, Model CP 01, Inglewood, CA, USA). A lead II ECG was obtained by use of nylon covered fluorocarbon leads attached to a shielded cable placed outside the radio frequency field. All measured variables were recorded continuously throughout experimentation on a Gould TA2000 recorder (Gould Inc., Cleveland, OH, USA).

Animals were also instrumented to monitor temperature at five sites: (i) left subcutaneous (lateral, mid thoracic, side facing the RFR antenna); (ii) right subcutaneous (lateral, mid thoracic, side facing away from the RFR source); (iii) right tympanic; (iv) colonic (5–6 cm post anus); and (v) tail (subcutaneous, dorsal, 1 cm from base). The tail was placed within a 4 cm diameter tube wrapped with two layers of RFR shielding material (Bekaert Steel Wire Corp., Dyersburg, TN, USA). Thus, the tail was shielded from direct exposure to RFR and yet air was allowed to flow freely around it. All temperature measurements were obtained via thermistor probes (BSD Medical Corporation, Salt Lake City, UT, USA) attached to a precision thermometry system (BSD Medical Corporation, Model BSD 200). All temperature and cardiovascular data were A/D (analogue to digital) converted by an IBM compatible custom designed Physiological Monitoring System (Berger, Frei & Jauchem, 1991) with real time graphics display and data analysis capabilities.

RFR equipment

Continuous wave 35 GHz fields were generated by a Millimeter Wave Exposure System (Applied Electromagnetics, Inc., Atlanta, GA, USA). Irradiation was conducted under far field conditions with the animal centred along the boresight, 110 cm from the antenna. The incident power density of the field was determined at the exposure site with an electromagnetic radiation monitor (Narda Microwave Corporation, Hauppauge, NY, USA). The generator power output was monitored throughout exposure with a Model 432 B Hewlett Packard powermeter. Irradiation

was conducted in an Eccosorb RF shielded anechoic chamber at the Radio Frequency Radiation Branch of the Air Force Research Laboratory, Brooks City Base, TX. Chamber temperature and relative humidity were maintained at 27.0 ± 0.5 °C and $20 \pm 5\%$, respectively, during experimentation.

Drugs

Diphenhydramine was solubilized in 0.9% NaCl immediately prior to administration. Cimetidine was solubilized in 50% ethanol and 50% D5W prior to injection.

Experimental procedure

After surgery, the rat was placed on a Plexiglas® holder in the anechoic chamber (Frei *et al.*, 1995). The animal was then instrumented with temperature probes and cardiovascular leads for data collection. During a 5 min control period, baseline temperatures were recorded and T_c was maintained between 37.0 ± 0.5 °C using a water perfused heating pad. After the control period, the animal was exposed in the H orientation (left lateral exposure, long axis of body parallel to electric field) to 35 GHz RFR at an incident power density of 75 mW cm^{-2} . This power density will result in a whole body specific absorption rate of approximately 13 W kg^{-1} in rats of the size used in this study (Frei *et al.*, 1995). Cardiovascular and temperature parameters were continuously recorded during the control period, throughout the RFR exposure, and following exposure.

Protocol 1. In this protocol, we determined whether combined pharmacological blockade of H1 and H2 receptors could reverse the hypotension induced by sustained RFR exposure and, consequently, increase survival time following RFR exposure. After a 5 min control period, the rats were exposed to RFR until MAP decreased to 75 mmHg. Previous experiments indicated that, if irradiation is discontinued at this point, MAP will continue to decline until death (Ryan, Frei, Berger & Jauchem, 1996; Jauchem, Ryan, Lovelace & Frei, 1997; Ryan, Frei & Jauchem, 1997). We therefore discontinued RFR exposure at this point and arbitrarily defined this as the point of circulatory shock induction. Immediately after the cessation of RFR exposure, the H1 and H2 receptor antagonists diphenhydramine and cimetidine, respectively, were administered intravenously in a single bolus injection. Group 1 ($n = 8$) received a low dose of both diphenhydramine (1 mg kg^{-1}) and cimetidine (5 mg kg^{-1}). Group 2 ($n = 8$) received a higher dose of both diphenhydramine (5 mg kg^{-1}) and cimetidine (10 mg kg^{-1}). Group 3 ($n = 8$) received a bolus injection

consisting of both vehicles. Each vehicle was administered in a volume equivalent to that used for drug dilution. Each rat received only one dose of either the drugs or the vehicle. Following administration of the drug or the vehicle, the rat was monitored until death.

Protocol 2. To determine whether pretreatment with H₁ and H₂ receptor antagonists could alter the onset of circulatory shock induction, animals in this protocol received diphenhydramine and cimetidine before initiation of RFR exposure. Each animal was first placed on the Plexiglas® holder in the exposure chamber and monitored for a 5 min control period. Following this, diphenhydramine and cimetidine were administered intravenously in a manner identical to that in protocol 1. Group 4 ($n = 8$) received the low doses of diphenhydramine (1 mg kg⁻¹) and cimetidine (5 mg kg⁻¹), while group 5 ($n = 8$) received the higher doses of diphenhydramine (5 mg kg⁻¹) and cimetidine (10 mg kg⁻¹). Group 6 ($n = 8$) received vehicles. Ten minutes following the injection, RFR exposure was initiated and was continued until death.

Data analysis

Data from this study are presented as mean $\pm$ SEM. The time of exposure to RFR necessary to reach the chosen endpoint (i.e. MAP = 75 mmHg) in protocol 1 varied across individual animals (46.9 $\pm$ 1.7 min). Cardiovascular and thermal data were therefore normalized for percentage of time to endpoint in order to make valid comparisons among animals with varying durations of RFR exposure. To determine differences between vehicle *vs.* histamine receptor antagonist treated groups, a two way analysis of variance (ANOVA) with repeated measures followed by the Student Newman Keuls (SNK) multiple comparison test was performed on the normalized data (related to amount of change from time '0' baseline).

Cardiovascular and thermal data from protocol 2 are presented in absolute time intervals of 2.5 min. To determine the differences among groups, a two way ANOVA with repeated measures was applied, followed by the SNK.

For protocols 1 and 2, temperatures at each measured site before and at cessation of RFR exposure were compared using a two way ANOVA, followed by the SNK when necessary.

Results

Protocol 1: RFR exposure until MAP of 75 mmHg followed by antihistamines

Temperatures at different sites before and at the end of RFR exposure are listed in Table 1. All postexposure temperatures were significantly higher than pre exposure values. Only one significant difference was found between the different treatment groups: postexposure tail temperature was higher in the low dose drug treated animals than in the vehicle treated group.

The time courses of left subcutaneous and colonic temperatures, and of MAP and HR are shown in Figs 1 & 2, respectively. (In the high dose group, animals survived only long enough for one measurement after cessation of RFR exposure.) For each parameter, there were significant changes over time in each treatment group. There were, however, no significant differences between changes (from baseline at time zero) between groups. Although initial HR was significantly lower in high dose drug treated animals, the pattern of HR increase was similar in all groups. It is possible, however, that the lower initial HR in that group affected subsequent changes during the experiments. Administration of diphenhydramine and cimetidine (either dose level) following RFR exposure failed to reverse the hyperthermia induced hypotension, as indicated by a continued decline in MAP after drug administration. Administration of the high doses

Table 1 Colonic (T_c), tympanic (T_{tym}), left (T_{sl}) and right (T_{sr}) subcutaneous, and tail (T_t) temperatures immediately before and at the end of RFR exposure

	T_c (°C)	T_{tym} (°C)	T_{sl} (°C)	T_{sr} (°C)	T_t (°C)
Pre-RFR					
Vehicle	36.9 $\pm$ 0.1	36.3 $\pm$ 0.2	34.4 $\pm$ 0.4	35.7 $\pm$ 0.4	27.5 $\pm$ 0.3
Low dose	37.1 $\pm$ 0.1	37.0 $\pm$ 0.1	34.3 $\pm$ 0.3	35.1 $\pm$ 0.5	29.1 $\pm$ 0.3
High dose	37.1 $\pm$ 0.1	37.0 $\pm$ 0.1	34.4 $\pm$ 0.4	35.7 $\pm$ 0.2	28.9 $\pm$ 0.4
Post-RFR					
Vehicle	41.9 $\pm$ 0.3	40.9 $\pm$ 0.3	48.6 $\pm$ 0.5	38.6 $\pm$ 0.7	31.4 $\pm$ 0.7
Low dose	41.4 $\pm$ 0.4	40.6 $\pm$ 0.6	48.9 $\pm$ 0.2	37.2 $\pm$ 0.9	33.6 $\pm$ 0.6*
High dose	40.8 $\pm$ 0.3	40.5 $\pm$ 0.2	48.9 $\pm$ 0.5	36.9 $\pm$ 0.5	32.1 $\pm$ 0.5

Rats were treated with diphenhydramine + cimetidine 10 min before RFR exposure. Low dose = 1 mg kg⁻¹ diphenhydramine + 5 mg kg⁻¹ cimetidine; high dose = 5 mg kg⁻¹ diphenhydramine + 10 mg kg⁻¹ cimetidine.

*Significantly different from vehicle.




Figure 1 Effects of combined administration (following RFR exposure until MAP dropped to 75 mmHg) of diphenhydramine and cimetidine on left subcutaneous and colonic temperatures. 'Low dose' = 1 mg kg^{-1} diphenhydramine and 5 mg kg^{-1} cimetidine. 'High dose' = 5 mg kg^{-1} diphenhydramine and 10 mg kg^{-1} cimetidine. In each group, $n = 8$. Time is represented as a percentage of the total RFR exposure time to account for differences in duration of individual experiments.

of antihistamines significantly decreased the post RFR survival time, compared with vehicle treated controls (Fig. 3).

Protocol 2: Pretreatment with antihistamines followed by RFR exposure until death

Temperatures at different sites before and at the end of RFR exposure are listed in Table 2. All temperatures at death were significantly higher than predrug and pre RFR values. There were no significant differences, however, between the different treatment groups. Thus, pretreatment with diphenhydramine and cimetidine did not significantly alter either control temperatures or temperatures at the end of RFR exposure.

The time courses of left subcutaneous and colonic temperatures, and of MAP and HR are shown in Figs 4 & 5, respectively. For each parameter, there were significant changes over time in each treatment group. Pretreatment with the antihistamines produced a decrease in MAP before RFR exposure began (Fig. 5). The change in MAP from time '0' baseline in animals receiving antihistamines at the higher doses remained significantly depressed compared with that of vehicle treated animals during the first 35 min of RFR exposure. Antihistamine pretreatment did not alter the total RFR exposure time required for death to occur (vehicle $59.8 \pm 2.2 \text{ min}$, low dose $60.0 \pm 1.3 \text{ min}$, high dose $63.0 \pm 3.7 \text{ min}$).




Figure 2 Effects of combined administration (following RFR exposure until MAP dropped to 75 mmHg) of diphenhydramine and cimetidine on mean arterial pressure (MAP) and heart rate. 'Low dose' = 1 mg kg⁻¹ diphenhydramine and 5 mg kg⁻¹ cimetidine. 'High dose' = 5 mg kg⁻¹ diphenhydramine and 10 mg kg⁻¹ cimetidine. In each group, $n = 8$. Time is represented as a percentage of the total RFR exposure time to account for differences in duration of individual experiments.





Figure 3 Effects of combined administration (following RFR exposure until MAP dropped to 75 mmHg) of diphenhydramine and cimetidine on RFR exposure time (i.e. time from beginning of exposure and drug until mean arterial pressure decreased to 75 mmHg) and subsequent survival time. 'Low dose' = 1 mg kg⁻¹ diphenhydramine and 5 mg kg⁻¹ cimetidine. 'High dose' = 5 mg kg⁻¹ diphenhydramine and 10 mg kg⁻¹ cimetidine. In each group, $n = 8$. *Significantly different from vehicle.

Table 2 Colonic (T_c), tympanic (T_{tym}), left (T_{sl}) and right (T_{sr}) subcutaneous, and tail (T_t) temperatures immediately before drug infusion and RFR exposure, and at death

	T_c (°C)	T_{tym} (°C)	T_{sl} (°C)	T_{sr} (°C)	T_t (°C)
Pre-drug					
Vehicle	37.0 ± 0.1	36.5 ± 0.1	34.0 ± 0.5	34.0 ± 0.5	27.9 ± 0.4
Low dose	37.2 ± 0.0	37.0 ± 0.1	34.6 ± 0.3	35.2 ± 0.6	28.8 ± 0.7
High dose	37.0 ± 0.1	36.7 ± 0.1	35.0 ± 0.4	35.0 ± 0.5	28.6 ± 0.3
Pre-RFR					
Vehicle	36.8 ± 0.1	36.3 ± 0.1	34.4 ± 0.6	33.7 ± 0.5	27.7 ± 0.2
Low dose	36.9 ± 0.1	36.6 ± 0.1	34.6 ± 0.3	34.6 ± 0.5	28.6 ± 0.5
High dose	36.5 ± 0.1	36.1 ± 0.1	33.7 ± 0.4	33.8 ± 0.6	29.8 ± 0.5
At death					
Vehicle	42.5 ± 0.3	41.5 ± 0.3	49.7 ± 0.6	36.7 ± 1.1	32.5 ± 0.8
Low dose	41.7 ± 0.2	40.9 ± 0.4	50.0 ± 0.3	36.6 ± 1.8	30.9 ± 0.3
High dose	42.5 ± 0.6	41.4 ± 0.4	48.4 ± 1.1	36.8 ± 1.8	32.2 ± 1.3

Rats were treated with diphenhydramine + cimetidine 10 min before RFR exposure. Low dose = 1 mg kg⁻¹ diphenhydramine + 5 mg kg⁻¹ cimetidine; high dose = 5 mg kg⁻¹ diphenhydramine + 10 mg kg⁻¹ cimetidine.

**Figure 4** Effects of combined administration of diphenhydramine and cimetidine, followed by RFR exposure, on left subcutaneous and colonic temperatures. 'Low dose' = 1 mg kg⁻¹ diphenhydramine and 5 mg kg⁻¹ cimetidine. 'High dose' = 5 mg kg⁻¹ diphenhydramine and 10 mg kg⁻¹ cimetidine. In each group, $n = 8$.




Figure 5 Effects of combined administration of diphenhydramine and cimetidine, followed by RFR exposure, on mean arterial pressure (MAP) and heart rate. 'Low dose' = 1 mg kg^{-1} diphenhydramine and 5 mg kg^{-1} cimetidine. 'High dose' = 5 mg kg^{-1} diphenhydramine and 10 mg kg^{-1} cimetidine. In each group, $n = 8$. *Significant difference between high dose and vehicle.

Discussion

Overall pattern of HR and MAP changes during RFR exposure and heating

The biphasic change (initial increase followed by decrease) in MAP, and the gradual rise in HR throughout the exposure period, were similar to changes previously found by other investigators during experiments of lethal environmental heat stress in anaesthetized rats (Kielblock, Strydom, Burger, Pretorius & Manjoo, 1982; Kregel & Gisolfi, 1990). In earlier studies of terminal exposures of anaesthetized rats to 2.45 GHz (Jauchem & Frei, 1994; Jauchem, Frei, Chang & Berger, 1995; Jauchem, Chang & Frei, 1996), 2.8 GHz (Jauchem, Frei & Heinmets, 1984), and 5.6 GHz RFR (Jauchem, Frei & Heinmets, 1983, 1988), the pattern of HR and MAP changes was also qualitatively comparable.

The pattern of HR and MAP changes during hyperthermia could be the result of diminished venous return as a result of massive venodilation, which could cause decreased stroke volume with a subsequent drop in cardiac output. Miki, Morimoto, Nose, Itoh & Yamada (1983) observed a prominent reduction in central venous pressure during severe hyperthermia. The mechanisms controlling such a decrease in pressure during heat stress are unclear. This diminished venous pressure, along with an accompanying decrease in cardiac filling, may be the limiting factor in survival during heat stress. Kregel & Gisolfi (1990) suggested that heating could prevent vasomotor stimuli from exerting an effect on the contractile apparatus of blood vessels. Takamata, Nose, Mack & Morimoto (1990) also suggested that the sudden decrease in MAP during hyperthermia may be due to a dysfunction in the peripheral circulation and a reduction in

venous return. Previously, Morimoto & Nose (1984) reported that vascular compliance is lower during hyperthermia.

A paradoxical bradycardia can develop in severe hypotensive hypovolaemic shock (Secher, Jensen, Werner, Warberg & Bie, 1984). In contrast to this, in experiments dealing with terminal heat stress, whether due to environmental exposure (Kielblock *et al.*, 1982; Kregel & Gisolfi, 1990) or RFR (as in the present experiments), tachycardia continued until death.

Effects of antihistamines

As the tail was shielded from direct exposure to RFR, any increase in tail temperature would be likely to be related to circulatory transfer of heat. During environmental heating, diversion of blood to the tail occurs with an increase in body temperature (Grant, 1963). The surface area to volume relationship of the tail allows for rapid dissipation of heat into the environment. Histamine induced relaxation of isolated rat tail arteries (synonymous with vasodilation) involves an endothelium independent mechanism that is sensitive to cimetidine (Grifoni & Brendhack, 2000). In our current study, postexposure tail temperature was elevated in the low dose (administered following RFR exposure) drug treated animals, compared with the vehicle treated group. Such an increase in temperature could be related to vasodilation in the tail. It is not known, however, why high dose antihistamine treated animals did not show an increase in tail temperature.

The actions of H 1 and H 2 antagonists on body temperature changes are dependent on the route of administration (Dey & Mukhopadhyaya, 1986). Cimetidine injection can inhibit hypothermia in rats (Brus, Krzeminski, Juraszczyk, Kurcok & Felinska, 1985). Conversely, chronic administration of diphenhydramine has been shown to block hyperthermia in rats (Nath, Patnaik, Saxena & Gupta, 1997). Dhawan, Shukla & Srimal (1982) presented a more complicated picture of effects of histamine on body temperature, with low and high doses of histamine causing hypothermia and hyperthermia, respectively. They concluded that H 1 receptors mediate hypothermia, while H 2 receptors mediate hyperthermia. Nowak (1982), however, reported that cimetidine, although blocking the hypothermic action of an H 2 agonist, had no effect on body temperature when administered alone. Although there were no significant differences in temperatures at death between groups in the present study, post RFR survival time was significantly decreased in the high dose (administered following RFR exposure) antihistamine group. This effect was unexpected. In fact, Brackett *et al.* (1985) found that combined treatment with both diphenhydramine and cimetidine resulted in

increased survival time during endotoxin induced shock.

In our experiments, the decrease in MAP in animals after pretreatment with antihistamines was consistent with findings by Paakkari, Totterman, Kupari, Karppanen & Paakkari (1982). Those investigators reported that i.v. cimetidine induced hypotension in anaesthetized rats; diphenhydramine pretreatment did not antagonize the hypotensive effect. It should be noted that the doses of antihistamines used in our experiments were similar to those in previous studies of diphenhydramine (e.g. Dimlich, 1985; Ulloa & Zaninovich, 1986; Carceller *et al.*, 1994; Wan, Wetmore, Sorensen, Greenberg & Nance, 1994) and cimetidine (e.g. Ezeamuzie, Nwankwoala & Emenike, 1986; Ulloa & Zaninovich, 1986; Couture, Gupta, K  rouac & Regoli, 1987) administered intravenously to rats by other investigators.

Other investigators have suggested several potential mechanisms of hypotensive effects of both H 1 and H 2 antagonists. In rats, H 1 antagonism can attenuate histamine induced release of noradrenaline (Bealer, 1993) or arginine vasopressin (Knigge, Willems, Kjaer, J  rgensen & Warberg, 1999). As noradrenaline and arginine vasopressin could each cause increases in blood pressure, inhibition of such release may contribute to hypotension, depending on interaction of other factors. In addition, humans with certain conditions react to administration of H 1 receptor antagonists with severe hypotension that is reversible by i.v. adrenaline (Roberts, Turk & Oates, 1982). Regarding H 2 antagonism, one potential mechanism of the hypotensive action of i.v. cimetidine is vasodilation (with a corresponding decrease in peripheral resistance), in both rats (Paakkari *et al.*, 1982) and humans (Totterman, Kupari, Paakkari & Nieminen, 1985). Due to the complex nature of possible mechanisms and interactions of different substances, it would be difficult to develop a unifying hypothesis for the hypotensive effects of high dose histamine antagonism in the present experiments.

The lack of a beneficial effect of antihistamines in our study is consistent with findings of Johnson *et al.* (1997), who used a different model of cardiovascular collapse: haemorrhagic shock in the Sprague Dawley rat. As in our study, those investigators used diphenhydramine and cimetidine as histamine antagonists. They concluded that histamine is not an important mediator of cardiovascular sequelae seen in the decompensatory phase of shock, although histamine levels rise coincidentally during this phase. The lack of a beneficial effect of antihistamines in our study does negate the possibility of some involvement of histamine in the overall responses. The model of hyperthermia was of a rather severe nature, and partial activation of some histamine receptors could still have occurred.

Possible role of other endogenous circulating factors during RFR exposure and heating

The role of other endogenous circulating factors in heat induced circulatory collapse (whether due to environmental or RFR heating) has been studied previously. McMeeken & Bell (1990) reported a pressor effect of RFR exposure and presumed that reflex adrenomedullary activation was involved, as plasma catecholamines were elevated. Gisolfi, Matthes, Kregel & Oppliger (1991) found that changes in sympathetic nerve activity and plasma catecholamines were not associated with mesenteric vasodilation that precedes heatstroke. Ryan, Jauchem, Tehrany & Boyle (2002) previously found that platelet activating factor was not associated with RFR induced circulatory failure. Nitric oxide was not associated with either RFR (Ryan *et al.*, 1996) or environmental heat induced (Ryan, Tehrany & Jauchem, 2001) circulatory collapse. McMeeken & Bell (1990) did not find any evidence that prostaglandins or kinins were involved in vasodilation that occurred during RFR exposure.

A marked increase in plasma levels of vasopressin occurs during heatstroke after heavy exercise (Aarseth, Eide, Skeie & Thaulow, 1986). In addition, circulating angiotensin converting enzyme activities in exertional heatstroke patients on admission were significantly lower than in normal subjects (Shieh, Shiang, Lin, Shiao & Wang, 1995). Effects of non exertional heating on vasopressin and the angiotensin system, however, have not been elucidated. It will be important to study these and other factors in future investigations of environmental and RFR induced heating.

Summary

In summary, pharmacological blockade of H₁ and H₂ receptors is not beneficial in anaesthetized rats made hypotensive by RFR exposure. This indicates that activation of H₁ and H₂ receptors by histamine does not occur to any significant extent and does not mediate the hypotensive response developed in this model of hyperthermia.

Acknowledgments

The authors wish to acknowledge the assistance of Michael C. Cook in producing the graphics in this paper. The views and opinions expressed in this paper are those of the authors and do not necessarily state or reflect those of the U.S. Government.

References

- AARSETH, H.P., EIDE, I., SKEIE, B. & THAULOW, E. (1986). Heat stroke in endurance exercise. *Acta Med. Scand.*, **220**, 279–283.

- BABA, M., NISHIBORI, M., OISHI, R., SAEKI, K. & KOSAKA, F. (1987). Effects of halothane, enflurane and pentobarbital on brain histamine dynamics in mice. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **335**, 686–691.
- BEALER, S.L. (1993). Histamine releases norepinephrine in the paraventricular nucleus/anterior hypothalamus of the conscious rat. *J. Pharmacol. Exp. Ther.*, **264**, 734–738.
- BERGER, R.E., FREI, M.R. & JAUCHEM, J.R. (1991). Computer based monitoring of thermoregulatory responses of anesthetized rats exposed to radiofrequency radiation. *Physiologist*, **34**, 246.
- BOYKIN, J.V., JR., ERIKSSON, E., SHOLLEY, M.M. & PITTMAN, R.N. (1980). Histamine mediated delayed permeability response after scald burn inhibited by cimetidine or cold water treatment. *Science*, **209**, 815–817.
- BRACKETT, D.J., SCHAEFER, C.F. & WILSON, M.F. (1985). The effects of H₁ and H₂ histamine receptor antagonists on the development of endotoxemia in the conscious, unrestrained rat. *Circ. Shock*, **16**, 141–153.
- BRUS, R., KRZEMINSKI, T., JURASZCZYK, Z., KURCOK, A. & FELINSKA, W. (1985). Influence of H₂ receptors blockade upon some central effects of prostacyclin in rats. *Pol. J. Pharmacol. Pharm.*, **37**, 831–839.
- CARCELLER, E., MERLOS, M., GIRAL, M., BALSÀ, D., ALMANSA, C., BARTOLÍ, J., GARCÍA-RAFANELL, J. & FORN, J. (1994). [(3 pyridylalkyl)piperi dylidene]benzocycloheptapyridine derivatives as dual antagonists of PAF and histamine. *J. Med. Chem.*, **37**, 2697–2703.
- COUTURE, R., GUPTA, A., KÉROUAC, R. & REGOLI, D. (1987). Studies on the cardiovascular effects produced by the spinal action of two substance P analogues in the rat: evidence for a central catecholaminergic mechanism. *Eur. J. Pharmacol.*, **135**, 345–354.
- DEY, P.K. & MUKHOPADHAYA, N. (1986). Involvement of histamine receptors in mediation of histamine induced thermoregulatory response in rats. *Indian J. Physiol. Pharmacol.*, **30**, 300–306.
- DHAWAN, B.N., SHUKLA, R. & SRIMAL, R.C. (1982). Analysis of histamine receptors in the central thermoregulatory mechanism of *mastomys natalensis*. *Br. J. Pharmacol.*, **75**, 145–149.
- DIMLICH, R.V. (1985). Histamine antagonists in the treatment of shock hyperglycemia in the rat. *Ann. Emerg. Med.*, **14**, 91–96.
- DUNNICK, C.A., GIBRAN, N.S. & HEIMBACH, D.M. (1996). Substance P has a role in neurogenic mediation of human burn wound healing. *J. Burn Care Rehabil.*, **17**, 390–396.
- EZEAMUZIE, I.C., NWANKWOALA, R.N.P. & EMENIKE, O. (1986). Dextran induced gastric secretion in rats. *Scand. J. Gastroenterol. (Suppl.)*, **124**, 145–148.
- FIGUEROA, A.H., STONE, R.H., COHLY, H.H.P., LEHAN, P.H. & MARKOV, A.K. (2001). Effect of fructose 1,6 diphosphate versus diphenhydramine on mortality in compound 48/80 induced shock. *Toxicol. Lett.*, **122**, 141–148.
- FREI, M.R. & JAUCHEM, J.R. (1989). Effects of 2.8 GHz microwaves on restrained and ketamine anesthetized rats. *Radiat. Environ. Biophys.*, **28**, 155–164.
- FREI, M.R., RYAN, K.L., BERGER, R.E. & JAUCHEM, J.R. (1995). Sustained 35 GHz radiofrequency irradiation induces circulatory failure. *Shock*, **4**, 289–293.
- GISOLFI, C.V., MATTHES, R.D., KREGEL, K.C. & OPPLIGER, R. (1991). Splanchnic sympathetic nerve activity and

- circulating catecholamines in the hyperthermic rat. *J. Appl. Physiol.*, **70**, 1821–1826.
- GRANT, R.T. (1963). Vasodilation and body warming in the rat. *J. Physiol. (London)*, **167**, 311–317.
- GRIFONI, S.C. & BRENDHACK, L.M. (2000). Relaxation induced by histamine is not endothelium dependent in tail arteries of L NAME treated rats. *Gen. Pharmacol.*, **34**, 435–441.
- HE, S. & WALLS, A.F. (1997). Human mast cell tryptase: a stimulus of microvascular leakage and mast cell activation. *Eur. J. Pharmacol.*, **328**, 89–97.
- JAUCHEM, J.R. & FREI, M.R. (1991). Cardiovascular changes in unanesthetized and ketamine anesthetized Sprague Dawley rats exposed to 2.8 GHz radiofrequency radiation. *Lab. Anim. Sci.*, **41**, 70–75.
- JAUCHEM, J.R. & FREI, M.R. (1992). Heart rate and blood pressure changes during radiofrequency irradiation and environmental heating. *Comp. Biochem. Physiol.*, **101A**, 1–9.
- JAUCHEM, J.R. & FREI, M.R. (1994). Cardiorespiratory changes during microwave induced lethal heat stress and β adrenergic blockade. *J. Appl. Physiol.*, **77**, 434–440.
- JAUCHEM, J.R., FREI, M.R. & HEINMETS, F. (1983). Terminal exposure to 5.6 GHz microwaves: a comparison of continuous wave and pulsed radiation. *Preprints of the 54th Annual Scientific Meeting of the Aerospace Medical Association*, pp. 241–242.
- JAUCHEM, J.R., FREI, M.R. & HEINMETS, F. (1984). Increased susceptibility to radiofrequency radiation due to pharmacological agents. *Aviat. Space Environ. Med.*, **55**, 1036–1040.
- JAUCHEM, J.R., FREI, M.R. & HEINMETS, F. (1988). Thermal responses to 5.6 GHz radiofrequency radiation in anesthetized rats: effect of chlorpromazine. *Physiol. Chem. Phys. Med. NMR*, **20**, 135–143.
- JAUCHEM, J.R., FREI, M.R., CHANG, K.D. & BERGER, R.E. (1995). Microwave induced lethal heat stress: effects of phentolamine, prazosin, and metoprolol. *Methods Find. Exp. Clin. Pharmacol.*, **17**, 241–248.
- JAUCHEM, J.R., CHANG, K.D. & FREI, M.R. (1996). Tolazoline decreases survival time during microwave induced lethal heat stress in anesthetized rats. *Proc. Soc. Exp. Biol. Med.*, **211**, 236–243.
- JAUCHEM, J.R., RYAN, K.L., LOVELACE, J.D. & FREI, M.R. (1997). Effects of esmolol on 35 GHz microwave induced lethal heat stress. *J. Auton. Pharmacol.*, **17**, 165–174.
- JOHANSSON, O., VIRTANEN, M. & HILLIGES, M. (1995). Histaminergic nerves demonstrated in the skin. A new direct mode of neurogenic inflammation? *Exp. Dermatol.*, **4**, 93–96.
- JOHNSON, K.B., CHARYA, R.V., ATKINS, J.L., WIESMANN, W.P. & PEARCE, F.J. (1997). The role of histamine in mediating the decompensatory phase of hemorrhagic shock in the rat. *Shock*, **8**, 444–449.
- KASZAKI, J., BOROS, M., SZABÓ, A. & NAGY, S. (1994). Role of histamine in the intestinal flow response following mesenteric ischemia. *Shock*, **2**, 413–420.
- KIELBLOCK, A.J., STRYDOM, N.B., BURGER, F.J., PRETORIOUS, P.J. & MANJOO, M. (1982). Cardiovascular origins of heatstroke pathophysiology: an anesthetized rat model. *Aviat. Space Environ. Med.*, **53**, 171–178.
- KNIGGE, U., WILLEMS, E., KJAER, A., JØRGENSEN, H. & WARBERG, J. (1999). Histaminergic and catecholaminergic interactions in the central regulation of vaso-pressin and oxytocin secretion. *Endocrinology*, **140**, 3713–3719.
- KOGURE, K. & SUZUKI, M. (1992). Effects of hepatic inflow occlusion on changes in plasma potassium, histamine, and norepinephrine in rats. *Circ. Shock*, **36**, 290–298.
- KREGEL, K.C. & GISOLFI, C.V. (1990). Circulatory responses to vasoconstrictor agents during passive heating in the rat. *J. Appl. Physiol.*, **68**, 1220–1227.
- MCMECKEN, J.M. & BELL, C. (1990). Effects of microwave irradiation on blood flow in the dog hindlimb. *Exp. Physiol.*, **75**, 367–374.
- MIKI, K., MORIMOTO, T., NOSE, H., ITOH, T. & YAMADA, S. (1983). Canine blood volume and cardiovascular function during hyperthermia. *J. Appl. Physiol.*, **55**, 300–306.
- MINK, S., BECKER, A., SHARMA, S., UNRUH, H., DUKE, K. & KEPRON, W. (1999). Role of autacoids in cardiovascular collapse in anaphylactic shock in anesthetized dogs. *Cardiovasc. Res.*, **43**, 173–182.
- MORIMOTO, T. & NOSE, H. (1984). Changes in vascular compliance during hyperthermia. *J. Thermal Biol.*, **9**, 149–151.
- NATH, C., PATNAIK, G.K., SAXENA, R.C. & GUPTA, M.B. (1997). Evaluation of inhibitory effect of diphenhydramine on benzodiazepine dependence in rats. *Indian J. Physiol. Pharmacol.*, **41**, 42–46.
- NOWAK, J.Z. (1982). Impromidine induced hypothermia in rats: effect of cimetidine and mianserine. *Agents Actions*, **12**, 148–152.
- PAAKKARI, I., TOTTERMAN, K.J., KUPARI, M., KARPPANEN, H. & PAAKKARI, P. (1982). Peripheral hypotensive and central hypertensive effects of cimetidine. *Agents Actions*, **12**, 152–155.
- REFINETTI, R. & CARLISLE, H.J. (1989). Thermoregulation during pentobarbital and ketamine anesthesia in rats. *J. Physiol. (Paris)*, **83**, 300–303.
- ROBERTS, L.J., II, TURK, J.W. & OATES, J.A. (1982). Shock syndrome associated with mastocytosis: pharmacologic reversal of the acute episode and therapeutic prevention of recurrent attacks. *Adv. Shock Res.*, **8**, 145–152.
- RYAN, K.R., FREI, M.R., BERGER, R.E. & JAUCHEM, J.R. (1996). Does nitric oxide mediate circulatory failure induced by 35 GHz microwave heating? *Shock*, **6**, 71–76.
- RYAN, K.R., FREI, M.R. & JAUCHEM, J.R. (1997). Circulatory failure induced by 35 GHz microwave heating: effects of chronic nitric oxide synthesis inhibition. *Shock*, **7**, 70–76.
- RYAN, K.R., D'ANDREA, J.A., JAUCHEM, J.R. & MASON, P.A. (2000). Radio frequency radiation of millimeter wave length: potential occupational safety issues relating to surface heating. *Health Phys.*, **78**, 170–181.
- RYAN, K.L., TEHRANY, M.R. & JAUCHEM, J.R. (2001). Nitric oxide does not contribute to the hypotension of heatstroke. *J. Appl. Physiol.*, **90**, 961–970.
- RYAN, K.L., JAUCHEM, J.R., TEHRANY, M.R. & BOYLE, H.L. (2002). Platelet activating factor does not mediate circulatory failure induced by 35 GHz microwave heating. *Methods Find. Exp. Clin. Pharmacol.*, **24**, 279–286.
- SECHER, N.H., JENSEN, K.S., WERNER, C., WARBERG, J. & BIE, P. (1984). Bradycardia during severe but reversible hypovolemic shock in man. *Circ. Shock*, **14**, 267–274.
- SHAKHANBEH, J. & ABO-GALYON, Y. (1996). Effects of inflammation on plasma extravasation and mast cell response in normal and denervated rat skin. *Skin Pharmacol.*, **9**, 203–210.

- SHARMA, H.S. & CERVOS-NAVARRO, J. (1991). Role of histamine in pathophysiology of heat stress in rats. *Agents Actions*, **33** (Suppl.), 97–102.
- SHIEH, S.D., SHIANG, J.C., LIN, Y.F., SHIAO, W.Y. & WANG, J.Y. (1995). Circulating angiotensin converting enzyme, von Willebrand factor antigen and thrombomodulin in exertional heat stroke. *Clin. Sci. (Lond.)*, **89**, 261–265.
- SHIMIZU, S., TANAKA, H., SAKAKI, S., YUKIOKA, T., MATSUDA, H. & SHIMAZAKI, S. (2002). Burn depth affects dermal interstitial fluid pressure, free radical production, and serum histamine levels in rats. *J. Trauma*, **52**, 683–687.
- SMITH, D.J., PEKOE, G.M., MARTIN, L.L. & COALGATE, B. (1980). The interaction of ketamine with the opiate receptor. *Life Sci.*, **26**, 789–795.
- TAKAMATA, A., NOSE, H., MACK, G.W. & MORIMOTO, T. (1990). Control of total peripheral resistance during hyperthermia in rats. *J. Appl. Physiol.*, **69**, 1087–1092.
- TOTTERMAN, K., KUPARI, M., PAAKKARI, I. & NIEMINEN, M.S. (1985). Acute cardiovascular effects of intravenous cimetidine. *Acta Med. Scand.*, **217**, 277–280.
- ULLOA, E.R. & ZANINOVICH, A.A. (1986). Effects of histamine H1 and H2 receptor antagonists on thyrotrophin secretion in the rat. *J. Endocrinol.*, **111**, 175–180.
- VALTONEN, E.J. (1966). The effects of microwave radiation on the cellular elements in the peritoneal fluid and peripheral blood of the rat. *Acta Rheum. Scand.*, **12**, 291–299.
- WAN, W., WETMORE, L., SORESENSEN, C.M., GREENBERG, A.H. & NANCE, D.M. (1994). Neural and biochemical mediators of endotoxin and stress induced c fos expression in the rat brain. *Brain Res. Bull.*, **34**, 7–14.
- WARD, P.A. & TILL, G.O. (1990). Pathophysiologic events related to thermal injury of skin. *J. Trauma.*, **30** (Suppl. 12), S75–S79.

(Received 20 October 2003

Revised 13 May 2004

Accepted 12 June 2004)

Copyright of Autonomic & Autacoid Pharmacology is the property of Blackwell Publishing Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.