

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP020036

TITLE: Photo-Oxidation From Mode-Locked Laser Exposure to
hTERT-RPE1 Cells

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: Laser Interaction with Tissue and Cells XV. Held in San Jose, Ca
on 26-28 January 2004.

To order the complete compilation report, use: ADA436676

The component part is provided here to allow users access to individually authored sections
of proceedings, annals, symposia, etc. However, the component should be considered within
the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP020007 thru ADP020056

UNCLASSIFIED

Photo-oxidation from Mode-locked Laser Exposure to hTERT-RPE1 Cells[‡]

Michael L. Denton^a, Debbie M. Eikum^a, Gary D. Noojin^a, David J. Stolarski^a, Randolph D. Glickman^b, and Benjamin A. Rockwell^c

^aNorthrop Grumman Information Technology, San Antonio, TX, USA, 78228-1330

^bDepartment of Ophthalmology, University of Texas Health Science Center,
San Antonio, TX, USA, 78284-6230

^cAir Force Research Laboratory, AFRL/HEDO,
Brooks Air Force Base, TX, USA, 78235-5214

ABSTRACT

Human retinal pigment epithelial (RPE) cells (hTERT-RPE1) were used to detect photo-oxidation products generated from chronic NIR (810 nm) laser exposure. Exposure of a discrete area within cell monolayers provided a means of distinguishing fluorescence above background levels. Oxidative stress was detected using the fluorescent dye H₂DCF-DA and its analog CM-H₂DCF-DA. Fluorescence was detected in cells exposed to mode-locked (76 MHz, ~160 femtoseconds) but not CW laser exposure. Detection of photo-oxidation from the mode-locked laser was dependent upon radiant exposure, but only if irradiance was greater than a threshold value. The CM-H₂DCF-DA dye proved a more sensitive indicator of oxidation than H₂DCF-DA, and the radiant exposure threshold for detection was dependent upon dye concentration. No oxidation was detected from CW exposures (using the most sensitive fluorescent dye conditions) when using 3 times the irradiance, and 10 times the radiant exposure needed to detect fluorescence from mode-locked exposure.

Keywords: photochemical oxidation, hTERT-RPE1 cells, mode-locked laser, fluorescence microscopy

INTRODUCTION

Retinal pigment epithelial (RPE) cells perform many functions in the eye. With a dense layer of intracellular melanosomes, the RPE act as a backstop for all light transmitted to the retina. They serve as a barrier between the neural retina and the blood supply of the choriocapillaris, filtering substrates and products of metabolism for the neural retinal cells. By nature of their phagocytic functions the RPE cells chemically recycle the membrane and vision pigment components of the rod and cone outer segments. These *in vivo* functions subject the RPE cells to chronic metabolic and oxidative stress. Through a combination of *in vivo* analysis and *in vitro* studies, knowledge of how this combination of factors leads to increased amounts of intracellular reactive oxygen intermediates (ROI), and how these compounds may lead to pathology,¹ is becoming increasingly clear.

Various *in vitro* models for studying oxidative stress in RPE cells have been established, utilizing either high doses of chemical oxidants,²⁻⁵ chronic mild hyperoxia,⁶ or blue-light illumination.^{7,8} For other cell types, there are indications that the damaging effects of violet/blue light is mediated by the production of hydrogen peroxide (H₂O₂), as a consequence of the photoreduction of flavins within oxidase enzymes.⁹

[‡] Opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the United States Air Force.

We are interested in laser bioeffects. Applications of mode-locked lasers have been increasing of late. For instance, in the field of fluorescence microscopy mode-locked lasers are used to generate multiphoton absorption of fluorescent probes, providing several advantages over single-photon excitation.¹⁰ Because mode-locked lasers generate beams with high peak-irradiance per pulse we have investigated their potential for causing damaging biological effects by comparison to CW lasers of the same average power. Our findings suggest that near-IR (NIR) mode-locked lasers do produce *in situ* production of photochemical oxidation in RPE cells, as detected by a fluorescent dye probe. We characterize the response of the cell line hTERT-RPE1 to laser exposure at 810 nm by measuring threshold irradiances and threshold radiant exposures for various concentrations (preloaded into the cells) of two dichlorofluorescein dyes.

METHODS AND MATERIALS

Cell Culture

All cell cultures were grown at 37°C in an atmosphere containing 5% CO₂. Stock cultures of the human RPE cell line hTERT-RPE1 (ClonTech) were maintained in DMEM/F12 medium supplemented with 10% fetal bovine serum (FBS; Atlanta Biochemicals) and were passaged at, or prior to confluence. At 2-9 days prior to exposure cells were plated into 48-well microtiter dishes (~90,000 cells/cm²).

Fluorescent dyes (Molecular Probes, Portland, OR) used in this study were dichlorodihydrofluorescein diacetate (H₂DCFDA; cat # D-399) and its 5-(and-6)-chloromethyl- derivative CM-H₂DCFDA (cat # C-6827) for detection of oxidation, and the combination of calcein-AM (cat# C-3099) with ethidium homodimer-1 (EthD-1, cat# E-1169) for viability staining.

Laser exposure

CW laser exposures (0.25 sec) at 405 nm were carried out¹² using a Coherent INNOVA-100 krypton laser where irradiance ranged from 2400 W/cm² to 15 W/cm². A Coherent MIRA900F titanium:sapphire laser provided CW and mode-locked beams at 810 nm. The repetition rate and pulse duration for the mode-locked beam were 76 MHz and ~200 fsec, respectively. Maximum irradiance from the MIRA was 27,000 W/cm² for the mode-locked beam and 35,000 W/cm² for the CW beam.

All laser exposures were performed in 48-well plates. Exposure durations at 810 nm ranged from 2 to 600 sec and beam diameters were set at 70 μm. The CW 405 nm beam was 140 μm in diameter. The physical formation of the laser exposure was a centrally located concentric ring formed by 7 individual laser exposures, creating an overall exposure area (figure 1) with diameters of about 340 μm (405 nm) and 170 μm (810 nm). This provided a defined region of fluorescence above background if the laser-induced oxidation was above the minimum level of detection when observed by microscopy.




Figure 1. Pattern of individual laser exposures forming the overall exposure area. Beams were separated by a 1/e spacing distribution.

Just prior to exposing, cells in experimental and mock wells were washed twice with Buffer (phosphate buffered saline [PBS] or Hank's balanced salt solution [HBSS]), allowed to pre-load with a fluorescent oxidation-sensitive dye (diluted in buffer) for 20 min in the culture incubator, and finally rinsed with 1 ml

fresh buffer. Control (no treatment) wells were treated in the same fashion except their cells were not subjected to fluorescent dye. Exposures were carried out in 0.1 ml buffer. Exposed cells were viewed directly using fluorescence microscopy. Fluorescent dye was either H₂DCFDA or CM-H₂DCFDA at varying concentrations from 10 μ M to 100 μ M.

Microscopy

After laser exposure cells were visualized and imaged on an inverted phase-contrast microscope set up for fluorescence (Olympus CK-40F; Leeds Instruments, Inc, Irving, TX). Photography was done with a Hamamatsu ORCA-100 digital camera with the resulting images processed using Simple PCI (C-Imaging Systems). Fluorescence from both DCF-based dyes was detected using a bandpass exciter of 460-490 nm and a bandpass emitter of either 515-555 nm or 490-530 nm.

RESULTS

Detecting laser-induced oxidation in hTERT-RPE1

Using the oxidation-sensitive fluorescent probe dichlorodihydrofluorescein diacetate (H₂DCFDA) we detected oxidative stress *in situ* in cultured RPE cells exposed (0.25 sec) to a CW laser at 405 nm.¹² The use of small exposure areas (see Figure 1) provided graphic representation of where laser-induced oxidation took place, within the detection limits of fluorescence using our dye conditions, microscope, and attached CCD camera. If laser exposure produced oxidation (presumably H₂O₂) within the cells of these targets, the H₂DCF is converted to the excitable species (DCF) that is visualized by excitation with blue light from the microscope. Targets with significant DCF appeared immediately during fluorescence microscopy on a black background. For exposures (0.25 sec) at 405 nm, cells were first preloaded with 10 μ M H₂DCFDA. Figure 2 shows a fluorescent "target" in the RPE monolayer corresponding to the laser exposed area. As further proof that the oxidation "target" is laser-induced, figure 2 also provides an image of "background" fluorescence due solely from blue light excitation at the microscope (Panel D). Threshold irradiance required to generate oxidized targets under these conditions (see ref. 12) was 38 W/cm² which correlated to 9.5 J/cm² delivered. Uncertainty was 8% for measuring both irradiance and radiant exposure.

Detecting NIR laser-induced oxidation in hTERT-RPE1

No definitive oxidation targets were generated from 810 nm exposures using the same conditions giving rise to the CW 405 nm oxidation results described above. By increasing both time of exposure and the concentration of dye during preloading of the cells we were able to detect oxidation products at 810 nm, but only for mode-locked beams. We again used a small exposure area to allow detection of specific laser-induced oxidation above background levels during fluorescence microscopy. Our analysis utilized data collected from several different experimental days. Prior to exposures each day, the laser was aligned to the center of wells to be exposed to facilitate quick identification of even faint regions of oxidation. Damage was scored as either a plus (initial fluorescence) or minus (no initial fluorescence) when observed on the fluorescent microscope. For scoring wells exposed to near threshold energies an experimentally "blind" approach was used. Figure 3 provides an example of an oxidation "target" generated by mode-locked NIR laser exposure (21,800 W/cm² for 100 sec) using 100 μ M H₂DCFDA. In addition to the H₂DCFDA, we successfully detected oxidation targets using its 5-(and-6)-chloromethyl- derivative CM-H₂DCFDA.

Although the damage targets from NIR mode-locked laser exposure were genuine (fluorescence above background in the shape of the expected target) they were not as sharply demarked as those generated by the (positive control) 405 nm laser.

The sensitivity of our system was clearly dependent upon the choice of dye and its concentration applied to the cells (Table 1). Due to its cellular retention properties (see Handbook for Molecular Probes; Molecular Probes, Portland, OR), CM-H₂DCFDA provided a much more sensitive detection system than the




Figure 2. 405 nm CW laser-induced DCF fluorescence in hTERT RPE cells.¹² A: DCF fluorescence image; B: phase-contrast image; C: image of DCF fluorescence overlaying phase-contrast; D: image of DCF fluorescence in unexposed cells upon prolonged excitation on microscope.




Figure 3. NIR (810 nm) mode-locked laser-induced DCF fluorescence in hTERT-RPE1 cells. A: image of DCF fluorescence overlaying phase-contrast; B: DCF fluorescence image.

H₂DCFDA. Using our most sensitive conditions for detecting cellular oxidation (cells preloaded with 100 μ M CM-H₂DCFDA) we have determined the approximate thresholds for irradiance (10,000 W/cm²) generating the oxidation event and the radiant exposure required for detecting the oxidation in our hTERT-RPE1 cells (200,000 J/cm²). Again, no targets of oxidation were observed from exposures of CW 810 nm beams, even at our most sensitive conditions (Table 1).

Table 1. Effect of dye and dye concentration on detection of laser-induced oxidation.

Dye preload into cells using HBSS unless otherwise specified	ML or CW	lowest E w/target (W/cm ²)	lowest H w/target (J/cm ²)
H ₂ DCFDA (50 μ M, PBS)	ML	20,500	820,000
H ₂ DCFDA (100 μ M)	ML	14,600	441,000
H ₂ DCFDA (100 μ M)	CW	no target	no target
CM-H ₂ DCFDA (12.5 μ M)	ML	no target	no target
CM-H ₂ DCFDA (25 μ M)	ML	no target	no target
CM-H ₂ DCFDA (50 μ M)	ML	19,900	662,400
CM-H ₂ DCFDA (100 μ M)	ML	9,920	180,900
CM-H ₂ DCFDA (100 μ M)	CW	no target	no target

Exposures of dye-embedded agarose

To test whether the oxidation observed in our mode-locked laser-exposed hTERT-RPE1 cells was due to an intracellular metabolic (oxidation) reaction, and not a consequence of direct interaction of the dye with the laser, we performed exposures in dye-embedded agarose samples. Here we chemically cleaved the diacetate group from the CM-H₂DCFDA using the manufacturers recommendations (NaOH treatment followed by neutralization) and incubated (at least 30 min at 37°C) the resulting reagent (163 μ M CM-H₂DCF) to wells of a 96-well microtiter plate that had 50 μ l solidified 4% agarose. We exposed the wells containing the dye-embedded agarose to mode-locked 810 nm at irradiances of greater than 19,500 W/cm² and for durations leading to over 1,200,000 J/cm² delivered. In all cases, no laser-specific "target" area was observed above background, represented in Figure 4. The blue exciter light of the fluorescence microscope (at extended excitation; Figure 4 Panel B) provided proof that the




Figure 4. Direct oxidation of CM-H₂DCFDA by blue light (B) but not mode-locked 810 nm light (A).

CM-H₂DCFDA was cleaved to CM-H₂DCF and that this chemical form was capable of being oxidized (by blue light) to CM-DCF.

DISCUSSION

We show that mode-locked NIR lasers can generate a multiphoton photochemical reaction at relatively low average fluences when exposed to living cells. Production of photo-oxidation at 810 nm as a linear event (single photon) is highly unlikely. We were unable to obtain evidence of oxidation when exposing hTERT-RPE1 cell to CW 810 nm laser under our most sensitive conditions, even at maximum output from the Mira laser, and at more than 10 fold greater radiant exposure than that leading to detection with mode-locked exposures.

With the new sensitive detection scheme (preloading cells with 100 μM CM-H₂DCFDA) damage targets from mode-locked 810 nm exposure was easily identifiable as being the result of laser exposures. As compared to the CW 405 nm laser, the oxidation efficiency for mode-locked 810 nm laser exposure is very low, even with the more sensitive conditions for detection. The oxidation from the CW 405 nm laser was detected from a 0.25 sec exposure using 10 μM H₂DCFDA while that from the mode-locked 810 nm laser was detected from a 20 sec (or more) exposure using 100 μM of the more sensitive dye CM-H₂DCFDA. However, the approximate peak irradiance ($8 \times 10^8 \text{ W/cm}^2$) for the threshold oxidation by mode-locked 810 nm laser exposure seems modest for a multiphoton mechanism. From the current results we now attribute the peculiar fluorescence staining reported previously¹² as an artifact of staining as we were unable to repeat any photo-oxidation at 810 nm using the prior conditions.

As judged by thresholds for radiant exposure required for effect in our hTERT-RPE1 cells, we found the general trend in detection efficiency by the fluorescent dyes to be 100 μM CM-H₂DCFDA as the most sensitive, followed by 100 μM H₂DCFDA (~2 fold lower), 50 μM CM-H₂DCFDA (~3 fold lower), and finally the 50 μM H₂DCFDA (~4 fold lower).

We examined potential cytotoxic effects from the mode-locked laser-induced photo-oxidation (results not shown). Upon assaying for viability post-exposure to near threshold fluences, we found that our cells were able to recover from the oxidative stress generated from the mode-locked 810 nm laser. This was not an unexpected result because our system for detection was very sensitive.

Finally, we tested for the possibility that the mode-locked laser was interacting directly with the fluorescent dyes, leading to the detected oxidation targets. Our exposure to agarose-embedded CM-H₂DCF yielded no targets at maximum irradiance and a total radiant exposure delivered that was 6 times above threshold.

In conclusion, we would caution the use of mode-locked lasers in applications for which continued survival of cells, such as extended fluorescent studies of cell cultures or the real-time detection of fluorescence in tissues *in vivo*, until safety issues can be resolved.

ACKNOWLEDGEMENTS

This research reported herein was supported from AFOSR (92HE0COR), AFRL/HEDO (7757B204), and Contract: F33615-92-0017.

REFERENCES

1. Winkler BS, Boulton ME, Gottsch JD, Sternberg P, "Oxidative damage and age-related macular degeneration", *Mol Vis*, **5**, 32-42, 1999.
2. Alizadeh M, Wada M, Gelfman CM, Handa JT, Hjelmeland LM, "Downregulation of differentiation specific gene expression by oxidative stress in ARPE-19 cells", *Invest Ophthalmol Vis Sci*, **42**, 2706-13, 2001.

3. Matsui H, Lin L-R, Singh KP, Shinohara T, Reddy VN, "Lens epithelium-derived growth factor: Increased survival and decreased DNA breakage of human RPE cells induced by oxidative stress", *Invest Ophthalmol Vis Sci*, **42**, 2935-41, 2001.
4. Cai J, Wu M, Nelson KC, Sternberg P, Jones DP, "Oxidant-induced apoptosis in cultured human retinal pigment epithelial cells", *Invest Ophthalmol Vis Sci*, **40**, 959-66, 1999.
5. Sternberg P Jr, Davidson PC, Jones DP, Hagen TM, Reed RL, Drew-Botsch C, "Protection of retinal pigment epithelium from oxidative injury by glutathione and precursors," *Invest Ophthalmol Vis Sci*, **34**, 3661-8, 1993.
6. Honda S, Hjelmeland LM, Handa JT, "The use of hyperoxia to induce chronic mild oxidative stress in RPE cells in vitro", *Mol Vis*, **7**, 63-70, 2000.
7. Sparrow, J. R., Nakanishi, K., Parish, C. A, "The lipofuscin fluorophore A2E mediates blue light-induced damage to retinal pigmented epithelial cells", *Invest Ophthalmol Vis Sci*, **41**, 1981-1989, 2000.
8. Brunk UT, Wihlmark U, Wrigstad A, Roberg K, Nilsson S-E, "Accumulation of lipofuscin within retinal pigment epithelial cells results in enhanced sensitivity to photooxidation", *Gerontology*, **41** (Suppl 2), 201-11, 1995.
9. Hockberger PE, Skimina TA, Centonze VE, Lavin C, Chu S, Dadras S, Reddy JK, White JG, "Activation of flavin-containing oxidases underlies light-induced production of H₂O₂ in mammalian cells", *Proc Natl Acad Sci USA*, **96**, 6255-60, 1999.
10. Xu C, Zipfel W, Shear JB, Williams RM, Webb W, "Multiphoton fluorescence excitation: New spectral windows for biological nonlinear microscopy", *Proc Natl Acad Sci USA*, **93**, 10763-68, 1996.
11. Hu J and Bok D, "A cell culture medium that supports the differentiation of human retinal pigment epithelium into functionally polarized monolayers", *Mol Vis*, **7**, 14-19, 2000.
12. Denton, ML, Eikum, DM, Stolarski, DJ, Noojin, GD, Thomas, RJ, Glickman, RD, Rockwell, BA. 2002. "Hydrogen peroxide production in cultured RPE cells exposed to near infrared lasers", In: *Laser-Tissue Interaction XII*, (S.L. Jacques, ed), SPIE **4617**, 150-155.